



Food and Agriculture
Organization of the
United Nations



World Health
Organization


Residue Monograph prepared by the meeting of the Joint FAO/WHO Expert
Committee on Food Additives (JECFA), 87th Meeting 2019

β -apo-8'-CAROTENAL

This monograph was also published in: Compendium of Food Additive Specifications. Joint
FAO/WHO Expert Committee on Food Additives (JECFA), 87th meeting 2019. FAO JECFA
Monograph 23 (2019)

β -apo-8'-CAROTENAL

Prepared at the 87th JECFA (2019) and published in FAO Monographs 23 (2019), superseding specifications prepared at the 74thJECFA (2011) and published in FAO Monographs 11 (2011). A group ADI of 0-5 mg/kg bw expressed as the sum of carotenoids including β -carotene, β -apo-8'-carotenal, and the methyl and ethyl esters of β -apo-8'-carotenoic acid was established at the 18th JECFA (1974).

SYNONYMS	CI Food Orange 6; CI (1975) No. 40820; INS No. 160e
DEFINITION	These specifications apply to β -apo-8'-carotenal which consists predominantly of all-trans- β -apo-8'-carotenal and may also contain minor quantities of other carotenoids such as all-trans-crocetindialdehyde, all-trans- β -apo-12'-carotenal and all-trans- β -carotene. Commercial preparations of β -apo-8'-carotenal intended for use in food are prepared from β -apo-8'-carotenal meeting these specifications and are formulated as suspensions in edible oil, emulsions and water dispersible powders. These preparations may also contain cis isomers.
Chemical names	β -Apo-8'-carotenal, 8'-apo- β -carotene-al 2E,4E,6E,8E,10E,12E,14E,16E)-2,6,11,15-tetramethyl-17-(2,6,6-trimethyl-1-cyclohexenyl)heptadeca-2,4,6,8,10,12,14,16-octaenal
C.A.S. number	1107-26-2
Chemical formula	C ₄₀ H ₄₀ O
Structural formula	All-trans- β -apo-8'-carotenal (main compound)
	
Formula weight	416.65
Assay	Not less than 96% total colouring matters
DESCRIPTION	Deep violet crystals with metallic lustre or crystalline powder; sensitive to oxygen and light and should therefore be kept in a light-resistant container under inert gas.
FUNCTIONAL USES	Colour

CHARACTERISTICS

IDENTIFICATION

- Solubility (Vol. 4) Insoluble in water; slightly soluble in ethanol; sparingly soluble in vegetable oils.
- Spectrophotometry (Vol. 4) Determine the absorbance of the diluted sample solution used in the Method of Assay at 460 nm and 488 nm. Determine the absorbance at 332 nm of a solution containing a ten-fold higher concentration as that of the diluted sample solution used in the Method of Assay. The ratio A_{488}/A_{460} is between 0.77 and 0.85. The ratio of A_{332}/A_{460} is between 0.63 and 0.75.

PURITY

- Sulfated ash (Vol. 4) Not more than 0.1%
Test 2 g of the sample (Method I)
- Carotenoids other than β -apo-8'-carotenal Not more than 3% of total colouring matters.
See description under TESTS
- Lead (Vol. 4) Not more than 2 mg/kg.
Determine using a method appropriate to the specified level. The selection of sample size and method of sample preparation may be based on the principles of the methods described in Volume 4 (under "General Methods, Metallic Impurities").

TESTS

PURITY TESTS

- Carotenoids other than β -apo-8'-carotenal Chromatographic system
- HPLC equipped with a UV/Vis detector or a photodiode array detector, refrigerated auto sampler and integrator
 - Detector wavelength: 463 nm
 - Column: Reverse phase C18; Suplex pkb-100 (250 x 4.6 mm, 5 μ m) from Supelco or equivalent
 - Mobile phase: In a 1000 ml volumetric flask, dissolve 50 mg BHT in 20 ml 2-propanol and add 0.2 ml N-ethyl-diisopropylamine, 25 ml 0.2% aqueous ammonium acetate solution, 455 ml acetonitrile, and 3pprox.. 450 ml methanol. Mixture cools and contracts. Allow to reach room temperature and dilute to volume with methanol. Discard after 2 days.
 - Isocratic elution
 - Column temperature: 30°
 - Flow rate: 0.6 ml/min
 - Injection volume: 10 μ l
 - Temperature of the autosampler: (3pprox.. 15°)

- Run time: 4 approx.. 35 min

Reagents

- Butylated hydroxytoluene (BHT), reagent grade
- 2-Propanol, HPLC grade
- N-ethyl-diisopropyl-amine, reagent grade
- Ammonium acetate, reagent grade
- Acetonitrile, HPLC grade
- Methanol, HPLC grade
- Ethanol, HPLC grade
- Tetrahydrofuran, HPLC grade

Sample solution

Weigh accurately (to ± 0.1 mg) 0.010 g of the sample and dissolve in tetrahydrofuran (stabilized with 0.025% BHT). Transfer to a 100 ml volumetric flask and bring to volume with tetrahydrofuran. Dilute to the ratio of 1:10 with ethanol.

Procedure

Inject the sample solution using the conditions detailed under *Chromatographic system*. The retention times for all-*trans*- β -apo-8'-carotenal is in the range of 7-9 min and corresponds to the largest peak in the chromatogram. The relative retention times of minor carotenoids with respect to the retention time of all-*trans*- β -apo-8'-carotenal are: all-*trans*-crocetin dialdehyde (0.54); all-*trans*- β -apo-12'-carotenal (0.84); all-*trans*- β -carotene (2.55).

Integrate the areas of the peaks in the chromatogram.

Calculation

Calculate the percentage of carotenoids other than β -apo-8'-carotenal (% w/w) using the following formula:

$$\text{Carotenoids other than } \beta - \text{apo} - 8' - \text{carotenal} \left(\%, \frac{W}{W} \right) = \frac{A_{\text{total}} - A_{\beta\text{-apo-8'-carotenal}}}{A_{\text{total}}}$$

where

A_{total} is the sum of the area of all the peaks in the chromatogram, excluding the solvent peak (area units); and

$A_{\beta\text{-apo-8'-carotenal}}$ is the area of the peak of β -apo-8'-carotenal in the chromatogram (area units).

METHOD OF ASSAY Total colouring matters content by spectrophotometry

Proceed as directed under Total Colouring Matters Content – Colouring Matters Content by Spectrophotometry, Procedure 2, using the following conditions:

Sample weight (W): 0.08 g (± 0.01 g);

Volume of the three volumetric flasks: $V_1 = V_2 = V_3 = 100$ ml;

Volume of the two pipets: $v_1 = v_2 = 5$ ml;

Specific absorbance of the standard: $A_{1\text{cm}}^{1\%} = 2640$;

Wavelength of maximum absorption: λ_{max} about 461 nm.

Calculation

Calculate the percentage of total colouring matters using the following formula:

$$\text{Total colouring matters (\%, w/w)} = \frac{A \times V_1 \times D}{A_{1\text{cm}}^{1\%} \times W}$$

where

A is the absorbance of the twice-diluted sample solution at 461 nm; and

D is the dilution factor $(V_2 \times V_3)/(v_1/v_2)$.