

Enhanced drought tolerance by expression of *hvDhn5* gene in poplar

M. S. Han¹, E. W. Noh²

¹mushan@korea.kr, Department of Variety of Examination, National Forest Seed & Variety Center,

²Korea Poplar Commission

Abstract

We transferred *Dhn5*(dehydrin5) gene from barley to poplar to determine the effect of its expression on the transgenic poplars. The results from northern blot analysis showed that the expression level of gene varied among the transgenic lines. During their culture on tissue culture media, the transgenic poplars formed vigorous growing callus in the presence of 5% PEG. When the transgenic poplars were growing in pots and withheld watering, they stayed much healthier than nontransgenic poplars. The transgenic poplars showed higher rates of photosynthetic rates, stomatal conductance and evaporation rates under the drought stress, although there was no significant difference in soil water content within the treatments. The relative electrical conductivity of the transgenic poplars after 20% PEG treatment was lower than that of nontransgenic poplars. The results provide evidence that the expression of *hvDhn5* gene conferred drought tolerance in the transgenic poplars.

Keywords : Dehydrin5, Drought tolerance, Photosynthetic rates, Relative electrical conductivity, Transgenic poplar

Introduction

Salt, drought and freezing stresses are the most critical abiotic stresses that affect productivity of annual crop species as well as perennial trees (Caruso et al. 2002). Recently rapid desertification has resulted in increased interest in the development of crop varieties that could tolerate the various environmental stresses. However, different plants adapt to their environment by different mechanisms. Therefore, it has been very difficult to select or develop varieties which withstand various environmental stresses via conventional breeding. The stacking of those traits into a selected variety requires cross fertility between the organisms that carry traits of interest (Flowers and Yeo 1995). This barrier could be overcome by genetic engineering (Singsit et al. 1997). It is now possible to selectively transfer only the traits of interest into a new plant variety so that we could avoid to transfer the unwanted portion of DNA in the donor genome (Cushman and Bohnert 2000).

A number of promising genes for salt and drought tolerance have been reported. Among them are LEA (late embryogenesis-abundant protein) genes (Brini et al. 2007b). Dehydrin belongs to LEA II group protein and known to be induced by environmental stresses including drought and cold temperature. Dehydrin is thought to play a role as a stabilizer of cellular macromolecules (Close 1996, 1997; Ingram & Bartels 1994). For example, transgenic plants expressing LEA genes from barley and corn showed increased tolerance against drought and salt stresses (Xu et al. 1996; Figueras et al. 2004). So far 13 dehydrin (*Dhn* 1 to 13) genes from barley have been reported (Rodriguez et al. 2005). Among them, *Dhn5* is known to be induced by low temperature stress and involved in freezing tolerance by protecting meristematic cells from losing water (Zhu et al. 2000). Han and Hwang (2003) transferred the barley *Dhn5* gene to *Artemisia adamsii* and reported that the calli from transgenic plants grew much better than those from nontransgenic plants in the presence of 20% PEG. Similarly, transgenic *Arabidopsis* carrying wheat *Dhn5* gene displayed enhanced tolerance against salt and osmotic stresses (Brini et al. 2007a). Many other cases of transgenic plants expressing LEA/Dhn proteins with increased tolerance against salt, drought and cold stresses have been reported (Xu et al. 1996; Babu et al. 2004; Sivamani et al. 2000; Honjoh et al. 1999).

We were interested in confirming whether the gene could confer poplar, a woody species, tolerance against such stresses. Therefore, in the present study, we introduced barley *Dhn5* gene to poplar and examined the phenotypes of the transgenic poplars under drought stress.

Materials and methods

Gene cloning and vector construction

Barley *Dhn5* gene was PCR cloned from barley genomic DNA using published sequence (GenBank accession No. M95810). A primer set was designed to amplify the gene: Forward: ggatcctaccctcgaacggcctataa, and Reverse: gagctctgaatgtgcatacgggtgga. An enzyme site (*Bam*H1 in F primer and *Sac*1 in R primer, respectively) was inserted to each primer to ease cloning process. The amplified PCR product was inserted into pBI121 after GUS gene was removed by *Bam*H1 and *Sac*1 digestion and transferred to an *Agrobacterium tumefaciens* vector, the strain GV3101.

Poplar transformation and regeneration

The chimeric *p35S-hvDhn5* gene was introduced to poplar by *A. tumefaciens* mediated transformation (Fillatti et al. 1987). Transformed cells were selected on MS medium (Murashige and Skoog 1962) containing 1.0 mg/L 2.4-D, 0.1 mg/L BAP, 0.01 mg/L NAA, 500 mg/L cefotaxime plus 50 mg/L kanamycin. Shoots were regenerated on WPM (Lloyd and McCown 1981) containing 1.0 mg/L zeatin, 0.1 mg/L BA, 0.01 mg/L NAA and 50 mg/L kanamycin.

Molecular analysis

Genomic DNA was extracted from the leaves of poplar plants using a MagExtractor-Plant Genome kit (Toyobo, Osaka, Japan). Ten µg of the genomic DNA were digested with a restriction enzyme *Pst* 1 overnight, run on 1% agarose gel and then transferred to Hybond-XL nylon membrane by capillary transfer method (Southern 1975). The full-length *hvDhn5* coding sequence labelled with 32P-dCTP was used as probe for hybridization for 12 h. The membrane was washed in 2 × SSC and 0.1% SDS (50°C) for 10 min and in 0.2 × SSC and 0.1% SDS (50°C) for 30 min followed by exposing to an X-ray film at -70°C.

For northern blot, the total RNA was isolated from young leaves using TRI Reagent (Molecular Research Center Inc., Cincinnati, OH, USA). Ten µg of the RNA were run on 1.2% formaldehyde agarose gel and then transferred to Hybond-XL nylon membrane. The methods for probe preparation and hybridization were the same as in Southern hybridization.

Drought tolerance assays

For drought tolerance assay, four different transgenic clones (*Dhn5*-1, 5-3, 5-9, and 5-10) showing different expression levels were used. To determine tolerance of the transgenic lines to drought stress, both *in vitro* and greenhouse pot trials were performed. For *in vitro* drought tolerance assay, leaf fragments (0.5 cm × 1.0 cm) were cultured on callus inducing media (MS with 1.0 mg/L 2.4-D and 0.1 mg/L BAP) containing various levels (0, 5, 10%) of polyethylene glycol (PEG). Fresh weight growth of the calli was determined after 5 weeks growth on culture medium. Drought tolerance of the two transgenic lines (*Dhn5*-1 and *Dhn5*-10) was assessed in a pot trial in the greenhouse. The plants were transferred to plastic pots filled with sand soil and fully watered. After 28 days, transgenic and nontransgenic poplars were subjected to drought treatment by withholding irrigation for 7 days. Reference plants were irrigated at 3 to 4 day interval. The photosynthesis of the poplars

was compared when damage symptoms were visible. A portable photosynthesis system (model LI-6400, LI-COR, Lincoln, NE) was used to determine photosynthetic rates. After photosynthesis measurements, soil water content at the end of 7 days was measured.

Ion leakage analysis

Ion leakage of the leaves was measured by relative electrical conductivity (REC) as described by Wu et al. (2008) with some modifications. Leaf tissue was cut into segments of 1 cm² with cork borer (No. 6). Five leaf segments were then drought stress for 48 hrs with 20% PEG solution at room temperature (25±2 °C). After the stress, the leaf segments were washed quickly five times with distilled deionized water and then put into 15 ml of distilled deionized water in falcon tube (50 ml) and shaken using a rotary shaker (170 rpm) at room temperature (25±2 °C). Electrical conductivity (EC) was measured at 1 hr later using an ion conductivity meter (Mettler co.) (E1). Then, the tubes containing the segments were autoclaved for 20 min at 121 °C and the conductivity of the effused was measured after cooling it to room temperature (E2). Relative electrical conductivity was measured by the formula $E1 / E2 \times 100$.

Results and discussion

Development and molecular analyses of transgenic poplars

Transgenic calli were first screened on callus inducing medium in the presence of 50 mg/L kanamycin. Shoots were also regenerated from the calli in the presence of 50 mg/L kanamycin. Southern blot analysis using barley *Dhn5* gene as a probe revealed that out of 5 transgenic poplars examined, two contained a single copy transgene and three contained multiple copies (Fig. 2). Northern blot using barley *Dhn5* gene showed that expression level varied much up to 10 fold difference among transgenic lines (Fig. 3). The difference in the transcription level among transgenic lines may be due to different copy number or different integration sites in the genome (Cooley et al. 1995; Sarria et al. 2000).




Fig. 1 Schematic illustration of a plant expression vector carrying *p35S-hvDhn5* gene (*Dhn5*: barley dehydrin5 gene, *p35S*: CaMV35S promoter, *pNOS*: nopaline synthase promoter, *npt II*: neomycin phosphotransferase II, *NOS3'*: nopaline synthase terminator)




Fig. 2 Southern blot analysis of poplars transformed by *p35S-hvDhn5*. Genomic DNA (10µg) from regenerated poplars was digested with *Pst* I and separated by electrophoresis in an 1.0% agarose gel. The gel was blotted onto nylon membrane, and hybridized with ³²P-labeled full-length *hvDhn5* gene. 1: nontransgenic poplar and 2 to 6: transgenic poplars




Fig. 3 Expression of *p35S-hvDhn5* gene revealed by Northern blot analysis. 10 μ g of total RNA extracted from leaf tissues were resolved on formaldehyde agarose gel and blotted onto nylon membrane. Hybridization was done with 32 P-labeled probe of full length *hvDhn5* gene. The top is EtBr-stained gel showing rRNA bands. C: nontransgenic poplar and 1 to 5: transgenic poplars

Drought tolerance test using test tube plants

To determine tolerance of the *hvDhn5* transgenic poplars to drought stress, calli were induced from the transgenic lines on callus inducing (and growing) medium containing 0, 5, 10% PEG. Although all the lines tested formed vigorously growing callus in the absence of PEG, they all showed growth suppression in the presence of PEG. However, transgenic lines showed less damage by PEG imposed stress than untransformed control. In the presence of 5% PEG nontransgenic poplars attained only 15% of callus growth (in fresh weight) that could be produced in the absence of PEG. In contrast, transgenic lines showed 26 and 37% of callus growth otherwise obtained in the absence of PEG. However, at 10% PEG, all the lines showed growth suppression and did not show any difference in the growth among the lines (Fig. 4). Transgenic *Artemisia* carrying the same *Dhn5* gene was reported to tolerate up to 20% PEG. The difference may be genetic nature of the plant materials. *Artemisia adamsii* is native to Gobi desert and thus thought to be highly tolerant to stress (Han and Hwang 2003). There might be other factors that protect *Artemisia* from drought or osmotic stresses.




Fig. 4 Fresh weight of callus of poplar with a barley *Dhn5* gene grown in media supplemented with PEG. Control is the nontransgenic poplars; *Dhn5-1*, *Dhn5-3*, *Dhn5-9*, and *Dhn5-10* are the transgenic poplars with the *hvDhn5* gene. All the plants were treated with PEG for 28 days. Vertical bars represent standard errors (n=3 per experiment)

Drought tolerance test using pot plants

hvDhn5 transgenic poplars withheld watering for a week did not show any wilting symptoms while nontransgenic poplars suffered from severe wilting damage (Fig. 5). There was no difference in photosynthetic rate between transgenic and nontransgenic poplars in normal conditions. However, at the time of wilting symptoms developed, the photosynthetic rate of nontransgenic poplars rapidly dropped to 33.7% of that in normal condition. In contrast, the transgenic lines *Dhn5-1* and *Dhn5-10* maintained the rates up to 84.6 and 92.7% of normal condition, respectively (Fig. 6-A).

Soil water content was also determined by the time when photosynthetic rates were measured. There was no difference in soil water content between the pots with transgenic lines and those with nontransgenic poplars (Fig. 7). Both drought and salt stress are known to cause the reduction in photosynthetic rates (Greenway and Munns 1980). When plants are under drought stress, they close stomata and thereby lowering CO₂ level in the mesophyll cells and reducing photosynthetic activity (Farquhar et al. 1987; Gimenez et al. 1992). Oh et al. (2005) also demonstrated that drought stress reduced plant growth and thus productivity of aspen plants by decreasing photosynthesis. Thus, it is critical to maintain photosynthetic structures and photosynthetic activity under drought stress for plant productivity (Tunnacliffe and Wise 2007).


Changes in stomatal conductivity under drought stress appeared to be similar to those in photosynthesis. Whereas nontransgenic poplars under drought stress displayed 8.2% activity of untreated poplars, the two transgenic lines were shown to maintain 23.3% and 32.5%, respectively of untreated poplars under the same conditions (Fig. 6-B). These results suggest that stomata closure play a role in the defense against drought stress and the transgenic lines seem to have better capacity to control stomatal closure. Similar observation was also made by other workers (Chaves et al. 2003).

As for changes in transpiration rate under drought stress, nontransgenic poplars showed markedly 93.2% decrease when compared to untreated poplars. In contrast, transgenic lines showed merely decreased rates of 64.3% and 62.8%, respectively when compared to untreated poplars (Fig. 6-C). Thus, the transgenic lines have higher activities in photosynthesis, stomatal conductivity, and transpiration than nontransgenic poplars. Therefore, overall, transgenic lines were more tolerant and thus showed less damage symptoms.



(A) (B) (C)

Fig. 5 Growth phenotypes of nontransgenic poplar vs. transgenic poplars over expressing *hvDhn5* gene after 7 days of withholding watering; (A) nontransgenic poplar, (B) and (C) transgenic poplars



(A) (B) (C)

Fig. 6 Photosynthetic rate (A), transpiration rate (B) and stomatal conductance (C) of *p35S-hvDhn5* transgenic poplars after 7 days of withholding watering. Control is the nontransgenic poplars; *Dhn5-1* and *Dhn5-10* are the transgenic poplars with the *hvDhn5* gene. All the plants were treated for 7 days. Vertical bars represent standard errors (n=3 per experiment)




Fig. 7 Relative soil water content after 7 days of withholding watering. Control is the nontransgenic poplars; *Dhn5-1* and *Dhn5-10* are the transgenic poplars with the *hvDhn5* gene. All the plants were treated for 7 days. Vertical bars represent standard errors (n=3 per experiment)

Ion leakage test

Figure 8 shows the relative ion conductivity of the solution exuded from leaf disks after 20% PEG treatment. There was no significant difference between transgenic and nontransgenic poplars in the absence of PEG. However, when treated with PEG, the relative ion conductivity was significantly lower in the solution from transgenic lines than that from nontransgenic poplars. Whereas the ion conductivity of nontransgenic poplars increased 153.3% by PEG treatment, those of transgenic lines increased to 138.4% and 136.5%, respectively. The result is consistent with the previous reports showing lower ion conductivity in both transgenic tobacco and rice plants expressing LEA or *HVA1* gene (Wang et al. 2006; Liu et al. 2009). Wang et al. (2006) conjectured that the low ion conductivity might be due to less damage of the transgenic plants since the LEA proteins protected the membrane under drought stress. Similarly, LEA proteins act as a stabilizer of cellular molecules and thus protect cellular components under the stress (Tunnacliffe and Wise 2007; Bajji et al. 2002). Therefore, the transgenic poplar lines are more tolerant to drought stress than nontransgenic poplars due to the expression of *hvDhn5* gene.




Fig. 8 The relative electrical conductivity in the leaf discs of nontransgenic and transgenic poplars after 20% PEG treatment for 48 hrs. Vertical bars represent standard errors (n=3 per experiment)

References

- Babu RC, Zhang JX, Blum A, Ho THD, Wu R, Nguyen HT. 2004. HVA1, a LEA gene from barley confers dehydration tolerance in transgenic rice (*Oryza sativa* L.) via cell membrane protection. *Plant Science*, 166:855-862.
- Bajji M, Kinet JM, Lutts S. 2002. The use of electrolyte leakage method for assessing cell membrane stability as a water stress tolerance test in durum wheat. *Plant Growth Regulation*, 36:61-70.
- Brini F, Hanin M, Lumbreras V, Amara I, Khoudi H, Hassairi A, Pagès M, Masmoudi K. 2007a. Overexpression of wheat dehydrin DHN-5 enhances tolerance to salt and osmotic stress in *Arabidopsis thaliana*. *Plant Cell Reports*, 26:2017-2026.
- Brini F, Hanin M, Lumbreras V, Irar S, Pagès M, Masmoudi K. 2007b. Functional characterization of DHN-5, a dehydrin showing a differential phosphorylation pattern in two Tunisian durum wheat (*Triticum durum* Desf.) varieties with marked differences in salt and drought tolerance. *Plant Science*, 172:20-28.
- Caruso A, Morabito D, Delmotte F, Kahlem G, Carpin S. 2002. Dehydrin induction during drought and osmotic stress in *Populus*. *Plant Physiology and Biochemistry*, 40:1033-1042.
- Chaves MM, Maroco JP, Pereira JS. 2003. Understanding plant responses to drought from genes to the whole plant. *Functional Plant Biology*, 30:239-264.
- Close TJ. 1996. Dehydrins: emergence of a biochemical role of a family of plant dehydration proteins. *Physiologia Plantarum*, 97:795-803.
- Close TJ. 1997. Dehydrins: a commonality in the response of plants to the dehydration and low temperature. *Physiologia Plantarum*, 100: 291-296.
- Cooley J, Ford T, Cristou P. 1995. Molecular and genetic characterization of elite transgenic rice plants produced by electric discharge particle acceleration. *Theoretical and Applied Genetics*, 90:97-104.
- Cushman JC, Bohnert HJ. 2000. Genomic approaches to plant stress tolerance. *Current Opinion in Plant Biology*, 3:117-124.
- Farquhar GD, Husic KT, Terashima I, Codon AG, Richards RA. 1987. Genetic variation in the relationship between photosynthetic CO₂ assimilation rate and stomata conductance to water loss. *Progress in Photosynthetic Research*, 4: 209-212.
- Figueras M, Pujal J, Saleh A, Savé R, Pagès M, Goday A. 2004. Mazie Rab17 over-expression in

Arabidopsis plants promotes osmotic stress tolerance. *Annals of Applied Biology*, 144:251-257.

Fillatti JJ, Sellmer J, McCown B, Hassig B, Comai L. 1987. Agrobacterium mediated transformation and regeneration of poplars. *Molecular and General Genetics*, 206:192-199.

Flowers TJ, Yeo AR. 1995. Breeding for salinity resistance in crop plants: where next?. *Austrian Journal of Plant Physiology*, 22:875-884.

Gimenez C, Mitchell VJ, Lawlor DW. 1992. Regulation of photosynthetic rate of two sunflower hybrids under water stress. *Plant Physiology*, 96:635-643.

Greenway H, Munns R. 1980. Mechanisms of salt tolerance in nonhalophytes. *Annual Review of Plant Physiology*, 123:149-190.

Han KH, Hwang CH. 2003. Transformation of *Artemisia adamsii*, Endemic to a gobi desert, with CLP, *Dhn5* to enhance environmental stress tolerance. *Journal of Plant Biotechnology*, 30:315-321.

Honjoh KI, Oda Y, Tkata R, Miyamoto T, Hatano S. 1999. Introduction of the hiC6 gene, which encodes a homologue of a late embryogenesis abundant (LEA) protein, enhances freezing tolerance of yeast. *J. Plant Physiology*, 155:509-512.

Ingram J, Bartels D. 1994. The molecular basis of dehydration tolerance in plants. *Annual Review of Plant Physiology*, 106:377-403.

Liu X, Wang Z, Wang L, Wu R, Phillips J, Deng X. 2009. LEA 4 group from the resurrection plant *Boea hygrometrica* confer dehydration tolerance in transgenic tobacco. *Plant Science*, 176: 90-98.

Lloyd D, McCown BH. 1980. Commercially feasible micropropagation of mountain laurel, *Kalmia latifolia*, by use of shoot tip culture. *Combined Proceedings, International Plant Propagators Society*, 30:421-427

Murashige T, Skoog F. 1962. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Plant Physiology*, 15: 473-497.

Oh CY, Han SH, Kim YY, Lee JC. 2005. Changes of drought tolerance and photosynthetic characteristics of *Populus davidiana* Doe according to PEG treatment. *Korean Journal of Agricultural and Forest Meteorology*, 7:296-302.

Rodriguez EM, Sevansson JT, Malatrasi M, Choi DW, Close TJ. 2005. Dhn13 encodes a KS-type dehydrin with constitutive and stress responsive expression. *Theoretical and Applied Genetics*, 110: 852-858.

Sarria R, Torres E, Angel F, Chavarriaga P, Roca WM. 2000. Transgenic plants of cassava (*Manihot esculenta*) with resistance to Basta obtained by Agrobacterium-mediated transformation. *Plant Cell Reports*, 19:339-344.

Singsit C, Adang MJ, Lynch RE. 1997. Expression of a *Bacillus thuringiensis* cryIA (c) gene in transgenic peanut plants and its efficacy against lesser cornstalk borer. *Transgenic research*, 6:169-170.

Sivamani E, Bahieldin A, Wraith JM, Al-Niemi T, Dyer WE, Ho TD, Wu R. 2000. Improved biomass productivity and water use efficiency under water-deficit conditions in transgenic wheat constitutively expressing the barley HVA1 gene. *Plant Science*, 155:1-9.

Southern EM. 1975. Detection of specific sequences among DNA fragments. *Journal of Molecular Biology*, 98:503-517.

Tunnacliffe A, Wise MJ. 2007. The continuing conundrum of the LEA proteins. *Naturwissenschaften*, 94:791-812.

Wang Y, Jiang J, Zhao X, Liu G, Yang C, Zhan L. 2006. A novel LEA gene from *Tamarix androssowii* confers drought tolerance in transgenic tobacco. *Plant Science*, 171:655-662.

Wu W, Su Q, Xia XY, Wang Y, Luan YS, An LJ. 2008. The *Suaeda liaotungensis* kitag betaine aldehyde dehydrogenase gene improves salt tolerance of transgenic maize mediated with minimum linear length of DNA fragment. *Euphytica*, 159:17-25.

Xu D, Duan X, Wang B, Hong B, Ho THD, Wu R. 1996. Expression of a late embryogenesis abundant protein gene, HVA1, from barley confers tolerance to water-deficit and salt-stress in transgenic rice. *Plant Physiology*, 110:249-257.

Zhu B, Choi DW, Fenton R, Close TJ. 2000. Expression of the barley dehydrin multigene family and the development of freezing tolerance. *Molecular and General Genetics*, 264:145-153.