

## **Part 1**

# **THE SCIENCE OF GENETIC MODIFICATION IN FOREST TREES**

# 1. Genetic modification as a component of forest biotechnology

*C. Walter and M. Menzies*

While the term “biotechnology” refers to a broad spectrum of modern tools and the application of those tools, it is frequently equated with genetic engineering by the lay public. FAO noted in their 2004 report *The State of Food and Agriculture* that “biotechnology is more than genetic engineering” (FAO, 2004a). In fact, 81% of all biotechnology activities in forestry over the past ten years were not related to genetic modification (Wheeler, 2004).

There are many definitions of biotechnology and they differ in their scope. FAO (2001) defines the term biotechnology as “any technological application that uses biological systems, living organisms, or derivatives thereof, to make or modify products or processes for specific use”. This definition, although accurate for the specific purposes for which it was intended, may contribute to the confusion surrounding the term. A simpler definition might be “the application of biological knowledge to practical needs such as technologies for altering reproduction, or technologies for locating, identifying, comparing or otherwise manipulating genes”.

In short, forest biotechnology is associated with a broad spectrum of modern methods applicable to agricultural and forest science, only some of which are related to genetic engineering. In forestry, the definition of biotechnology covers all aspects of tree breeding and plant cloning, DNA genotyping and gene manipulation, and gene transfer.

Forest biotechnologies can be classified in many ways (Yanchuk, 2001; Wheeler, 2004), but here they are grouped under five major, though undoubtedly overlapping, categories (Henderson and Walter, 2006; Trontin *et al.*, 2007; El-Kassaby, 2003, 2004):

- propagation;
- molecular markers;
- marker-assisted selection (MAS) and marker-assisted breeding (MAB);
- genomics, metabolomics and proteomics;
- genetic modification or genetic engineering.

This chapter provides a brief discussion of these technologies in the context of existing or proposed deployment in commercial forestry. However, this should be read only as an introduction, and the reader is referred to the vast literature available on those subjects.

## PROPAGATION

Plant cloning has been used for centuries for tree breeding and propagation using grafts and cuttings. Chinese fir (*Cunninghamia lanceolata*) has been propagated by cuttings for clonal forestry in China for more than 800 years (Li and Ritchie, 1999) and Japanese cedar (*Cryptomeria japonica*) has been propagated clonally by cuttings in Japan for plantations since the beginning of the fifteenth century (Toda, 1974). Some tree species are easier than others to propagate by cuttings. Easy-to-root hardwood species, such as poplars (*Populus* spp.), willows (*Salix* spp.) and some eucalypt (*Eucalyptus*) species, and conifer species, such as spruces (*Larix* spp.), redwood (*Sequoia sempervirens*), and some pines (*Pinus* spp.), are widely planted as cuttings in family or clonal plantations (Ritchie, 1991; Ahuja and Libby, 1993; Assis, Fett-Neto and Alfenas, 2004; Menzies and Aimers-Halliday, 2004). In the future, the use of vegetatively propagated trees for intensively managed, high-yielding plantations is expected to increase in all regions of the world.

While the main use of propagation technologies has been for forest establishment of genetically-improved families or clones, there is also a conservation use for those species that are at risk, rare, endangered or of special cultural, economic or ecological value (Benson, 2003). Integrating traditional methods such as *in situ* conservation and seed storage with biotechnologies such as micropropagation and cryopreservation can provide successful solutions.

### Micropropagation

Micropropagation refers to the *in vitro* vegetative multiplication of selected plant genotypes, using organogenesis and/or somatic embryogenesis. Approximately 34% of all biotechnology activities reported in forestry over the past ten years related to propagation (Chaix and Monteuuis, 2004; Wheeler, 2004). Micropropagation is used to multiply (bulk-up) desirable genotypes or phenotypes to create large numbers of genetically identical individuals of clones or varieties. These techniques are gaining increased attention by foresters and tree breeders because vegetative propagation offers a unique opportunity to bypass the genetic mixing associated with sexual reproduction.

### Organogenesis

While macropropagation methods, such as cuttings, involve comparatively large pieces of tissue, micropropagation by organogenesis involves *in vitro* culture of very small plant parts, tissues or cells, particularly meristems from germinating embryos or juvenile plant apices. There are a number of stages in organogenesis, involving sterilization and shoot initiation, shoot elongation and multiplication, rooting and acclimatization. Sterilization is typically done with a diluted bleach solution, followed by initiation of shoots on an appropriate tissue culture medium. Shoots can develop from existing axillary meristems or from meristems of adventitious origin. Adventitious meristems can be stimulated from plant tissue, such as cotyledons or leaves, by exposure to a pulse of the plant hormone, cytokinin. Plants arising from shoots of adventitious origin may show undesirable

advanced maturation characteristics (Frampton and Isik, 1987). There have been many different media developed for organogenesis, depending on the species (McCown and Sellmer, 1987). Following shoot initiation, shoots are elongated on a medium without cytokinin. The addition of 0.5–1.0% activated charcoal may be beneficial. Once shoots have elongated sufficiently, they can be cut into nodal sections or topped to stimulate lateral side shoot or shoot clump development, which can then be separated and elongated. When sufficient multiplication has been achieved, the shoots can be stimulated to form roots by transferring them to a medium containing auxin. Rooting may be done *in vitro* or *ex vitro*, depending on the species. Venting of the culture container by using a hole in the container lid covered with a permeable membrane or cotton wool during the time in auxin medium may help acclimatization for transfer *ex vitro*. Similarly, the container lid may be left loosened or unwrapped to allow some gaseous exchange and exposure to ambient humidity. Once shoots are transferred *ex vitro* and have rooted, the humidity may be gradually reduced to ambient conditions in an acclimatization phase.

There are a number of methods available for maintaining or storing of clones in tissue culture by organogenesis, including repeated subculture (serial propagation), minimal growth media, cool storage and cryopreservation. Radiata pine clones have been maintained as shoots for more than ten years with repeated subculture every 6–8 weeks (Horgan, Skudder and Holden, 1997). However, long-term success at halting ageing is uncertain and the costs are high because of the requirement for regular transfers and a controlled environment. Using diluted nutrient concentrations in the media does reduce the need for regular subculturing, and radiata pine shoots have been maintained successfully for four years at 20–22 °C with annual subculturing (Horgan, Skudder and Holden, 1997). Successful cryopreservation of organogenic material has proved to be more difficult. Cotyledons from radiata pine zygotic embryos have been successfully frozen and thawed (Hargreaves *et al.*, 1999). Cryopreservation of axillary meristems is also being attempted (Hargreaves *et al.*, 1997) and results are now very promising (Hargreaves and Menzies, 2007). Organogenesis methods have been developed for a large number of forestry species for large-scale production, including hardwoods such as poplars, willows and eucalypts, and for conifers such as coast redwoods, radiata pine (*Pinus radiata*), loblolly pine (*Pinus taeda*) and Douglas fir (*Pseudotsuga menziesii*). More detailed protocols for various hardwoods and conifers can be found in Bonga and Durzan (1987a, b) and Bajaj (1986, 1989, 1991).

### Embryogenesis

Another micropropagation technology that has been more recently developed and has promising applications for clonal forestry is somatic embryogenesis. Successful embryogenesis was first reported for sweetgum (*Liquidambar styraciflua*) in 1980 (Sommer and Brown, 1980) and for spruce (*Picea abies*) in the mid-1980s (Hakman and von Arnold, 1985; Chalupa 1985). Since then, somatic embryogenesis has been

investigated for many forestry species, including hardwoods such as poplars, willows and eucalypts, and conifers such as spruces, larch (*Larix* spp.), pines and Douglas fir. Embryogenesis differs from organogenesis in that somatic embryos are formed from embryogenically competent somatic cells *in vitro*, with both shoot and root axes, and these embryos will germinate, whereas with organogenesis shoots are developed, and these must be rooted as mini-cuttings.

As in organogenesis, there are a number of stages for embryogenesis, involving initiation of embryogenic tissue, multiplication, development and maturation, germination and acclimatization. Typically, embryogenic tissue is established from immature seeds, just after fertilization, using either embryos within intact megagametophytes or excised embryos. Tissue can be maintained or multiplied in a relatively undifferentiated state. However, by changing the medium, embryos can be stimulated to develop into bullet-stage embryos with suspensors. Further medium changes, including the addition of abscisic acid, increasing the osmotic potential, and controlled desiccation using water-vapour-permeable plastic film, stimulate the embryos to develop and mature into the cotyledonary stage. These embryos can be harvested and, after germination under sterile conditions, transferred to containers in a greenhouse. The somatic seedlings are transferred to larger containers or lined out in a nursery bed when they are large enough. More detailed protocols for various hardwoods and conifers can be found in Bajaj (1989, 1991), Jain, Gupta and Newton (1999, 2000) and Jain and Gupta (2005).

An important advantage of embryogenesis is the ability to maintain or store clones through cryopreservation. Reliable cryogenic storage of embryogenic tissue at  $-196\text{ }^{\circ}\text{C}$  has been possible for many years (Cyr, 1999; Gupta, Timmis and Holmstrom, 2005). Typically, free water is removed by the use of a higher osmoticum medium, followed by the addition of a cryoprotectant, such as sorbitol and dimethylsulphoxide (DMSO). This avoids the formation of the ice crystals that cause cell disruption and death. Similarly, thawing is done rapidly to avoid ice crystal formation.

The efficiency of embryogenesis needs further improvement, but the technology has the potential to produce unlimited quantities of embryos of desirable genotypes at costs cheaper than current control-pollinated seed prices. These benefits will be achieved once genotype capture is improved, automation technology is designed and artificial seed is developed. Micropropagation, and in particular embryogenesis, is the gateway to genetic engineering (Henderson and Walter, 2006). While *Agrobacterium tumefaciens* transformation is most successful with hardwood species, using organogenic or embryogenic technologies, biolistic transformation can be used most successfully with embryogenic cultures of both softwoods and hardwoods. This means that the development of genetically modified trees is dependant on the availability of a reliable, reproducible propagation system (Campbell *et al.*, 2003).

### Choosing the appropriate system

A range of propagation systems are available for clonal deployment and they each have advantages and disadvantages. Micropropagation systems have the advantages of high potential multiplication rates, potentially reliable cooled storage or cryopreservation, and amenability to genetic modification. However, major disadvantages are that the techniques may not work for a considerable proportion of genotypes, plant quality may be poor and costs are high. Nursery cuttings systems have lower multiplication rates and allow short-term clonal storage through stool-bed systems, but can reliably produce good quality plants at lower cost than current micropropagation systems. A hybrid system might be the best option. For example, organogenesis or embryogenesis could be used initially to capture and cryopreserve genotypes and to produce sufficient plants for clonal testing. Once clones had been selected for clonal production, sufficient individuals could be produced by micropropagation to be planted as stock plants for the production of cuttings, producing more robust and cheaper plants for outplanting (Menzies and Aimers-Halliday 1997). Also, if embryogenesis is producing low numbers of germinating somatic seedlings for some clones, the germinating plants can be transferred to an organogenesis multiplication system while still sterile to increase plant numbers before transfer *ex vitro*.

### MOLECULAR MARKERS, MAS, QTL DETECTION AND FINGERPRINTING

The introduction of biochemical (e.g. terpenes and flavanoids) and Mendelian-inherited protein (allozymes) markers in the latter quarter of the past century drove a rapid increase in evolutionary biology studies in forestry. These markers also found valuable application in seed orchard management (Wheeler, Adams and Hamrick, 1993; El-Kassaby, 2000). In the past decade, the development of molecular markers based directly on DNA polymorphisms has largely replaced allozymes for most practical and scientific applications. This replacement was accelerated by the development of the polymerase chain reaction (PCR) technique. Molecular markers come in many forms, each with an array of benefits and drawbacks (Ritland and Ritland, 2000). The utility of these molecular markers and the analytical methods used differ according to the type of question asked and the nature of the markers (dominant vs co-dominant).

Molecular markers are routinely used for a number of research and development and practical applications in forestry, the most common of which is the estimation of genetic diversity in natural and artificial populations. According to Chaix and Monteuis (2005), over 25% of all biotechnology activity reported in the past ten years related to marker application, predominately focused on measures of diversity. Other applications include the study of gene flow and mating systems, tracking clonal and seedling materials in breeding programmes, paternity studies, gene conservation, and construction of genetic linkage maps. Recently, a new approach to tree breeding that relies on molecular markers for full pedigree reconstruction following polycross mating was proposed (Lambeth *et al.*, 2001). This technology allows for making greater gains while reducing breeding and

testing costs. The use of markers for MAS and MAB will be discussed in the next section. In short, the application of molecular marker technology in forestry is extensive and likely to expand in the years ahead.

### Marker-assisted selection and marker-assisted breeding

MAS and MAB refer to approaches to tree improvement that rely on the statistical association of molecular markers with desirable genetic variants. With the development of new and easily obtained molecular markers in the 1990s, the prospect for practising MAS/MAB was bright. Fifteen years of research around the globe has both tempered and rejuvenated this prospect.

Initially, MAS was attempted by creating genetic linkage maps using molecular markers in segregating populations (pedigrees or crosses), and placing quantitative trait loci (QTLs) that explained some portion of the variation in a trait of interest (e.g. wood density) on those maps. Markers are identified as being in close genetic linkage with the genes responsible for the trait of interest, and can be used to select for the desired alleles of those genes. In addition to MAS, potential applications for QTL maps include the genetic dissection of complex quantitative traits, and the provision of guidance for selection and prioritization of candidate genes (Wheeler *et al.*, 2005). QTL maps have been created for over two dozen forest tree species (Sewell and Neale, 2000). Though highly informative, QTL maps are difficult and costly to produce, and have utility limited largely to the pedigrees for which they were created. Use of this technology for MAS is modest, but finds strong advocates for selected applications in North America, Europe and New Zealand.

Currently, research on another approach to identifying QTLs using natural populations rather than pedigrees is receiving increasing attention in forestry and agriculture. This technology, called association genetics, proposes finding markers that tag the actual genetic variants that cause a phenotypic response (i.e. markers occurring within the gene of interest) (Neale and Savolainen, 2004). This approach holds great promise for MAS and MAB, and applications within forestry are possible within the next ten years.

### Genomics

Genomics is a recent field, with many subdisciplines (Krutovskii and Neale, 2001). Over the past six years, substantial resources have been invested in the genomics sciences of humans, agronomic crops and forest trees. Genomics encompasses a wide range of activities, including gene discovery, gene space and genome sequencing, gene function determination, comparative studies among species, genera and families, physical mapping and the burgeoning field of bio-informatics. The ultimate goal of genomics is to identify every gene and its related function in an organism.

The completion of a whole-genome sequence for *Populus trichocarpa* (Tuskan *et al.*, 2006) has laid the foundation for reaching this goal for a model species. Efforts follow to replicate this deed in *Eucalyptus* sp. and *Pinus* sp., though

progress may be slower due to larger genome sizes, in particular for pines. Gene and expressed sequence tag (EST) (cDNA) libraries for conifers by far exceed one million entries; however, not all entries are readily available to the scientific community due to private ownership. The immediate applications of genomics include identification of candidate genes for association studies and targets for genetic modification studies. Also, comparative studies of genes from different trees have revealed the great similarity among taxa throughout the conifers, and raise hope that what is learned from one species will benefit many others.

Genomic sciences, like the other '-omics', namely metabolomics and proteomics, require substantial investment and are done on a very large scale, primarily by commercial entities with highly-trained laboratory staff, technology protected by intellectual property rights (IPR) and vast bio-informatics and associated statistical capacity. In general, genomics currently represents the most rapidly expanding area of biotechnological research; however, in forestry, most of the activities are concentrating on high throughput gene discovery and function elucidation. Characterization of genetic components of disease or pest resistance is a rapidly expanding field (Ellis *et al.*, 2001; Gartland, Kellison and Fenning, 2002). Other applications are expected to increase to complement traditional tree improvement through association genetics (Neale and Savolainen, 2004).

### Proteomics

Proteomics is the large-scale study of the proteins expressed by an organism, particularly protein structure and function. The term 'proteomics' was coined to make an analogy with genomics, the study of the genes. The proteome of an organism is the set of proteins it produces during its life, and the genome of the organism is the set of genes it contains.

Proteomics is often considered the next step in the study of biological systems, after genomics. It is much more complicated than genomics, mostly because while an organism's genome is fairly constant, a proteome differs from cell to cell and constantly changes through its biochemical interactions with the genome and the environment. Another major difficulty is the complexity of proteins relative to nucleic acids. For example, in the human body there are about 25 000 identified genes, but an estimated >500 000 proteins are derived from these genes. This increased complexity derives from mechanisms such as alternative splicing, protein modification (glycosylation, phosphorylation) and protein degradation.

Proteomics has attracted much interest because it yields information that is potentially more complex and informative in comparison with that gained from genomic studies. The level of transcription of a gene provides an approximate estimate of its level of expression into a protein. An mRNA produced in abundance may be degraded rapidly, modified or translated inefficiently. This could result in reduced amounts or types of protein being produced. In addition, many transcripts give rise to more than one protein, through alternative splicing or alternative post-translational modifications. Many proteins form complexes with other proteins or RNA molecules, and only function in the presence of these other molecules.



Proteomic studies require significant analytical and biocomputing capability, including instrumentation such as electrophoresis, crystallography, infrared and mass spectroscopy, and matrix-assisted laser desorption/ionization – time-of-flight mass spectrometer (MALDI-TOF) equipment.

Proteomics can be of value to forestry in a number of ways. For example, a proteomic study with somatic embryogenesis in *Picea glauca* identified a number of differentially expressed proteins across different stages of embryogenesis (Lippert *et al.*, 2005). The knowledge gained from such experiments may help to better understand and manipulate the process of embryogenesis.

### Metabolomics

Metabolomics is the “systematic study of the unique chemical fingerprints that specific cellular processes leave behind” - specifically, the study of their small-molecule metabolite profiles. The metabolome represents the collection of all metabolites in a biological organism, which are the end products of its gene expression. Thus, while mRNA gene expression data and proteomic analyses do not tell the whole story of what might be happening in a cell, metabolic profiling can give an instantaneous snapshot of the physiology of that cell. One of the challenges of systems biology is to integrate proteomic, transcriptomic and metabolomic information to provide a more complete picture of living organisms. The typical technical approach to metabolomics is through mass spectroscopy.

Metabolomics can be an excellent tool for determining the phenotype caused by a genetic manipulation, such as gene deletion or insertion. Sometimes this can be a sufficient goal in itself, such as to detect any phenotypic changes in a genetically modified tree, and to compare this with the naturally occurring variation in a tree population. It can also be used to understand variation that is induced by various factors such as genetic or environmental factors. For example, a metabolomic study with field-planted Douglas fir found that environmental variation was greater than genetic variation (Robinson *et al.*, 2007).

### GENETIC ENGINEERING

Biotechnological advancements in crop improvement through genetic engineering have attracted great attention from both the scientific and lay communities. This is as true for forestry as it is for agriculture. In fact, genetic modification is so embedded in the public conscientiousness that it is often considered synonymous with the term biotechnology. However, genetic engineering represents only one-fifth of the total biotechnology activities published in the past ten years (Walter and Killerby, 2004). Genetic modification is frequently seen as the most controversial use of biotechnology (Dale, 1999; Stewart, Richards and Halfhill, 2000; Thompson-Campbell, 2000; Dale, Clarke and Fontes, 2002; Conner, Glare and Nap, 2003; Burdon and Walter, 2004; Walter, 2004a, b; Walter and Fenning, 2004).

A major apprehension with genetic modification is the possible widespread gene transfer via escapes and hybridization and/or introgression with related native species. This concern is particularly felt in areas where inter-fertile species

are present in the vicinity of a plantation of genetically modified plants and when measures to prevent gene flow are not considered. Various approaches have been considered to ensure containment of genetically modified organisms (GMOs) through sterility (Brunner *et al.*, 2007).

Compared with the advances made in agricultural biotechnology, which can now be seen through looking back at more than ten years of successful commercial application, forest genetic engineering has lagged behind. This is mainly due to much fewer resources, longer rotation times of the crop and significant hurdles to overcome with regard to efficient tissue culture and propagation technologies. The more recent development of efficient plant tissue culture techniques has allowed forestry to emulate what has been achieved for agricultural and horticultural species. While there have been major advances with conventional tree breeding, there are some desirable traits that are not available in the tree species of choice. Possible traits of interest include herbicide and insect resistance, and modified lignin and cellulose content (Hu *et al.*, 1999; Bishop-Hurley *et al.*, 2001; Pilate *et al.*, 2002; Grace *et al.*, 2005). Also, more recently, research has focused on traits that are associated with the wood secondary cell wall and that have the potential to make transformational changes to wood-based products (Wagner *et al.*, 2007; Li *et al.*, 2003; Moeller *et al.*, 2005). Of increasing interest is the current trend towards a bio-based economy that derives resource materials from plant matter rather than petrochemicals.

## GENETIC MODIFICATION TECHNOLOGIES

Two main technologies are available to transfer foreign DNA into plant cells, and then regenerate plants from these transformed cells. These technologies are the use of bacterium, typically *Agrobacterium tumefaciens* (Gelvin, 2003), or biolistics (gene gun) (Klein *et al.*, 1987). *A. tumefaciens* is a bacterium that causes crown gall disease in some, particularly dicotyledonous, plants. The bacterium characteristically infects a wound, and incorporates a segment of Transfer-DNA (T-DNA) (syn. Ti [Tumour inducing] DNA) into the host genome. This DNA codes for the production of plant hormones and its expression in the host plant cell leads to undifferentiated growth. The T-DNA resides on a bacterial plasmid that also carries other genes (virulence or *vir* genes), which are responsible for the transfer of the T-DNA into the plant cells. The *A. tumefaciens* T-DNA can be replaced by any gene(s) of interest, which will then be transferred to plant cells during *A. tumefaciens* infection. Poplar was the first hardwood species to be transformed using this technology, with a herbicide resistance gene in 1987 (Fillatti *et al.*, 1987). Conifer species are difficult to transform using *A. tumefaciens*, although successful transformations of larch (*Larix decidua*) (Huang, Diner and Karnosky, 1991), pine (*Pinus radiata*) (Grant, Cooper and Dalr, 2004; Charity *et al.*, 2005) and spruce (*Picea* spp.) (Klimaszewska *et al.*, 2001; Le *et al.*, 2001) species has been reported (Henderson and Walter, 2006).

Biolistic techniques have now been developed to stably transform species that are difficult to transform using *A. tumefaciens* (Walter *et al.*, 1998, 1999; Find

*et al.*, 2005; Henderson and Walter, 2006; Trontin *et al.*, 2007). For this technology, the DNA is coated onto small metal particles (tungsten or gold) and these are propelled by various methods fast enough to puncture target cells. Typically, a pulse of pressurized helium is used to inject the particles into the target cells. Provided that the cell is not irretrievably damaged, the DNA can be taken up by the cell and integrated into its genome. Any transformed cells need to be actively selected from non-transformed cells, so that chimeric cell lines are avoided. This can be achieved by including a selectable marker gene in the transferred DNA, such as for antibiotic resistance. Following the transformation event, the cells are cultured on a medium containing the antibiotic. Over time, only stably-transformed cells will survive this exposure to an antibiotic, and so transformed cell lines can be established and tested for the presence of the new DNA. The efficiency of transclone production using biolistic techniques is usually slightly higher than when *A. tumefaciens* is used as a vector for gene transfer. However, recent modifications to the biolistic process (Walter, unpublished) have increased the efficiency significantly, so that more than 200 transclones can be produced by one operator in a single day. Transgenic plants can be regenerated from these cell lines and evaluated in greenhouse and field tests.

The successful expression of genes that are of commercial interest has already been demonstrated in laboratory and field experiments. These include the modification of lignin and cellulose biosynthesis (Hu *et al.*, 1999; Pilate *et al.*, 2002), herbicide resistance (Bishop-Hurley *et al.*, 2001), and insect resistance (Grace *et al.*, 2005). Field tests of transgenic pine plants produced through biolistic techniques have also demonstrated the long-term stability of the introduced gene, up eight years of age (Walter, in preparation).

Genetic modification technology is still new to forestry. However, relatively numerous (124) introduced traits of transgenic trees have been under regulatory examination in the United States of America (McLean and Charest, 2000), and a commercial plantation of genetically-modified poplar trees has been reported in China (Su *et al.*, 2003). A new wave of transgenic trees with improved secondary cell wall characteristics (improved pulpability, increased cellulose content, better stability) will soon be available for field testing and subsequent commercial deployment in plantation forestry. In many cases, particularly where interfertile species are present, reproductive sterility will be required to prevent introgression of transgenes into native populations (Brunner *et al.*, 2007; Höfig *et al.*, 2006).

Forestry genetic modification activities are taking place in at least 35 countries, 16 of which host some form of experimental field trials (Wheeler, 2004). These field trials are generally small (12 to 2 850 plants in reported studies) and typically of short duration. In many countries, such trials must be destroyed before seed production occurs. In other countries, experimentation is restricted to laboratories or greenhouses. To date, only China (Wang, 2004) has reported the establishment of approved, commercial plantations of genetically modified trees. While the majority of activities on genetic modification are experimental and regulated

under very strict conditions, concerns about genetically modified trees are similar to those about agricultural crops.

## REFERENCES

- Ahuja, M.-R. & Libby, W.J. (editors). 1993. *Clonal forestry II: conservation and application*. Springer-Verlag, Berlin and Heidelberg, Germany.
- Assis, T.F., Fett-Neto, A.G. & Alfenas, A.C. 2004. Current techniques and prospects for the clonal propagation of hardwoods with emphasis on Eucalyptus. pp. 303–333, in: C. Walter & M. Carson (editors). *Plantation forest biotechnology for the 21st century*. Research Signpost, Trivandrum, India.
- Bajaj, Y.P.S. (editor). 1986. *Biotechnology in agriculture and forestry 1: Trees I*. Springer-Verlag, Berlin, Germany.
- Bajaj, Y.P.S. (editor). 1989. *Biotechnology in agriculture and forestry 5: Trees II*. Springer-Verlag, Berlin, Germany.
- Bajaj, Y.P.S. (editor). 1991. *Biotechnology in agriculture and forestry 16: Trees III*. Springer-Verlag, Berlin, Germany.
- Benson, E. 2003. Conserving special trees: integrating biotechnological and traditional approaches. pp. 23–24, in: S. McCord & K. Gartland (editors). *Forest biotechnology in Europe: impending barriers, policy and implication*. Institute of Forest Biotechnology, Edinburgh, UK.
- Bishop-Hurley, S.L., Zabkiewicz, J.A., Grace, L., Gardner, R.C., Wagner, A. & Walter, C. 2001. Conifer GE: transgenic *Pinus radiata* (D. Don) and *Picea abies* (Karst) plants are resistant to the herbicide Buster. *Plant Cell Reports*, 20: 235–243.
- Bonga, J.M. & Durzan, D.J. 1987a. *Cell and tissue culture in forestry, Vol. 2: Specific principles and methods: growth and developments*. Martinus Nijoff Publishers, Dordrecht, Netherlands.
- Bonga, J.M. & Durzan, D.J. 1987b. *Cell and tissue culture in forestry, Vol. 3: Case histories: Gymnosperms, Angiosperms and Palms*. Martinus Nijoff Publishers, Dordrecht, Netherlands.
- Brunner, A., Li, J., DiFazio, S.P., Shevchenko, O., Montgomery, B.E., Mohamed, R., Wei, H., Ma, C., Elias, A.A., VanWormer, K. & Strauss, S.H. 2007. Genetic containment of forest plantations. *Tree Genetics and Genomes*, 3: 75–100.
- Burdon, R.D. & Walter, C. 2004. Exotic pines and eucalypts: perspectives on risks of transgenic plantations. In: S.H. Strauss & H.D. Bradshaw (editors). *The bioengineered forest: challenges for science and society*. Resources for the Future, Washington, DC, USA.
- Campbell, M.M., Brunner, A.M., Jones, H.M. & Strauss, S.H. 2003. Forestry's Fertile Crescent: the application of biotechnology to forest trees. *Plant Biotechnology Journal*, 1: 141–154.
- Chaix, G. & Monteuis, O. 2004. Biotechnology in the forestry sector. In: FAO, 2004b, q.v.
- Chalupa, V. 1985. Somatic embryogenesis and plantlet regeneration from cultured immature and mature embryos of *Picea abies* (L.) Karst. *Communications of the Institute for Forestry of the Czech Republic*, 14: 57–63.
- Charity, J.A., Holland, L., Grace, L.J. & Walter, C. 2005. Consistent and stable expression of the *nptII*, *uidA* and *bar* genes in transgenic *Pinus radiata* after *Agrobacterium*-mediated transformation using nurse cultures. *Plant Cell Reports*, 23: 606–616.
- Conner, A.J., Glare, T.R. & Nap, J.-P. 2003. The release of genetically modified crops into the environment. *The Plant Journal*, 33: 19–46.
- Cyr, D.R. 1999. Cryopreservation of embryogenic cultures of conifers and its application to clonal forestry. pp. 239–261, in: Jain, Gupta & Newton, 1999, q.v.
- Dale, P.D. 1999. Public concerns over transgenic crops. *Genome Research*, 9: 1159–1162.
- Dale, P.J., Clarke, B. & Fontes, M.G. 2002. Potential for the environmental impact of transgenic crops. *Nature Biotechnology*, 20: 567–574.
- El-Kassaby, Y.A. 2000. Effect of forest tree domestication on gene pools. pp. 197–213, in: A. Young, D. Boshier & T. Boyle (editors). *Forest conservation genetics: principles and practice*. Commonwealth Scientific and Industrial Research Organisation (CSIRO) Publishing-CABI Publishing, Canberra, Australia.

- El-Kassaby, Y.A. 2003. *Feasibility and proposed outline of a global review of forest biotechnology*. Discussion Paper of the Forest Resources Division, Forestry Department. FAO, Rome.
- El-Kassaby, Y.A. 2004. Anticipated contribution to and scale of impact of biotechnology in forestry. *In*: FAO, 2004b, q.v.
- Ellis, D., Meillan, R., Pilate, G. & Skinner, J.S. 2001. Transgenic trees: where are we now? pp. 113–123, *in*: S.H. Strauss & H.D. Bradshaw (editors). *Proceedings 1st International Symposium on Ecological and Societal Aspects of Transgenic Plantations*. College of Forestry, Oregon State University, Corvallis, Oregon, USA.
- FAO. 2001. Glossary of biotechnology for food and agriculture: a revised and augmented edition of the glossary of biotechnology and genetic engineering. *FAO Research and Technology Paper*, No. 9. Rome.
- FAO. 2004a. The State of Food and Agriculture 2003–2004. Agricultural biotechnology: meeting the needs of the poor? *FAO Agriculture Series*, No. 35. Rome.
- FAO. 2004b. *Preliminary review of biotechnology in forestry, including genetic modification*. Forest Genetic Resources Working Paper FGR/59E. Forest Resources Development Service, Forest Resources Division. Rome (available at [www.fao.org/docrep/008/ae574e/ae574e00.htm](http://www.fao.org/docrep/008/ae574e/ae574e00.htm)).
- Fillatti, J.J., Sellmer, J., McCown, B., Haissig, B. & Comai, L. 1987. *Agrobacterium*-mediated transformation and regeneration of *Populus*. *Molecular and General Genetics*, 206: 192–199.
- Find, J.I., Charity, J.A., Grace, L.J., Kristensen, M.M.M.H., Krogstrup, P. & Walter, C. 2005. Stable genetic transformation of embryogenic cultures of *Abies nordmanniana* (Nordman fir) and regeneration of transgenic plants. *In vitro Cellular & Developmental Biology–Plant*, 41: 725–730.
- Frampton, L.J. Jr & Isik, K. 1987. Comparison of field growth among Loblolly pine seedlings and three plant types produced *in vitro*. *Tappi*, 70(7): 119–123.
- Gartland, K.M.A., Kellison, R.C. & Fenning, T.M. 2002. Forest biotechnology and Europe's forests of the future. *In*: *Proceedings, Forest biotechnology in Europe: impending barriers, policy and implications*. Edinburgh, UK.
- Gelvin, S.B. 2003. *Agrobacterium*-mediated plant transformation: the biology behind the 'gene jockey' tool. *Microbiology and Molecular Biology Reviews*, 67(1): 16–37.
- Grace, L.J., Charity, J.A., Gresham, B., Kay, N. & Walter, C. 2005. Insect-resistant transgenic *Pinus radiata*. *Plant Cell Reports*, 24(2): 103–111.
- Grant, J.E., Cooper, P.A. & Dalr, T.M. 2004. Transgenic *Pinus radiata* from *Agrobacterium tumefaciens* mediated transformation of cotyledons. *Plant Cell Reports*, 108(6): 1177–1181.
- Gupta, P.K., Timmis, R. & Holmstrom, D. 2005. Cryopreservation of embryonal cells. pp. 567–572, *in*: Jain & Gupta, 2005, q.v.
- Hakman, I. & von Arnold, S. 1985. Plantlet regeneration through somatic embryogenesis in *Picea abies* (Norway spruce). *Journal of Plant Physiology*, 121: 149–158.
- Hargreaves, C.L. & Menzies, M.I. 2007. Organogenesis and cryopreservation of juvenile radiata pine. pp. 51–65, *in*: S. Jain & M. Haggman (editors). *Protocols for micropropagation of woody trees and fruits*. Springer, Berlin, Germany.
- Hargreaves, C.L., Smith, D.R., Foggo, M.N. & Gordon, M.E. 1997. Cryopreservation of zygotic embryos of *Pinus radiata* and subsequent plant regeneration. pp. 281–284, *in*: R.D. Burdon & J.M. Moore (editors). *IUFRO '97 Genetics of radiata pine*. Proceedings of conference, 1–4 December 1997, Rotorua, New Zealand. NZ Forest Research Institute, FRI Bulletin, No. 203.
- Hargreaves, C.L., Foggo, M.N., Smith, D.R. & Gordon, M.E. 1999. Development of protocols for the cryopreservation of zygotic embryos of *Pinus radiata* and subsequent plant regeneration. *NZ Journal of Forest Science*, 29(1): 54–63.
- Henderson, A.R. & Walter, C. 2006. Genetic engineering in conifer plantation forestry. *Silvae Genetica*, 55(6): 253–262.
- Höfig, K.P., Möller, R., Donaldson, L., Putterill, J. & Walter, C. 2006. Towards male-sterility in *Pinus radiata* – a stilbene synthase approach to genetically engineer nuclear male sterility. *Plant Biotechnology Journal*, 4: 333–343.



- Horgan, K., Skudder, D. & Holden, G. 1997. Clonal storage and rejuvenation. pp. 273–280, in: R.D. Burdon & J.M. Moore (editors). *IUFRO '97 Genetics of radiata pine*. Proceedings of conference, 1–4 December 1997, Rotorua, New Zealand. NZ Forest Research Institute, FRI Bulletin, No. 203.
- Hu, W.-J., Harding, S.A., Lung, J., Popko, J.L., Ralph, J., Stokke, D.D., Tsai, C.-J. & Chiang, V.L. 1999. Repression of lignin biosynthesis promotes cellulose accumulation and growth in transgenic trees. *Nature Biotechnology*, 17: 808–815.
- Huang, Y., Diner, A.M. & Karnosky, D.F. 1991. *Agrobacterium rhizogenes* mediated genetic transformation and regeneration of a conifer: *Larix decidua*. In vitro *Cellular & Developmental Biology–Plant*, 27: 201–207.
- Jain, S.M., Gupta, P.K. & Newton, R.J. (editors). 1999. *Somatic embryogenesis in woody plants*, Vol. 4. Kluwer Academic Publishers, Dordrecht, Netherlands.
- Jain, S.M., Gupta, P.K. & Newton, R.J. (editors). 2000. *Somatic embryogenesis in woody plants*, Vol. 6. Kluwer Academic Publishers, Dordrecht, Netherlands.
- Jain, S.M. & Gupta, P.K. (editors). 2005. *Protocol for somatic embryogenesis in woody plants*. Springer, Dordrecht, Netherlands.
- Klein, T.M., Wolf, E.D., Wu, R. & Sanford, J.C. 1987. High-velocity microprojectiles for delivering nucleic acids into living cells. *Nature*, 327: 70–73.
- Klimaszewska, K., Lachance, D., Pelletier, G., Lelu, A.M. & Seguin, A. 2001. Regeneration of transgenic *Picea glauca*, *P. mariana* and *P. abies* after co-cultivation of embryogenic tissue with *Agrobacterium tumefaciens*. In vitro *Cellular & Developmental Biology–Plant*, 37(6): 748–755.
- Krutovskii, K.V. & Neale, D.B. 2001. *Forest genomics for conserving adaptive genetic diversity*. Forest Genetics Working Paper FGR/3. Forest Resources Development Service, Forest Resources Division, FAO. Rome.
- Lambeth, C., Lee, B.-C., O'Malley, D.M. & Wheeler, N.C. 2001. Polymix breeding combined with paternal analysis (PMX/WPA) of progeny: an alternative to full-sib breeding and testing systems. *Theoretical and Applied Genetics*, 103: 930–943.
- Le, V.Q., Belles-Isles, J., Dusabenyagasani, M. & Tremlay, F.M. 2001. An improved procedure for production of white spruce (*Picea glauca*) transgenic plants using *Agrobacterium tumefaciens*. *Journal of Experimental Botany*, 364: 2089–2095.
- Li, M. & Ritchie, G.A. 1999. Eight hundred years of clonal forestry in China: I. Traditional afforestation with Chinese fir (*Cunninghamia lanceolata* (Lamb.) Hook). *New Forests*, 18: 131–142.
- Li, L., Zhou, Y., Cheng, X., Sun, J., Marita, J.M., Ralph, J. & Chianf, V.L. 2003. Combinatorial modification of multiple lignin traits in trees through multigene co-transformation. *Proceedings of the National Academy of Sciences of the United States of America*, 100(8): 4939–4944.
- Lippert, D., Zhuang, J., Ralph, S., Ellis, D.E., Gilbert, M., Olafson, R., Ritland, K., Ellis, B., Douglas, C.J. & Bohlmann, J. 2005. Proteome analysis of early somatic embryogenesis in *Picea glauca*. *Proteomics*, 5(2): 461–473.
- McCown, B.H. & Sellmer, J.C. 1987. General media and vessels suitable for woody plant culture. pp. 4–16, in: J.M. Bonga & D.J. Durzan (editors). *Cell and tissue culture in forestry, Vol. 1: General principles and biotechnology*. Martinus Nijhoff Publishers, Dordrecht, Netherlands.
- McLean, M.A. & Charest, P.J. 2000. The regulation of forest trees in North America. *Silvae Genetica*, 49(6): 233–239.
- Menzies, M.I. & Aimers-Halliday, J. 1997. Propagation options for clonal forestry with *Pinus radiata*. pp. 256–263, in: R.D. Burdon & J.M. Moore (editors). *IUFRO '97 Genetics of radiata pine*. Proceedings of conference, 1–4 December 1997, Rotorua, New Zealand. NZ Forest Research Institute, FRI Bulletin, No. 203.
- Menzies, M.I. & Aimers-Halliday, J. 2004. Propagation options for clonal forestry with conifers. pp. 255–274, in: C. Walter & M. Carson (editors). *Plantation forest biotechnology for the 21st century*. Research Signpost, Trivandrum, Kerala, India.
- Moeller, R., Steward, D., Phillips, L., Flint, H. & Wagner, A. 2005. Gene silencing of cinnamyl alcohol dehydrogenase in *Pinus radiata* cell cultures. *Plant Physiology and Biochemistry*, 43: 1061–1066.

- Neale, D.B. & Savolainen, O. 2004. Association genetics of complex traits in conifers. *Trends in Plant Science*, 9(7): 325-330.
- Pilate, G., Guiney, E., Holt, K., Petit-Conil, M., Lapierre, C., Leple, J.C., Ppillet, B.; Mila, I., Webster, E.A., Marstrop, H.G., Hopkins, D.W., Jouanin, L., Boerjan, W., Schuch, W., Cornu, D. & Halpin, C. 2002. Field and pulping performances of transgenic trees with altered lignification. *Nature Biotechnology*, 20(6): 558-560.
- Ritchie, G.A. 1991. The commercial use of conifer rooted cuttings in forestry: a world overview. *New Forests*, 5: 247-275.
- Ritland, C. & Ritland, K. 2000. DNA-fragment markers in plants. pp. 208-234, in: A.J. Baker (editor). *Molecular methods in ecology*. Blackwell Scientific, Oxford, UK.
- Robinson, A.R., Ukrainetz, N.K., Kang, K.-Y. & Mansfield, S. 2007. Metabolite profiling of Douglas fir (*Pseudotsuga menziesii*) field trials reveals strong environmental and weak genetic variation. *New Phytologist*, 174(4): 762-773.
- Sewell, M.M. & Neale, D.B. 2000. Mapping quantitative traits in forest trees. pp. 407-424, in: S.M. Jain & S.C. Minocha (editors). *Molecular biology of woody plants*. Kluwer Academic Publishers, Dordrecht, Netherlands.
- Sommer, H.E. & Brown, C.L. 1980. Embryogenesis in tissue cultures of sweetgum. *Forest Science*, 26(2): 257-260.
- Stewart, C.N., Richards, H.A. & Halfhill, M.D. 2000. Transgenic plants and biosafety: science, misconceptions and public perceptions. *BioTechniques*, 29: 832-843.
- Su, X.-H., Zhang, B.-Y., Huang, Q.-J., Huang, L.-J. & Zhang, X.-H. 2003: Advances in tree genetic engineering in China. Paper submitted to the XII World Forestry Congress.2003, Quebec City, Canada (available at [www.fao.org/DOCREP/ARTICLE/WFC/XII/0280-B2.HTM](http://www.fao.org/DOCREP/ARTICLE/WFC/XII/0280-B2.HTM)).
- Thompson-Campbell, F.T. 2000. *Genetically engineered trees: questions without answers*. American Lands Alliance. Washington, DC, USA.
- Toda, R. 1974. Vegetative propagation in relation to Japanese forest tree improvement. *NZ Journal of Forestry Science*, 4(2): 410-417.
- Trontin, J.-F., Walter, C., Klimaszewska, K., Park, Y.-S. & Walter, M.-A. 2007. Recent progress in genetic transformation of four *Pinus* spp. *Transgenic Plant Journal*, 1(2): 314-329.
- Tuskan, G.A., DiFazio, S., Jansson, S. *et al.* 2006. The genome of black cottonwood, *Populus trichocarpa* (Torr & Gray). *Science*, 313(5793): 1596-1604.
- Walter, C. 2004a. Stability of novel gene expression in transgenic conifers: An issue of concern? In: A. Mujib, M.J. Cho, S. Predieri & S. Banerjee (editors). *In vitro application in crop improvement*. Science Publishers Inc., Enfield, New Hampshire, USA.
- Walter, C. 2004b. Genetic engineering in conifer forestry: technical and social considerations. *In vitro Cellular & Developmental Biology-Plant*, 40(5): 434-441.
- Walter, C. & Fenning, T. 2004. Deployment of genetically-engineered trees in plantation forestry - An issue of concern? The science and politics of genetically-modified tree plantations. pp. 423-446, in: C. Walter & M. Carson (editors). *Plantation forest biotechnology for the 21st century*. Research Signpost, Trivandrum, India.
- Walter, C. & Killerby, S. 2004. A global study on the state of forest tree genetic modification. In: FAO, 2004b, q.v.
- Walter, C., Grace, L.J., Wagner, A., White, D.W.R., Walden, A.R., Donaldson, S.S., Hinton, H., Gardner, R.C. & Smith, D.R. 1998. Stable transformation and regeneration of transgenic plants of *Pinus radiata* D Don. *Plant Cell Reports*, 17: 460-468.
- Walter, C., Grace, L.J., Donaldson, S.S., Moody J., Gemmell, J.E., Van Der Maas, S., Kwaalen, H. & Loenneborg, A. 1999. An efficient biolistic transformation protocol for *Picea abies* (L.) Karst. embryogenic tissue and regeneration of transgenic plants. *Canadian Journal of Forestry Research*, 29: 1539-1546.
- Wagner, A., Ralph, J., Akiyama, T., Flint, H., Phillips, L., Torr, K., Nanayakkara, B. & Te Kiri, L. 2007. Exploring lignification in conifers by silencing hydroxycinnamoyl-CoA:shikimate hydroxycinnamoyltransferase in *Pinus radiata*. *Proceedings of the National Academy of Sciences of the United States of America*, 104(28): 11856-11861.

- Wang, H.** 2004. The state of genetically modified forest trees in China. *In*: FAO, 2004b, q.v.
- Wheeler, N.** 2004. A snapshot of the global status and trends in forest biotechnology. *In*: FAO, 2004b, q.v.
- Wheeler, N.C., Adams, W.T. & Hamrick, J.L.** 1993. Pollen distribution in wind-pollinated seed orchards. pp. 25–31, *in*: D.L. Bramlett, G.A. Askew, T.D. Blush, F.E. Bridgwater & J.B. Jett (editors). *Advances in pollen management. USDA Forestry Service Agricultural Handbook*, No. 698.
- Wheeler, N., Jermstad, K.D., Krutovsky, K., Aitken, S.N., Howe, G.T., Krakowski, J. & Neale, D.B.** 2005. Mapping of quantitative trait loci controlling adaptive traits in coastal Douglas fir. IV. Cold hardiness QTL verification and candidate gene mapping. *Molecular Breeding*, 15: 145–156.
- Yanchuk, A.D.** 2001. The role and implications of biotechnological tools in forestry. *Unasylva*, 204: 53–61.



## 2. Biotechnology techniques

*R. Meilan, Z. Huang and G. Pilate*

Biotechnology can be divided into two broad areas: conventional breeding and molecular genetics. The former has been used for centuries to improve plant and animal species to satisfy human needs. Advances in molecular genetics have been rapidly adopted by the scientific community over the last two decades, and they complement tools already available to conventional breeders.

Molecular genetics can itself be subdivided into two distinct categories. In the first, which could be called ‘non-controversial technologies’, the plant genome is not altered. This category comprises molecular markers, which are used for DNA fingerprinting and MAS (e.g. QTL mapping and association genetics); sequence analysis (genomic DNA, cDNA libraries [ESTs], and bacterial artificial chromosome [BAC] clones), which aid in gene discovery; and *in vitro* propagation (e.g. somatic embryogenesis). The benefits of research using these technologies are increased genetic gain per generation through improved selection in conventional breeding programmes, faster deployment of genetically improved material to plantations, and a deeper understanding of the genes controlling commercially important traits.

The second major subdivision of molecular genetics, termed ‘controversial technologies’, includes recombinant DNA and gene-transfer techniques. These are the basis for genetic engineering, which is defined as the stable, usually heritable, modification of an organism’s genetic makeup via asexual gene transfer, regardless of the origin and nature of the introduced gene. The product of this process is generally referred to as a genetically modified organism (GMO). Genetic engineering offers the opportunity to add new genes to existing, elite genotypes. Although much progress has been made, genetically engineered forest species are not likely to be deployed commercially in much of the world for several more years. One reason for this delay is our limited understanding of the key genes that contribute to the control of commercially important traits, such as wood properties, flowering control and pest resistance. Research in these areas will broaden our knowledge of the genetic and physiological mechanisms that govern tree growth and development. In addition, it will allow the assessment of risks associated with these controversial technologies—assessments that will be required if we are to produce genetically improved material for meeting the growing societal demands for high-quality wood and fibre (Farnum, Lucier and Meilan, 2007).

To make more rapid progress with tree biotechnology, certain innovations are needed, including improved regeneration protocols, alternative *in vitro* selection strategies, dependable excision mechanisms and reliable confinement strategies. One limitation is in our understanding of the roles played by genes controlling key aspects of tree development. Poplar is widely accepted as the model tree for

forest biology owing to its small genome, expanding molecular resources, fast growth, and the relative ease with which it can be clonally propagated *ex vitro* and transformed and regenerated *in vitro* (Bradshaw *et al.*, 2000; Wullschleger, Jansson and Taylor, 2002). The recently released *Populus trichocarpa* genome sequence (Tuskan *et al.*, 2006) and newly developed genomics approaches have already and will continue to expedite gene discovery. The knowledge gained through our work with poplar can then be applied to other tree species.

## TECHNIQUES

### Recombinant DNA

The application of a variety of techniques collectively referred to as ‘recombinant DNA technology’ permits the study of gene structure and function, gene transfer to various species, and the efficient expression of their products. Using microbiological methods, it is possible to combine genetic material from various organisms in novel ways. Through these techniques it has been possible to expand our knowledge concerning the way in which genes are regulated, eukaryotes synthesize proteins, and eukaryotic genomes are organized. With regard to genetic engineering, recombinant DNA techniques are essential for:

- identifying genes responsible for specific traits;
- isolating these genes;
- creating genetic constructs harbouring both these genes and flanking regulatory sequences needed for expression in the host organism (in our case, a tree);
- selecting transgenic cells (generally by using an antibiotic or herbicide resistance gene).

Once genetically modified plants have been produced, this technology also allows us to select the best individuals with preferred levels of integration and expression and to monitor, at the molecular level, whether transgene integration and expression are maintained from one growing season to the next, after sexual reproduction, and in various environments.

### Transformation

The main steps required for the production of GMOs are:

- stably introducing a novel piece of DNA into the genome of a cell (i.e. transformation);
- isolating transgenic plant cells on a medium containing a selection agent (e.g. the antibiotic or herbicide against which the selectable marker gene imparts resistance);
- regenerating whole plants from the transformed cells through *in vitro* culture;
- screening various transgenic lines that result from independent transformation events on the basis of insert copy number and configuration, and expression.

To date, much of the research on genetic engineering of trees has concentrated on optimizing transformation. Three gene-transfer techniques are commonly

utilized here: protoplast transformation, biolistics and *Agrobacterium*-mediated transformation. Historically, angiosperms were transformed primarily through the use of *Agrobacterium tumefaciens*. Because of early difficulties encountered when transforming conifers with common *Agrobacterium* strains, gymnosperms were initially transformed using particle bombardment (Pena and Seguin, 2001). These problems have now largely been resolved, and several different species are being efficiently transformed via standard *Agrobacterium* strains (e.g. Pilate *et al.*, 1999; Tang, Newton and Weidner, 2007; Tereso *et al.*, 2006). However, except for larch (*Larix kaempferi* × *L. decidua*) (Levee *et al.*, 1997), much work remains to be done on the other steps leading to the production of genetically modified trees, particularly with regard to the regeneration of whole plants from transgenic cells. Plants are regenerated through one of two methods: direct organogenesis or somatic embryogenesis. The latter leads to the production of embryos from somatic tissues, whereas the former involves the generation of organs, such as shoots and roots, from various mature tissues or undifferentiated cell masses derived therefrom. No matter which approach is used, *in vitro* regeneration is often a genotype-dependent process.

### Protoplast transformation

Protoplasts are derived by enzymatically digesting the walls of plant cells that are usually isolated from the leaf mesophyll, and are often grown in a liquid suspension culture. Frequently, protoplasts can be transformed either by direct DNA uptake, following polyethylene glycol pre-treatment, or by electroporation. Although many studies have resulted in successful transient expression of a transgene in cell-derived protoplasts (Bekkaoui, Tautorius and Dunstan, 1995), very few have described the regeneration of transgenic trees (e.g. Chupeau, Pautot and Chupeau, 1994). This is probably due to difficulties in regenerating whole plants from protoplasts.

### Biolistics

Particle bombardment relies on the delivery of DNA-coated tungsten or gold microprojectiles, which are accelerated variously by ignited gunpowder, compressed gases (helium, nitrogen or carbon dioxide) or electrical discharge (Hansen and Wright, 1999). Although this technique was used to produce some of the first transgenic plants from recalcitrant coniferous or monocotyledonous species (Klein *et al.*, 1988; Ellis *et al.*, 1993), such transformation efficiency remains generally low, and usually results in a high number of transgene inserts in the genome. For these reasons, direct DNA transfer techniques have been avoided in favour of *Agrobacterium*-mediated protocols.

### **Agrobacterium-mediated transformation.**

*Agrobacterium tumefaciens* is a soil-borne bacterium responsible for crown gall, a disease of dicotyledonous plants that causes chaotic cell proliferation at the infection site, ultimately leading to the development of a plant tumour. During

the complex infection process, bacterial DNA is stably incorporated into the plant genome. Today *A. tumefaciens* co-cultivation is the most widely used and preferred method for transforming many types of plants (reviewed by Gelvin, 2003).

*A. tumefaciens* harbours a large, tumour-inducing (Ti) plasmid, which encodes several products needed to transfer a piece of its DNA into the host-plant genome. This transferred sequence, called T-DNA, contains a region delimited by two borders, and carries genes that are responsible for tumour development and for the synthesis of opines (molecules that serve as a carbon and nitrogen source for the bacterium, and which result from an association between amino acids and sugars). The virulence genes (*Vir*), located outside the T-DNA region on the Ti plasmid, facilitate T-DNA transfer.

This naturally occurring mechanism for DNA transfer has been exploited by plant biotechnologists, who have demonstrated that the bacterium recognizes the DNA to be transferred to the plant cell genome by its unique borders. An *A. tumefaciens* strain is said to be disarmed when the genes within those T-DNA borders are removed. Another plasmid, a binary vector that contains the genes of interest between the border sequences, is then transformed into the disarmed strain of *A. tumefaciens*. The *Vir* genes located on the disarmed vector are able to act in trans.

The transfer of T-DNA into the host-plant genome takes place following the co-cultivation of explants (generally leaf disks, petioles, stem internodes or root segments) with the bacterium. The explants are then extensively washed to remove excess bacterium before being maintained on media containing bacteriostats (e.g. cefotaxime or timentin) and the appropriate selection agent. Transgenic cells are multiplied then transferred to a series of media that have been optimized to contain the proper amounts of nutrients and plant growth regulators so that the various phases of plant regeneration are induced through either somatic embryogenesis or organogenesis.

The first genetically modified tree, a poplar, was produced 20 years ago (Fillatti *et al.*, 1987). Today, the number of forest tree species for which transformation and regeneration techniques have been optimized remains low; they include aspen, cottonwood, eucalyptus and walnut. Recently, transformation and regeneration protocols have been developed for several gymnosperms, mostly species within the genera *Pinus*, *Larix* and *Picea*. Within each of these species, only a few genotypes have been amenable to the recovery of transgenic plants. In general, for a wide range of genotypes, effective plant regeneration has been more difficult to achieve through organogenesis than through somatic embryogenesis.

### Transgene type and its control

A gene comprises a coding sequence that is preceded by a promoter, which controls where, when and to what extent it will be expressed in a plant. This coding sequence might originate from a different species and therefore may not be present in the host plant. For example, Bt genes, which confer resistance to insects, are derived from a bacterium, *Bacillus thuringiensis*. Alternatively, the

transgene may already exist in the host plant (i.e. an endogene). For example, ferulate-5-hydroxylase (F5H) is an enzyme specific for the synthesis of syringyl lignins; homologues of this gene are found in angiosperm trees. In general, foreign genes are relatively easy to express in the host plant. Depending on the configuration of the genetic construct (e.g. the orientation of the coding sequence or the occurrence of an inverted repeat), expression of the introduced gene may be ectopic (e.g. expressed in a tissue or at a stage not ordinarily seen in the wild-type plant), elevated or down-regulated (e.g. RNA interference (RNAi)). Moreover, a promoter could be fused to a reporter gene, such as  $\beta$ -glucuronidase (GUS) (Jefferson, Burgess and Hirsch, 1986) or to the green fluorescence protein (GFP) gene from jellyfish (*Aequoria victoria*) (Haseloff *et al.*, 1997), which can be used to reveal the pattern of expression conferred by a given promoter.

## IDENTIFYING CANDIDATE GENES

### Mutation analysis

Several experimental approaches have been taken to isolate genes that either confer a commercially useful trait or control a key aspect of plant development. The first, mutation analysis, involves screening thousands and possibly millions of seedlings for rare mutations that might aid in identifying desirable genes. This is a random, hit-or-miss approach that is slow, labour-intensive and sporadic when applied to tree species. In addition, because trees have long generation times, mate by cross-pollination and are highly heterozygous, rare recessive mutations are difficult to detect. A directed programme of inbreeding could be employed to expose recessive mutations, but inbreeding can also result in trees with poor form and low vigour owing to their high genetic loads, confounding attempts to identify valuable alleles. Tree improvement through these conventional means could require many decades, even with rapid advances in the area of plant genetics and the ease with which biotechnological tools can be applied to certain tree species (e.g. poplar; Bradshaw and Strauss, 2001).

### *In silico* cloning

A second method for identifying candidate genes involves utilizing information from other model plants, such as the herbaceous annual *Arabidopsis thaliana*, to identify tree orthologs. An example of this approach is the identification of the *NAC1* gene, a root-specific member of a family of transcriptional regulators in plants. A mutation in *NAC1* diminishes lateral root formation and perturbs expression of *AIR3* (Xie *et al.*, 2000), a downstream gene associated with the emergence of lateral roots (Neuteboom *et al.*, 1999a, b). Furthermore, transgenic complementation with a functional *NAC1* gene restores lateral root formation, and overexpression results in a proliferation of lateral roots. Thus, the *NAC1* gene product appears to be both necessary and sufficient for lateral root formation. In this case, both sequence and functional information are being tested for functionality via transgenesis (B. Goldfarb, personal communication, North Carolina State University).

### Forward genomics

A third way to facilitate gene discovery relies on the use of direct, random mutagenesis. Gene and enhancer trapping are methods for insertion-based gene discovery that both reference genome sequence data and result in a dominant phenotype (Springer, 2000). In short, gene-trap vectors carry a reporter gene lacking a functional promoter, while enhancer-trap constructs contain a minimal promoter preceding a reporter gene. In each case, the reporter gene is expressed in a fashion that imitates the normal expression pattern of the native gene at the insertion site, as has been demonstrated for *Arabidopsis* gene- and enhancer-trap lines (e.g. Springer *et al.*, 1995; Gu *et al.*, 1998; Pruitt *et al.*, 2000). The genomic region flanking the insertion site is amplified using PCR and sequenced; alignment of the flanking sequence with the genome sequence allows immediate mapping of insertions (Sundaresan *et al.*, 1995). This technique has recently been applied to identify genes likely to be involved in vascularization (Groover *et al.*, 2004). A similar strategy, using a luciferase-based promoter-trap vector, has allowed the identification of tissue- or cell-specific promoters (Johansson *et al.*, 2003).

Another forward genomics approach, namely activation tagging, utilizes a strong enhancer element that is randomly inserted into the genome and can be effective some distance from a promoter (Weigel *et al.*, 2000). Elevated expression of the nearby native gene may result in an aberrant phenotype. Lines exhibiting an obvious difference (early flowering, modifications in crown form, adventitious root development, etc.) are then analysed for the causative gene. Overexpression of some native genes (e.g. those affecting wood quality) may not give rise to a visually apparent change. In such cases, high throughput analyses are needed for screening a population of transgenics. The feasibility of this approach has already been demonstrated in poplar (Busov *et al.*, 2003). The recent release of the annotated draft of the *Populus trichocarpa* genome ([www.phytozome.net/poplar.php](http://www.phytozome.net/poplar.php)) is facilitating the isolation and characterization of loci underpinning mutations found in similar ways.

### Microarrays

A fourth approach to identifying candidate genes utilizes differential gene expression. The development of microarray technology has provided biologists with a powerful tool for studying the effects of gene expression on development and environmental responses (Brown and Botstein, 1999; Rishi, Nelson and Goyal, 2002). Expression levels of entire suites of genes, of both known and unknown function, can be measured simultaneously rather than one or a few genes at a time. This approach has already been successful in many systems. For root formation, a screen of loblolly pine shoots given a rooting treatment (auxin pulse) yielded a putative membrane transport protein that was induced by auxin treatment in juvenile (rooting) but not in mature (non-rooting) stem bases (Busov *et al.*, 2004). This gene shows homology to a large multigene family in *Arabidopsis*, members of which are similar to what was first classified as a nodulin from alfalfa.

### PCR-based techniques

The fifth molecular technique to identify candidate genes is based on PCR, and includes suppression subtractive hybridization (SSH), differential display PCR (DD-PCR), and cDNA-AFLP (amplified fragment length polymorphism).

SSH is a PCR-based technique that was developed for the generation of subtracted cDNA libraries, and combines normalization and subtraction in a single procedure. Diatchenko *et al.* (1996) demonstrated that SSH could result in the enrichment of rare sequences by over 1000-fold in one round of subtractive hybridization. This technique has been a powerful tool for many molecular genetic and positional cloning studies to identify developmental, tissue-specific and differentially expressed genes (Matsumoto, 2006). For example, using SSH, bract-specific genes have been successfully identified in the ornamental tree *Davidia involucreata* (Li *et al.*, 2002), and genes responsive to benzothiadiazole (BTH; used to induce systemic acquired resistance) in the tropical fruit tree papaya (Qiu *et al.*, 2004). Genes involved in flowering have also been isolated from carnation (*Dianthus caryophyllus*; Ok *et al.*, 2003) and black wattle (*Acacia mangium*; Wang, Cao and Hong, 2005) using this method.

DD-PCR is another widely used method for detecting altered gene expression between samples, often derived from the treated and untreated individuals from the same genotype or species. An amplification is done using a primer that hybridizes to the poly(A) tail and an arbitrary 5' primer. The first application of this technology was reported by Liang and Pardee (1992), and has since been used with a wide variety of organisms, including bacteria, plants, yeast, flies and higher animals, to expedite gene discovery. A Myb transcription factor HbMyb1 associated with a physiological syndrome known as tapping panel dryness has been identified and characterized from rubber trees using differential display reverse transcriptase PCR (DDRT-PCR) (Chen *et al.*, 2002). Transcriptional profiling of gene expression from leaves of apricot (*Prunus armeniaca*) was conducted by DDRT-PCR and up- or down-regulated genes in response to European stone fruit yellows phytoplasma infection were identified (Carginale *et al.*, 2004). A significant disadvantage of this technique is its high percentage of false-positives (Zegzouti *et al.*, 1997).

cDNA-AFLP was first used by Bachem *et al.* (1996) to analyse differential gene expression during potato tuber development and was subsequently modified by Breyne *et al.* (2003). It too is a PCR-based method, which starts with cDNA synthesis, using random hexamer primers and total or mRNA as a template. Following digestion with two different restriction enzymes, adapters are ligated before amplification via PCR. This method has proven to be an efficient tool for differential quantitative transcript profiling and a useful alternative to microarrays (Breyne *et al.*, 2003). cDNA-AFLP was used to identify transcripts that accumulated in mature embryos and in *in vitro*-cultured plantlets subjected to desiccation or abscisic acid (ABA) treatment in almond (*Prunus amygdalus*; Campalans, Pages and Messeguer, 2001). Using this approach a novel gene, designated *Mal-DDNA*, was cloned and confirmed to play an important role in lowering the acidity of apple fruit (Yao *et al.*, 2007).



### RNA interface

Double-stranded RNA-mediated gene suppression, also known as RNA interference (RNAi), was first reported in *Caenorhabditis elegans* a decade ago (Fire *et al.*, 1998). It is currently the most widely used method to down-regulate gene expression. It can be used to knock out all copies of a given gene, thus providing insight into its functionality. However, it does not always result in complete inhibition of a gene's expression. Recent advances in targeted gene mutagenesis and replacement using the yeast *RAD54* gene (Shaked, Melamed-Bessudo and Levy, 2005) or zinc-finger nucleases (Lloyd *et al.*, 2005; Wright *et al.*, 2005) may eventually lead to efficient methods for engineering null alleles in trees.

## IMPROVEMENTS NEEDED

### Regeneration

Regeneration protocols are typically optimized for a single genotype by conducting complex, labour-intensive, complete-factorial experiments. A more universal protocol has not been developed because of a lack of fundamental understanding of how plant cells acquire the competence to regenerate *in vitro*. Using rapidly advancing genomics tools, it is now possible to unravel this mystery. The research community now has access to a chip on which sequence information for all poplar genes has been spotted. Using this microarray, it is possible to identify genes that interfere with or promote regeneration by evaluating expression levels for all genes in tissues that differ in their regeneration potential, before and after being induced to regenerate. In addition, gene expression profiling that is done on tissues gathered during the juvenility-to-maturity transition could help identify genes affecting regeneration, in a similar manner to the approach described by Brunner and Nilsson (2004) to identify genes involved in flowering control.

### Selection systems

As described above, a selectable marker gene is linked to the gene of interest that is being inserted. Transformed cells can then be isolated on a medium containing the appropriate selection agent. While this method is convenient, it is often problematic. First, performing subsequent rounds of transformation may not be possible because only a limited number of selectable marker genes are available. Second, various selection agents can have dramatic and negative effects on regeneration. Finally, the presence of a selectable marker gene is usually an impediment to gaining public acceptance of genetically engineered plants.

Recently, alternative selection systems have been developed. These are based on a growth medium that lacks a substance needed for metabolic activity or proper development. A particularly attractive option exploits the inability of a cell to regenerate a whole plant without the addition of a phytohormone, or its derivative, to the culture medium at a precise step in the regeneration process. For example, most regeneration protocols rely on an exogenous supply of cytokinin to induce differentiation of adventitious shoots or embryos from transgenic calli.



The *GUS* gene, a common reporter, encodes an enzyme that cleaves glucuronide residues. The glucuronide derivative of benzyladenine is biologically inactive; if it is the sole cytokinin incorporated in the induction medium, regeneration will not occur. However, upon hydrolysis by  $\beta$ -glucuronidase, a biologically active cytokinin is liberated to induce regeneration (Okkels, Ward and Joersbo, 1997). This supplement must necessarily be transitory because cytokinin can inhibit subsequent steps in development.

Another positive selection strategy involves inserting a gene whose product imparts a metabolic advantage to the transformed cell. Mannose is a sugar that plants are unable to metabolize; cells starve when grown on a medium containing mannose as the sole carbon source. When taken up by the cells, this sugar is phosphorylated by a native hexokinase. However, plants lack a native phosphomannose isomerase gene, which encodes an enzyme that catalyses the conversion of mannose to a usable six-carbon sugar (Joersbo *et al.*, 1998). Similarly, xylose isomerase, another enzyme that plants lack, is able to convert xylose to a sugar that can be utilized (Haldrup, Petersen and Okkels, 1998). Regeneration protocols that exploit positive-selection strategies such as these can be up to ten fold more efficient than those that rely on more traditional, negative-selection strategies.

### Excision systems

The ability to delete unwanted pieces of DNA reliably is a valuable tool for both basic and applied research. Excision systems can remove selectable marker genes, thereby alleviating public concern and allowing for easy re-transformation using vectors derived from a common backbone. Moreover, some alternative regeneration methods (e.g. MAT, discussed below) depend on excision for their success. Because transposons have proven too unreliable, alternative systems, such as *Cre/lox* (Russell, Hoopes and Odell, 1992), *FLP/FRT* (Lyznik, Rao and Hodges, 1996) and *R/RS* (Onouchi *et al.*, 1995), have been utilized. Excision vectors typically include a recombinase gene, usually under the control of an inducible promoter, and recognition sites that flank the DNA being targeted for removal. However, these systems have not proven to be reliable in certain plants. Thus, it is necessary to determine which is the most appropriate for use with various tree species. For each system, one must ascertain the efficacy of the recombinase and how cleanly it excises the target sequence. Moreover, it is imperative to have an inducible promoter that functions reliably in the plant being transformed.

### Producing marker-free plants

The recently developed multiautonomous transformation system (MAT) allows for the production of transgenic plants lacking selectable marker genes from a variety of species (e.g. tobacco, aspen, rice, snapdragon) (Ebinuma *et al.*, 1997; Ebinuma and Komamine, 2001). These vectors harbour *Agrobacterium* genes (*ipt* or *rol*) that control sensitivity to or the biosynthesis of phytohormones. Cells transformed with these vectors regenerate into plants with either a 'shooty'

or ‘hairy-root’ phenotype. MAT vectors also contain a site-specific, inducible recombinase for excision of both the recombinase and the oncogenes. This alternative production system is attractive because it has the potential to increase both the yield and speed with which transgenic plants can be produced, and may eliminate the need for specific selection and regeneration conditions, making it possible to transform a wider array of genotypes. Such a system will also be useful for stacking genes in forest trees, as described by Halpin and Boerjan (2003).

### Mitigating transgene spread

The Coordinated Framework of the United States Animal and Plant Health Inspection Service (APHIS) now gives consideration to transgenic woody perennials. It is likely that before such trees can be deployed commercially, a method to mitigate the risk of transgene spread in the environment will be required, particularly in the cases when the introduced gene will improve the fitness of the genetically engineered tree. Many researchers are investigating ways to modify floral development to satisfy this need. The two most common approaches are to engineer trees that are either reproductively sterile or have delayed flowering. The latter may be particularly useful for short-rotation intensive culture (SRIC), where trees are harvested before the onset of maturation. Nevertheless, the main techniques being employed to modify floral development are:

- cell ablation (floral-specific expression of a cytotoxin gene);
- RNAi (silencing native genes via short, interfering RNAs);
- dominant negative mutations (DNMs), which lead to the production of a dysfunctional version of a gene product, such as a transcription factor (reviewed by Meilan *et al.*, 2001).

Because of functional redundancy, suppression of more than one floral regulatory gene is likely to be needed to achieve complete sterility. Where redundancy is obvious, RNAi constructs can be designed to silence effectively several members of a multigene family (Waterhouse and Helliwell, 2003). It is also advisable to utilize multiple techniques (e.g. cell ablation, RNAi or DNM, alone or in combination) to alter the expression of genes in more than one family to increase the likelihood of developing a durable confinement strategy. Transgene expression has been found to be unstable under various conditions (Brandle *et al.*, 1995; Köhne *et al.*, 1998; Metz, Jacobsen and Stiekema, 1997; Neumann *et al.*, 1997; Scorza *et al.*, 2001). Matrix attachment regions (MARs) have been used to enhance and stabilize transgene expression (Han, Ma and Strauss, 1997; Allen, Spiker and Thompson, 2000); however, there is some question about their utility (Li *et al.*, 2008). Given the potential for instability, it will be imperative to conduct multiyear field studies, in a variety of environments, and extending past the onset of maturity, in order to ensure the reliability of a given confinement system.

Progress in this area has been hampered by the inherent, delayed maturation of trees. Even the five- to seven-year juvenile period for poplar is a serious impediment. There is a report of a *Populus alba* genotype (6K10) that can be

induced to flower precociously, but it is of limited practical use (Meilan *et al.*, 2004). Its induction regime is lengthy and complex, and specialized equipment is required. In addition, not every plant in a population responds to induction. Moreover, the efficiency with which the genotype can be transformed and regenerated is very low. Because both male and female sterility will be needed, poplar is dioecious and 6K10 is a female, confinement systems will need to be tested in another poplar genotype. Early-flowering genotypes are rare and many trees do not respond well to treatments that induce precocious flowering (Meilan, 1997). Thus, there is a need for alternative genotypes that can be reliably and efficiently induced to flower.

## BIO-INFORMATICS TECHNOLOGY

Bio-informatics is an interdisciplinary approach that utilizes computational and statistical techniques to aid in solving biological problems at the molecular level. Initially, bio-informatic tools were merely used to store, retrieve and analyse nucleic acid and protein sequence information. The field is now evolving rapidly, and being employed in newly emerging disciplines such as comparative genomics, transcriptomics, functional genomics and structural genomics. Below we briefly discuss some of the basic bio-informatics applications that are commonly used today.

### Sequence analysis

One of the fundamental goals of sequence analysis is to determine the similarity of unknown or 'query' sequences to those previously identified and stored in various databases. A commonly used algorithm known as BLAST (basic local alignment search tool) provides a way to rapidly search nucleotide and protein databases. Since BLAST performs both local and global alignments, regions of similarity embedded in other, seemingly unrelated, proteins can be detected. Sequence similarity can provide important clues concerning the function of uncharacterized genes and the proteins they encode.

Other sequence-analysis tools are available to aid in determining the biological function and structure of genes and proteins, or to cluster them into related families based on their sequence information. Some software packages need to be purchased, others are available at no cost. The European Molecular Biology Open Software Suite (EMBOSS) is free, open-source software that can be downloaded from <http://emboss.sourceforge.net/>. It integrates many bio-informatics tools for sequence analysis into a single environment and can be used to analyse DNA and protein sequence in a variety of formats. Within EMBOSS there are hundreds of applications covering areas such as sequence alignment, rapid database searching for sequence patterns (e.g. to identify islands or repeats), protein motif identification (domain analysis), codon usage analysis for small genomes, and rapid identification of sequence patterns in large sequence sets. In addition, because extensive libraries are provided with this package, it is possible for users to develop and release software of their own. An example of another integrated

bio-informatics software can be found at <http://ca.expasy.org/tools>. As with EMBOSS, this package is helpful for characterizing and predicting the function of biomolecules of interest. Other commonly used sequence analysis applications include ClustalW and IMAGE.

### Structure prediction

There are also software packages that can predict protein structure based on its sequence information or that of the gene encoded by it. Understanding protein structure is the key to revealing its function. Currently there are many programs for performing primary, secondary and tertiary structural analyses. ProtParam is a tool that computes physical or chemical parameters for a protein, such as molecular weight, amino acid and atomic composition, isoelectric point, extinction coefficient, estimated half-life, stability index and aliphatic index, based on user-entered sequence information. RasMol is an excellent graphics tool for visualizing macromolecular structure in order to help elucidate function. Other structure-prediction programs include Dowser, FastDNAMl, LOOPP, MapMaker/QTL and PAML.

### THE -OMICS

The 'omics' suffix is used to describe disciplines in which researchers analyse biological interactions on a genome-wide scale. The associated prefix indicates the object of study in each field. Examples include genomics, transcriptomics, metabolomics and proteomics. These encompass the study of the genetic make-up, the complete set of mRNA produced, the collection of metabolites, and protein function and interaction, respectively, in organisms, tissues or cells. The main focus of -omics is on gathering information at a given level and using computer-based tools to identify relationships in order to understand heterogeneous, biological networks, often with the ultimate goal of manipulating regulatory mechanisms. Omics require a multidisciplinary approach, bringing scientists together from a variety of fields to interpret the data collected.

### APPLICATIONS

Rapidly emerging biotechnological tools can be used to help us better understand how biological systems function. The resulting discoveries allow us to introduce novel or alter existing traits that are useful to humans. Chapter 4 by McDonnell *et al.* in this volume provides a description of some commercially important and environmentally beneficial traits that have been incorporated into trees.

### REFERENCES

- Allen, G.C., Spiker, S.L. & Thompson, W.F. 2000. Use of matrix attachment regions (MARs) to minimize transgene silencing. *Plant Molecular Biology*, 43: 361–376.
- Bachem, C.W.B., van der Hoeven, R.S., de Bruijn, S.M., Vreugdenhil, D., Zabeau, M. & Visser, R.G.F. 1996. Visualization of differential gene expression using a novel method of RNA fingerprinting based on AFLP: analysis of gene expression during potato tuber development. *Plant Journal*, 9: 745–753.

- Bekkaoui, F., Tautorius, T.E. & Dunstan, D.I. 1995. Gymnosperm protoplasts. pp. 167–191, in: Jain, S.M., Gupta, P.K. & Newton, R.J. (editors). *Somatic embryogenesis in woody plants*, Vol. 1. Kluwer Academic Publishers, Dordrecht, Netherlands.
- Bradshaw, H.D. Jr & Strauss, S.H. 2001. Breeding strategies for the 21st century: domestication of poplar. pp. 383–394, in: D.I. Dickmann, J.G. Isebrands, J.E. Eckenwalder & J. Richardson (editors). *Poplar culture in North America*. Part 2. NRC Research Press, National Research Council of Canada, Ottawa, Canada.
- Bradshaw, H.D. Jr, Ceulemans, R., Davis, J. & Stettler, R.F. 2000. Emerging model systems: poplar (*Populus*) as a model forest tree. *Journal of Plant Growth Regulation*, 19: 306–313.
- Brandle, J.E., McHugh, S.G., James, L., Labbe, H. & Miki, B.L. 1995. Instability of transgene expression in field grown tobacco carrying the *csr1-1* gene for sulfonylurea herbicide resistance. *Bio-Technology*, 13(9): 994–998.
- Breyne, P., Dreese, R., Cannoot, B., Rombaut, D., Vandepoele, K., Rombauts, S., Vanderhaeghen, R., Inze, D. & Zabeau, M. 2003. Quantitative cDNA-AFLP analysis for genome-wide expression studies. *Molecular Genetics and Genomics*, 269: 173–179.
- Brown, P.O. & Botstein, D. 1999. Exploring the new world of the genome with DNA microarrays. *Nature Genetics*, 21: 33–37.
- Brunner, A.M. & Nilsson, O. 2004. Revisiting tree maturation and floral initiation in the poplar functional genomics era. *New Phytologist*, 164(1): 43–51.
- Busov, V.B., Meilan, R., Pearce, D.W., Ma, C., Rood, S.B. & Strauss, S.H. 2003. Activation tagging of a dominant gibberellin catabolism gene (*GA 2-oxidase*) from poplar that regulates tree stature. *Plant Physiology*, 132: 1283–1291.
- Busov, V.B., Johannes, E., Whetten, R.W., Sederoff, R.R., Spiker, S.L., Lanz-Garcia, C. & Goldfarb, B. 2004. An auxin-inducible gene from loblolly pine (*Pinus taeda* L.) is differentially expressed in mature and juvenile-phase shoots and encodes a putative transmembrane protein. *Planta*, 218(6): 916–927.
- Campalans, A., Pages, M. & Messegue, R. 2001. Identification of differentially expressed genes by the cDNA-AFLP technique during dehydration of almond (*Prunus amygdalus*). *Tree Physiology*, 21: 633–643.
- Carginale, V., Maria, G., Capasso, C., Ionata, E., Cara, F.L., Pastore, M., Bertaccini, A. & Capasso, A. 2004. Identification of genes expressed in response to phytoplasma infection in leaves of *Prunus armeniaca* by messenger RNA differential display. *Gene*, 332: 29–34.
- Chen, S., Peng, S., Huang, G., Wu, K., Fu, X. & Chen, Z. 2002. Association of decreased expression of a Myb transcription factor with the TPD (tapping panel dryness) syndrome in *Hevea brasiliensis*. *Plant Molecular Biology*, 51: 51–58.
- Chupeau, M.C., Pautot, V. & Chupeau, Y. 1994. Recovery of transgenic trees after electroporation of poplar protoplasts. *Transgenic Research*, 3(1): 13–19.
- Diatchenko, L., Lau, Y.F.C., Campbell, A.P., Chenchik, A., Mogadam, F., Huang, B., Lukyanov, S., Lukyanov, K., Gurskaya, N., Sverdlov, E.D. & Siebert, P.D. 1996. Suppression subtractive hybridization: a method for generating differentially regulated or tissue-specific cDNA probes and libraries. *Proceedings of the National Academy of Sciences of the United States of America*, 93(12): 6025–6030.
- Ebinuma, H. & Komamine, A. 2001. MAT (Multi-Auto-Transformation) vector system. The oncogenes of *Agrobacterium* as positive markers for regeneration and selection of marker-free transgenic plants. In vitro *Cellular & Developmental Biology-Plant*, 37(2): 103–113.
- Ebinuma, H., Sugita, K., Matsunaga, E. & Yamakado, M. 1997. Selection of marker-free transgenic plants using the isopentenyl transferase gene. *Proceedings of the National Academy of Sciences of the United States of America*, 94(6): 2117–2121.
- Ellis, D.D., McCabe, D.E., McInnis, S., Ramachandran, R., Russell, D.R., Wallace, K.M., Martinell, B.J., Roberts, D.R., Raffa, K.F. & McCown, B.H. 1993. Stable transformation of *Picea glauca* by particle acceleration. *Bio-Technology*, 11(1): 84–89.
- Farnum, P., Lucier, A. & Meilan, R. 2007. Ecological and population genetics research imperatives for transgenic trees. *Tree Genetics and Genomes*, 3(2): 119–133.

- Fire, A., Xu, S., Montgomery, M.K., Kostas, S.A., Driver, S.E. & Mello, C.C. 1998. Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegans*. *Nature*, 391: 806–811.
- Fillatti, J.J., Sellmer, J., McCown, B., Haissig, B. & Comai, L. 1987. *Agrobacterium*-mediated transformation and regeneration of *Populus*. *Molecular Genomics and Genetics*, 206: 192–196.
- Gelvin, S.B. 2003. *Agrobacterium*-mediated plant transformation: the biology behind the “gene-jockeying” tool. *Microbiology and Molecular Biology Reviews*, 67(1): 16–37.
- Groover, A., Fontana, J., Dupper, G., Ma, C., Martienssen, R., Strauss, S. & Meilan, R. 2004. Gene and enhancer trap tagging of vascular-expressed genes in poplar trees. *Plant Physiology*, 134: 1742–1751.
- Gu, Q., Ferrandiz, C., Yanofsky, M. & Martienssen, R. 1998. The FRUITFULL MADS-box gene mediates cell differentiation during *Arabidopsis* fruit development. *Development*, 125: 1509–1517.
- Haldrup, A., Petersen, S.G. & Okkels, F.T. 1998. Positive selection: a plant selection principle based on xylose isomerase, an enzyme used in the food industry. *Plant Cell Reporter*, 18(1-2): 76–81.
- Halpin, C. & Boerjan, W. 2003. Stacking transgenes in forest trees. *Trends in Plant Science*, 8: 363–365.
- Han, K.-H., Ma, C. & Strauss, S.H. 1997. Matrix attachment regions (MARs) enhance transformation frequency and transgene expression in poplar. *Transgenic Research*, 6(6): 415–420.
- Hansen, G. & Wright, M.S. 1999. Recent advances in the transformation of plants. *Trends in Plant Science*, 4: 226–231.
- Haseloff, J., Siemering, K.R., Prasher, D.C. & Hodge, S. 1997. Removal of a cryptic intron and subcellular localization of green fluorescent protein are required to mark transgenic *Arabidopsis* plants brightly. *Proceedings of the National Academy of Sciences of the United States of America*, 94: 2122–2127.
- Jefferson, R.A., Burgess, S.M. & Hirsch D. 1986.  $\beta$ -glucuronidase from *Escherichia coli* as a gene-fusion marker. *Proceedings of the National Academy of Sciences of the United States of America*, 83: 8447–8451.
- Joersbo, M., Donaldson, I., Kreiberg, J., Petersen, S.G., Brunstedt, J. & Okkels, F.T. 1998. Analysis of mannose selection used for transformation of sugar beet. *Molecular Breeding*, 4(2): 111–117.
- Johansson, A.-M., Wang, C., Stenberg, A., Hertzberg, M., Little, C.H.A. & Olsson, O. 2003. Characterization of a PttRPS18 promoter active in the vascular cambium region of hybrid aspen. *Plant Molecular Biology*, 52(2): 317–329.
- Klein, T.M., Fromm, M., Weissinger, A., Tomes, D., Schaaf, S., Sletten, M. & Sanford, J.C. 1988. Transfer of foreign genes into intact maize cells with high-velocity micro-projectiles. *Proceedings of the National Academy of Sciences of the United States of America*, 85: 4305–4309.
- Köhne, S., Neumann, K., Pühler, A. & Broer, I. 1998. The heat-treatment-induced reduction of the *pat* gene encoded herbicide resistance in *Nicotiana tabacum* is influenced by the transgene sequence. *Plant Physiology*, 153: 631–642.
- Levee, V., Lelu, M.A., Jouanin, L., Cornu, D. & Pilate, G. 1997. *Agrobacterium tumefaciens*-mediated transformation of hybrid larch (*Larix kaempferi*  $\times$  *L. decidua*) and transgenic plant regeneration. *Plant Cell Reports*, 16(10): 680–685.
- Li, J., Brunner, A.M., Shevchenko, O., Meilan, R., DiFazio, S.P., Ma, C., Skinner, J.S. & Strauss, S.H. 2008. Efficient and stable transgene suppression via RNAi in field-grown poplars. *Transgenic Research*, 17(4): 679–694.
- Li, Y.X., Chen, L., Juan, L., Li, Y. & Chen, F. 2002. Suppression subtractive hybridization cloning of cDNAs of differentially expressed genes in Dovetree (*Davidia involucrata*) bracts. *Plant Molecular Biology Reports*, 20: 231–238.
- Liang, P. 2005. A decade of differential display. *Biotechniques*, 33: 338–346.
- Liang, P. & Pardee, A.B. 1992. Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science*, 257(5072): 967–971.



- Lloyd A., Plaisier, C.L., Carroll, D. & Drews, G.N. 2005. Targeted mutagenesis using zinc-finger nucleases in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America*, 102: 2232–2237.
- Lyznik, L.A., Rao, K.V. & Hodges, T.K. 1996. FLP-mediated recombination of FRT sites in the maize genome. *Nucleic Acids Research*, 24: 3784–3789.
- Matsumoto T.K. 2006. Genes uniquely expressed in vegetative and potassium chlorate induced floral buds of *Dimocarpus longan*. *Plant Science*, 170(3): 500–510.
- Meilan, R. 1997. Floral induction in woody angiosperms. *New Forests*, 14(3): 179–202.
- Meilan, R., Brunner, A., Skinner, J. & Strauss, S. 2001. Modification of flowering in transgenic trees. pp. 247–256, in: A. Komamine & N. Morohoshi (editors). *Molecular breeding of woody plants*. (Progress in Biotechnology series.) Elsevier Science BV, Amsterdam, Netherlands.
- Meilan, R., Sabatti, M., Ma, C. & Kuzminsky, E. 2004. An early-flowering genotype of *Populus*. *Journal of Plant Biology*, 47(1): 52–56.
- Metz, P.L.J., Jacobsen, E. & Stiekema, W.J. 1997. Occasional loss of expression of phosphinothricin tolerance in sexual offspring of transgenic oilseed rape (*Brassica napus* L.). *Euphytica*, 98(3): 189–196.
- Neumann, K., Dröge-Laser, W., Köhne, S. & Broer, I. 1997. Heat treatment results in a loss of transgene-encoded activities in several tobacco lines. *Plant Physiology*, 115: 939–947.
- Neuteboom, L.W., Ng, J.M., Kuyper, M., Clijdesdale, O.R., Hooykaas, P.J. & van der Zaal, B.J. 1999a. Isolation and characterization of cDNA clones corresponding with mRNAs that accumulate during auxin-induced lateral root formation. *Plant Molecular Biology*, 39: 27–278.
- Neuteboom, L.W., Veth-Tello, L.M., Clijdesdale, O.R., Hooykaas, P.J. & van der Zaal, B.J. 1999b. A novel subtilisin-like protease gene from *Arabidopsis thaliana* is expressed at sites of lateral root emergence. *DNA Research*, 6: 13–19.
- Ok, S.H., Park, H.M., Kim, J.Y., Bahn, S.C., Bae, J.M., Suh, M.C., Jeung, J.U., Kim, K.N. & Shin, J.S. 2003. Identification of differentially expressed genes during flower development in carnation (*Dianthus caryophyllus*). *Plant Science*, 165: 291–295.
- Okkels, F.T., Ward, J.L. & Joersbo, M. 1997. Synthesis of cytokinin glucuronides for the selection of transgenic plant cells. *Phytochemistry*, 46: 801–804.
- Onouchi, H., Nishihama, R., Kudo, M., Machida, Y. & Machida, C. 1995. Visualization of site-specific recombination catalyzed by a recombinase from *Zygosaccharomyces rouxii* in *Arabidopsis thaliana*. *Molecular Genomics and Genetics*, 247: 653–660.
- Pena, L. & Seguin, A. 2001. Recent advances in the genetic transformation of trees. *Trends in Biotechnology*, 19(12): 500–506.
- Pilate, G., Leplé, J.C., Cornu, D. & Lelu, M.A. 1999. Transgenic larch (*Larix* species). pp. 125–141, in: Y.P.S. Bajaj (editor). *Transgenic trees*. Vol. 44. Springer-Verlag, Berlin and Heidelberg, Germany.
- Pruitt, R., Vielle-Calzada, J., Ploense, S., Grossniklaus, U. & Lolle, S. 2000. FIDDLEHEAD, a gene required to suppress epidermal cell interactions in *Arabidopsis*, encodes a putative lipid biosynthetic enzyme. *Proceedings of the National Academy of Sciences of the United States of America*, 97: 1311–1316.
- Rishi, A.S., Nelson, N.D. & Goyal, A. 2002. DNA micro-arrays: gene expression profiling in plants. *Reviews in Plant Biochemistry and Biotechnology*, 1: 81–100.
- Russell, S.H., Hoopes, J.L. & Odell, J.T. 1992. Directed excision of a transgene from the plant genome. *Molecular Genomics and Genetics*, 234: 49–59.
- Qiu, X., Guan, P., Wang, M.L., Moore, P.M., Zhu, Y.J., Hu, J., Borth, W. & Albert, H.H. 2004. Identification and expression analysis of BTH induced genes in papaya. *Physiological and Molecular Plant Pathology*, 65(1): 21–30.
- Scorza, R., Callahan, A., Levy, L., Damsteegt, V., Webb, K. & Ravelonandro, M. 2001. Post-transcriptional gene silencing in plum pox virus-resistant transgenic European plum containing the plum pox potyvirus coat protein gene. *Transgenic Research*, 10(3): 201–209.

- Shaked, H., Melamed-Bessudo, C. & Levy, A.A. 2005. High-frequency gene targeting in *Arabidopsis* plants expressing the yeast *RAD54* gene. *Proceedings of the National Academy of Sciences of the United States of America*, 102: 12265–12269.
- Springer, P. 2000. Gene traps: tools for plant development and genomics. *Plant Cell*, 12: 1007–1020.
- Springer, P., McCombie, W., Sundaresan, V. & Martienssen, R. 1995. Gene trap tagging of Prolifera, an essential MCM2-3-5-like gene in *Arabidopsis*. *Science*, 268: 877–880.
- Sundaresan, V., Springer, P., Volpe, T., Haward, S., Jones, J., Dean, C., Ma, H. & Martienssen, R. 1995. Patterns of gene action in plant development revealed by enhancer trap and gene trap transposable elements. *Genes and Development*, 9: 1797–1810.
- Tang, W., Newton, R.J. & Weidner, D.A. 2007. Genetic transformation and gene silencing mediated by multiple copies of a transgene in eastern white pine. *Journal of Experimental Botany*, 58(3): 545–554.
- Tereso, S., Miguel, C., Zoglauer, K., Valle-Piquera, C. & Oliveira, M.M. 2006. Stable *Agrobacterium*-mediated transformation of embryogenic tissues from *Pinus pinaster* Portuguese genotypes. *Plant Growth Regulation*, 50(1): 57–68.
- Tuskan, G.A., DiFazio, S., Jansson, S. *et al.* 2006.: The genome of black cottonwood, *Populus trichocarpa* (Torr & Gray). *Science*, 313(5793): 1596–1604.
- Wang, X.J., Cao, X.L. & Hong, Y. 2005. Isolation and characterization of flower-specific transcripts in *Acacia mangium*. *Tree Physiology*, 25: 167–185.
- Waterhouse, P.M. & Helliwell, C.A. 2003. Exploring plant genomes by RNA-induced gene silencing. *Nature Reviews Genetics*, 4: 29–38.
- Weigel, D., Ahn, J.H., Blazquez, M.A., Borevitz, J.O., Christensen, S.K., Fankhauser, C., Ferrandiz, C., Kardailsky, I., Malancharuvil, E.J., Neff, M.M., Nguyen, J.T., Sato, S., Wang, Z.Y., Xia, Y.J., Dixon, R.A., Harrison, M.J., Lamb, C.J., Yanofsky, M.F. & Chory, J. 2000. Activation tagging in *Arabidopsis*. *Plant Physiology*, 122(4): 1003–1013.
- Wright, D.A., Townsend, J.A., Winfrey R.J. Jr, Irwin, P.A., Rajagopal, J., Lonosky, P.M., Hall, B.D., Jondle, M.D. & Voytas, D.F. 2005. High-frequency homologous recombination in plants mediated by zinc-finger nucleases. *Plant Journal*, 44: 693–705.
- Wullschleger, S.D., Jansson S. & Taylor G. 2002. Genomics and forest biology: *Populus* emerges as the perennial favorite. *Plant Cell*, 14: 2651–2655.
- Xie, Q., Frugis, G., Colgan, D. & Chua, N.-H. 2000. *Arabidopsis* *NAC1* transduces auxin signal downstream of *TIR1* to promote lateral root development. *Genes & Development*, 14(23): 3024–3036.
- Yao, Y.X., Li, M., Liu, Z., Hao, Y.J. & Zhai, H. 2007. A novel gene, screened by cDNA-AFLP approach, contributes to lowering the acidity of fruit in apple. *Plant Physiology and Biochemistry*, 45(2): 139–145.
- Zegzouti, H., Marty, C., Jones, B., Bouquin, T., Latché, A., Pech, J. & Bouzayen, M. 1997. Improved screening of cDNA generated by mRNA differential display enables the selection of true positives and the isolation of weakly expressed messages. *Plant Molecular Biology Reports*, 15: 238–245.



## 3. Genetic containment of forest plantations

*A.M. Brunner, J. Li, S.P. DiFazio, O. Shevchenko, B.E. Montgomery, R. Mohamed, H. Wei, C. Ma, A.A. Elias, K. VanWormer and S.H. Strauss*

“It is essential that new molecular gene-containment strategies... be developed and introduced.”

Editorial, *Nature Biotechnology*, 20: 527 (2002)

### CONTEXT FOR GENE CONTAINMENT APPROACHES

In an ideal world, industrial forest plantations would operate in harmony with, and in isolation from, natural ecosystems. Plantations would occur within a landscape designed to maintain biodiversity and minimize ecological impacts of plantations on external ecosystems, and economic goals would be the primary consideration within plantations. However, the reality is that plantations have multiple ecological connections with other managed and wild ecosystems and operate in a social milieu where their actual and perceived impacts may or may not be tolerated. Regulations, laws, and marketplace mechanisms such as certification systems set limits on the kinds of activities that may occur within plantations and on the impacts that these activities may have outside of plantations. All of these mechanisms strongly constrain research and commercial application of genetically engineered trees (reviews in Strauss and Bradshaw, 2004). Genetically engineered, genetically modified or transgenic organisms, as used in this review paper, are defined as those that have been modified using recombinant DNA and asexual gene transfer methods – regardless of the source of the DNA employed.

Forest certification systems represent a growing mechanism for expression of social preferences in the marketplace (Cashore, Auld and Newsom, 2003). One major forestry certification system aimed at environmental and social compliance, that of the Forest Stewardship Council, bans all forms of genetically engineered trees on certified lands. This rule is absolute; it applies regardless of the level of containment, whether the genes are from the same or different species, whether the goal is purely scientific research vs application, or whether the primary aim is the solution of substantial environmental problems rather than economic benefits (Strauss *et al.*, 2001a, b). Such a broad ban, which covers even contained research with environmental goals, is difficult to justify on scientific grounds, especially given the long-standing scientific consensus that “product not process” should dominate risk assessment for genetically engineered organisms (Snow *et al.*, 2005). It shows that social considerations can overwhelm technical innovations.

---

This review was first published in *Tree Genetics & Genomes*, 3: 75–100, © Springer-Verlag 2007, and is reproduced here by kind permission of Springer Science and Business Media. A.M. Brunner and S.H. Strauss contributed equally to this paper.

Thus, containment systems may be required even for genes where no significant biological impact, or even a positive environmental effect, is expected to occur. By allowing effective isolation of trees produced in different ways on the landscape, containment systems should provide a mechanism whereby different social values can more easily co-exist.

However, genetic mechanisms for isolation have never before been required even when highly bred or exotic species have been used in agriculture or forestry; their novelty, therefore, creates new forms of social controversy. Although genetic containment systems have long been called for by ecologists and other scientists to reduce a number of undesired effects of genetically engineered crops (NRC, 2004; Snow *et al.*, 2005), there has been strong pressure on companies and governments against use of any forms of ‘Terminator-like’ containment technology (ETC, 2006). For example, a law against the use of such technology in Brazil (Law 11,105/05, banning “...the commercialization of any form of Gene Use Restriction Technology (GURTs)”) delayed approval of a field trial of a reduced-lignin, putatively sterile eucalypt (ISAAA, 2006). In agriculture, these concerns primarily are about control of intellectual property and the forced repurchase of seed by farmers. But in the forestry area, there has also been activism against containment technology because of a lack of confidence that it will be fully effective, concerns about loss of biodiversity associated with modification or loss of floral tissues (Cummins and Ho, 2005), and legal uncertainties and liability risks from the dispersal of patented genes. These biological concerns occur despite the intention to use such technology mainly in plantations that, due to breeding, high planting density and short life spans, already produce few flowers and seeds compared with long-lived and open-grown trees. The powerful inverse association between forest stand density and degree of tree reproduction is widely known (Daniel, Helms and Baker, 1979). There is also an abundance of means to avoid and mitigate such effects at gene to landscape levels (Johnson and Kirby, 2004; Strauss and Brunner, 2004). Government regulations against the dispersal of genes from research trials also pose very substantial barriers to field research to study the efficiency of containment mechanisms (Strauss *et al.*, 2004; Valenzuela and Strauss, 2005). Thus, genetic containment technology is, itself, difficult and highly controversial, requiring special social conditions even to carry out research.

From a biological viewpoint, however, there are good reasons to employ containment technologies to control some forms of highly domesticated, exotic or genetically engineered organisms. Once genes or organisms move beyond plantation boundaries, the risks to external ecosystems are virtually impossible to control, and as with other biological introductions of mobile organisms, may be irreversible. Novel organisms of all kinds may impair the health of some wild ecosystems or create management problems for human-dominated ecosystems (James *et al.*, 1998). If we could confidently segregate intensely domesticated trees by control of reproduction, it would avoid the need for much of the complex, imprecise and costly ecological research that would otherwise be required to try to understand and predict impacts of spread. The costs and obstacles to conducting

commercially relevant environmental research with genetically engineered trees are great and occur for a number of reasons:

- laboratory cost of genetically engineered tree production, including production and study of many kinds of gene constructs and gene transfer events;
- ecological complexity in space and time and high stochastic variance in gene flow and related ecological processes, requiring many sites, environmental conditions, long time frames and large spatial scales;
- cost of needed patents, licenses, publication agreements, and transactions for access to genes intended for commercial use (required if results are to be directly relevant to regulatory decisions);
- cost of record keeping and compliance with regulations, which can be very demanding and legally risky for complex programmes that span many years and sites;
- uncertainty over what data regulators will require due to vagueness in regulatory standards and political volatility creating substantial changes in regulations or their interpretations over time;
- risk of spread into the environment during research, including costly steps to prevent any spread (e.g. premature termination of trials, bagging all flowers in test plantings, use of non-commercial but sterile genotypes, or use of geographically distant planting environments);
- disincentives to undertaking costly and risky research, as a result of possible marketplace rejection and separation costs; other significant disincentives result from primary ownership of the genes and gene transfer methods generally being out of the hands of the tree breeders and producers that bear most of the risks and costs of field testing.

These very formidable obstacles, many of which have substantial similarities in many other crop species, have forced companies and governments to ask whether these obstacles do more harm than good by blocking economically and environmentally beneficial technologies. It has also prompted calls for regulations that would place genetically engineered organisms into risk categories that call for dramatically different levels of research and containment depending on the novelty and risk of the new traits (Bradford *et al.*, 2005). For example, it has been suggested that ‘genomics guided transgenes (GGTs)’, where the expression of native or functionally homologous genes are altered in a manner analogous to conventional breeding, and ‘domestication transgenes’ that encode traits highly likely to reduce fitness in the wild, should be put into a low risk category or exempted from regulation entirely (Strauss, 2003). In contrast, new types of genetically engineered plants that are more likely to produce ecologically novel traits, or produce hazardous forms of pharmaceutical or industrial compounds, would be regulated with increased stringency. The Animal and Plant Health Inspection Service (APHIS) of the United States Department of Agriculture (USDA), which regulates all field research in the United States of America, is currently undergoing a major review, with one goal being the creation of risk categories. The obstacles to

field research have also called for increased emphasis on ecogenetic models, where the spread and impacts of transgenes with different properties, and under different environmental and social conditions, can be studied over decades as they spread within the containment of a computer (reviewed below).

The sense for a mandate to use containment technologies was also inspired by the creation of genetically engineering-based male and female sterility mechanisms during the early 1990s (Mariani *et al.*, 1990, 1992), when the possibilities of plant biotechnology seemed limitless, public acceptance was not an issue, and regulatory hurdles appeared modest (reviewed below). It was also stimulated by the suggestion of ‘mitigation’ genes that can both increase value in managed environments and reduce competitive ability in the wild (Gressel, 1999). If gene spread creates irreversible risks and social discomfort, and technology exists to greatly reduce these risks, is it not the ethical responsibility of scientists and companies to act to minimize these risks? The incorporation of biosafety features into genetically engineered organisms during their design has been promoted as key elements of good stewardship (Doering, 2004).

Unfortunately, as discussed above and in genetic detail below, applying containment technology to trees is an extremely costly and difficult endeavour. Caution is therefore warranted in assuming that containment systems – even the use of genes with a neutral or negative effect on fitness – present good stewardship. If genetic containment were incomplete, genes that provide a significant and evolutionarily highly stable selective advantage (should such transgenes be feasible to create and deploy), could eventually spread widely. Even neutral or deleterious genes can persist and even become fixed in wild populations in situations where transgenes numerically swamp native genes (Haygood, Ives and Andow, 2003). Obtaining licences to the set of patents that cover all of the elements of the best containment technology can also be very costly or impossible. At the same time, it is also likely that the spread of fitness-improving transgenes could, in some cases, provide ecological benefits. A gene for resistance against a serious exotic pest of trees such as the chestnut blight or Asian longhorn beetle might provide large ecological benefits by maintaining or restoring healthy ecological dominants and their dependent communities. Genes for general pest or abiotic stress resistance, including against native herbivores or pathogens, might also provide net ecological benefits by increasing the vigour of a native organism like poplar, which provides habitat for myriad dependent organisms (Whitham *et al.*, 2006), even if some introduced herbivores or plant species were disadvantaged as a consequence. It is therefore essential that containment technology is not indiscriminately required by regulations or used when its net benefits are questionable.

The goal of the remainder of this paper is to review the state of sterility technology that might be useful for sexual containment of trees used in clonal forestry and ornamental horticulture. We previously reviewed the many options for sex-specific sterility and inducible sterility/fertility (Strauss *et al.*, 1995) that might be used to enable continued seed propagation. Here, we focus on complete sterility under some form of vegetative propagation. Only after a simple method for strong

and bisexual sterility is shown to be effective and socially accepted is it likely that more sophisticated methods for fertility control will be developed and deployed.

## TECHNICAL APPROACHES AND THEIR ADVANTAGES AND DISADVANTAGES

Below, we discuss the main approaches to engineering containment relevant to forest trees. In addition, via electronic searches, we have scanned the recent (2000 to time of writing) scientific and patent (United States Patent and Trademark Office (US PTO)) literature and presented representative examples of developments. Tables 3-1 and 3-2 summarize the kinds of approaches being taken, nearly all of which are relevant to one kind of tree species or another.

TABLE 3-1

### Selected literature on genetic engineering of sterility published from 2000 onwards

Phenotype	Mechanism	Promoter	Active gene	Plant species	Reference
<b>Delayed flowering</b>					
Late flowering	Overexpression of <i>FLM</i>	35S CaMV	<i>Flowering Locus M</i>	<i>Arabidopsis</i>	Scortecci, Michaels and Amasino, 2001
	<i>AGL20</i> /shoot apical meristem	35S CaMV	<i>AGAMOUS LIKE 20</i>	<i>Arabidopsis</i>	Borner <i>et al.</i> , 2000
<b>Cell ablation</b>					
Male sterility	Altered pollen development	Endosperm specific promoter, <i>AGP2</i>	Fission yeast <i>cdc25</i>	Wheat	Chrimes <i>et al.</i> , 2005
	Pollen sterility	Rice tapetum promoter ( <i>TAP</i> )	Barnase/rice tapetum gene <i>rts</i>	Creeping bentgrass	Luo <i>et al.</i> , 2005
	Alteration in tapetal cells	<i>Tapetum A9</i> promoter	Chimeric gene in transgenic plant	<i>Arabidopsis</i>	Guerineau <i>et al.</i> , 2003
	Abnormal pollen	<i>BcA9</i>	<i>DTx-A</i>	<i>Brassica</i>	Lee <i>et al.</i> , 2003
	Tapetal dysfunction	<i>TA29</i> promoter	<i>RIP</i>	Tobacco	Cho <i>et al.</i> , 2001
	Reduced pollen viability	Pollen specific promoter <i>G9</i>	Chimeric genes <i>G9 uidA</i> and <i>G9-RNase</i>	Tobacco	Bernd-Souza <i>et al.</i> , 2000
Male and female sterility	Floral organ ablation with otherwise normal growth	<i>PopulusPTD</i>	<i>DTA</i>	Tobacco, poplar, <i>Arabidopsis</i>	Skinner <i>et al.</i> , 2003
Recoverable block of function (RBF)	Inducible fertility	Sulfhydryl endopeptidase, heat-shock promoter	<i>Barnase</i> (the blocking construct) and <i>barstar</i> (recovering construct)	Tobacco	Kuvshinov <i>et al.</i> , 2001
<b>Gene suppression</b>					
Male sterility	Distorted pollen morphology	Various	<i>AtMYB32 AtMYB4</i>	<i>Arabidopsis</i>	Preston <i>et al.</i> , 2004
	Temperature sensitive male sterility due to silencing choline biosynthesis	S-adenosyl-L-methionine	Phosphoethanolamine N-methyltransferase (PEAMT)	<i>Arabidopsis</i>	Mou <i>et al.</i> , 2002
	Mitochondrial dysfunction	Tapetum specific promoter	Antisense pyruvate dehydrogenase E1 $\alpha$ subunit	Sugar beet	Yui <i>et al.</i> , 2003
	Abnormal pollen	<i>Nin88</i> promoter	Antisense <i>Nin88</i>	Tobacco	Goetz <i>et al.</i> , 2001
	Abnormal pollen	Glutenin subunit gene promoter	Antisense sucrose non-fermenting-1-related (SnRK1) protein kinase	Barley	Zhang <i>et al.</i> , 2001
Restoration of fertility	Glucanase gene suppression	<i>pA9</i>	Sense and antisense PR glucanase	Tobacco	Hird <i>et al.</i> , 2000

TABLE 3-2  
Selected patents on genetic engineering of sterility published from 2000 onwards

Phenotype	Mechanism	Promoter	Active gene/Protein	Species	Reference
<b>Time of flowering</b>					
Altered floral development	Expression of floral meristem identity protein	Modified native promoter	<i>CAULIFLOWER (CAL)</i> , <i>APETELA 1 (AP1)</i> , <i>LEAFY (LFY)</i>	Angiosperm or gymnosperm	Yanofsky, 2000
<b>Cytotoxin ablation</b>					
Suicide gene to ablate gamete	Any of several cytotoxic genes expressed in gametes	Male- or female-specific promoter expressed in gamete	Various "suicide" genes ( <i>barnase</i> , <i>tasselseed2</i> , <i>diphtheria toxin A</i> )	Rice	Dellaporta and Moreno, 2004
Female sterility	Enhance fruit development or induce sterility	<i>DefH9</i> promoter	DNases, RNases, proteases, glucanases, lipases, toxins, etc.	Many	Spena <i>et al.</i> , 2002
<b>Gene suppression</b>					
Male sterility	Calcium/calmodulin-dependent protein kinase (CCaMK) expression	Developmental stage-specific anther promoter	Antisense RNA	Tobacco	Poovalah, Patil and Takezawa, 2002
Reversible male sterility	Biosynthesis of amino acids inhibited in male reproductive organs, reversible by application of those amino acids	Male organ-specific promoter	Antisense RNA	<i>Arabidopsis</i> , tobacco	Dirks <i>et al.</i> , 2001
Male sterility	Suppression of <i>ATH1</i> gene to control flowering time	35S CaMV	Antisense <i>ATH1</i>	<i>Arabidopsis</i>	Smeekeens, Weisbeek and Proveniers, 2005
Delayed flowering time	Loss of function of <i>SIN1</i> by RNAi	35S CaMV	Short integuments 1 protein	Unspecified	Ray and Golden, 2004
	RNAi construct	Constitutive, inducible, or tissue-specific promoter	Sequence similar to transgene or endogenous gene	Unspecified	Waterhouse and Wang, 2002
<b>Floral promoters</b>					
Male sterility	Anther development-specific genes and promoters	Tapetum, pollen	Antisense RNA or any gene that compromises pollen viability	<i>Brassica</i> , <i>Arabidopsis</i> , tobacco	Knox, Singh and Xu, 2004
Female sterility	Regulatory region of corn silk/pistil genes	C3 promoter	Silk-specific gene, C3	Maize	Ouellet <i>et al.</i> , 2003
Restoration of fertility to cytoplasmic male sterile plants	Wild-type <i>atp6</i>	<i>AP3</i> promoter	Wild-type <i>atp6</i> gene fused to a mitochondrial transit peptide	<i>Brassica</i>	Brown, 2002
Conditional male sterility	Upon application of acetylated toxin	Stamen-selective promoters	Deacetylase	Wheat	Quandt, Bartsch and Knittel, 2002
Male and female sterility	Poplar floral homeotic genes and promoters	Native promoters	<i>PTLF</i> , <i>PTD</i> , <i>PTAG-1</i> , <i>PTAG-2</i>	Poplar	Strauss <i>et al.</i> , 2002
Male sterility	Recessive mutant causes sterility	<i>Ms41-A</i> promoter	<i>Ms41-A</i>	<i>Arabidopsis</i> , maize,	Baudot <i>et al.</i> , 2001
Male sterility	Absence of a functional callase enzyme	<i>MsMOS</i> promoter	<i>msMOS</i>	Soy	Davis, 2000

TABLE 3-2 (CONTINUED)

Phenotype	Mechanism	Promoter	Active gene/Protein	Species	Reference
<b>Protein interference</b>					
Reversible male sterility	Dominant negative genes under anther-promoter reversed by expression of a repressor	Anther-specific promoter and <i>lexA</i> operator	Any cytotoxic methylase or growth-inhibiting gene	Maize	Cigan and Albertsen, 2002
Cytoplasmic male sterility	ATP synthesis in mitochondria inhibited	Ubiquitin promoter	Unedited <i>Nad 9</i> gene	Rice, wheat, maize, soybean	Patell <i>et al.</i> , 2003
Male sterility	Biotin-binding polypeptide ablates male gamete tissue, fertility can be restored	Promoter regulated by the <i>LexA</i> operon expressed in anther	Biotin-binding polypeptide and inhibitory proteins	<i>Arabidopsis</i> and tobacco	Albertsen and Huffman, 2002
Male sterility	Repressor protein under male promoter repressed by antisense RNA	Male flower specific promoter	Repressor protein	Multiple	Bridges <i>et al.</i> , 2001
Male sterility	Protein that disturbs metabolism, development and gene for reversibility	Stamen-specific promoter	A sterility RNA, protein or polypeptide	<i>Brassica</i> , maize, rice	Michiels, Botterman and Cornelissen, 2000
<b>Mitigation</b>					
Male sterile and dwarf	Unknown	Native promoter	<i>df11</i> gene	Safflower	Weisker, 1995
Dwarf plants	GA insensitive	Native promoter	Mutant of <i>GA1</i>	<i>Arabidopsis</i>	Harberd <i>et al.</i> , 2004
Dwarf plants	Rht mutant dominant allele causes GA-insensitivity	Native promoter	Mutant of <i>Rht</i> (D8)	Rice	Harberd, Richards and Peng, 2004

There are five major approaches to containment. One approach, mitigation (e.g. Al-Ahmad, Galili and Gressel, 2004), is a directed form of plant domestication such that the fitness benefits of transgenes are effectively cancelled by tight linkage to a gene that is beneficial within farms or plantations, but deleterious elsewhere. It has the advantage of being applicable to vegetative and sexual dispersal, which is useful for species like poplars that can spread vegetatively. Mitigation genes could also be combined with sterility genes to provide a second layer of containment. Genes that reduce the rate of height growth in forest trees, especially for shade-intolerant species like poplars (Daniel, Helms and Baker, 1979), are expected to provide a very powerful competitive disadvantage in competition with wild trees (Strauss *et al.*, 2004). Only two patents for dwarfism genes are shown under mitigation in Table 3-3 (Harberd, Richards and Peng, 2004; Harberd *et al.*, 2004), though there are a number of such genes now reported in both the scientific and patent literature. It is unclear, however, if such genes could be used and still maintain or improve yield and adaptability in plantation grown trees, but such studies are underway (e.g. Strauss *et al.*, 2004; Busov *et al.*, 2006).

The other forms of containment affect sexual reproduction, which is overwhelmingly the most important means for large-scale propagule spread in most tree species. There are basically four genetic engineering approaches: ablation, where floral tissues are effectively destroyed or made non-functional



TABLE 3-3  
Summary of studies on stability of transgene expression in plants

Taxa	Gene	Number of events (unstable) <sup>1</sup>	Environment	Propagation	Generations or years	Associated factors	Non-associated factors	Reference
Chrysanthemum	<i>35S-gus</i>	17(0)	Greenhouse	Vegetative	1 generation			Pavingerová <i>et al.</i> , 1994
Citrus	<i>35S-uidA</i> , <i>NOS-nptII</i>	70 (0)	Screenhouse	Vegetative	4–5 years	Copy number	T-DNA rearrangements	Cervera <i>et al.</i> , 2000
Poplar	<i>FMV-cp4</i> , <i>FMV-gox</i>	40 (1)	Field	Vegetative	4 years			Meilan <i>et al.</i> , 2002
Poplar	<i>35S-rolC</i>	6–22 (2–6)	<i>In vitro</i> , greenhouse, field	Vegetative	5–6 years	T-DNA repeat formation, flanking AT-rich sequence		Kumar and Fladung, 2001
Poplar	<i>35S-uidA</i> , <i>EuCAD-uidA</i>	44 (0)	<i>In vitro</i> , greenhouse, field	Vegetative	6 years		Copy number, extra vector sequence	Hawkins <i>et al.</i> , 2003
Poplar	<i>35S-ASCAD</i> <i>35S-ASCOMT</i>	4	Field	Vegetative	4 years			Pilate <i>et al.</i> , 2002
Potato	<i>Gus</i> , <i>nptII</i>	2	<i>In vitro</i> , greenhouse	Vegetative	2 years			Borkowska <i>et al.</i> , 1995
Potato	<i>NptII</i> , <i>gus</i> , <i>ocs</i> , <i>rolA</i> , and <i>C</i>	4	Greenhouse	Vegetative	3 generations			Ottaviani, Hanisch ten Cate and Doting, 1992
Sugar cane	<i>Ubi-bar</i>	1	Greenhouse	Vegetative	3 generations		Contained five copies	Gallo-Meagher and Irvine, 1996
Sugar cane	<i>Pat</i>	1	Field	Vegetative	3 generations		Contained nine copies	Leibbrandt and Snyman, 2003
Tall fescue	<i>Actin1-gus</i>	2	Growth room	Vegetative	5 generations			Bettany <i>et al.</i> , 1998
<i>Arabidopsis</i>	<i>NOS-nptII</i>	7	<i>In vitro</i>	Sexual	4 generations	Promoter methylation		Kilby, Leyser and Furner, 1992
<i>Arabidopsis</i>	<i>35S-hpt</i>	28 (14)	<i>In vitro</i>	Sexual	1 generations	Copy number		Scheid, Paszkowski and Potrykus, 1991
<i>Arabidopsis</i>	<i>NOS-nptII</i>	111 (62)	<i>In vitro</i> , growth chamber	Sexual	3 generations	Construct configuration, temperature	Copy number	Meza <i>et al.</i> , 2001
<i>Arabidopsis</i>	<i>Fpl-dsFAD2</i>	1	Greenhouse	Sexual	4 generations			Stoutjesdijk <i>et al.</i> , 2002
Petunia	<i>35S-A1</i>	1	Field	Sexual	1 year	Promoter methylation, temperature, endogenous factors		Meyer <i>et al.</i> , 1992
Rice	<i>35S-bar</i> , <i>35S-gusA</i>	12 (0–2)		Sexual	3 generations	Presence of truncated transgene sequences	Copy number, position effect	Kohli <i>et al.</i> , 1999
Rice	<i>Ltp2-gus</i>	3	Greenhouse	Sexual	5 generations	Partial rearranged transgene		Morina, Olsen and Shimamoto, 1999
Tobacco	<i>NOS-nptII</i>	2	<i>In vitro</i>	Sexual	3 generations			Müller <i>et al.</i> , 1987
Tobacco	<i>NOS-nptII</i>	18 ( $5 \times 10^{-6}$ – $5.9 \times 10^{-4}$ ) <sup>2</sup>	<i>In vitro</i>	Sexual	1 generation	Environmental stress	MAR	Conner <i>et al.</i> , 1998
Tobacco	<i>35S-hpt</i> , <i>35S-cat</i>	4	<i>In vitro</i>	Sexual	8 generations	T-DNA flanking sequences, position effect, extra vector sequence		Iglesias <i>et al.</i> , 1997

<sup>1</sup> Unstable events given in parentheses only where data on ten or more independent events reported.


<sup>2</sup> Frequency of kanamycin-sensitive seedlings derived from each event.



by a cytotoxin; excision, where some or all functional transgenes are removed from gametes before their release; gene suppression, where the activity of one or more genes essential for reproduction are impaired at the DNA, RNA or protein levels; and repression, where the onset of flowering is postponed by modifying the expression of genes that promote vegetative growth or repress the transition to reproductive growth.


### Ablation approaches

Genetic ablation methods employ promoters active in specific cells to control the expression of a deleterious gene, usually encoding a cytotoxin (e.g. Burgess *et al.*, 2002). However, many kinds of deleterious genes may be employed, as demonstrated by the patent applications of Dellaporta and Moreno (2004) and Spena *et al.* (2002), which cite in addition to the widely used RNases and protein synthesis inhibitors (Table 3-1), DNases, proteases, glucanases and lipases. Höfig *et al.* (2006) recently reported that targeted expression of stilbene synthase, which interferes with pollen function, gave a high rate of male sterility. For engineering reproductive sterility, a floral predominant promoter has been used to control the expression of a cytotoxin such as the ribonuclease barnase (Mariani *et al.*, 1990). Ideally, cytotoxin expression will be confined to floral cells; however, it appears that many floral promoters are not expressed exclusively in floral tissues (e.g. Brunner *et al.*, 2000; Rottmann *et al.*, 2000), and even low levels of unintended cytotoxin expression may impair tree growth (Skinner *et al.*, 2000). Thus, great care is needed in selection of promoters and cytotoxins. Skinner *et al.*, (2003) showed how the promoter of the poplar floral homeotic gene PTD, used to drive the cytotoxin DTA, gave rise to high levels of sterility in tobacco and *Arabidopsis* and did not impair vegetative growth in a greenhouse trial. The tapetal specific promoter TA29 from tobacco, when fused to barnase, caused very high levels of male sterility in field-grown poplars (Figures 3-1 and 3-2). However, Wei *et al.* (2007), studying poplar, and Lemmetyinen, Keinonen and Sopenan (2004) and Lännepää *et al.* (2005), studying birch, found that many transgenic events with floral homeotic promoter::barnase fusions showed abnormal growth or morphology in the greenhouse. In an attempt to avoid deleterious effects on growth seen with the poplar *LEAFY* (*PTLF*) promoter driving barnase, barstar, a specific inhibitor of barnase, was co-expressed in transgenic poplars using various promoters, and it was found that gene insertion events with low ratios of barstar to barnase activity had abnormal growth and morphology (Figure 3-3), and that even among plants with normal growth and morphology in the greenhouse, those events with barnase grew slower in the field than events with only barstar or that lacked both genes (Wei *et al.*, 2007). We found that we were unable to regenerate any transgenic poplars containing an intact *pAPETALA1::DTA* transgene, a likely result of leaky expression (root and leaf) seen with this promoter in transgenic poplars with *pAPETALA1::GUS* fusion genes (data not shown). Thus, ablation-based systems need to be carefully engineered in trees via judicious choice of promoters, cytotoxins and vectors, and then carefully field tested.




### Gene excision approaches

There have been considerable efforts to develop more precise means for manipulation of transgenes and their genomic locations via the use of site-specific recombinase systems such as *cre/lox* from bacteriophage P1 (reviewed in Gilbertson, 2003). Although the primary goals have been the removal of selectable marker genes and the targeting of transgenes to defined locations, a more recent application has been to use them to selectively remove transgenes before the release of seeds and pollen. By flanking transgenes with recombinase recognition sites and placing the recombinase under the control of a floral predominant promoter, it appears that very high levels of transgene excision can be obtained. Mlynárová, Conner and Nap (2006) used the microspore-predominant NTM19 promoter to control expression of an intron-containing *cre* gene to successfully excise GUS encoding transgenes from tobacco pollen at a rate above 99.98%. No



excision activity was detected other than in target tissues. Li and Pei (2006 and personal communication) used the promoter of the bisexually expressed PAB5 gene (Belostotsky and Meagher, 1996) to drive either or both the *cre* or FLP recombinase genes, targeting loxP-FRT fusion recognition sites. Based on GUS activity examined in more than 25 000  $T_1$  progeny per transgenic event, they reported a 100% rate of transgene removal from both male and female gametes of tobacco in 18 of 45 events studied. Although this is a promising system for transgene containment in vegetatively propagated plants, its effectiveness in the long term under field conditions is unknown, and predicting and verifying that gametes will lack transgenes in large trees when they begin flowering will be difficult. It is also distinct from the other approaches in that it does not impair fertility, and thus would provide containment of only the excised transgenes – not of exotic or highly domesticated organisms. However, reproductive transgene excision could be used in combination with a sterility transgene to provide a more robust containment system.



### Gene suppression approaches

The activity of genes essential for fertility can be suppressed by transcriptional gene suppression, posttranscriptional gene suppression, blocking the activity of the encoded protein, or by directed mutation or deletion. As shown in Tables 3.2 and 3.3, there have been a great variety of genes and approaches in various plant species that have been successfully used to impart sterility and/or restore fertility. This includes targeting of signal transduction proteins (Zhang *et al.*, 2001; Poovaiah, Patil and Takezawa, 2002), amino acid metabolism (Dirks *et al.*, 2001), choline biosynthesis (Mou *et al.*, 2002), transcription factors (Preston *et al.*, 2004; Smeekens, Weisbeek and Proveniers, 2005), methylases or methyltransferases (Cigan and Albertsen, 2002; Luo *et al.*, 2005) and mitochondrial genes (Patell *et al.*, 2003; Yui *et al.*, 2003).

### RNA interference and related methods

Double-stranded RNA (dsRNA) can induce a variety of sequence-specific gene suppression processes in plants, animals and fungi (reviewed in Baulcombe, 2004; Matzke and Birchler, 2005). RNA-mediated gene suppression, also called RNA interference (RNAi), is now widely exploited to reduce the expression of specific genes (reviewed in Watson *et al.*, 2005). Virus-induced gene silencing (VIGS)

vectors are one option for inducing sequence-specific suppression and have great potential for functional genomics (Burch-Smith *et al.*, 2004 and discussed below), but are not suited to stable introduction of a biosafety trait.

Stable transformation of transgenes containing an inverted repeat or hairpin sequence corresponding to a transcribed region of the target gene has been effective in a variety of plants, and post-transcriptional suppression has been shown to be stably inherited over several generations (Chuang and Meyerowitz, 2000; Wesley *et al.*, 2001). However, stability through rounds of vegetative propagation and across multiple years in field environments has not been extensively studied (discussed below). Inverted-repeat transgenes of promoter regions can induce methylation and transcriptional gene suppression of endogenous plant promoters, and this approach was used to engineer male sterility in maize (Cigan, Unger-Wallace and Haug-Collet, 2005). Nonetheless, there have been relatively few studies, and thus its utility as a gene suppression approach is uncertain. Moreover, it appears that promoters vary in their sensitivity to different types of cytosine methylation, depending on their sequence composition (Matzke *et al.*, 2004).

Multiple genes can be silenced by using a conserved region or by joining sequence segments of multiple genes together to create a compound RNAi transgene (reviewed in Watson *et al.*, 2005). This capability is especially important for sterility systems where a redundant approach is desirable to produce a highly robust and reliable biosafety trait. Because of genetic redundancy in the regulation of flowering and many taxon-specific gene duplications and losses (Irish and Litt, 2005), the extent and configuration of redundancy required for robust and effective RNAi suppression will vary between species.

A population of transgenic events carrying the same RNAi transgene typically exhibit highly diverse levels of suppression. Although RNAi transgenics that phenocopy null mutations in floral regulatory and other genes have been obtained, strong suppression can be infrequent (Chuang and Meyerowitz, 2000; Stoutjesdijk *et al.*, 2002). In addition, the level of endogene suppression appears to be target-specific (Kerschen *et al.*, 2004). The endogenous expression level of the target gene appears to influence the effectiveness of RNA-mediated silencing, but does not appear to be the only gene-specific determinant of RNAi effectiveness (Han, H. Griffiths and D. Grierson, 2004; Kerschen *et al.*, 2004; Wagner *et al.*, 2005).

Possible additional determinants include spatiotemporal expression, RNA turnover and sequence composition. Single-copy RNAi transgenics are preferable because multicopy events appear more variable with respect to level of suppression and stability, perhaps because multicopy transgenes are more susceptible to transcriptional gene suppression (Kerschen *et al.*, 2004). For practical application, successful transformation events (i.e. those exhibiting strong suppression) must be identifiable via molecular tests when trees are still juvenile. This potentially limits the utility of this approach because many target genes are specifically or predominantly expressed in floral tissues. We have produced transgenic poplars carrying RNAi transgenes targeting various genes regulating floral onset and floral organ development. Using vegetative tissue from poplar transgenics still in tissue

culture or the greenhouse, we have been able to identify events exhibiting strong target endogene suppression using quantitative reverse transcriptase-polymerase chain reaction (qRT-PCR; Figure 3-4), suggesting that RNAi transgenic trees with greatly reduced fertility can be selected at an early, non-flowering stage.

Pleiotropic effects of RNAi methods can be significant. Non-target effects of dsRNAs are well-known in animal systems (Jackson and Linsley, 2004). However, this does not appear to be a common problem in plants for well-targeted dsRNAs, perhaps because both siRNAs and miRNAs require high levels of complementarity with their target (Watson *et al.*, 2005; Schwab *et al.*, 2005). Transitive suppression, whereby suppression spreads from the initiator sequence to an adjacent region, could potentially cause pleiotropic effects in plants. However, several plant studies have shown that transitive suppression occurred when the target was a transgene, but did not occur when an endogene was the target (Vaistij, Jones and Baulcombe, 2002; Petersen and Albrechtsen, 2005; Miki, Itoh and Shimamoto, 2005). Why transitive silencing appears to commonly occur with transgenes, but not endogenes, is unknown. However, to date, a few studies have looked for transitive silencing with endogene targets.


### **DOMINANT NEGATIVE PROTEINS**

Alternative approaches to repressing floral genes include introduction of dominant negative mutant forms of the target endogene and artificial transcription factors. Several studies have identified dominant negative mutant forms of plant signal transduction proteins and transcription factors, including MADS box genes regulating floral development (e.g. Jeon *et al.*, 2000; Dievart *et al.*, 2003; Ferrario *et al.*, 2004). Most dominant negative forms appear to exploit the modular nature of these proteins and that they often form multiprotein complexes. For example, a dominant negative protein might be able to interact with other proteins, but the protein complex cannot bind DNA. Based on studies of rice and mammalian MADS-box genes, we used site-specific mutagenesis to alter amino acids predicted to be necessary for dimerization and/or DNA binding in *AG* and *APE-TALAI(API)*. Constitutive expression induced strong loss-of-function phenotypes at a frequency of approximately 30% in primary *Arabidopsis* transformants, and these transgenes are now being evaluated in poplar and sweetgum (data not shown).


Another option for dominant repression of transcription factor activity is the introduction of chimeric transgenes that are translational fusions of the selected transcription factor coding region and a repression domain such as the ERF amphiphilic repressor (EAR) motif (Hiratsu *et al.*, 2003). Expression of EAR chimeras has proven to be useful for producing phenocopies of double knockouts in *Arabidopsis* and thus, can overcome the problem of genetic redundancy among gene duplicates. Recently, Mitsuda *et al.* (2006) used this chimeric repressor approach with *AP3*, *AG*, *LEAFY*, and a floral expressed *MYB* gene, and reported very high levels of sterility in *Arabidopsis* and/or rice. Recent studies have also shown that synthetic zinc-finger domains fused to a transcriptional activation or repression domain are highly effective for manipulating the expression of specific



FIGURE 3-4  
Range of RNAi gene suppression (top) and repeatability among biological replicates (bottom) for floral genes expressed in vegetative tissues



Relative expression level of native PTLF gene in selected poplar PTLF-RNAi transgenic trees and non-transgenic controls of poplar clone 353-53 (*Populus tremula* × *tremuloides*). Expression was determined by qRT-PCR analysis of native transcripts in vegetative shoots (a ubiquitin gene served as an internal control). Each datum represents a pool of total RNA from four to five ramets per transgenic event; error bars are standard deviations over three PCR technical replicates.



Relative expression level of native *Poplar SOC1* (*PSOC1*) gene in pairs of biological replicates (RNA extraction from different ramets) of selected *PSOC1*-RNAi transgenic trees and non-transgenic controls. qRT-PCR methods as in top graph. Data are means of independent qRT-PCR runs for two different ramets for single transgenic events; error bars are standard deviations over the average of two PCR technical replicates ( $r^2=0.41$ ). Pairs (shading) show biological replicates per event.



genes (reviewed in Segal, Stege and Barbas, 2003). By combining pre-defined zinc-finger modules appropriately, three- or six-finger domains can be created that specifically bind to a selected 12 to 18 bp DNA sequence. For example, a transgene containing a human repression domain, fused to a zinc-finger module designed to bind to a site in the *AP3* promoter, was able to repress endogenous *AP3* expression and induce a loss-of-function phenotype (Guan *et al.*, 2002).

It remains to be determined how these different methods of gene suppression compare with respect to frequency of transformants exhibiting strong repression or loss-of-function phenotypes, and stability over multiple years, in the field. It is also important to investigate whether pleiotropic effects are more common with certain methods. As discussed above, deleterious side-effects are not always evident under controlled conditions, but may appear as a cumulative effect of tree development, especially in the field. Although most studies have used strong constitutive promoters, tissue-specific promoters have been successfully used for RNAi and other repression methods. Promoters directing more restricted expression could reduce the occurrence of pleiotropic effects. However, they might be less effective at inducing strong, stable sterility.

### Targeted gene mutagenesis and replacement

The long-sought-after goal of routinely creating precise deletions, insertions or mutations with plant genes has been elusive, largely due to the propensity for random rather than homologous DNA recombination in plants. However, recent studies have demonstrated new strategies that achieve substantial improvements in the rate of targeted mutagenesis and gene replacement. By constitutively expressing the yeast *RAD54* gene, a member of the *SWI2/SNF2* chromatin remodelling gene family, Shaked, Melamed-Bessudo and Levy (2005) achieved gene targeting frequencies of 3 to 17% in *Arabidopsis*. Another approach employs the zinc-finger modules discussed above for targeted gene repression. In this case, the zinc-finger domain is fused to a nuclease to introduce double-strand breaks at specific genomic sites. In one study, zinc-finger nucleases (ZFN) were expressed in *Arabidopsis* to create breaks that were subsequently repaired by non-homologous end joining, resulting in site-specific insertion/deletion mutations at frequencies of 2–20% (Lloyd *et al.*, 2005). Using a ZFN to facilitate gene replacement via homologous recombination, Wright *et al.* (2005) achieved 10% gene targeting efficiency. Both ZFN and donor genes had been introduced into tobacco protoplasts via electroporation. In four of seven tobacco plants that were homozygous for the target reporter gene, the desired gene replacement occurred on both chromosomes; such a capability is critical for induction of sterility as loss of function effects are expected to be recessive, and breeding for homozygosity in trees is generally not feasible.

Genetic redundancy further complicates introducing sterility via gene targeting (e.g. both alleles of two or more genes might need to be replaced or mutated). However, replacement of only one allele of one gene with a dominant suppression transgene might be more effective in achieving reliable sterility than random

integration of the sterility transgene because it would reduce wild-type gene dosage and may avoid position effects that can occur with random transgene integration. A key factor limiting the use of gene targeting is ease and efficiency of transformation in the species or genotype of interest. The feasibility of gene targeting is dependent of the combined frequencies of transformation and gene targeting and ease of transformation, regeneration and selection. *In planta* transformation is routine for *Arabidopsis* and that allows production and screening of a large number of transgenics with little effort; no similar system exists for trees.

One caveat to gene mutation or deletion is that recent studies suggest the possibility that there might be cases where it is not permanent. *Arabidopsis* hothead (*hth*) mutants can inherit allele-specific DNA sequences at multiple loci that were not present in the genomes of their parents, but were present in an earlier ancestor (Lolle *et al.*, 2005). Under certain environmental conditions, varieties of flax exhibit highly specific DNA changes at multiple loci from parents to progeny, including a large insertion that is found in natural populations, but is not present in the genome of the progenitor (Chen, Schneeberger and Cullis, 2005). To explain the non-Mendelian inheritance of *hth* mutants, Lolle *et al.* (2005) proposed that a cache of stable RNA serves as the template for extra-genomic DNA sequence reversion; however, others have posited alternative explanations (e.g. Comai and Cartwright, 2005). It is unclear whether this type of reversion could occur somatically in trees (e.g. during vegetative propagation or under certain stressful conditions). Rates of transgene instability under vegetative growth appear to be considerably lower than under sexual reproduction (discussed below).

### Repressors of flowering

The activities of some strong repressors of the transition to flowering are directly correlated with their expression level (reviewed in Boss *et al.*, 2004). Thus, constitutive expression or overexpression of a floral repressor in appropriate tissues may be effective at long-term postponement of flowering. Because of the multiple pathways promoting flowering, this approach might delay, rather than prevent, the transition to flowering, but if flowering were delayed until long after harvest age, it still could be an effective biosafety approach. In addition, a floral repressor transgene could be combined with a different sterility transgene, such as one suppressing genes necessary for reproductive organ development, to provide redundancy. Overexpression of a floral repressor might be more likely to induce pleiotropic effects that, as discussed above, might not be apparent until trees are field-tested. Maintaining trees in a purely vegetative phase throughout their rotation cycle, whether by overexpression of a floral repressor, suppression of a floral promoter, or both, is highly desirable because this would completely prevent resource allocation to reproductive structures. However, depending on the tree taxon and environment, development of sterile reproductive structures might not be desirable if, for example, the plantation provides important habitat for birds or beneficial insects that feed on flower parts.

## REPRODUCTIVE GENE MOLECULAR BIOLOGY AND GENOMICS IN TREES

### Analysis of floral gene homologs


Most published studies of genes controlling flowering in trees have described the isolation and gene expression patterns of homologs of genes known to control various stages of flowering in *Arabidopsis* (e.g. Southerton *et al.*, 1998; Sheppard *et al.*, 2000; Cseke, Zheng and Podila, 2003). Results from heterologous overexpression in *Arabidopsis* and tobacco have also been reported, and these studies have usually shown a phenotype similar to that induced by overexpression of the *Arabidopsis* homolog (e.g. Kyojuka *et al.*, 1997; Rutledge *et al.*, 1998; Elo *et al.*, 2001). Functional gene studies of flowering in trees are rare because of the lack of sufficiently efficient transformation systems to produce multiple-event transgenic populations for large numbers of target genes. In addition, the multiple-year non-flowering phase of trees requires long and costly time spans and large areas for field research. *LFY* and *AP1* and tree orthologs of *FT*, which accelerate flowering when overexpressed in *Arabidopsis*, have been shown to induce early flowering in poplar and/or citrus, potentially bypassing the long time delays to flowering (Weigel and Nilsson, 1995; Rottmann *et al.*, 2000; Pena *et al.*, 2001; Endo *et al.*, 2005; Böhlenius *et al.*, 2006; Hsu *et al.*, 2006). In some cases, however, the inflorescences have been abnormal or gametes inviable (Rottmann *et al.*, 2000; Hsu *et al.*, 2006); induction of at least some *FT* homologs may bypass this problem (Böhlenius *et al.*, 2006).

Both overexpression and antisense constructs of the silver birch genes, BpMADS1 and BpMADS6, homologs of *SEPALLATA3* and *AG*, were transformed into an early flowering birch genotype (Lemmetäinen *et al.*, 2004). Although mutant phenotypes were somewhat inconsistent or rare, suppression of BpMADS1 appeared to cause some inflorescences to partially revert to vegetative shoots, and in two BpMADS6 transgenics, some male inflorescences lacked stamens, suggesting functions similar to their *Arabidopsis* counterparts. In *PTLF* antisense poplar transgenics that flowered after several years in the field, some male transgenic events produced mutant flowers with homeotic conversion similar to *lfy* mutants (data not shown). Phenotypes were consistent between catkins from a single transgenic event, but catkins typically displayed a basal to tip gradient with flowers at the tip having a more severe mutant phenotype; thus, basal flowers often produced stamens that were wild-type in appearance. However, in the transgenic event with the most severe mutant phenotype, few flowers with stamens were observed. RNAi transgenes have been reported to be more efficient at inducing suppression than antisense constructs (Wesley *et al.*, 2001), suggesting that RNAi versions of *PTLF* now entering field trials (data not shown) might give a higher rate of sterility both within and between events.

Encouraging results were found with RNAi studies of *PCENL1*, a poplar homolog of the *Arabidopsis* floral repressor, *TERMINALFLOWER 1*. Transgenic events that showed strong reduction in target endogene expression as determined by qRT-PCR initiated flowering earlier than wild-type in the field (Mohamed, 2006); the extent of precocious flowering was significantly correlated with the

level of endogene suppression (Figure 3-5). These studies suggest that RNAi suppression of orthologs of *Arabidopsis* genes that promote flowering, and do not appear to have any role in vegetative development, can be an effective method for introducing biosafety traits. They also suggest that transgenic events will need to be carefully screened to select lines exhibiting strong suppression. Where vegetative tissue expression is detectable, it should be possible to screen for desirable events during seedling growth, saving years of study and reducing the costs and issues of screening large numbers of field-grown trees.

The extent of overlap in genes and pathways regulating reproductive development in angiosperms and gymnosperms is poorly known. Most studies have focused on MADS-box genes. For example, studies have identified *Picea*, *Ginkgo*, *Gnetum* and *Cycas* genes belonging to the AG subfamily (Rutledge *et al.*, 1998; Shindo *et al.*, 1999; Jager *et al.*, 2003; Zhang *et al.*, 2004). The expression patterns of the gymnosperm AG homologs and phenotypes induced by heterologous ectopic expression or complementation of an *Arabidopsis ag* mutant support a conserved function in controlling reproductive organ development. Conifer homologs of the MADS-box B-class floral organ identity genes, the flowering time gene, *SOC1*, and *LEAFY* have also been identified (Tandre *et al.*, 1995; Sundstrom *et al.*, 1999; Mellerowicz *et al.*, 1998; Mouradov *et al.*, 1998). The Norway spruce gene *DAL10* belongs to a MADS-box subgroup that is possibly gymnosperm-specific



and is specifically expressed in pollen and seed cones (Carlsbecker *et al.*, 2003). Another spruce MADS-box gene, *DAL1*, belongs to the *AGL6* subfamily and its expression correlates with maturation to the adult or flowering phase (Carlsbecker *et al.*, 2004).

### Forward-looking genomics approaches

Although comparative studies indicate that similar genes and pathways control reproductive development in angiosperms and to an extent in gymnosperms, taxon-specific gene duplications and losses, and subsequent subfunctionalization and neofunctionalization, make predictions of gene function based solely on orthology or expression patterns problematic (Irish and Litt, 2005). The poplar genome sequence and an increasing number of large expressed sequence tags (EST) datasets for various tree taxa greatly facilitates identification of tree homologs to various *Arabidopsis* genes regulating flowering and their lineage-specific gene duplications and losses (Brunner and Nilsson, 2004). Moreover, the Floral Genome Project ([www.floralgenome.org](http://www.floralgenome.org)) (Albert *et al.*, 2005) and other projects (e.g. Brenner *et al.*, 2005) have developed extensive floral EST datasets from diverse plants including phylogenetically important eudicots, non-grass monocots, basal angiosperms and gymnosperms. Although many of the floral EST sets are not from trees, comparative floral genomics studies are still informative because tree taxa occur in almost all eudicot orders (Groover, 2005). These extensive sequence resources are beginning to reveal patterns of conservation and divergence of families of floral regulatory genes (e.g. Zahn *et al.*, 2006).

Genomic platforms for analysing gene networks controlling flowering in trees will enable selection of genes and design of sterility strategies with greater precision and effectiveness. Global expression analyses of *Arabidopsis* development, responses to floral induction stimuli and spatial patterns in flowers of *Arabidopsis* mutants, have revealed tissue-predominant expression patterns and components of gene networks controlling floral initiation and floral organ development (Schmid *et al.*, 2003, 2005; Wellmer *et al.*, 2004). Bio-informatic analyses of co-expressed genes, chromatin immunoprecipitation studies and comparison of regulatory regions of orthologous genes can identify cis-regulatory elements associated with a particular response or process (e.g. Li, Zhong and Wong, 2005; Kreiman, 2004; Rombauts *et al.*, 2003). Yeast two-hybrid screens were used to develop a comprehensive interaction map of all *Arabidopsis* MADS domain proteins (de Folter *et al.*, 2005). Combined with global expression analysis, protein interaction studies would be especially useful for selecting genes and sterility methods unlikely to have pleiotropic effects. Similar strategies are beginning to be applied to poplar, and a new United States of America National Science Foundation Plant Genome Project is studying the transition to flowering in poplar. This includes use of overexpression and RNAi poplar transgenics for transcriptome analyses.

Comprehensive study of gene expression is more difficult in trees than annuals due to complex developmental phase changes and increasing size and tissue complexity across years. We have observed that some genes showing floral-

predominant expression in poplar show levels of vegetative expression that vary in intensity across an annual cycle of growth and dormancy (data not shown). Furthermore, trees are exposed to very variable abiotic and biotic conditions over many years that can markedly affect gene expression. For example, galling insects appear to induce ectopic organ developmental programmes that are similar to reproductive development; *LEAFY*, *API* and C-class MADS-box genes directing carpel development, but not B-class genes, are expressed during development of galls on grape vine leaves (J.C. Shultz, personal communication). This is especially problematic for ablation sterility systems where selection criteria for appropriate promoters are most stringent.

In addition to not having complete genome sequences, studies in most tree taxa are generally limited by lack of efficient transformation systems. Development of VIGS vectors for trees could be particularly valuable for studying genes controlling flowering. A VIGS vector has recently been developed for poplar (Naylor *et al.*, 2005), but unfortunately a poplar genotype that reliably flowers in the greenhouse in the absence of *FT* overexpression is not currently available. Some other tree species, such as eucalypts and apple, can be reliably induced to flower via use of plant hormones and cultural treatments.

As tree genomics tools and knowledge of candidate genes for flowering advance, it should be possible to clone genes that control onset of flowering using high-resolution quantitative trait locus (QTL) or association genetics approaches. This approach potentially allows discovery of mechanisms of reproductive development that are unique to trees, rather than relying on studies of herbaceous annual model plants for target gene identification. Liebhard *et al.* (2003) reported QTLs for juvenile phase in apple. Missiaggia, Piacuzzi and Grattapaglia (2005) identified a QTL for very early flowering in eucalypts. For these studies, it will be essential to have large populations ready that include segregants with rare precocious flowering. To prevent flowering, these genes could then be suppressed or mutated, as discussed above.

### STABILITY OF TRANSGENE EXPRESSION

It is well known that newly produced transgenic plants often exhibit instability in expression of transgenes, related endogenes and their encoded traits. It is also widely known that the level of instability varies widely among constructs, species and gene transfer methods. However, after field screening, gene insertion events with strong and stable expression are generally identified, and these are the ones focused upon during research and commercial development. The ability to identify highly stable transgenic events has been firmly established by the hundreds of millions of hectares of genetically engineered crops that have been grown by farmers, which contain a variety of genetic constructs in a variety of genotypes and species. These include commercialized trees (papaya, poplar), with traits induced via RNAi (papaya, tomato, squash) and with conventional transgene expression.

Questions remain, however, about the long-term stability of specific traits in vegetatively propagated crops, including containment traits and to what extent



stable expression can be identified and delivered in an efficient manner in breeding programmes with transgenics. It is also unclear how strong and stable a sterility phenotype must be to confer an adequate level of containment. A high level of stability of a leaf-expressed gene for herbicide resistance, imparted by genes derived from other species, does not guarantee that a native gene designed to suppress a floral meristem identity gene via RNAi will be sufficiently reliable for stringent, long-term containment goals. Because of the importance of stability of gene expression for genetic containment in trees, we review both what has been learned from studies in other vegetatively propagated crops, and then in the following section consider how a modelling approach can help to identify how much trait instability (i.e. reversion to fertility) might be biologically acceptable.

Due to the long life cycles of forest trees and the complex environments they experience, stability of expression of genetically engineered-introduced traits in trees has received considerable debate (Fladung, 1999; Hoenicka and Fladung, 2006a). In addition, possible genome instability due to effects of the gene transfer process and interaction with plant genome sequences adds to scientific uncertainties about long-term performance of primary transformants in the field. In an AFLP study with four *Agrobacterium*-transformed aspen transgenic lines carrying a *rolC* gene, 886 out of 889 (99.9%) of the amplified bands were common between the control and transgenics, suggesting very limited genetic engineering-associated genomic change compared with extensive wild AFLP polymorphism in poplar and most other tree species (reviewed in Hoenicka and Fladung, 2006b). In agronomic crops, it also appears that genomic variation imparted by transformation is modest compared to the extensive genomic variation present in traditionally bred and wild plants (Bradford *et al.*, 2005).

A number of factors have been implicated in transgene silencing, including insert number, chromosomal environment (position effect), T-DNA structure, environmental stress and endogenous factors (Table 3-3). Unfortunately, most of these factors do not seem to be consistent predictors of long-term stability. For example, there appears to be little association between insert number and instability, even though single-copy transgenes are widely assumed to be important for obtaining stable gene expression. Where transgene structure was studied, however, instability was often associated with transgene repeat structure, truncation, or other re-arrangements at or near transgene insertion sites (Table 3-3).

Transgene stability under vegetative propagation has been studied in poplar, citrus, tall fescue, sugar cane, chrysanthemum and potato. Transgene expression appears far less stable over sexually propagated generations than over vegetatively propagated generations (Table 3-3). Unfortunately, most studies have used a small number of transgenic events (<20), and are thus of limited relevance to commercial transformation and breeding programmes, which often screen many dozens or hundreds of events. Moreover, many of the published studies on stability of transgene expression have focused on unstable events observed in preliminary screens, and are thus biased with respect to the levels of instability expected in commercial programmes.




In a study similar to what a tree breeding programme might address, Meilan *et al.* (2002) reported high stability of herbicide resistance genes in 40 independent poplar transgenic events over four years in the field. Hawkins *et al.* (2003) reported stable expression of a GUS reporter gene in 44 independent poplar transgenic events over a period of six years under *in vitro*, greenhouse and field conditions. Histological GUS analysis in 70 transgenic events showed similar patterns of GUS expression over a period of four to five years in citrus (Cervera *et al.*, 2000). In contrast, in a study of 22 transgenic events carrying the morphological marker gene, *rolC*, phenotypic alteration or reversion was observed for up to one-third of the events during vegetative growth in either *in vitro*, greenhouse or field conditions (Kumar and Fladung, 2001). In biolistically transformed pine, Wagner *et al.* (2005) reported that the level of silencing of a cinnamyl alcohol dehydrogenase (CAD) gene during embryogenic propagation was associated with expression level.

Variation in stability of transgene expression among studies can result from uncontrolled differences in experimental protocols, as demonstrated by James *et al.* (2004). Because native and introduced genes show stochastic (Raser and O'Shea, 2004) and developmental variation in expression, it is important to pick a suitable control gene. For example, the strong and deleterious effects of variable expression of the *rolC* gene discussed above might be similar to the normal variation that occurs with many endogenes and transgenes, but its gene product is so powerful and toxic that its effect on development is amplified. In contrast, no such consequence, nor possibly any phenotypic effect at all, would be expected for similar levels of variation in a transgene encoding insect or herbicide resistance.

We have performed three stability studies using different transgene constructs (unpublished data). In one study, the *BAR* herbicide resistance gene was transferred into two poplar clones, and 32 transgenic events produced. The expression of the *BAR* gene was monitored on 384 plants over a period of eight years of repeated coppicing in the field. No instability or loss of the initial resistance phenotype was observed based on visualized herbicide damage and protein enzyme-linked immunosorbent assays (ELISA). In another study, the reporter genes *GFP* and *BAR* were assembled in the same binary vector, and transferred into two poplar clones. The expression levels were measured on 2 256 transgenic poplar trees generated from 404 independent transgenic events over three years in the greenhouse and the field. The expression of both genes was highly stable over three years, with no cases of gene silencing observed. However, the physical loss of transgene sequences was observed in three of the 80 transgenic events after they were regenerated via a second round of organogenesis in tissue culture.

In a third study, we examined the stability of RNAi silencing of a resident *BAR* gene in transgenic poplars that had been re-transformed with inverted repeats (IR) of either a section of the coding sequence or the promoter sequence of the *BAR* gene. RNAi silencing efficiency and stability were studied in 56 RNAi transgenic events over two years in the field. The results suggested that dsRNA of the *BAR* coding sequence was highly efficient in suppressing *BAR* expression; 80% of



the events showed more than 90% gene suppression. However, dsRNA of the *BAR* promoter sequence was much less efficient; only 6% of the events showed more than 90% suppression. Most importantly for gene containment, the degree of RNAi suppression appeared to be stable for both constructs over two years (Figure 3-6). These studies, plus the reporter gene studies described above, suggest that instability of gene expression may only rarely be a problem in vegetatively propagated trees, though longer-term studies are desirable.

### **STERILITY AS A QUANTITATIVE TRAIT: HOW MUCH DO WE NEED?**

Complete prevention of sexual reproduction with 100% certainty is a daunting technical and social challenge. The long time frames and large numbers of potential reproductive meristems in transgenic tree plantations provide many opportunities for reversion to fertility, such that rare events become probable. Furthermore, transgenic approaches to sterility will incur added economic and regulatory costs and social resistance (discussed above). It is therefore critical to define if sterility is needed at all for biological or social reasons, and if so, what level and form is required. However, there does not seem to have been any serious field studies, in any crop, sufficient to estimate the operational effectiveness of containment genes (Ellstrand, 2003). Until many such studies are published, it would be unwise to assume that genes can be fully and safely contained in the near future. Conventional approaches to fertility reduction, including the use of hybrids or aneuploid germplasm (Bradshaw and Stettler, 1993), also generally do not provide complete containment. However, they could provide an option for deployment of some

transgenes in breeding programmes that use ploidy-modified trees. However, such genotypes are rare in most forest tree breeding programmes. Poplar and some other tree species are capable of dispersal and establishment of vegetative propagules, thereby potentially bypassing most containment measures based on sexual sterility. Though local spread from plantings can usually be managed, some degree of long-distance vegetative spread can occur through adventitious rooting from broken or abscised branches (Rood *et al.*, 2003). If transgene containment is an important goal, it is important to explore the consequences of all of the different modes and levels of reproduction under realistic ecological scenarios. This is best addressed in the context of a risk assessment and is facilitated by the use of ecological modelling.

Risk assessment includes hazard identification, exposure assessment, consequence assessment, risk characterization and delineation of mitigation options (Hill, 2005). Risk from transgene dispersal is sometimes treated as synonymous with the exposure portion of the process, and demonstrations of potential distributions of transgenic propagules are treated as examples of the inherent risks of forest biotechnology (e.g. Williams, 2005). However, the mere presence of transgenic propagules does not automatically constitute a negative endpoint (Stewart, Halfhill and Warwick, 2003). Production and dispersal of transgenic seed and pollen constitute the first steps in a network of processes contributing to introgression of transgenes to wild populations. Even with the extensive dispersal distances expected for forest trees (Nathan *et al.*, 2002), realized transgene introgression could still be extremely low due to sexual incompatibility with wild trees, lack of availability of safe sites for establishment, negative fitness effects of transgenes or domestication genes in a wild setting, and extensive dilution from non-transgenic planted and wild stands (Pilson and Prendeville, 2004; Hails and Morley, 2005). As discussed above, transgene dispersal could also have large net ecological benefits.

Trees create special challenges for generating the data necessary for assessing potential introgression. Very large temporal and spatial scales must be considered for movement of tree pollen and seeds (Nathan *et al.*, 2002; Smouse and Sork, 2004). Furthermore, long-distance gene flow is a disproportionately important determinant of rates of spread of introduced organisms or genes (Higgins and Richardson, 1999), and this process is subject to stochastic influences that make accurate measurement extremely challenging, if not impossible (Clark *et al.*, 2003). This difficulty is magnified when one considers the network of interacting, highly variable factors that determine establishment and spread in wild systems (Parker and Kareiva, 1996; Pilson and Prendeville, 2004). Therefore, realistic, replicated experiments cannot be performed at appropriate scales and time frames for predicting introgression of transgenes (Parker and Kareiva, 1996). However, data from non-transgenic populations can be used in simulations to provide useful estimates of what is likely to occur under various deployment situations and environments (Dunning *et al.*, 1995; Pilson and Prendeville, 2004).

Simulation approaches have been used successfully to investigate factors affecting the spread of transgenic insect-resistant oilseed rape varieties (Kelly *et al.*, 2005)

and to investigate factors affecting fitness of transgenic fish with enhanced growth (Howard, DeWoody and Muir, 2004). However, many of these kinds of studies have not taken into account realistic spatial distributions of transgenic organisms on the landscape relative to wild and managed habitats. The spatial dimensions of gene flow are an essential component of introgression because habitat availability and competition from wild relatives are likely to be two of the primary factors inhibiting spread of partially fertile transgenic trees, and these will be determined by management regimes and locations of wild populations on the landscape.

Many different types of models have been used for simulating dispersal and gene flow across a landscape (Nathan *et al.*, 2003). One approach is to devise mechanistic models of pollen and seed dispersal based on the physical properties of the propagules and the environment (Katul *et al.*, 2005; Nathan *et al.*, 2002; Clark *et al.*, 2003). Such models have a distinct advantage in that they are easily parameterized for a large number of species because flight characteristics of pollen and seeds are readily measured, detailed microclimatic data can be obtained for many sites, and the physics of dispersal by abiotic agents are fairly well characterized. Disadvantages include the large number of parameters that require estimation (particularly if realized gene flow is to be modelled) and the high computational requirements that limit the extent of the area and time frame that can be modelled (Nathan *et al.*, 2002).

An alternative approach is to model gene flow phenomenologically based on field observations of dispersal and demographic processes. A common method is to use reaction-diffusion models to depict the movement of an 'invasion front' using a diffusion approximation and logistic growth models (Fisher, 1937; Shigesada and Kawasaki, 1997). Alternatively, probability density functions of propagule movement and/or reproductive success can be used to determine the probability of dispersal between points on a lattice of habitat cells (Higgins and Cain, 2002; Lavorel, Smith and Reid, 1999). This approach has the advantage of being easily parameterized from historical data (e.g. a chronosequence of air photos or survey data) and readily integrated with geographical information systems (GIS). A major disadvantage is the difficulty of measuring contemporaneous realized gene flow on appropriate space and time scales to parameterize the models.

As an example of the latter approach, we developed a spatially explicit model of gene flow from hybrid poplar plantations based on observations of realized gene flow in wild populations (DiFazio, 2002; Slavov, DiFazio and Strauss, 2004). The model, called Simulation of Transgene Effects in a Variable Environment (STEVE), was applied to a landscape grid in northwest Oregon (23 km × 37 km, 100 m<sup>2</sup> cells) containing information about elevation, habitat type and poplar populations. The simulation has an annual time step, with modules to simulate creation and conversion of poplar patches, growth, reproduction, dispersal and competition within poplar cohorts. The primary objective of this model was to produce a framework for virtual experiments that could accommodate the diverse silvicultural, agronomic and ecological settings in which transgenic trees might be released, and to incorporate many different types of transgenic traits.

The findings of the STEVE model most germane to discussions of reproductive sterility come from simulations with different levels of innate fertility of transgenics and with various probabilities of reversion to fertility. Relative pollen production was calculated for each genotype within each sexually mature cohort of trees in each poplar cell. Representation of pollen and seed was entirely relative because the most important quantity is the ratio of transgenic to conventional genotypes in the propagule pools. Therefore, pollen production was directly proportional to the basal area of each genotype in a particular location on the landscape.


Relative fertility varied annually based on a user-defined standard deviation determined from annual field observations of flowering in plantations. In addition, transgenics with reduced fertility could have their fertility partially restored according to a user-defined probability. Vegetative propagule production was also stochastic and proportional to basal area. Pollen was dispersed within the immediate vicinity of male trees and across the landscape according to empirically determined dispersal kernels (Slavov, DiFazio and Strauss, 2004), and transgenic and conventional seed production was determined by the proportion of pollen of each genotype dispersed to female trees, modified by relative fertility factors.

As expected, fertility of transgenic trees had a strong effect on rate of gene flow from transgenic plantations. With highly reduced fertility, gene flow was at some of the lowest levels observed for all scenarios tested: between 0.1 and 0.2%, compared with approximately 5% for fully fertile transgenic plantations. In addition, transgene flow rates were not distinguishable within the range of 0 to 1% of wild fertility, indicating that complete sterility was not required to attain maximum gene containment (Figure 3-7a). Thus, the reductions in fertility of approximately  $10^5$  that we have observed in the field (Figure 3-1) would appear to be far in excess of the level needed for effective mitigation in this scenario. (In practice, only the pollenless events might be chosen for commercial purposes.) The low level of gene flow that we observed for fully sterile plantations was due to movement of vegetative propagules in the vicinity of plantations. However, transgenic gene flow remained very low under a wide range of rates of vegetative establishment (Figure 3-7b), and gene flow rates were insensitive to changes in rates of vegetative establishment and shapes of vegetative dispersal curves (data not shown). Sexual fertility was therefore much more important than vegetative establishment in controlling gene flow in this system. Nearly 50% of the gene flow with low-fertility transgenics (fertility  $<0.1$ ) was due to sexual reproduction, as demonstrated by simulations with vegetative establishment eliminated (Figure 3-7b).


Other investigations have also identified fertility as a major factor limiting plant spread. For example, a reduction of fertility of as little as 75% was projected to limit the spread of scotch broom (*Cytisus scoparius* L.), based on insect-protection assays and simulations (Rees and Paynter, 1997). Density of pines spreading from plantations in South Africa was sensitive to fecundity and age of reproductive maturity in spatially explicit simulations (Higgins, Richardson and Cowling, 1996). Spread of feral oilseed rape was hypothesized to be limited by seed input

FIGURE 3-7  
**Simulated effects of transgenic fertility on transgene flow based on the STEVE model**


Simulations were conducted over a 50-year time period, and gene flow was indexed by the proportion of 100 m<sup>2</sup> *Populus* cohorts greater than ten years of age that contained at least one transgenic tree outside of plantations (Mean Area of Mature Transgenics). Responses were averaged over the final 25 years of the simulation to simplify presentation of results (responses stabilized by age 25 for the simulations shown).



(a) Effects of fertility of transgenic trees relative to non-transgenics.



(b) Interaction between vegetative establishment and fertility. Vegetative establishment is the proportion of established individuals in a new cohort that are derived from vegetative propagules. Variation in vegetative establishment had little overall effect on transgene flow, although a minor effect is apparent at low levels of fertility.



(c) Effects of unstable sterility on transgene flow. Probability of sterility breakdown is the probability of a reversion to fertility (x-axis), which is then restored with a fertility level of 0.1 or 0.5, sampled from a normal distribution with a standard deviation of 0.05 or 0.25, respectively. Reversion was permanent and cumulative (Cumul.) for each tree through time, or fertility was transient and reset to the original value each succeeding year of the simulation (Noncum.). Low values of instability had little effect on gene flow; a cumulative reversion rate of about 20%, with 50% fertility restoration, would be required for gene flow levels to approach those of fully fertile transgenic trees.

Source: DiFazio, 2002.

based on patterns of establishment along shipping (i.e. dispersal) routes (Crawley and Brown, 1995), and simulation modelling implicated seed viability as a major factor limiting spread of transgenic oilseed rape (Kelly *et al.*, 2005). Therefore, the effectiveness of partial sterility in attenuating gene flow is not surprising, but the model is useful in demonstrating the importance of different modes of reproduction (vegetative vs various degrees of sexual reproduction).

The model was also useful for exploring implications of unstable sterility. We simulated this by allowing some restoration of fertility for trees that began the simulations with highly reduced fertility (fertility level of 0.01 compared to wild-type trees) (Figure 3-7c). These simulations had three important parameters: the probability of reversion to fertility (sampled from a normal distribution), the level of fertility restoration for each reversion event (10 or 50%, sampled from a normal distribution), and the duration of the restoration (cumulative or permanent restoration vs non-cumulative or transient restoration, with reversion to the original fertility level each year). With a permanent restoration level of 50% per reversion event, a 20% probability of reversion was required for gene flow levels to approach those of fully fertile trees. With a permanent restoration level of 10%, gene flow was considerably less than full fertility, and this was true even with reversion rates as high as 60%. Gene flow with reversion rates up to 3% were nearly indistinguishable from that of trees with stable sterility. If reversion was not cumulative (i.e. fertility was reset to 0.01 each year for each tree), gene flow was still greatly reduced compared to wild trees and was marginally greater than for trees with stable sterility. These results were manifested across a broad range of probabilities of reversion. Reversion rates that we have observed under vegetative propagation for transgenic *Populus* (reported above) appear to be considerably below the rates required for significant effects on modelled transgene flow. In addition, such high rates of reversion would likely be detected with moderate pre-commercial screening and post-release monitoring efforts. The simulations discussed above dealt with sterility in relation to spread of neutral transgenes. Transgenes that enhanced the competitiveness of trees in wild settings caused greatly enhanced gene flow for fully fertile transgenic trees, but a tightly linked sterility gene was very effective at attenuating spread, even in the face of a strong selective advantage and incomplete sterility (DiFazio, 2002).

## CONCLUSIONS

“In theory, there is no difference between theory and practice. In practice, there is.”

Andrew S. Tannenbaum, TIGR (The Institute for Genomic Research)

There are many genes of interest for commercial purposes that are likely to present very low risks, either because they are very similar to native genes, because they will reduce fitness or be neutral in the wild, or because their benefits outweigh their detriments. At the same time, there may be crops, such as forms of bio-industrial crops that encode novel and potentially ecotoxic compounds, for which very strong biological containment would be clearly warranted. Nonetheless,



the loudest social resistance seems to focus not on the products, traits and their benefits vs risks, but on perception of ‘contamination’ by GMOs generally. Indeed, because of the long-known propensity for long distance movement of pollen and/or seed from most tree species, if complete containment is the social goal, there is unlikely to be any place for genetically engineered trees in forestry plantation or horticulture – at least not for many decades. The technologies and simulations presented assume that some level of transgene dispersal could be socially and biologically acceptable – much like dispersal of new or modified genes and chromosomes introduced by breeding continues to have high social acceptance.

It has often been said that plant sterility should be an easy trait to engineer; after all, there are dozens of ways to damage a motor so it does not work. Unfortunately, motors do not have the redundancy and resilience of biological systems that have evolved to reproduce ‘at all costs’, nor do vandalism-leaning auto mechanics face the large biological and social obstacles that researchers and companies do when trying to conduct field-relevant research with genetically engineered trees. To arrive at efficient, reliable, effective sterility systems, we make the following suggestions:

1. *Functional genomics in trees.* Much more basic functional genomics is required in model taxa that represent the major forestry species. In this research, the main candidate genes based on studies in *Arabidopsis* and other model plant species, combined with newly discovered genes from trees identified in QTL, EST or microarray studies of trees, would be repressed or over-expressed and their functions identified in the field or the greenhouse, hopefully under conditions of accelerated flowering. This should allow the most important genes and promoters to be identified, thus, informing efforts to combine genes in redundant, reliable systems. It is hoped that inducible systems that make use of the *FT* gene might provide the much needed acceleration in production of normal flowers (Böhlenius *et al.*, 2006).
2. *Transformation technology improvements.* Gene transfer, gene targeting and highly specific recombinase technology needs to be greatly improved if mutagenesis of floral genes, and efficient addition or removal of sterility genes in many genotypes, is to become feasible. This requires much basic research on innovative transformation, excision, and homologous recombination methods – first in model plant species; but then, considerable work will be required to transfer these systems to trees.
3. *Regulatory and intellectual property constraints.* Candidate sterility cassettes based on the results of suggestions 1 and 2 need to be designed to meet regulatory standards and have freedom to operate with respect to intellectual property. They must then be tested in a diversity of commercially relevant environments and genotypes for stability and pleiotropic effects. These should be combined with predictive assays where possible to enable their effectiveness and pleiotropy to be forecast from a young age. The current ‘anti-commons’ (Boettiger and Bennett, 2006), where the licences for each genetic and construct

element, and basic transformation technology, are owned by parties different from those bearing the costs and risks of this long-term research, appear to provide large disincentives to moving forward. High regulatory and licensing costs and market stigmas impede the ‘adaptive management’ approaches so common in forestry (where research and commercial development go hand-in-hand, a result of the high costs and long time frames for forestry research).

4. *Transparency*. Containment research, due to its cost, long time frame and high level of scrutiny from society, should ideally be conducted by non-commercial third parties. A similar model is applied for all environmental research by Weyerhaeuser Company because of the need for independent validation of results for social acceptance (P. Farnum, personal communication). It is doubtful that company-based research, where only selected results are presented to the public, will be trusted, yet this model continues to be followed by some biotechnology companies. Ironically, the “eco”-vandalism that is still common in Europe, and continues to be a concern in North America, limits the extent to which the details of field and laboratory research can be safely disclosed. It appears that both vandalism risks to companies and Forest Stewardship Council exclusion of genetically engineered trees from field trials – both motivated by ecological concerns over appropriate uses of forest biotechnology – are delaying, rather than promoting, the development of ecologically sound genetic engineering technologies.

Because of the rapid rate of growth of genetic information and technological innovations, we believe that highly efficient containment systems can be developed and their reliability established. Without such systems, which will require testing over many years, it appears that many kinds of transgenes may never obtain regulatory or social approval in many countries, greatly limiting the benefits that transgenic biotechnologies are likely to be capable of providing.

## REFERENCES

- Al-Ahmad, H., Galili, S. & Gressel, J. 2004. Tandem constructs to mitigate transgene persistence: tobacco as a model. *Molecular Ecology*, 13: 697–710.
- Albert, V.A., Soltis, D.E., Carlson, J.E., Farmerie, W.G., Wall, P.K., Ilut, D.C., Solow, T.M., Mueller, L.A., Landherr, L.L., Hu, Y., Buzgo, M., Kim, S., Yoo, M.J., Frohlich, M.W., Perl-Treves, R., Schlarbaum, S.E., Bliss, B.J., Zhang, X., Tanksley, S.D., Oppenheimer, D.G., Soltis, P.S., Ma, H., DePamphilis, C.W. & Leebens-Mack, J.H. 2005. Floral gene resources from basal angiosperms for comparative genomics research. *BMC Plant Biology*, 5: Art. no. 5.
- Albertsen, M. & Huffman, G. 2002. Biotin-binding compounds for induction of sterility in plants. United States Patent and Trademark Office 20020129399
- Baudot, G., Garcia, D., Hodge, R. & Perez, P. 2001. DNA sequences coding for a protein conferring male sterility. United States Patent and Trademark Office 6,207,883.
- Baulcombe, D. 2004. RNA silencing in plants. *Nature*, 431: 356–363.
- Belostotsky, D.A. & Meagher, R.B. 1996. A pollen-, ovule-, and early embryo-specific poly (A) binding protein from *Arabidopsis* complements essential functions in yeast. *Plant Cell*, 8: 1261–1275.
- Bernd-Souza, R.B., de Sa, M.F.G., Ellis, D.D. & McCown, B.H. 2000. A rat ribonuclease fused to late cotton pollen promoter severely reduces pollen viability in tobacco plants. *Genetics and Molecular Biology*, 23(2): 435–443.

- Bettany, A.J.E., Dalton, S., Timms, E. & Morris, P. 1998. Stability of transgene expression during vegetative propagation of protoplast-derived tall fescue (*Festuca arundinacea* Schreb), *Journal of Experimental Botany*, 49: 1797–1804.
- Boettiger, S. & Bennett, A.B. 2006. Bayh-Dole: if we knew then what we know now. *Nature Biotechnology*, 24: 320–323.
- Böhlenius, H., Huang, T., Charbonnel-Campaa, L., Brunner, A.M., Jansson, S., Strauss, S.H. & Nilsson, O. 2006. The conserved CO/FT regulatory module controls timing of flowering and seasonal growth cessation in trees. *Science*, 312: 1040–1043.
- Borkowska, M., Kleczkowski, K., Pawelczak, A. & Wielgat, B. 1995. Transformation of diploid potato with an *Agrobacterium tumefaciens* binary vector system. II. Stability of transformation in tubers, micropropagated and greenhouse grown plants. *Acta Physiologia Plantarum*, 17: 275–280.
- Borner, R., Kampmann, G., Chandler, J., Gleibner, R., Wisman, E., Apel, K. & Melzer, S. 2000. A MADS domain gene involved in the transition to flowering in *Arabidopsis*. *Plant Journal*, 24: 591–599.
- Boss, P.K., Bastow, R.M., Mylne, J.S. & Dean, C. 2004. Multiple pathways in the decision to flower: enabling, promoting, and resetting. *Plant Cell*, 16: 18–31.
- Bradford, K., Van Deynze, A., Gutterson, N., Parrott, W. & Strauss, S.H. 2005. Regulating transgenic crops sensibly: lessons from plant breeding, biotechnology and genomics. *Nature Biotechnology*, 23: 439–444.
- Bradshaw, H.D. Jr & Stettler, R.F. 1993. Molecular genetics of growth and development in *Populus*. I. Triploidy in hybrid poplars. *Theoretical and Applied Genetics*, 86: 301–307.
- Brenner, E.D., Katari, M.S., Stevenson, D.W., Rudd, S.A., Douglas, A.W., Moss, W.N., Twigg, R.W., Runko, S.J., Stellari, G.M., McCombie, W.R. & Coruzzi, G.M. 2005. EST analysis in *Ginkgo biloba*: an assessment of conserved developmental regulators and gymnosperm specific genes. *BMC Genomics*, 6: Art. no. 143.
- Bridges, I., Bright, S., Greenland, A. & Schuch, W. 2001. Plant gene construct encoding a protein capable of disrupting the biogenesis of viable pollen. United States Patent and Trademark Office 6,172,279.
- Brown, G. 2002. Method for enhancement of naturally occurring cytoplasmic male sterility and for restoration of male fertility and uses thereof in hybrid crop production. United States Patent and Trademark Office 20020133851, 2 patents.
- Brunner, A.M. & Nilsson, O. 2004. Revisiting tree maturation and floral initiation in the poplar functional genomics era. *New Phytologist*, 164: 43–51.
- Brunner, A.M., Rottmann, W.H., Sheppard, L.A., Krutovskii, K., DiFazio, S.P., Leonardi, S. & Strauss, S.H. 2000. Structure and expression of duplicate AGAMOUS orthologues in poplar. *Plant Molecular Biology*, 44: 619–634.
- Burch-Smith, T.M., Anderson, J.C., Martin, G.B. & Dinesh-Kumar, S.P. 2004. Applications and advantages of virus-induced gene silencing for gene function studies in plants. *Plant Journal*, 39: 734–746.
- Burgess, D.G., Ralston, E.J., Hanson, W.G., Heckert, M., Ho, M., Jenq, T., Palys, J.M., Tang, K. & Gutterson, N. 2002. A novel, two-component system for cell lethality and its use in engineering nuclear male-sterility in plants. *Plant Journal*, 31: 113–125.
- Busov, V., Meilan, R., Pearce, D.W., Rood, S.B., Ma, C., Tschaplinski, T.J. & Strauss, S.H. 2006. Transgenic modification of *gai* or *rgl1* causes dwarfing and alters gibberellins, root growth, and metabolite profiles in *Populus*. *Planta*, 24: 288–299.
- Carlsbecker, A., Sundstrom, J., Tandré, K., Englund, M., Kvarnheden, A., Johanson, U. & Engstrom, P. 2003. The *DAL10* gene from Norway spruce (*Picea abies*) belongs to a potentially gymnosperm-specific subclass of MADS-box genes and is specifically active in seed cones and pollen cones. *Evolution & Development*, 5(6): 551–561.
- Carlsbecker, A., Tandré, K., Johanson, U., Englund, M. & Engstrom, P. 2004. The MADS-box gene *DAL1* is a potential mediator of the juvenile-to-adult transition in Norway spruce (*Picea abies*). *Plant Journal*, 40: 546–557.

- Cashore, B., Auld, G. & Newsom, D. 2003. Forest certification (eco-labeling) programs and their policy-making authority: explaining divergence among North American and European case studies. *Forest Policy and Economics*, 5(3): 225–247.
- Cervera, M., Pina, J.A., Juárez, J., Navarro, L. & Pena, L. 2000. A broad exploration of a transgenic population of citrus: stability of gene expression and phenotype. *Theoretical and Applied Genetics*, 100: 670–677.
- Chen, Y., Schneeberger, R.G. & Cullis, C.A. 2005. A site-specific insertion sequence in flax genotrophs induced by environment. *New Phytologist*, 167: 171–180.
- Cho, H., Kim, S., Kim, M. & Kim, B. 2001. Production of transgenic male sterile tobacco plants with the cDNA encoding a ribosome inactivating protein in *Dianthus sinensis*. *Molecules and Cells*, 11(3): 326–333.
- Chrimes, D., Rogers, H.J., Francis, D., Jones, H.D. & Ainsworth, C. 2005. Expression of fission yeast *cdc25* driven by the wheat ADP-glucose pyrophosphorylase large subunit promoter reduces pollen viability and prevents transmission of the transgene in wheat. *New Phytologist*, 166: 185–192.
- Chuang, C.F. & Meyerowitz, E.M. 2000. Specific and heritable genetic interference by double-stranded RNA in *Arabidopsis thaliana*. *Proceedings of the National Academy of Sciences of the United States of America*, 97: 4985–4990.
- Cigan, A. & Albertsen, M. 2002. Reversible nuclear genetic system for male sterility in transgenic plants. United States Patent and Trademark Office 6,399,856. 8 patents.
- Cigan, A.M., Unger-Wallace, E. & Haug-Collet, K. 2005. Transcriptional gene silencing as a tool for uncovering gene function in maize. *Plant Journal*, 43: 929–940.
- Clark, J.S., Lewis, M., McLachlan, J.S. & HilleRisLambers, J. 2003. Estimating population spread: what can we forecast and how well? *Ecology*, 84: 1979–1988.
- Comai, L. & Cartwright, R.A. 2005. A toxic mutator and selection alternative to the non-Mendelian RNA cache hypothesis for hothead reversion. *Plant Cell*, 17: 2856–2858.
- Conner, A.J., Mlynrov, L., Stiekema, W.J. & Nap, J. 1998. Meiotic stability of transgene expression is unaffected by flanking matrix-associated regions. *Molecular Breeding*, 4: 47–58.
- Crawley, M.J. & Brown, S.L. 1995. Seed limitation and the dynamics of feral oilseed rape on the M25 motorway. *Proceedings of the Royal Society of London Series B-Biological Sciences*, 259(1354): 49–54.
- Cseke, L.J., Zheng, J. & Podila, G.K. 2003. Characterization of PTM5 in aspen trees: a MADS-box gene expressed during woody vascular development. *Gene*, 318: 55–67.
- Cummins, J. & Ho, M.W. 2005. Terminator Trees. *Institute of Science in Society (ISIS) Report* 01/03/05 (available at [www.i-sis.org.uk/full/TerminatorTreesFull.php](http://www.i-sis.org.uk/full/TerminatorTreesFull.php), 29 May 2010).
- Daniel, T.W., Helms, J.A. & Baker, F.S. 1979. *Principles of silviculture*. McGraw-Hill, New York, USA.
- Davis, W. 2000. Mutant male sterile gene of soybean. United States Patent and Trademark Office 6,046,385.
- de Folter, S., Immink, R.G., Kieffer, M., Parenicova, L., Henz, S.R., Weigel, D., Busscher, M., Kooiker, M., Colombo, L., Kater, M.M., Davies, B. & Angenent, G.C. 2005. Comprehensive interaction map of the *Arabidopsis* MADS Box transcription factors. *Plant Cell*, 17: 1424–1433.
- Dellaporta, S. & Moreno, M. 2004. Methods and compositions to reduce or eliminate transmission of a transgene. United States Patent and Trademark Office 20040154054.
- Dievart, A., Dalal, M., Tax, F.E., Lacey, A.D., Huttly, A., Li, J. & Clark, S.E. 2003. CLAVATA1 dominant-negative alleles reveal functional overlap between multiple receptor kinases that regulate meristem and organ development. *Plant Cell*, 15: 1198–1211.
- DiFazio, S.P. 2002. Measuring and modeling gene flow from hybrid poplar plantations: implications for transgenic risk assessment. Ph.D. Thesis, Oregon State University, Corvallis, Oregon, USA.
- Dirks, R., Trinks, K., Uijtewaal, B., Bartsch, K., Peeters, R., Hofgen, R. & Pohlenz, H.-D. 2001. Process for generating male sterile plants. United States Patent and Trademark Office 6,262,339.

- Doering, D.S. 2004. *Designing genes: aiming for safety and sustainability in U.S. agriculture and biotechnology*. World Resources Institute, Washington, DC, USA. 43 p.
- Dunning, J.B., Stewart, D.J., Danielson, B.J., Noon, B.R., Root, T.L., Lamberson, R.H. & Stevens, E.E. 1995. Spatially explicit population models: current forms and future uses. *Ecological Applications*, 5(1): 3–11.
- Ellstrand, N.C. 2003. *Dangerous liaisons? When cultivated plants mate with their wild relatives*. The Johns Hopkins University Press, Baltimore, Maryland, USA. 244 p.
- Elo, A., Lemmetyinen, J., Turunen, M.L., Tikka, L. & Sopanen, T. 2001. Three MADS-box genes similar to *APETALA1* and *FRUITFUL* from silver birch (*Betula pendula*). *Physiologia Plantarum*, 112(1): 95–103.
- Endo, T., Shimada, T., Fujii, H., Kobayashi, Y., Araki, T. & Omura, M. 2005. Ectopic expression of an *FT* homolog from citrus confers an early flowering phenotype on trifoliolate orange (*Poncirus trifoliata* L. Raf.). *Transgenic Research*, 14: 703–712.
- ETC [Action Group on Erosion, Technology and Concentration]. 2006. Terminator Threat Looms: Intergovernmental meeting to tackle suicide seeds issue. ETC Group News Release, 20 January 2006 (available at [www.etcgroup.org/en/node/36](http://www.etcgroup.org/en/node/36), accessed 24 May 2010).
- Ferrario, S., Busscher, J., Franken, J., Gerats, T., Vandebussche, M., Angenent, G.C. & Immink, R.G. 2004. Ectopic expression of the petunia MADS box gene *UNSHAVEN* accelerates flowering and confers leaf-like characteristics to floral organs in a dominant-negative manner. *Plant Cell*, 16: 1490–1505.
- Fisher, R.A. 1937. The wave of advance of advantageous genes. *Annals of Eugenics*, 7: 355–369.
- Fladung, M. 1999. Gene stability in transgenic aspen (*Populus*). I. Flanking DNA sequences and T-DNA structure. *Molecular Genomics and Genetics*, 260: 574–581.
- Gallo-Meagher, M. & Irvine, J.E. 1996. Herbicide resistant transgenic sugarcane plants containing the bar gene. *Crop Science*, 36: 1367–1374.
- Gilbertson, L. 2003. *Cre-lox* recombination: creative tools for plant biotechnology. *Trends in Biotechnology*, 21: 550–555.
- Goetz, M., Godt, D.E., Giuvarch, A., Kahmann, U., Chriqui, D. & Roitsch, T. 2001. Induction of male sterility in plants by metabolic engineering of the carbohydrate supply. *Proceedings of the National Academy of Sciences of the United States of America*, 98: 6522–6527.
- Gressel, J. 1999. Tandem constructs: preventing the rise of super-weeds. *Trends in Biotechnology*, 17: 361–366.
- Groover, A.T. 2005. What genes make a tree a tree? *Trends in Plant Science*, 10: 210–214.
- Guan, X., Stege, J., Kim, M., Dahmani, Z., Fan, N., Heifetz, P., Barbas, C.F. 3rd & Briggs, S.P. 2002. Heritable endogenous gene regulation in plants with designed polydactyl zinc finger transcription factors. *Proceedings of the National Academy of Sciences of the United States of America*, 99: 13296–13301.
- Guerineau, F., Sorensen, A., Fenby, N. & Scott, R.J. 2003. Temperature-sensitive diphtheria toxin confers conditional male-sterility in *Arabidopsis thaliana*. *Plant Biotechnology Journal*, 1: 33–42.
- Hails, R.S. & Morley, K. 2005. Genes invading new populations: a risk assessment perspective. *Trends in Ecology & Evolution*, 20(5): 245–252.
- Han, Y., Griffiths, A., Li, H. & Grierson, D. 2004. The effect of endogenous mRNA levels on co-suppression in tomato. *FEBS Letters*, 563: 123–128.
- Harberd, N., Peng, J., Carol, P. & Richards D. 2004. Nucleic acid encoding GAI gene of *Arabidopsis thaliana*. United States Patent and Trademark Office 6,830,930.
- Harberd, N., Richards, D.E. & Peng, J. 2004. Genetic control of plant growth and development. United States Patent and Trademark Office 6,762,348.
- Hawkins, S., Lep le, J., Cornu, D., Jouanin, L. & Pilate, G. 2003. Stability of transgene expression in poplar: a model forest tree species. *Annals of Forest Science*, 60: 427–438.
- Haygood, R., Ives, A.R. & Andow, D.A. 2003. Consequences of recurrent gene flow from crops to wild relatives. *Proceedings of the Royal Society of London – Series B – Biological Sciences*, 270(1527): 1879–1886.



- Higgins, S.I. & Cain, M.L. 2002. Spatially realistic plant metapopulation models and the colonization-competition trade-off. *Journal of Ecology*, 90: 616–626.
- Higgins, S.I. & Richardson, D.M. 1999. Predicting plant migration rates in a changing world: the role of long-distance dispersal. *American Naturalist*, 153: 464–475.
- Higgins, S.I., Richardson, D.M. & Cowling, R.M. 1996. Modelling invasive plant spread: the role of plant-environment interactions and model structure. *Ecology*, 77: 2043–2054.
- Hill, R.A. 2005. Conceptualizing risk assessment methodology for genetically modified organisms. *Environmental Biosafety Research*, 4(2): 67–70.
- Hiratsu, K., Matsui, K., Koyama, T. & Ohme-Takagi, M. 2003. Dominant repression of target genes by chimeric repressors that include the EAR motif, a repression domain, in *Arabidopsis*. *Plant Journal*, 34: 733–739.
- Hird, D.L., Paul, W., Hollyoak, J.S. & Scott, R.J. 2000. The restoration of fertility in male sterile tobacco demonstrates that transgene silencing can be mediated by T-DNA that has no DNA homology to the silenced transgene. *Transgenic Research*, 9: 91–102.
- Hoenicke, H. & Fladung, M. 2006a. Biosafety in *Populus* spp. and other forest trees: from non-native species to taxa derived from traditional breeding and genetic engineering. *Trees*, 20: 131–144.
- Hoenicke, H. & Fladung, M. 2006b. Genome instability in woody plants derived from genetic engineering. pp. 301–321, in: M. Fladung & D. Ewald (editors). *Tree transgenesis – recent developments*. Springer, Berlin, Heidelberg and New York.
- Höfig, K.P., Möller, R., Donaldson, L., Putterill, J. & Walter, C. 2006. Towards male-sterility in *Pinus radiata* – a stilbene synthase approach to genetically engineer nuclear male sterility. *Plant Biotechnology Journal*, 4: 333–343.
- Howard, R.D., DeWoody, J.A. & Muir, W.M. 2004. Transgenic male mating advantage provides opportunity for Trojan gene effect in a fish. *Proceedings of the National Academy of Sciences of the United States of America*, 101: 2934–2938.
- Hsu, C.Y., Liu, Y., Luthe, D.S. & Yuceer, C. 2006. Poplar FT2 shortens the juvenile phase and promotes seasonal flowering. *Plant Cell*, 18(8): 1846–1861.
- Iglesias, V.A., Moscone, E.A., Papp, I., Neuhuber, F., Michalowski, S., Phelan, T., Spiker, S., Matzke, M. & Matzke, A.J.M. 1997. Molecular and cytogenetic analyses of stably and unstably expressed transgene loci in tobacco. *Plant Cell*, 9: 1251–1264.
- Irish, V.F. & Litt, A. 2005. Flower development and evolution: gene duplication, diversification and redeployment. *Current Opinion in Genetic Development*, 15: 454–460.
- ISAAA [International Service for the Acquisition of Agri-biotech Applications]. 2006. Brazil: curtailing research and technological innovation. CropBiotech Update, ISAAA, SEAsia Center, 14 July 2006 (available at [www.cedab.it/newsletter\\_ISAAA.asp?IDnews=127#an\\_cora2](http://www.cedab.it/newsletter_ISAAA.asp?IDnews=127#an_cora2), accessed 29 July 2006).
- Jackson, A.L. & Linsley, P.S. 2004. Noise amidst the silence: off-target effects of siRNAs? *Trends in Genetics*, 20: 521–524.
- Jager, M., Hassanin, A., Manuel, M., Le Guyader, H. & Deutsch, J. 2003. MADS-box genes in *Ginkgo biloba* and the evolution of the AGAMOUS family. *Molecular Biology and Evolution*, 20(5): 842–854.
- James, R.R., DiFazio, S.P., Brunner, A.M. & Strauss, S.H. 1998. Environmental effects of genetic engineering of woody biomass crops. *Biomass Bioenergy*, 14: 403–414.
- James, V., Worland, B., Snape, J.W. & Vain, P. 2004. Strategies for precise quantification of transgene expression levels over several generations in rice. *Journal of Experimental Botany*, 55: 1307–1313.
- Jeon, J.S., Jang, S., Lee, S., Nam, J., Kim, C., Lee, S.H., Chung, Y.Y., Kim, S.R., Lee, Y.H., Cho, Y.G. & An, G. 2000. Leafy hull sterile1 is a homeotic mutation in a rice MADS box gene affecting rice flower development. *Plant Cell*, 12: 871–884.
- Johnson, B. & Kirby, K. 2004. Potential impacts of genetically modified trees on biodiversity of forestry plantations: a global perspective. pp. 190–207, in: S.H. Strauss & H.D. Bradshaw (editors). *The bio-engineered forest: challenges to science and society*. Resources for the Future. Washington DC, USA.

- Katul, G.G., Porporato, A., Nathan, R., Siqueira, M., Soons, M.B., Poggi, D., Horn, H.S. & Levin, S.A. 2005. Mechanistic analytical models for long-distance seed dispersal by wind. *American Naturalist*, 166: 368–381.
- Kelly, C.K., Bowler, M.G., Breden, F., Fenner, M. & Poppy, G.M. 2005. An analytical model assessing the potential threat to natural habitats from insect resistance transgenes. *Proceedings of the Royal Society B – Biological Sciences*, 272(1574): 1759–1767.
- Kerschen, A., Napoli, C.A., Jorgensen, R.A. & Muller, A.E. 2004. Effectiveness of RNA interference in transgenic plants. *FEBS Letters*, 566: 223–228.
- Kilby, N.J., Leyser, H.M.O. & Furner, I.J. 1992. Promoter methylation and progressive transgene inactivation in *Arabidopsis*. *Plant Molecular Biology*, 20: 103–112.
- Knox, R., Singh, M. & Xu, H. 2004. Developmental regulation in anther tissue of plants. United States Patent and Trademark Office 6,740,748.
- Kohli, A., Gahakwa, D., Vain, P., Laurie, D.A. & Christou, P. 1999. Transgene expression in rice engineered through particle bombardment: molecular factors controlling stable expression and transgene silencing. *Planta*, 208: 88–97.
- Kreiman, G. 2004. Identification of sparsely distributed clusters of cis-regulatory elements in sets of co-expressed genes. *Nucleic Acids Research*, 32: 2889–2900.
- Kumar, S. & Fladung, M. 2001. Gene stability in transgenic aspen (*Populus*). II. Molecular characterization of variable expression of transgene in wild and hybrid aspen. *Planta*, 213: 731–740.
- Kuvshinov, V.V., Koivu, K., Kanerva, A. & Pehu, E. 2001. Molecular control of transgene escape from genetically modified plants. *Plant Science*, 160: 517–522.
- Kyozuka, J., Harcourt, R., Peacock, W.J. & Dennis, E.S. 1997. Eucalyptus has functional equivalents of the *Arabidopsis AP1* gene. *Plant Molecular Biology*, 35: 573–584.
- Lännepää, M., Hassinen, M., Ranki, A., Hölttä-Vuora, M., Lemmetyinen, J., Keinonen, K. & Sapanen, T. 2005. Prevention of flower development in birch and other plants using a *BPFULL1::BARNASE* construct. *Plant Cell Reports*, 24: 69–78.
- Lavorel, S., Smith, M.S. & Reid, N. 1999. Spread of mistletoes (*Amyema preissii*) in fragmented Australian woodlands: a simulation study. *Landscape Ecology*, 14: 147–160.
- Lee, Y.H., Chung, K.H., Kim, H.U., Jin, Y.M., Kim, H.I. & Park, B.S. 2003. Introduction of male-sterile cabbage using a tapetum-specific promoter from *Brassica campestris* L. ssp. *pekinensis*. *Plant Cell Reports*, 22: 268–273.
- Leibbrandt, N.B. & Snyman S.J. 2003. Stability of gene expression and agronomic performance of a transgenic herbicide-resistant sugarcane line in South Africa. *Crop Science*, 43: 671–677.
- Lemmetyinen, J., Keinonen, K. & Sapanen, T. 2004. Prevention of flowering of a tree, silver birch. *Molecular Breeding*, 13: 243–249.
- Lemmetyinen, J., Hassinen, M., Elo, A., Porali, I., Keinonen, K., Makela, H. & Sapanen, T. 2004. Functional characterization of *SEPALLATA3* and *AGAMOUS* orthologues in silver birch. *Physiologia Plantarum*, 121: 149–162.
- Li, X., Zhong, S. & Wong, W.H. 2005. Reliable prediction of transcription factor binding sites by phylogenetic verification. *Proceedings of the National Academy of Sciences of the United States of America*, 102: 16945–16950.
- Li, Y. & Pei, Y. 2006. Biotech approaches to improve biomass production of poplar and to produce genetically modified gene free pollen and seed from genetically modified plants. p. 13 (abstracts), in: International Poplar Symposium IV. Nanjing, China, 5–9 June 2006.
- Liebhart, R., Kellerhals, M., Pfammatter, W., Jertmini, M. & Gessler, C. 2003. Mapping quantitative physiological traits in apple (*Malus × domestica* Borkh.). *Plant Molecular Biology*, 52: 511–526.
- Lloyd, A., Plaisier, C.L., Carroll, D. & Drews, G.N. 2005. Targeted mutagenesis using zinc-finger nucleases in *Arabidopsis*. *Proceedings of the National Academy of Sciences of the United States of America*, 102: 2232–2237.
- Lolle, S.J., Victor, J.L., Young, J.M. & Pruitt, R.E. 2005. Genome-wide non-Mendelian inheritance of extra-genomic information in *Arabidopsis*. *Nature*, 434: 505–509.



- Luo, H., Kausch, A.P., Hu, Q., Nelson, K., Wipff, J.K., Fricker, C.C., Owen, T., Moreno, M.A., Lee, J. & Hodges, T.K. 2005. Controlling transgene escape in GM creeping bentgrass. *Molecular Breeding*, 16: 185–188.
- Mariani, C., Beuckeleer, M.D., Truettner, J., Leemans, J. & Goldberg, R.B. 1990. Induction of male sterility in plants by a chimaeric ribonuclease gene. *Nature*, 347: 737–741.
- Mariani, C., Gossele, V., Beuckeleer, M.D., Block, M.D., Goldberg, R.B., Greef, W.D. & Leemans, J. 1992. A chimaeric ribonuclease-inhibitor gene restores fertility to male sterile plants. *Nature*, 357: 384–387.
- Matzke, M.A. & Birchler, J.A. 2005. RNAi-mediated pathways in the nucleus. *Nature Reviews Genetics*, 6: 24–35.
- Matzke, M., Aufsatz, W., Kanno, T., Daxinger, L., Papp, I., Mette, M.F. & Matzke, A.J. 2004. Genetic analysis of RNA-mediated transcriptional gene silencing. *Biochimica et Biophysica Acta-Gene Structure and Expression*, 1677(1-3): 129–141.
- Meilan, R., Auerbach, D.J., Ma, C., DiFazio, S.P. & Strauss, S.H. 2002. Stability of herbicide resistance and GUS expression in trans-genic hybrid poplars (*Populus* sp.) during four years of field trials and vegetative propagation. *HortScience*, 37: 277–280.
- Mellerowicz, E.J., Horgan, K., Walden, A., Coker, A. & Walter, C. 1998. *PRFLL*-a *Pinus radiata* homologue of *FLORICAULA* and *LEAFY* is expressed in buds containing vegetative shoot and undifferentiated male cone primordia. *Planta*, 206: 619–629.
- Meyer, P., Linn, F., Heidmann, I., Meyer, H., Niedenhof, I. & Saedler, H. 1992. Endogenous and environmental factors influence 35S promoter methylation of a maize A1 gene construct in transgenic petunia and its colour phenotype. *Molecular Genomics and Genetics*, 231: 345–352.
- Meza, T.J., Kamfjord, D., Hakelien, A., Evans, I., Godager, L.H., Mandal, A., Jakobsen, K.S. & Aalen, R.B. 2001. The frequency of silencing in *Arabidopsis thaliana* varies highly between progeny of siblings and can be influenced by environmental factors. *Transgenic Research*, 10: 53–67.
- Michiels, F., Botterman, J. & Cornelissen, M. 2000. Method to obtain male sterile plants. United States Patent and Trademark Office 6,025,546.
- Miki, D., Itoh, R. & Shimamoto, K. 2005. RNA silencing of single and multiple members in a gene family of rice. *Plant Physiology*, 138: 1903–1913.
- Missiaggia, A.A., Piacuzzi, A.L. & Grattapaglia, D. 2005. Genetic mapping of *Eef1*, a major effect QTL for early flowering in *Eucalyptus grandis*. *Tree Genetics & Genomes*, 1(2): 79–84.
- Mitsuda, N., Hiratsu, K., Todaka, D., Nakashima, K., Yamaguchi-Shinozaki, K. & Ohme-Takagi, M. 2006. Efficient production of male and female sterile plants by expression of a chimeric repressor in *Arabidopsis* and rice. *Plant Biotechnology Journal*, 4: 325–332.
- Mlynárová, L., Conner, A.J. & Nap, J.-P. 2006. Directed microspore-specific recombination of transgenic alleles to prevent pollen-mediated transmission of transgenes. *Plant Biotechnology Journal*, 4: 445–452.
- Mohamed, R. 2006. Expression and function of *Populus* homologs to *TERMINAL FLOWER 1* genes: roles in onset of flowering and shoot phenology. Ph.D. Thesis, Oregon State University, Corvallis, Oregon, USA. 136 p. (available at [www.fsl.orst.edu/tgerc/strauss/Rozi\\_Mohamed\\_PhD\\_Thesis\\_January\\_2006.pdf](http://www.fsl.orst.edu/tgerc/strauss/Rozi_Mohamed_PhD_Thesis_January_2006.pdf), accessed 30 May 2010).
- Morina, K., Olsen, O. & Shimamoto, K. 1999. Silencing of an aleurone-specific gene in transgenic rice is caused by a rearranged transgene. *Plant Journal*, 17: 275–285.
- Mou, Z., Wang, X., Fu, Z., Dai, Y., Han, C., Ouyang, J., Bao, F., Hu, Y. & Li, J. 2002. Silencing of phosphoethanolamine N-methyltransferase results in temperature-sensitive male sterility and salt hypersensitivity in *Arabidopsis*. *Plant Cell*, 14: 2031–2043.
- Mouradov, A., Glassick, T., Hamdorf, B., Murphy, L., Fowler, B., Marla, S. & Teasdale, R.D. 1998. *NEEDLY*, a *Pinus radiata* ortholog of *FLORICAULA/LEAFY* genes, expressed in both reproductive and vegetative meristems. *Proceedings of the National Academy of Sciences of the United States of America*, 95: 6537–6542.

- Müller, A.J., Mendel, R.R., Schiemann, J., Simoens, C. & Inzé, D. 1987. High meiotic stability of a foreign gene introduced into tobacco by *Agrobacterium*-mediated transformation. *Molecular Genomics and Genetics*, 207: 171–175.
- Nathan, R., Katul, G.G., Horn, H.S., Thomas, S.M., Oren, R., Avissar, R., Pacala, S.W. & Levin, S.A. 2002. Mechanisms of long-distance dispersal of seeds by wind. *Nature*, 418: 409–413.
- Nathan, R., Perry, G., Cronin, J.T., Strand, A.E. & Cain, M.L. 2003. Methods for estimating long-distance dispersal. *Oikos*, 103: 261–273.
- Naylor, M., Reeves, J., Cooper, J.L., Edwards, M.L. & Wang, H. 2005. Construction and properties of a gene-silencing vector based on Poplar mosaic virus (genus Carlavirus). *Journal of Virology Methods*, 124: 27–36.
- NRC [National Research Council]. 2004. *Biological confinement of genetically engineered organisms*. The National Academies, Washington, DC, USA. 256 p.
- Ottaviani, M.P., Hanisch ten Cate, C.H. & Doting, L.V. 1992. Expression of introduced genes after tuber propagation of transgenic potato plants. *Plant Breeding*, 109: 89–96.
- Ouellet, T., Singh, J., Tao, T. & Simmonds, J. 2003. Corn silk gene and regulatory region. United States Patent and Trademark Office 6,515,204.
- Parker, I.M. & Kareiva, P. 1996. Assessing the risks of invasion for genetically engineered plants: acceptable evidence and reasonable doubt. *Biological Conservation*, 78(1-2): 193–203.
- Patell, V., Rayapuram, N., Venkataramiah, M., Joma, J. & Goswami, S. 2003. Process for generating cytoplasmic male sterile line in rice and other crops by RNA editing. United States Patent and Trademark Office 20030163856.
- Pavingerová, D., Dostál, J., Bísková, R. & Benetka, V. 1994. Somatic embryogenesis and *Agrobacterium*-mediated transformation of chrysanthemum. *Plant Science*, 97: 95–101.
- Pena, L., Martin-Trillo, M., Juarez, J., Pina, J.A., Navarro, L. & Martinez-Zapater, J.M. 2001. Constitutive expression of *Arabidopsis* *LEAFY* or *APETALA1* genes in citrus reduces their generation time. *Nature Biotechnology*, 19: 263–267.
- Petersen, B.O. & Albrechtsen, M. 2005. Evidence implying only unprimed RdRP activity during transitive gene silencing in plants. *Plant Molecular Biology*, 58(4): 575–583.
- Pilate, G., Guiney, E., Holt, K., Petit-Conil, M., Lapierre, C., Lep'l'e, J., Pollet, B., Mila, I., Webster, E.A., Marstorp, H.G., Hopkins, D.W., Jouanin, L., Boerjan, W., Schuch, W., Cornu, D. & Halpin, C. 2002. Field and pulping performances of transgenic trees with altered lignification. *Nature Biotechnology*, 20: 607–612.
- Pilson, D. & Prendeville, H.R. 2004. Ecological effects of transgenic crops and the escape of transgenes into wild populations. *Annual Review of Ecology, Evolution and Systematics*, 35: 149–174.
- Poovalah, B., Patil, S. & Takezawa, D. 2002. Compositions and methods for production of male-sterile plants. United States Patent and Trademark Office 6,403,352.
- Preston, J., Wheeler, J., Heazlewood, J., Li, S.F. & Parish, R.W. 2004. AtMYB32 is required for normal pollen development in *Arabidopsis thaliana*. *Plant Journal*, 40: 979–995.
- Quandt, J., Bartsch, K. & Knittel, N. 2002. Conditional sterility in wheat. United States Patent and Trademark Office 6,384,304.
- Raser, J.M. & O'Shea, E.K. 2004. Control of stochasticity in eukaryotic gene expression. *Science*, 304: 1811–1814.
- Ray, A. & Golden, T. 2004. Gene encoding short integuments and uses thereof. United States Patent and Trademark Office 6,737,561.
- Rees, M. & Paynter, Q. 1997. Biological control of Scotch broom: modelling the determinants of abundance and the potential impact of introduced insect herbivores. *Journal of Applied Ecology*, 34: 1203–1221.
- Rombauts, S., Florquin, K., Lescot, M., Marchal, K., Rouze, P. & van de Peer, Y. 2003. Computational approaches to identify promoters and cis-regulatory elements in plant genomes. *Plant Physiology*, 132: 1162–1176.

- Rood, S.B., Kalischuk, A.R., Polzin, M.L. & Braatne, J.H. 2003. Branch propagation, not cladogenesis, permits dispersive, clonal reproduction of riparian cottonwoods. *Forest Ecology and Management*, 186(1-3): 227–242.
- Rottmann, W.H., Meilan, R., Sheppard, L.A., Brunner, A.M., Skinner, J.S., Ma, C., Cheng, S., Jouanin, L., Pilate, G. & Strauss, S.H. 2000. Diverse effects of over expression of LEAFY and PTLF, a poplar (*Populus*) homolog of LEAFY/FLOICAULA, in transgenic poplar and *Arabidopsis*. *Plant Journal*, 22: 235–245.
- Rutledge, R., Regan, S., Nicolas, O., Fobert, P., Cote, C., Bosnich, W., Kauffeldt, C., Sunohara, G., Seguin, A. & Stewart, D. 1998. Characterization of an *AGAMOUS* homologue from the conifer black spruce (*Picea mariana*) that produces floral homeotic conversions when expressed in *Arabidopsis*. *Plant Journal*, 15: 625–634.
- Scheid, O.M., Paszkowski, J. & Potrykus, I. 1991. Reversible inactivation of a transgene in *Arabidopsis thaliana*. *Molecular Genomics and Genetics*, 228: 104–112.
- Schmid, M., Uhlentaut, N.H., Godard, F., Demar, M., Bressan, R., Weigel, D. & Lohmann, J.U. 2003. Dissection of floral induction pathways using global expression analysis. *Development*, 130: 6001–6012.
- Schmid, M., Davison, T.S., Henz, S.R., Pape, U.J., Demar, M., Vingron, M., Scholkopf, B., Weigel, D. & Lohmann, J.U. 2005. A gene expression map of *Arabidopsis thaliana* development. *Nature Genetics*, 37: 501–506.
- Schwab, R., Palatnik, J.F., Riester, M., Schommer, C., Schmid, M. & Weigel, D. 2005. Specific effects of microRNAs on the plant transcriptome. *Developmental Cell*, 8(4): 517–527.
- Scortecci, K.C., Michaels, S.D. & Amasino, R.M. 2001. Identification of a MADS-box gene, *FLOWERING LOCUS M*, that represses flowering. *Plant Journal*, 26: 229–236.
- Segal, D.J., Stege, J.T. & Barbas, C.F. 3rd. 2003. Zinc fingers and a green thumb: manipulating gene expression in plants. *Current Opinion in Plant Biology*, 6: 163–168.
- Shaked, H., Melamed-Bessudo, C. & Levy, A.A. 2005. High-frequency gene targeting in *Arabidopsis* plants expressing the yeast RAD54 gene. *Proceedings of the National Academy of Sciences of the United States of America*, 102:12265–12269
- Sheppard, L.A., Brunner, A.M., Krutovskii, K.V., Rottmann, W.H., Skinner, J.S., Vollmer, S.S. & Strauss, S.H. 2000. A *DEFICIENS* homolog from the dioecious tree *Populus trichocarpa* is expressed in both female and male floral meristems of its two-whorled, unisexual flowers. *Plant Physiology*, 124: 627–640.
- Shigesada, N. & Kawasaki, K. 1997. *Biological invasions: theory and practice*. Oxford University Press, New York, USA.
- Shindo, S., Ito, M., Ueda, K., Kato, M. & Hasebe, M. 1999. Characterization of MADS genes in the gymnosperm *Gnetum parvifolium* and its implication on the evolution of reproductive organs in seed plants. *Evolution and Development*, 1: 180–190.
- Skinner, J.S., Meilan, R., Brunner, A.M. & Strauss, S.H. 2000. Options for genetic engineering of floral sterility in forest trees. pp. 135–153, in: S.M. Jain & S.C. Minocha (editors). *Molecular biology of woody plants*. Kluwer, Dordrecht, the Netherlands.
- Skinner, J.S., Meilan, R., Ma, C. & Strauss, S.H. 2003. The *Populus* PTD promoter imparts floral-predominant expression and enables high levels of floral-organ ablation in *Populus*, *Nicotiana* and *Arabidopsis*. *Molecular Breeding*, 12: 119–132.
- Slavov, G.T., DiFazio, S.P. & Strauss, S.H. 2004. Gene flow in forest trees: gene migration patterns and landscape modeling of transgene dispersal in hybrid poplar. pp. 89–106, in: H.C.M. den Nijs, D. Bartsch & J. Sweet (editors). *Introgression from genetically modified plants into wild relatives*. CAB International, Wallingford, UK.
- Smeekens, S., Weisbeek, P. & Proveniers, M. 2005. Plant gene constructs and their use. United States Patent and Trademark Office 6,864,051.
- Smouse, P.E. & Sork, V.L. 2004. Measuring pollen flow in forest trees: an exposition of alternative approaches. *Forest Ecology and Management*, 197: 21–38.

- Snow, A.A., Andow, D.A., Gepts, P., Hallerman, E.M., Power, A., Tiedje, J.M. & Wolfenbarger, L.L. 2005. Genetically engineered organisms and the environment: current status and recommendations. *Ecological Applications*, 15(2): 377–404.
- Southerton, S.G., Marshall, H., Mouradov, A. & Teasdale, R.D. 1998. Eucalypt MADS-box genes expressed in developing flowers. *Plant Physiology*, 118: 365–372.
- Spena, A., Saedler, H., Sommer, H. & Rotino, G. 2002. Methods for producing parthenocarpic or female sterile transgenic plants and methods for enhancing fruit setting and development. United States Patent and Trademark Office 6,483,012.
- Stewart, C.N., Halfhill, M.D. & Warwick, S.I. 2003. Transgene introgression from genetically modified crops to their wild relatives. *Nature Reviews Genetics*, 4: 806–817.
- Stoutjesdijk, P.A., Singh, S.P., Liu, Q., Hurlstone, C.J., Waterhouse, P.A. & Green, A.G. 2002. hpRNA-mediated targeting of the *Arabidopsis* *FAD2* gene gives highly efficient and stable silencing. *Plant Physiology*, 129: 1723–1731.
- Strauss, S.H. 2003. Genomics, genetic engineering, and domestication of crops. *Science*, 300: 61–62.
- Strauss, S.H. & Bradshaw, H.D. (editors). 2004. *The bioengineered forest: challenges to science and society*. Resources for the Future, Washington, DC, USA. 245 p.
- Strauss, S.H. & Brunner, A.M. 2004. Tree biotechnology in the 21st century: transforming trees in the light of comparative genomics. pp. 76–97, in: Strauss & Bradshaw, 2004, q.v.
- Strauss, S.H., Rottmann, W.H., Brunner, A.W. & Sheppard, L.A. 1995. Genetic engineering of reproductive sterility in forest trees. *Molecular Breeding*, 1: 5–26.
- Strauss, S.H., Coventry, P., Campbell, M.M., Pryor, S.M. & Burley, J. 2001a. Certification of genetically modified forest plantations. *International Forestry Review*, 3: 87–104.
- Strauss, S.H., Campbell, M.M., Pryor, S.N., Coventry, P. & Burley, J. 2001b. Plantation certification and genetic engineering: FSC's ban on research is counterproductive. *Journal of Forestry*, 99(12): 4–7.
- Strauss, S., Rottmann, W., Brunner, A. & Sheppard, L. 2002. Floral homeotic genes for manipulation of flowering in poplar and other plant species. United States Patent and Trademark Office 6,395,892.
- Strauss, S.H., Brunner, A.M., Busov, V.B., Ma, C. & Meilan, R. 2004. Ten lessons from 15 years of transgenic *Populus* research. *Forestry*, 77: 455–465.
- Sundstrom, J., Carlsbecker, A., Svensson, M.E., Svenson, M., Johanson, U., Theissen, G. & Engstrom, P. 1999. MADS-box genes active in developing pollen cones of Norway spruce (*Picea abies*) are homologous to the B-class floral homeotic genes in angiosperms. *Developmental Genetics*, 25(3): 253–266.
- Tandre, K., Albert, V.A., Sundas, A. & Engstrom, P. 1995. Conifer homologues to genes that control floral development in angiosperms. *Plant Molecular Biology*, 27: 69–78.
- Vaistij, F.E., Jones, L. & Baulcombe, D.C. 2002. Spreading of RNA targeting and DNA methylation in RNA silencing requires transcription of the target gene and a putative RNA-dependent RNA polymerase. *Plant Cell*, 14: 857–867.
- Valenzuela, S. & Strauss, S.H. 2005. Lost in the woods. *Nature Biotechnology*, 23: 532–533.
- Wagner, A., Phillips, L., Narayan, R.D., Moody, J.M. & Geddes, B. 2005. Gene silencing studies in the gymnosperm *Pinus radiata*. *Plant Cell Reports*, 24: 95–102.
- Waterhouse, P. & Wang, M.-B. 2002. Methods for obtaining modified phenotypes in plant cells. United States Patent and Trademark Office 6,423,885.
- Watson, J.M., Fusaro, A.F., Wang, M. & Waterhouse, P.M. 2005. RNA silencing platforms in plants. *FEBS Letters*, 579(26; Special issue S1): 5982–5987.
- Wei, H., Meilan, R., Brunner, A.M., Skinner, J.S., Ma, C., Gandhi, H.T. & Strauss, S.H. 2007. Field trial detects incomplete barstar attenuation of vegetative cytotoxicity in *Populus* trees containing a poplar LEAFY promoter::barnase sterility transgene. *Molecular Breeding*, 19(1): 69–85.
- Weigel, D. & Nilsson, O. 1995. A developmental switch sufficient for flower initiation in diverse plants. *Nature*, 377: 495–500.

- Weisker, A. 1995. Hybrid safflower production utilizing genetic dwarf male sterility. United States Patent and Trademark Office 5,436,386.
- Wellmer, F., Riechmann, J.L., Alves-Ferreira, M. & Meyerowitz, E.M. 2004. Genome-wide analysis of spatial gene expression in *Arabidopsis* flowers. *Plant Cell*, 16: 1314–1326.
- Wesley, S.V., Helliwell, C.A., Smith, N.A., Wang, M.B., Rouse, D.T., Liu, Q., Gooding, P.S., Singh, S.P., Abbott, D., Stoutjesdijk, P.A., Robinson, S.P., Gleave, A.P., Green, A.G. & Waterhouse, P.M. 2001. Construct design for efficient, effective and high-throughput gene silencing in plants. *Plant Journal*, 27: 581–590.
- Whitham, T.G., Bailey, J.K., Schweitzer, J.A., Shuster, S.M., Bangert, R.K., LeRoy, C.J., Lonsdorf, E.V., Allan, G.J., DiFazio, S.P., Potts, B.M., Fischer, D.G., Gehring, C.A., Lindroth, R.L., Marks, J.C., Hart, S.C., Wimp, G.M. & Wooley, S.C. 2006. A framework for community and ecosystem genetics: from genes to ecosystems. *Nature Reviews Genetics*, 7(7): 510–523.
- Williams, C.G. 2005. Framing the issues on transgenic forests. *Nature Biotechnology*, 23: 530–532.
- Wright, D.A., Townsend, J.A., Winfrey, R.J. Jr, Irwin, P.A., Rajagopal, J., Lonosky, P.M., Hall, B.D., Jondle, M.D. & Voytas, D.F. 2005. High-frequency homologous recombination in plants mediated by zinc-finger nucleases. *Plant Journal*, 44: 693–705.
- Yanofsky, M. 2000. Seed plants exhibiting early reproductive development and methods of making same. United States Patent and Trademark Office 6,025,543.
- Yui, R., Iketani, S., Mikami, T. & Kubo, T. 2003. Antisense inhibition of mitochondrial pyruvate dehydrogenase E1 alpha subunit in anther tapetum causes male sterility. *Plant Journal*, 34(1): 57–66.
- Zahn, L.M., Leebens-Mack, J.H., Arrington, J.M., Hu, Y., Landherr, L.L., Depamphilis, C.W., Becker, A., Theissen, G. & Ma, H. 2006. Conservation and divergence in the *AGAMOUS* subfamily of MADS-box genes: evidence of independent sub- and neo-functionalization events. *Evolution and Development*, 8: 30–45.
- Zhang, Y., Shewry, P.R., Jones, H., Barcelo, P., Lazzeri, P.A. & Halford, N.G. 2001. Expression of antisense Sn RK1 protein kinase sequence causes abnormal pollen development and male sterility in transgenic barley. *Plant Journal*, 28: 431–441.
- Zhang, P., Tan, H.T., Pwee, K.H. & Kumar, P.P. 2004. Conservation of class C function of floral organ development during 300 million years of evolution from gymnosperms to angiosperms. *Plant Journal*, 37: 566–577.

## 4. Engineering trees with target traits

*L.M. McDonnell, H.D. Coleman, D.G. French, R. Meilan and S.D. Mansfield*

Despite their unparalleled importance both ecologically and economically, very little is known about the molecular mechanisms that underpin the development, growth and health of forest trees. However, the past decade has yielded remarkable progress in elucidating the biochemical and genetic mechanisms controlling the growth and survival of annual plants. An understanding of these processes will inform efforts aimed at ensuring the long-term maintenance and sustainability of global forests. Much of this progress has been made through the application of what is collectively known as functional genomics.

Functional genomics entails the analysis of an organism's genetic material (the genome), and relates this to its form and function. Genomic analysis of a tree or model plant may identify gene(s), or in some cases spontaneous or induced mutations, which offer opportunities for directed modification of the corresponding trait(s). Using new forward genomics techniques (e.g. activation tagging and gene/enhancer traps), it is now possible to induce useful 'mutations'. The resulting phenotype may reflect the underlying gene's function and may point to further, desirable alterations. Newly available, high-throughput screening techniques may also identify useful, but cryptic, mutations that are not normally manifested as an obvious phenotype (Davis *et al.*, 2006; Ehling *et al.*, 2005; Kelley *et al.*, 2004; Kent *et al.*, 2006; Labbe *et al.*, 2005; Tsuchikawa, 2007; Tuskan *et al.*, 1999; Wiklund *et al.*, 2005). Ultimately, functionality must be demonstrated by either silencing or overexpressing the putative gene.

Along with the great progress with the model plant *Arabidopsis thaliana*, recent publication of the first genomic sequence for a tree (*Populus trichocarpa*; Tuskan *et al.*, 2006), is revolutionizing our understanding of tree biology and permitting the exploitation of several powerful biotechnological techniques that can aid in the domestication of other tree species. The next decade should prove extremely exciting for tree researchers as they exploit the genomic sequence, and combine postgenomic technologies (e.g. cell biology, bio-informatics, transcriptome and proteome analyses, and metabolic profiling) to unravel many of the mysteries surrounding genotype-phenotype relationships in trees. Consideration of costs and benefits, including unintended effects, will influence the ultimate choice of a target trait. This is especially true when comparing directed target trait modification to alternative approaches (e.g. conventional breeding or cultural treatments).



## BIOTIC STRESS

Damage to forest trees caused by both native and introduced pests is of global importance. These biotic stresses significantly affect forest growth and productivity, with substantial economic consequences. For example, in China in 1989, damage to hybrid poplar plantations by common defoliators such as the poplar lopper (*Apochemia cineraria*) and the gypsy moth (*Lymantria dispar*) resulted in substantial (40%) stand loss (Hu *et al.*, 2001). Similarly, coniferous trees such as loblolly pine (*Pinus taeda*) are often damaged by the insect pests *Dendrolimus punctatus* and *Crypyothelea formosicola* (Tang and Tian, 2003), while white spruce is often negatively affected by defoliating insects such as the spruce budworm (*Choristoneura fumiferana*; Lachance *et al.*, 2007). In addition to defoliating insects, there are fungal, bacterial and viral pathogens that can affect forest health and productivity (Table 4-1). The following sections highlight the results of genetic modifications aimed at improving tree defences against damaging pests.

TABLE 4-1

**Summary of genetic modifications in trees, targeting resistance to various pathogens**

Gene	Modification	Effects	Reference
Bt <i>Cry3Aa</i>	Expressed in <i>E. coli</i>	Using <i>E. coli</i> as a preliminary system to test the effectiveness of a variety of Bt toxin, the protein caused high mortality in long-horned beetle larvae	Chen <i>et al.</i> , 2005
Bt <i>Cry1A</i>	Expressed in <i>Picea glauca</i> embryogenic callus	Low toxicity to spruce budworm when fed embryonic callus tissue	Ellis <i>et al.</i> , 1993
Synthetic Bt <i>Cry3Aa</i>	Expressed in hybrid poplar, <i>P. tremula</i> × <i>tremuloides</i>	Transgenic leaves toxic when fed to <i>Chrysomela tremulae</i> beetle	Genissel <i>et al.</i> , 2003
Bt <i>Cry1Ac</i>	Expressed in <i>Pinus radiata</i> , ubiquitin promoter (pr)	Some lines had higher resistance to painted apple moth; mature needles were more toxic compared with young needles	Grace <i>et al.</i> , 2005
Bt <i>Cry1Aa</i>	Expressed in <i>P. alba</i> × <i>grandidentata</i>	Gypsy moth fed more on mature leaves among transgenics	Kleiner <i>et al.</i> , 2003
Bt <i>Cry1Ab</i>	Expressed in <i>P. glauca</i> , ubiquitin pr.	Embryogenic, young somatic tissue and needles from 5 year field trial trees were toxic to spruce budworm	Lachance <i>et al.</i> , 2007
Synthetic <i>Cry1Ac</i>	Expressed in <i>Pinus taeda</i> , 35S pr.	64 to 75% mortality of <i>D. punctatus</i> Walker and <i>C. formosicola</i> Staud larvae after 7 days of feeding	Tang and Tian, 2003
Various forms of Bt toxin	Expressed in <i>P. nigra</i> , 2x35S pr.	Low expression of toxin, but high mortality of various pests and reduced damage to leaves on transgenic plants	Wang <i>et al.</i> , 1996
Bt toxin	Expressed in <i>P. alba</i> × <i>grandidentata</i> and <i>P. nigra</i> × <i>trichocarpa</i> , 35S pr.	Leaf feeding assay: high mortality to forest tent caterpillar and gypsy moth	McCown <i>et al.</i> , 1991
Polyphenol oxidase (PPO)	Overexpressed in <i>P. tremula</i> × <i>alba</i>	Increased levels of PPO in leaves, but no effect on feeding caterpillars	Barbehenn <i>et al.</i> , 2007
Kunitz trypsin proteinase inhibitor	Expressed in <i>P. nigra</i>	Feeding assays: no change to <i>Lymantria dispar</i> and <i>Clostera anastomosis</i> larval mortality or pupal weight	Confalonieri <i>et al.</i> , 1998
Cysteine proteinase inhibitor ( <i>Atcys</i> )	Overexpressed in <i>P. alba</i>	Higher mortality to <i>Chrysomela populi</i> beetle larvae when fed transgenic leaves	Delledonne <i>et al.</i> , 2001

Note: Bt = *Bacillus thuringiensis*

### Transgenic trees expressing Bt toxins

Insect pests are a major problem for poplar plantation managers. The two main classes of poplar pests are chrysomelid beetles and lepidopteran caterpillars, both of which are susceptible to microbial pesticides derived from different strains of *Bacillus thuringiensis* (Bt). This bacterium synthesizes polypeptides that are activated within the gut of certain insects, causing lesions and eventually insect death (Knowles and Dow, 1993). The insecticidal proteins, collectively referred to as Bt toxins, have been used safely as microbial pesticides in numerous crops (Carozzi and Koziel, 1997) for many years, both exogenously and endogenously. These toxins are relatively selective insecticides that have very few non-target effects (James, 1997). Several Bt strains have been identified, each affecting a select group of insects that are usually closely related phylogenetically (Thompson, Schnepf and Feitelson, 1995). Genetically modifying trees to produce forms of Bt toxin offers an appealing alternative for establishing plantations that are resistant to damage from a broad range of pests (Table 4-1).

Trees expressing Bt transgenes may be preferable to use of spray applications for several reasons. First, vegetation, soil and water surrounding the crop are not exposed to spray drift. Susceptible, non-target, insects in areas adjacent to the transgenic crop would not be exposed, reducing the potential for development of Bt resistance. Second, spray applications quickly degrade, persisting on leaves for, at most, a few days (Thompson, Schnepf and Feitelson, 1995; James, Croft and Strauss, 1999). Genetically engineered trees, however, can produce the toxin continuously, thereby avoiding sensitivities to application timing and the costs associated with repeated applications. Finally, because transgenic trees produce the toxin within plant tissues, it is possible to affect insects residing in the plant, such as wood borers and leaf folders. For some of these pests, no insecticides are currently available that target the life stage(s) most responsible for damage.

McCown *et al.* (1991) were the first to report on poplars that were stably transformed with a Bt toxin gene. One transgenic line in particular showed high levels of resistance to two pests: the forest tent caterpillar and the gypsy moth, as observed through leaf feeding experiments. Ellis *et al.* (1993) then described the first stable transformation of a conifer, white spruce (*Picea glauca*), with the *CryIA* gene (a gene coding for a form of Bt toxin). Spruce budworms fed a diet of the transgenic embryonic tissues showed few signs of toxicity. The lack of toxicity was attributed to low transgene expression in embryogenic tissues. More recently, transgenic Monterey pine (*Pinus radiata*) expressing a Bt toxin gene showed variable resistance to damage from painted apple moth larvae (*Teia anartoides*), depending on the age (maturity) of needles (Grace *et al.*, 2005). These studies emphasize the importance of transgene expression levels and tissue specificity.

One example of successful engineering of poplar trees to combat an insect pest is found in the cottonwood leaf beetle (CLB, *Chrysomela scripta* Fabricius), the primary insect pest in poplar plantations. The CLB is a multivoltine insect that has a wide distribution, which can culminate in outbreaks causing severe defoliation, particularly in young plantations (Hart *et al.*, 1996). James, Croft and Strauss

(1999) have demonstrated that a Cry3A Bt toxin is highly effective against the CLB. In this study, a binary vector containing a *Cry3A* gene under the control of the cauliflower mosaic virus (CaMV) 35S promoter was used to produce 51 insect-resistant lines in four genotypes of *Populus* sp. The transgenic trees were field-tested in eastern Washington State, United States of America. This trial relied on insect pressure from the surrounding, commercial stands to evaluate insect resistance. Trees were evaluated for damage, basal diameter and height at various stages during the growing season. Virtually all of the Bt transgenics showed very low feeding damage, whereas the non-transgenic lines sustained significantly higher levels of defoliation. Moreover, in most cases, the mean growth for transgenic lines was greater than that for the non-transgenic controls within each clone (Meilan *et al.*, 2000).

Others have explored the impacts of transgenic expression of synthetic *Cry1A* genes. Leaves of transgenic *Populus tremula* × *P. tremuloides* expressing a synthetic form of Bt toxin (*Cry3Aa*) proved to be highly effective in resisting damage by the phytophagous beetle *Chrysomela tremulae* (Genissel *et al.*, 2003). The synthetic Bt gene was modified to possess dicotyledonous codon usage and no AT-rich regions, and beetles feeding on high- and low-expressing *Cry3Aa*-leaves died within days. Synthetic *Cry1Ac* expressed in loblolly pine under the control of the 35S promoter resulted in a nearly eight-fold increase in toxicity to various insect larvae (e.g. *Dendrolimus punctatus* and *Crypythoelea formosicola*), in laboratory feeding experiments. Although verification of resistance in both short- and long-term field trials is needed, stable expression of synthetic Bt toxin appears to be an effective method to reduce damage and, hence, lost growth or mortality caused by various insect pests.

The potential for insects to develop resistance to genetically engineered crops is a major concern (DiCosty and Whalon, 1997; James, 1997; Roush and Shelton, 1997). Before insect-resistant transgenics can be commercialized, a resistance management plan must be developed. Many management strategies have been proposed based on prior experiences with pesticide resistance (e.g. Luttrell and Caprio, 1996; Roush, 1997; Gould, 1998; McGaughey, Gould and Gelernter, 1998). Combining resistance genes (pyramiding or stacking) is one way of reducing the risk of insects becoming resistant to Bt gene products (Nwanze *et al.*, 1995; Maredia and Mihm, 1997; Roush 1997). This approach has proven to be an effective strategy for resistance management with many insects, including the cotton bollworm (*Helicoverpa armigera*; Zhao *et al.*, 1997). However, in the United States of America, for example, in order to obtain approval from the United States Environmental Protection Agency (USEPA) to deploy trees containing a gene encoding a Bt toxin, additional studies of beetle dispersal will be needed.

### Non-Bt modifications

Despite the relief offered by Bt toxins, both native and synthetic, against damage caused by insect pests, other research has targeted herbivore resistance using different compounds. For example, Confalonieri *et al.* (1998) generated *P. nigra*

expressing a soybean trypsin proteinase inhibitor (Kunitz proteinase inhibitor, *KTi3*). Although the transgenic Kunitz protein inhibited digestive proteinases of the polyphagous moths *Lymantria dispar* and *Clostera anastomosis in vitro*, leaf feeding bio-assays showed no increase of larval mortality as a result of the transgenic expression. Perhaps increased expression would improve resistance, or a different proteinase inhibitor would be more detrimental to the digestion processes of larvae. In contrast, greater success was achieved in white poplar (*P. alba*) expressing an *Arabidopsis* cysteine proteinase inhibitor (*Atcys*), which resulted in up to 100% mortality of chrysomelid beetle (*Chrysomela populi*) larvae after only 16 days of feeding on transgenic leaf tissue (Delledonne *et al.*, 2001). Interestingly, expression of the scorpion neurotoxin, AaIT, in hybrid poplars appears to impart resistance against the gypsy moth (*Lymantria diaper*; Wu *et al.*, 2000).

### Fungal pathogens

Fungal infections can be equally damaging to forest trees. Genetic modifications using a variety of genes from several plants have been evaluated to improve fungal pathogen resistance, and have met with varying success. Expression of the bacterio-opsin gene in tobacco suggested that defence mechanisms would be elicited by its expression and pathogen resistance therefore increased (Mittler, Shulaev and Lam, 1995). However, the expression of a synthetic bacterio-opsin gene in hybrid poplars did not elicit a significant increase in defence-response against a variety of fungal pathogens, such as leaf rust, leaf and shoot blight, and stem canker (Mohamed *et al.*, 2001). Similarly, white poplar expressing grapevine stilbene synthase (*StSy*), which has been implicated in the production of resveratrol compounds, did not significantly affect the efficacy of resistance against a rust disease (*Melampsora pulcherrima*; Giorcelli *et al.*, 2004). Although these transgenic trees offered an interesting opportunity to produce pharmacologically valuable compounds, they were not a viable option to reduce loss due to rust (Giorcelli *et al.*, 2004). In contrast, transgenic poplars expressing a rabbit defensin gene (*NP-1*; Zhao *et al.*, 1999) or a chitinase (*CH5B*) appear to have increased resistance to a broad spectrum of fungal pathogens (Meng *et al.*, 2004). Hybrid poplars expressing a wheat germin-like oxalate oxidase gene, directed at metabolizing the oxalic acid produced by fungal pathogens, showed signs of delayed infection by *Davidiella populorum* (syn. *Septoria musiva*) (Liang *et al.*, 2001). Liang *et al.* (2002) have also investigated the ability of transgenic poplars expressing antimicrobial peptides to alter resistance against *Davidiella populorum*. Two-year-old transgenic trees expressing a combination of antimicrobial peptides did, indeed, show greater resistance in leaf disc assays, and initial field trials showed less frequent *Davidiella* cankers on transgenic trees (unpublished, but reported in Powell *et al.*, 2006). It is evident that increasing resistance against fungal pathogens requires the use of a variety, and perhaps a combination, of transgenic products.

### Bacterial pathogens

Reports of genetic modifications resulting in increased resistance to bacterial pathogens are less common. Although bacterial damage occurs less frequently, serious infections of *Xanthomonas* spp. have been reported (Haworth and Spiers, 1988; De Kam, 1984). Transgenic poplar expressing the antimicrobial peptide, D4E1, showed mixed resistance against *Agrobacterium* and *Xanthomonas* infection (Mentag *et al.*, 2003). In particular, the transgenic line displaying the highest transgene transcript abundance showed a significant increase in resistance, as defined by reduced tumour formation after *Agrobacterium* inoculation or the development of smaller cankers following *Xanthomonas* infection. However, D4E1-transformants did not show improved resistance against fungal pathogens. It should be noted that resistance against one strain of *Agrobacterium*, C58, was not improved; therefore, D4E1 appears to have limited specificity (Mentag *et al.*, 2003).

### Field trials

The value of transgenic trees will only be realized after the completion of many extensive field trials. In the area of pest and pathogen resistance, a number of field trials have been reported, and some trials have yielded contrasting findings. In one case, resistance appeared to be lower in the field than in laboratory tests, and resistance levels can vary depending on the tissue tested. For example, a three-year field trial of transgenic birch expressing sugar beet chitinase revealed that although plants showed greater resistance in greenhouse trials, in the field the birch were equally, if not more, susceptible to fungal diseases such as leaf spot, caused by *Pyrenopeziza betulicola* (Pasonen *et al.*, 2004). In contrast, Hu *et al.* (2001) conducted a three-year trial of Bt-transgenic *P. nigra* and showed a decrease in damage from defoliators: 10% leaf damage compared to 80 to 90% damage on control plants. This study had other significant implications, as it demonstrated that there was a concurrent decrease in the abundance of insect pupae in the soil on transgenic plots, and that non-transgenic, wild-type trees were more protected when grown near or amongst transgenics. In conifers, although Cry1Ab levels in needles from field-grown trees were lower than levels in embryonic tissues or somatic seedlings, Bt-transgenic spruce had improved resistance against spruce budworm (Lachance *et al.*, 2007). Mortality of larvae feeding on tissues from field-tested plants ranged from 44 to 100% for transgenic trees, compared with approximately 37% for controls. Five-year-old trees also appeared phenotypically normal. This study illustrates the inherent variation of genetic modifications, and emphasizes the need for long-term trials. Efforts to understand the broad changes caused by genetic modifications were also evaluated by Davis *et al.* (2006), who looked at the effects of a Bt transgene on wood properties in hybrid poplars, and found no significant difference in chemical composition.

### Considerations

As with chemical sprays, it is extremely important to consider the potential for pests to become resistant to the transgene product. Although laboratory

experiments suggested that feeding on transgenic Bt tissues enhanced the development of Bt-resistant pests, field tests report no increase in Bt-resistant pests either on or near Bt crops (Tabashnik *et al.*, 2003). Natural variation present in the field may prevent the rapid evolution of resistant pests. Results from Hu *et al.* (2001) demonstrate the value of planting stands mixed with both non-transgenic and transgenic trees. Genetic modification in other non-forest trees may be relevant when assessing the long-term effects of field-grown, genetically modified trees. One such example is the production of virus-resistant papaya, which was genetically modified to resist infection by the papaya ring-spot virus (Lius *et al.*, 1997). The genetically modified papaya trees have been grown commercially since the mid-1990s, and continue to be of benefit in Hawaii (NASS, 2005). However, the movement of transgenes from engineered trees to other organisms is still of concern (Fuchs and Gonsalves, 2007).

New research has highlighted the production of antibodies within transgenic plant cells, which could aid in combating infection from various pathogens (Powell *et al.*, 2006). For example, the expression of a recombinant antibody in citrus trees may reduce the infectious nature of pathogen proteins and thereby reduce disease progression. Such new technologies will surely change the direction of some research programmes, and continued efforts to understand the genes involved in plant defence mechanisms will lead to new avenues to improve resistance (e.g. Ralph *et al.*, 2006). Although commercial deployment of transgenic trees is several years away, using genetic modifications to increase tree survival, reduce the impact of chemical sprays to the environment and increase economic value of forests has significant potential.

## NUTRITION

Forest nutrition is a key factor affecting tree growth. Nitrogen and sulphur are two elements that are essential for normal growth and development, and are often in short supply. With afforestation and reforestation efforts increasingly occurring on marginal agricultural land, tree nutrition is a key target area for genetic improvements.

### Nitrogen

Nitrogen availability is a common limiting factor in forest tree growth (Suárez *et al.*, 2002). Development depends not only on the inorganic nitrogen available in the soil, but also on recycling within the plant, particularly in situations with limited nitrogen. Glutamine synthetase (GS) plays a significant role in both nitrogen uptake and recycling, as it catalyses the incorporation of ammonium into glutamine, the precursor to glutamate. Glutamine is also the precursor for all other plant N-containing compounds (Mifflin and Lea, 1980). In an attempt to alter tree growth, GS has been an important target for genetic engineering (Fu *et al.*, 2003).

There are two iso-enzymes of GS, one localized in the cytosol (GS1) and the other in the chloroplast (GS2; Gallardo *et al.*, 1999). In angiosperms, GS2 is thought to function in the assimilation of ammonium from nitrate and respiration,



while GS1 has been suggested to be involved in glutamine generation for transport within the plant (Lam *et al.*, 1996). In conifers, GS2 has not been identified; GS1 is expressed in the photosynthetic cells and has been proposed to be involved in the primary assimilation in roots and re-assimilation from other metabolic processes (Jing *et al.*, 2004). GS1 has been found to co-localize with QTLs associated with yield (Hirel *et al.*, 2001; Obara *et al.*, 2001).

Suárez *et al.* (2002) overexpressed pine GS1 in poplar, and the plants were shown to form GS protein that is different from the native version, and elevated levels of chlorophyll (Gallardo *et al.*, 1999). Transgenic plants grew significantly faster than non-transformed controls. The transgenic plants also show increased early vegetative growth, increased leaf area, and a greater number of internodes (Fu *et al.*, 2003). GS activity was increased by 66%, chlorophyll by 33% and protein content by 21%. The results suggest that GS activity in young leaves is an effective marker for vegetative development (Fu *et al.*, 2003), even under low-nitrogen conditions. GS activity was strongly correlated with height growth, more so than chlorophyll or protein content. It is possible that the GS expression could affect additional pathways involved in vegetative growth other than through enhanced nitrogen, as was seen in tobacco, where changes in photosynthetic and photorespiratory capacities resulted in improved growth (Fuentes *et al.*, 2001).

These same poplars trees were field-tested for three years. The plants were again shown to be taller than the corresponding controls and had increased protein, total GS activity/protein and ferredoxin-dependent glutamate synthase (Fd-GOGAT), but showed no change in the Rubisco large subunit or in water content (Jing *et al.*, 2004). No significant differences were seen in polysaccharide or lignin content in the stems, but stem diameter and bark protein content suggest that nitrogen reserves accumulated to a greater extent in the stems of transgenics (Jing *et al.*, 2004). This increased growth and nitrogen cycling in poplars with ectopic pine cytosolic GS expression may be of particular importance given that marginal lands are being increasingly reclaimed for tree plantations, a trend that is likely to increase with the demand for bio-energy.

## Sulphur

Sulphur is an essential element found mostly in its reduced form as the amino acids cysteine and methionine. In plants, cysteine is used either in the synthesis of proteins, or can be further metabolized to methionine, glutathione (GSH) and phytochelatins. GSH plays several crucial roles in plants, including acting as an antioxidant, protecting against reactive oxygen species (Foyer *et al.*, 1995); as a substrate for glutathione S-transferases, enabling detoxification of xenobiotics (Marrs, 1996); as a precursor in the synthesis of phytochelatins, which participate in the detoxification of heavy metals (Cobbett, 2000); in the regulation of gene expression (Wingate, Lawton and Lamb, 1988); and in the storage and transport of reduced sulphur (Herschbach, Jouanin and Rennenberg, 1998).

GSH is synthesized in two steps. Initially,  $\gamma$ -glutamylcysteine synthetase ( $\gamma$ -ECS) catalyses the fusion of cysteine to glutamate, producing  $\gamma$ -glutamylcysteine,

which then reacts with glycine in the synthesis of GSH via GSH synthetase (GSS; Herschbach and Kopriva, 2002). There are three factors that control the rate of GSH synthesis in leaves: the availability of cysteine (Stroh *et al.*, 1995), feedback inhibition of  $\gamma$ -ECS by GSH (Schneider and Bergmann, 1995) and the availability of the enzyme  $\gamma$ -ECS (Fargo and Brunold, 1994). Glycine may also be a limiting factor during prolonged darkness, as its synthesis is affected by photorespiration (Noctor *et al.*, 1999).

Given the biological significance of sulphur in plant development, it has been a key target for genetic engineering in trees, particularly GSH formation. Numerous strategies have been used in designing transgenics over the years. Originally,  $\gamma$ -ECS and GSS were up-regulated in the cytosol (Foyer *et al.*, 1995; Stroh *et al.*, 1995; Noctor *et al.*, 1996; Arisi *et al.*, 1997). Overexpression of GSS did not yield an increase in GSH, despite the supplemental supply of cysteine. In subsequent feeding studies, an exogenous supply of  $\gamma$ -EC overcame this limitation, leading to increases in GSH, suggesting that  $\gamma$ -ECS is the rate-limiting step (Stroh *et al.*, 1995). Overexpression of  $\gamma$ -ECS resulted in an increase in GSH without affecting the redox state, as well as a significant increase in  $\gamma$ -EC (Stroh *et al.*, 1995; Noctor *et al.*, 1996). An exogenous supply of cysteine resulted in further increases in GSH. Similar results were observed in transgenic tobacco (Creissen *et al.*, 1996); however, in tobacco there was increased sensitivity to oxidative stress, and necrotic lesions formed.

Additional studies revealed an increase in GSH in the phloem of poplars overexpressing  $\gamma$ -ECS, confirming its role as a major transport form of sulphur (Herschbach, Jouanin and Rennenberg, 1998). Furthermore, overexpression of  $\gamma$ -ECS in the chloroplast, using the 35S promoter along with the pea *rbcS* chloroplast transit peptide or *rbcS* itself, resulted in increased concentrations of foliar valine, leucine, isoleucine, tyrosine and lysine. These findings suggest that there is an additional indirect effect on nitrogen metabolism (Noctor *et al.*, 1998).

Similarly, the overexpression of glutathione reductase (GR) in the chloroplasts of poplar led to increased GSH pools (Foyer *et al.*, 1995). Although there were no quantifiable changes in photosynthetic rates under normal growing conditions, when photo-inhibition was induced by subjecting the transgenics to low temperatures and high light levels, the plants proved to be much less sensitive to stress compared with the corresponding controls. The improved stress tolerance was attributed to a number of factors, including: increased cellular recycling of GSH and ascorbate pools, helping to combat the production of harmful free radicals; and elevated GSH levels that may aid in stabilizing enzymes that require reduced thiol groups for activity, and could also exert indirect actions on protein synthesis and gene expression (Foyer *et al.*, 1995).

The significance of these findings extends beyond the bounds of general plant nutrition, as sulphur is a key component of compounds involved in stress and herbicide resistance. The implications of alteration of sulphur nutrition are discussed further below.

### Herbicide resistance

Generally, there are three main breeding and improvement objectives, including improved growth rates, improved wood characteristics, and improved resistances to herbicides, pests and disease (Chupeau, Pautot and Chupeau, 1994). Improvements in herbicide resistance would allow for lower total herbicide use as well as application of more environmentally benign active ingredients, not to mention more flexibility with regard to the timing of application.

### Glyphosate

Fillatti *et al.* (1987a, b) reported the first successful insertion of a herbicide-tolerance gene in trees, generating transgenic poplar (*P. alba* × *P. grandidentata*) expressing the *Salmonella typhimurium aroA* gene. A mutation in this gene resulted in the formation of a 5-enolpyruvyl-3-phosphoshikimate synthase (EPSPS) that is resistant to inhibition by glyphosate, the active ingredient in Roundup® herbicide (Comai, Sen and Stalker, 1983). Ectopic expression of *aroA* under the mannopine synthase promoter yielded surprisingly low levels of herbicide tolerance (Riemenschneider *et al.*, 1988), which was thought to be the result of low cytosolic expression of the gene. When the same poplar was transformed with *aroA* under the control of the 35S promoter and a chloroplast transit peptide, the resulting plants showed higher levels of glyphosate tolerance (Riemenschneider and Haissig, 1991; Donahue *et al.*, 1994).

Although use of the 35S promoter led to higher expression levels than mannopine synthase, and the transit peptide directed transgene product to the chloroplast, performance of these *aroA*-containing lines still fell short of expectations (Karnosky *et al.*, 1997). Following greenhouse spray tests, chlorophyll content in all transgenic lines was inversely correlated to glyphosate concentration, height growth was arrested following herbicide treatment, and only one line retained live leaves six weeks after treatment (Donahue *et al.*, 1994). It was clear from this work that the production of a herbicide-tolerant poplar line would not be as straightforward as originally thought.

More recently, Meilan *et al.* (2002a) tested a construct containing a glyphosate-tolerance gene, *CP4*, for its ability to impart glyphosate tolerance in poplar. *CP4* is an alternative form of EPSPS that originates from *Agrobacterium tumefaciens* strain CP4, and glyphosate has a low affinity for this form of the enzyme. Using an *Agrobacterium*-mediated transformation protocol (Leple *et al.*, 1992; Han *et al.*, 2000), Meilan *et al.* (2000) generated transgenic plants in 12 genotypes of hybrid poplar. The resultant transgenic plants were field-tested for two years on the west and east sides of the Cascade Mountains in Oregon, United States of America. Growth of the lines expressing *CP4* was significantly better than controls and lines expressing both the *CP4* and *GOX*, another gene implicated in herbicide tolerance. In addition, the lines expressing *CP4* only exhibited less damage in response to glyphosate treatment. This was the first report of transgenic poplars exhibiting high levels of glyphosate tolerance when grown under field conditions

(Meilan *et al.*, 2002b). Herbicide resistance has remained stable for over eight years in trees grown under field conditions (Li *et al.*, 2008).

### Chlorsulphuron

Chlorsulphuron [chlorsulfuron] is a sulphonylurea herbicide that acts on acetolactate synthase and blocks the biosynthesis of valine and isoleucine (Ray, 1984). A mutant acetolactate synthase gene (*crs1-1*) from *Arabidopsis*, which confers resistance to chlorsulphuron, has been expressed in hybrid poplar (*Populus tremula* × *P. alba*) under the control of both the native and 2×35S promoter. Both promoters led to transgenic lines that were completely resistant to higher-than-normal field application rates of chlorsulphuron in greenhouse tests. Control poplar trees died within two to three weeks of treatment, whereas transgenic lines survived. Following slightly delayed growth and root development, the transgenic plants returned to normal growth following the treatment (Brasileiro *et al.*, 1992).

### Chloroacetanilide

Acetochlor and metochlor are active ingredients in chloroacetanilide herbicides, which are detoxified by GSH-dependent reactions (Gullner, Komives and Rennenberg, 2001). GSH and the glutathione S-transferase (GST) family play crucial roles in the degradation of several herbicides. GSTs are able to catalyse conjugation reactions between a number of xenobiotics and GSH. Herbicide:GSH conjugates are less toxic and more water soluble than the herbicide molecules alone (Edwards, Dixon and Walbot, 2000). When poplars expressing  $\gamma$ -ECS in chloroplast or cytosol were exposed to acetochlor and metochlor dispersed in soil (Gullner, Komives and Rennenberg, 2001), the growth and biomass of all lines was markedly reduced, but the reduction was less dramatic in the transgenic lines relative to the non-transformed trees, and the growth rate of cytosol expressers was less affected than the chloroplast expressers. Foliar  $\gamma$ -EC and GSH levels increased in all lines, but more so in the transgenics poplar (Gullner, Komives and Rennenberg, 2001).

### Glufosinate

Glufosinate (phosphinothricin, PPT) is the active ingredient in herbicides known as Basta<sup>®</sup> and Buster<sup>®</sup>, and is a structural analogue of glutamate. It inhibits glutamine synthetase, causing ammonium to accumulate, which at elevated concentrations is lethal (Bishop-Hurley *et al.*, 2001). This interaction causes irreversible inactivation of GS, which also blocks photorespiration, resulting in the depletion of leaf amino acid pools (Pascual *et al.*, 2008). Plants respond to PPT by developing necrosis, usually initiating at or near the apical meristem and spreading throughout the plant (Pascual *et al.*, 2008).

The *BAR* gene encodes for phosphinothricin acetyltransferase (PAT), which inactivates glufosinate by acetylating its free ammonium group (Thompson *et al.*,

TABLE 4-2  
Summary of genetic modifications in trees, targeting resistance to herbicides

Gene	Modification	Effects	Reference
<i>aroA</i> (EPSP gene)	Overexpression in <i>Populus alba</i> × <i>P. grandidentata</i>	First record of insertion and expression of a foreign gene of agronomic importance in woody plants; slight resistance to glyphosate	Fillatti <i>et al.</i> , 1987
<i>bar</i> (PAT gene)	Overexpressed in <i>P. alba</i> × <i>P. tremula</i> and <i>P. trichocarpa</i> × <i>deltoides</i>	Transgenics did not accumulate NH <sub>4</sub> <sup>+</sup> when treated with Basta®	De Block, 1990
<i>crs 1-1</i>	Overexpressed in <i>P. tremula</i> × <i>P. alba</i>	Resistance to chlorsulphuron in greenhouse tests	Brasileiro <i>et al.</i> , 1992
<i>aroA</i>	Overexpressed in <i>P. alba</i> × <i>P. grandidentata</i>	Conferred resistance to glyphosate	Donahue <i>et al.</i> , 1994
<i>als, pat</i>	Overexpressed in <i>P. tremula</i> × <i>P. alba</i>	Conferred herbicide resistance to calli	Chupeau, Pautot and Chupeau, 1994
<i>aroA</i>	Overexpressed in <i>Larix decidua</i>	Conferred resistance to glyphosate at moderate treatment levels	Shin <i>et al.</i> , 1994
<i>bar</i>	Overexpressed in <i>Eucalyptus camaldulensis</i>	Conferred resistance to herbicide at over twice the normal field application rate	Harcourt <i>et al.</i> , 2000
<i>bar</i>	Overexpressed in <i>Pinus radiata</i> and <i>Picea abies</i>	Conferred resistance to glufosinate in both species in greenhouse testing	Bishop-Hurley <i>et al.</i> , 2001
γ-ECS	Overexpression in <i>Populus tremula</i> × <i>P. alba</i>	Conferred resistance to acetochlor and metolachlor present in soil. Plants with cytosolic expression were more resistant than those with chloroplastic expression	Gullner, Komives and Renneberg, 2001
GOX, <i>CP4</i>	Overexpressed in various poplar hybrids, including <i>P. trichocarpa</i> × <i>P. deltoides</i>	Genes shown to confer resistance up to 66% at low glyphosate levels, with 25% of lines showing increased growth following herbicide treatment. Lack of damage was attributed to <i>CP4</i> , as GOX was suspected to cause undesirable side effects. Twelve lines expressing only <i>cp4</i> had similar herbicide tolerance, but grew better and had less damage in response to treatment	Meilan <i>et al.</i> , 2002a Meilan <i>et al.</i> , 2002b
γ-ECS	Overexpressed in <i>P. tremula</i> × <i>P. alba</i>	No change in tolerance to paraquat	Bittsanszky <i>et al.</i> , 2006
GS	Overexpressed in <i>P. tremula</i> × <i>P. alba</i>	Conferred resistance to PPT	Pascual <i>et al.</i> , 2008
<i>bar</i>	Overexpressed in <i>P. alba</i>	Conferred resistance to Basta® at normal field dosage; poplar still tolerant at twice the normal field dosage	Confalonieri <i>et al.</i> , 2000

1987). The effectiveness of *BAR* in conferring herbicide resistance has been shown in many tree species. Poplar explants transformed with *BAR* were able to survive and grow on glufosinate-containing medium at the callus phase, proving it to be an effective selectable marker gene (Chupeau, Pautot and Chupeau, 1994). Similar experiments have been carried out in other species (see Table 4-2).

Poplar overexpressing the pine GS gene showed increased resistance to PPT. Resistance was measured at 5, 25 and 100 µM PPT. At 5 µM, there was limited effect on all plants, while at 25 µM, 75% of control plants died and 50–100% of the transgenic plants in each line remained viable. At 100 µM all of the wild-type trees were dead within eight days, whereas 20–45% of the transgenics survived (Pascual *et al.*, 2008).

### Abiotic stress

Environmental stresses can significantly affect productivity. Low temperatures and high salt concentrations during the growing season can damage seedlings, leading

to impaired growth or even death (Blennow and Lindkvist, 2000). Because certain plants and bacteria are able to survive these harsh conditions, genomic tools have been used to identify target genes within the protective pathways and impart stress tolerance through genetic modification (Cushman and Bohnert, 2000).

Increased resistance to many types of stress has already been achieved for several plant species. Expression of anti-freeze or ice nucleation genes has been shown to improve freeze tolerance in tobacco and maize protoplasts (Baertlein *et al.*, 1992; Murata *et al.*, 1992; Georges, Saleem and Cutler, 1990). Two antifreeze genes, glucose-6-phosphate dehydrogenase (PsG6PDH) and anti-freeze protein (PsAFP), have been introduced into poplar and freeze-resistance tests are currently underway (Lin and Zhang, 2004).

Control mechanisms for abiotic stresses are based on the activation or regulation of specific stress-related genes, which can be involved in controlling transcription, cell signalling, protecting membranes and proteins, or scavenging free-radicals, or a combination (Wang, Vinocur and Altman, 2003). Several different abiotic stresses often activate similar signalling pathways and cellular responses (Knight and Knight, 2001; Zhu, 2002) and therefore result in similar plant phenotypes. As such, transcription factors hold particular interest because of their potential to increase tolerance to multiple stresses via the overexpression of a single transcription factor. An ERF/AP2-type transcription factor has been of particular interest because when overexpressed in *Arabidopsis*, it caused the simultaneous up-regulation of pathogen- and cold-response (*COR*) genes (Yi *et al.*, 2004). Similarly, the overexpression of the dehydration response element 1A (*DREB1A*) transcription factor in *Arabidopsis* caused increased tolerance to multiple abiotic stresses by the increased *COR* expression (Kasuga *et al.*, 1999; Jaglo-Ottosen *et al.*, 1998). Gains in the tolerance of abiotic stress have also been seen in forest trees.

### Ozone stress

Ozone is formed by photochemical reactions between nitrogen oxides, hydrocarbons and carbon monoxide, and is highly phytotoxic (Lelieveld and Crutzen, 1990). At elevated concentrations it elicits changes in plant biochemical and physiological processes, resulting in foliar injury, increased senescence, decreased growth rates (Kress and Skelly, 1982; Sandermann, 1996) and increased production of reactive oxygen species (ROS; Foyer *et al.*, 1994).

The ascorbate-glutathione pathway plays an important role in protecting plants from ROS (Foyer *et al.*, 1994). GSH acts as an antioxidant that can directly scavenge ROS, and also protect thiol-containing enzymes and reduce dehydroascorbate, as it is oxidized to glutathione disulfide (GSSG). The reduced GSH pool is maintained by the activity of GR, and many plant species have exhibited gains in resistance to photo-oxidative stress, herbicides or drought, or a combination, through the up-regulation of GR or superoxide dismutase (Foyer *et al.*, 1994).

When transgenic poplar overexpressing GSS in the cytosol or GR in the cytosol or chloroplast were exposed to various levels of ozone stress, there was no



apparent difference in the phenotypic response between the transgenic and control trees, despite elevated activities of GSS and GR (Strohm *et al.*, 1995). Rather, sensitivity to ozone stress appeared to be directly related to leaf developmental stage (Strohm *et al.*, 1995). Poplar with GR overexpressed in the chloroplast did, however, recover more quickly when exposed to high light levels and low temperatures than did wild-type trees, or trees expressing GR or GSS in the cytosol (Foyer *et al.*, 1995; Strohm *et al.*, 1995).

Plants with increased peroxidase activity also display increased resistance to abiotic stress (Hiriga *et al.*, 2001). Hybrid aspen (*P. sieboldii* × *P. grandidentata*) overexpressing the horseradish peroxidase gene (*prxC1a*) showed increased growth and elevated peroxidase activity. Transgenic callus tissue and plantlets were resistant to oxidative stress imposed by hydrogen peroxide, although the growth rate was decreased (Kawaoka *et al.*, 2003).

### Salt stress

Salt stress is an increasingly important issue throughout the world, and it is imposed by two factors: water deficit due to osmotic stress, and the accumulation of ions that negatively affect biochemical processes (Tang, Charles and Newton, 2005). A number of genes have been tested in attempts to increase salt tolerance in trees. Poplar transformed with the *E. coli* mannitol-1-phosphate dehydrogenase gene (*mt1D*) grew faster and had a higher survival rate than non-transformed controls (Liu *et al.*, 2000). Hu *et al.* (2005) found that the up-regulation of *mt1D* in poplar led to increased mannitol levels, and, under salt stress, all lines had higher stomatal conductance, transpiration and photosynthetic rates. Under non-salt-stressed conditions, transgenic plant growth was about 50% that of controls (Hu *et al.*, 2005). Other species have also demonstrated gains in salt tolerance when transformed with *mt1D*, glucitol-6-phosphate dehydrogenase (*gutD*), choline dehydrogenase (*betA*) and choline oxidase (*codA*) genes (see Table 4-3).

### Drought stress

Drought stress is primarily osmotic stress, which causes the disruption of homeostasis and ion distribution in the cell (Serrano *et al.*, 1999; Zhu, 2001). Poplar transformed with a pine cytosolic GS (GS1) was shown to be more tolerant to drought stress than wild-type trees (El-Khatib *et al.*, 2004). At all levels of water availability, the transgenic trees had higher photosynthetic assimilation rates and stomatal conductance than the corresponding controls. All GS1-containing lines also showed an irreversible decline in photosystem II (PSII) antennae transfer efficiency after drought and during recovery, but the increased photo-assimilation capacity of the transgenic poplar allowed more resources to be allocated to photoprotective mechanisms. Gains in drought stress were also reported in hybrid eucalypts (*Eucalyptus grandis* × *E. urophylla*) transformed with the DREB1A transcription factor (Kawazu, 2004).

Efforts have been made to engineer tolerance to multiple stresses. Hybrid larch (*Larix* × *leptoeuropaea*) expressing a pyrroline 5-carboxylate synthase gene

TABLE 4-3

## Summary of genetic modifications in trees, targeting resistance to abiotic stress

Gene	Modification	Effects	Reference
<i>GSS</i>	Overexpression in <i>Populus tremula</i> × <i>alba</i>	No changes in response to ozone stress; ozone sensitivity related to leaf development stage	Strohm <i>et al.</i> , 1995
<i>FeSOD</i>	Overexpression in <i>Populus tremula</i> × <i>alba</i>	No change in response to high light and photo-inhibition of PSII; suggests rate of conversion of superoxide to hydrogen peroxide is not a rate limiting factor in protection against or repair of photo-inhibition	Tyystjarvi, 1999
<i>mtlD</i>	Overexpression in <i>Populus</i> sp.	Transgenic plants grew significantly better with a higher survival rate	Liu <i>et al.</i> , 2000; Liu <i>et al.</i> , 2002
<i>Bet-A</i>	Overexpression in <i>Populus</i> sp.	Conferred salt resistance	Yang <i>et al.</i> , 2001
<i>Phospholipase D</i>	Antisense expression in <i>Populus</i> sp.	Conferred salt tolerance	Liu <i>et al.</i> , 2002
<i>PsG6PDH</i> and <i>PsAFP</i>	Overexpression in <i>Populus</i> sp.	Freezing resistance tests under way	Lin and Zhang, 2004
<i>mt1D</i> and <i>gutD</i>	Overexpression in <i>Pinus taeda</i>	Increased salt tolerance at both calli and plantlet stage; accumulated mannitol and glucitol	Tang, Peng and Newton, 2005
<i>codA</i>	Overexpressed in <i>Eucalyptus camaldulensis</i>	Increased salt stress tolerance	Yamada-Watanabe, Kawaoka and Matsunaga, 2003
<i>mt1D</i>	Overexpression in <i>Populus tomentosa</i>	Increased mannitol; increased salt tolerance <i>in vitro</i> and in hydroponic culture; decreased growth in the absence of salt	Hu <i>et al.</i> , 2005
<i>prxC1a</i>	Overexpression in <i>P. sieboldii</i> × <i>P. grandidentata</i>	Increased growth rate; elevated peroxidase activities; calli resistant to oxidative stress imposed by hydrogen peroxide	Kawaoka <i>et al.</i> , 2003
<i>vhb</i>	Overexpression in <i>P. alba</i>	No change in growth pattern, or chlorophyll and protein contents; no change in stress resistance	Zelasco <i>et al.</i> , 2006
<i>dreb1a</i> and citrate synthase	Overexpression in <i>Eucalyptus grandis</i> × <i>E. urophylla</i>	Conferred resistance to drought and acid soil tolerance	Kawazu, 2004
<i>GS1</i>	Overexpression in <i>P. tremula</i> × <i>P. alba</i>	Higher photosynthetic assimilation and stomatal conductance at all levels of water availability; increased photo-assimilation allows increased allocation of resources to photoprotective mechanisms	El-Khatib <i>et al.</i> , 2004
<i>P5CS</i>	Overexpression in <i>Larix leptoeuropaea</i>	Increased proline; increased resistance to cold, salt and freezing stress	Gleeson, Lelu-Walter and Parkinson, 2005
<i>CaPF1</i>	Overexpressed in <i>Pinus strobus</i>	Increase in tolerance to drought, freezing and salt stress	Tang, Newton and Weidner, 2007

(*P5CS*), which functions as a rate-limiting step in proline synthesis, had increased proline content and were shown to be more resistant to cold, salt and freezing stresses (Gleeson, Lelu-Walter and Parkinson, 2005). Proline, which is produced via the glutamic acid pathway (Delauney and Verma, 1993; Kavi Kishor *et al.*, 1995), has been shown to play a role in protecting trees against stresses (Gleeson, Lelu-Walter and Parkinson, 2004). When *P5CS* was up-regulated in tobacco, rice, lettuce and wheat, increased proline and biomass production were observed, despite environmental stresses (Kavi Kishor *et al.*, 1995; Zhu *et al.*, 1998; Pileggi *et al.*, 2001; Sawahel and Hassan, 2002). Similarly, Tang *et al.* (2007a) overexpressed

paprika (*Capsicum annuum*) pathogen and freezing tolerance-related protein 1 (CaPF1) in eastern white pine (*Pinus strobus*). The result was a dramatic increase in tolerance to drought, freezing, and salt stress. This was related to polyamine biosynthesis, as putrescine, spermidine and spermine levels were maintained in the transgenic lines, while they decreased in stress-treated controls (Tang *et al.*, 2007).

### Phytoremediation

The use of plants to remove contaminants from the environment is known as phytoremediation (Schnoor *et al.*, 1995). This technology has recently been applied to several environmental problems, including disposal of municipal wastewater, biofiltration of farm and industrial runoff, and the remediation of soils spoiled by industrial processes (Che *et al.*, 2003, 2006; Lee, Isenhardt and Schultz, 2003; Strand *et al.*, 2005). Because this technology is less costly, less invasive, more aesthetic, and often yields a usable product (e.g. biomass), it has many advantages over traditional, engineering-based methods. Phytoremediation plantings can provide additional environment benefits, such as a means to sequester carbon (i.e. carbon credits), erosion control, wildlife habitat maintenance and the creation of buffers against noise, garbage and harmful dust (Rockwood *et al.*, 2004).

Using transgenic plants for improved phytoremediation is a relatively new, but highly successful, technological advance. For example, transgenic plants have been developed that are more effective in translocating arsenic (Dhankher *et al.*, 2002) and selenium (LeDuc *et al.*, 2006) from soil to the plant. Similarly, transgenics have been generated that are more tolerant of and better able to degrade explosives (French *et al.*, 1999; Hannink *et al.*, 2001), and that can detoxify sites contaminated with mercury (Bizily, Rugh and Meagher, 2000; Meagher, 2000).

Heavy metal contamination is a major problem globally. Anthropogenic uses of mercury, zinc, cadmium, selenium, lead and arsenic have led to problems in many terrestrial and aquatic ecosystems. These harmful pollutants are generated from numerous sources, and although many have recently been the target of stricter regulation, historically, contaminated areas have not been reclaimed due to extremely high cost and the destructive nature of available methods. Many metallic compounds can be taken up by plants or microbes and thus enter the food chain, potentially causing significant problems for animals and humans. Although the concept of using plants to remediate contaminated sites is not new (Baker and Brooks, 1989), heavy metals are multisite inhibitors of several metabolic pathways and are therefore generally phytotoxic. Genetic manipulation can overcome these obstacles. Several plant species have been considered for phytoremediation efforts, but trees have most recently been identified as particularly useful vehicles because they produce large amounts of biomass, have far-reaching roots and are perennial, although leaves may need to be collected for incineration (Bittsanszky *et al.*, 2005). There is the added benefit of employing the woody biomass for fuel or value-added products. Despite the development of successful transformation techniques for numerous tree species, research in dendroremediation is relatively new, but results are consistent with what has been seen in other plant species.

The volatilization of mercury by tobacco and *Arabidopsis* transformed with the mercuric ion reductase (*mer*) genes from bacteria inhabiting contaminated sites has been particularly successful (Summers, 1986). The *mer* gene catalyses the conversion of ionic mercury Hg(II) to its volatile derivative Hg(0). Yellow-poplar (*Liriodendron tulipifera*) was transformed with *merA18* from *E. coli* (Rugh *et al.*, 1998), and the resulting plantlets grew vigorously on a medium containing levels of mercury that were roughly ten fold those known to be toxic. They also released elemental mercury at ten times the rate of wild-type trees over a six-day trial in mercury vapour sampling tubes, with no apparent effect on growth.

Eastern cottonwood (*Populus deltoides*) has been identified as a key dendroremediant, as its native growth habitat includes riparian areas, which are similar to many contaminated sites. In 2003, Che *et al.* transformed eastern cottonwood with the *merA9* and *merA18* genes from *E. coli*. The transgenic trees grew normally and rooted in medium containing 25  $\mu$ M Hg(II), while wild-type trees did not survive. When exposed to lower levels of Hg(II), which were not lethal to wild-type trees, the transgenic lines emitted two- to four-fold more Hg(0) than the corresponding wild-type trees. Transgenic plants (*merA18*) also accumulated significantly higher biomass than wild-type trees when grown in soil containing 40 ppm Hg(II). However, in soils with lower levels of contamination, there was no difference in biomass accumulation, and in soil that was not contaminated, wild-type trees grew faster (Che *et al.*, 2003).

More recently, eastern cottonwood harbouring both *merA* and *merB*, which encode for organomercury lyases, were produced for use in mercury-contaminated soils (Lyyra *et al.*, 2007). *In vitro*-grown plants were highly resistant to phenylmercuric acetate, and were able to detoxify organic mercury compounds at two to three times the rate of controls or trees containing one of the two *mer* genes. Only trees transformed with both genes were capable of rooting in media supplemented with mercury, although their roots were shorter and thicker than in mercury-free media. The plants expressing both transgenes probably convert mercury first to Hg<sup>2+</sup> and then to elemental mercury, with lower toxicity in a coupled reaction, as was previously shown in both *Arabidopsis* and tobacco (Bizily, Rugh and Meagher, 2000; Ruiz *et al.*, 2003). It is expected that when grown in soil with no mercury contamination, the *merA/merB* poplar will be less productive than the wild-type trees, based on a previous report of *merA* alone in poplar (Che *et al.*, 2003).

Another heavy metal of interest is zinc, which can cause reduced foliage and dry-mass accumulation (Di Baccio *et al.*, 2003). When the cytosol and chloroplasts of *P. canescens* were transformed with the *E. coli gsh1* gene encoding for  $\gamma$ -glutamylsysteine synthetase ( $\gamma$ -ECS), the resulting trees clearly contained elevated levels of glutathione (Bittsanszky *et al.*, 2005). It is expected that higher GSH levels will result in enhanced phytochelatin production (Cobbett, 2000). When these transgenic and wild-type trees were subjected to varying levels of zinc, similar results were observed. At 10<sup>-1</sup> M, the symptoms were necrosis and severe phytotoxicity, while at 10<sup>-2</sup> M the leaves bleached, but continued growing.

At lower levels ( $10^{-3}$  to  $10^{-5}$  M), there were no toxic effects. Leaf zinc content increased with increasing treatment concentration, but did not significantly differ between transgenics and controls. However, trees expressing  $\gamma$ -ECS in the cytosol accumulated significantly more Cd, Cr and Cu than the wild-type or other transgenic lines, which is consistent with previously published results in poplar and *Arabidopsis* (Koprivova *et al.*, 2002). While GST activity in the wild-type and chloroplast-expressing lines increased, there was no observable change in lines with cytosolic expression, suggesting a lower stress response. GST is known to possess GSH peroxidase activity and can contribute to detoxification of active oxygen species (AOS). Therefore, increased GST levels are thought to contribute to the improved detoxification capacity in Zn-treated poplar leaves, but the mechanism is still unknown (Bittsanszky *et al.*, 2005).

Cadmium, another significant pollutant, can accumulate in soils and be phytotoxic due to its reactivity with O-, N- and S-containing ligands. In short, it inhibits photosynthesis and increases respiration, as carbohydrate metabolism (e.g. TCA cycle) is induced by increased leaf Cd content. In addition, Cd induces the synthesis of phytochelatin, which form complexes with Cd that can then be sequestered in the vacuole (Arisi *et al.*, 2000).

Phytochelatin are synthesized by  $\gamma$ -glutamylcysteine ( $\gamma$ -EC) dipeptidyl transpeptidase from reduced GSH, which is formed by two sequential reactions catalysed by  $\gamma$ -ECS and GS in the chloroplasts and cytosol of plant cells (Arisi *et al.*, 2000). Cd tolerance is related to GSH accumulation in leaves and increased capacity for GSH synthesis. Hybrid poplar (*P. tremula*  $\times$  *P. alba*) overexpressing bacterial  $\gamma$ -ECS in the cytosol displayed 30-fold increases in foliar  $\gamma$ -ECS activity in the absence of Cd (Arisi *et al.*, 2000). Foliar  $\gamma$ -EC was increased ten fold and foliar GSH accumulation increased 2.5 to 3.5-fold relative to controls grown in the absence of Cd. All transgenic trees also had higher leaf cysteine concentrations in the absence of Cd. In the presence of Cd, foliar  $\gamma$ -ECS,  $\gamma$ -EC, glutathione and cysteine all increased in both transgenics and control trees. However, the transgenic lines were able to accumulate more Cd than control at all concentrations, and there was also less GSH accumulation in the leaves of controls. Cd-induced changes in enzyme activities were less pronounced in the leaves of transgenic lines. Despite being able to accumulate more Cd in the leaves, the transgenic plants did not have a greatly increased tolerance to Cd (Arisi *et al.*, 2000). Koprivova *et al.* (2002) also examined the effects of overexpression of a bacterial  $\gamma$ -ECS on Cd accumulation in poplar. They initially looked at CdCl<sub>2</sub> in a hydroponic system, and the transgenics were able to accumulate significantly more Cd in root tissue than plants overexpressing GS or wild-types. At low Cd concentrations there was no difference in accumulation, but at higher Cd concentrations the transgenics were able to accumulate 2.5 to 3 times more Cd in the leaf tissue than the wild-type trees (Koprivova *et al.*, 2002).

Many sites with heavy metal contamination are also contaminated with organic components such as trichloroethylene (TCE; Meagher *et al.*, 1998). Chlorinated compounds, such as TCE, are among the most widespread groundwater

contaminants in the United States of America. TCE was commonly used as a dry cleaning solvent and as a metal-degreasing agent, especially by the military and the electronics industry. The USEPA classifies TCE as a suspected human carcinogen and gives high priority to its clean-up (Doty *et al.*, 2000). About 40% of all its Superfund sites are contaminated with this substance. It has also been shown that exposure can result in depression of the central nervous system (Costa, Katz and Ivanetich, 1980). Moreover, TCE persists in the environment for decades. Existing remediation techniques (e.g. pumping or air-stripping) are labour-intensive, expensive and wasteful. While some plants have an innate ability to absorb and metabolize TCE, those that have been genetically engineered to contain the appropriate gene possess enhanced ability to metabolize TCE. For example, tobacco plants transformed with the gene encoding a mammalian cytochrome P450 2E1 (CYP2E1) are able to metabolize TCE at a rate that is 640-fold greater than plants without this gene (Doty *et al.*, 2000). The encoded enzyme also metabolizes a wide range of other harmful pollutants, including ethylene dibromide (used as a gasoline additive and as a soil fumigant to control nematodes), carbon tetrachloride, chloroform and vinyl chloride (Doty *et al.*, 2007). This gene is a prime candidate for genetic engineering trees to remediate contaminated sites.

As is the case with heavy metals, a disadvantage of phytoremediation is that when the plants remove pollutants from the groundwater and soil, they either transpire them, unaltered, into the atmosphere, or sequester them in various tissues. Thus, plants that could metabolize the pollutant would be more desirable. Although significant gains in the phytoremediative capacity of trees have been shown in controlled environments, limited field-testing has been conducted to date.

## Hormones

Much research has been carried out in lignin modification, flowering, and abiotic and biotic resistances. Genes that control hormone synthesis and sensitivity are potential candidates for producing trees that have altered wood properties and other desirable characteristics. These include reduced axillary bud break, high density, long fibres, better rooting and improved growth rate, all of which are influenced by hormones.

Cytokinins are important as mediators of growth and differentiation in plants. The isopentenyltransferase gene (*IPT*) from *Agrobacterium tumefaciens* catalyses the conversion of adenosine-5'-monophosphate and isopentenylpyrophosphate to isopentenyladenosine-5'-monophosphate (Akiyoshi *et al.*, 1984), which is then converted to isopentenyl- and zeatin-type cytokinins (Von Schwartzberg *et al.*, 1994). Poplar overexpressing *IPT* showed increased formation of branching, with short internodes that were unable to root. The calli were able to regenerate buds in the absence of exogenous cytokinins, and contained high concentrations of zeatin, zeatin riboside and isopentenyladenosine (Von Schwartzberg *et al.*, 1994). The effects of the transformation were noticeable even at a very early stage, as transgenic explants developed green calli with significantly more buds than



non-transformed controls on medium lacking thidiazuron (TDZ). When grown on media containing TDZ, fewer buds formed on the transgenic explants. Isolated transgenic shoots showed reduced apical dominance with frequent branching, shorter internodes and the inability to root (Von Schwartzenberg *et al.*, 1994).

Another phytohormone, indole-3-acetic acid (IAA), is important for maintaining the structure and integrity of the vascular cambium. Exogenous application of auxins or auxin transport inhibitors affect many aspects of cambial growth, including xylem production, cell size and thickness, and reaction wood and vessel density (reviewed by Little and Savidge, 1987). Enzymes encoded by *iaaM* (trp-2-mono-oxygenase) and *iaaH* (indole-3-acetamide hydrolase) catalyse the two-step formation of IAA from tryptophan, and the overexpression of these two genes in plants has led to increased IAA levels (Sitbon *et al.*, 1991, 1992). Poplar transformed with *Agrobacterium iaaM* and *iaaH* showed phenotypic alterations (Tuominen *et al.*, 1995), as might be expected. All transgenic lines were smaller than controls, but there was variation in extent of stunting. Some lines showed elevated levels of free and conjugated IAA in the mature leaves and roots, relative to controls. Decapitated transgenic lines had a lower bud release rate than controls, with some plants showing no release at all. Many of these plants survived and continued to grow, with large increases in the growth of the lower trunk. Transgenics also had alterations in xylem formation, with decreased width and altered structure, and also had large fibres and small vessels, which appeared more uniform in size and rounded.

Tuominen *et al.* (2000) engineered poplar (*P. tremula* × *P. tremuloides*) to overexpress *iaaM* fused to the *GUS* reporter gene under the control of the cambial region-specific *Agrobacterium rhizogenes rolC* promoter. While IAA levels were increased, the radial distribution pattern remained unchanged, and no changes were seen in the developmental pattern of cambial derivatives or in cambial second growth, suggesting that the distribution pattern of IAA holds a more important role in wood formation than changes in the amount of IAA (Uggla *et al.*, 1996). Despite a 35% increase in IAA (with only minor changes in conjugate pools), phenotypic changes were relatively minor. Transgenic lines had increased internode length, and decreased occurrence of axillary bud break following decapitation. Some lines also showed decreased leaf size or height, but this response was not consistent.

Plants overexpressing the *rolC* gene showed significant alteration in growth and development. This has been observed in various species, including tobacco (Nilsson *et al.*, 1993) and potato (Fladung, 1990). The plants are dwarfed and have reduced apical dominance, shorter internodes and smaller leaves, suggesting that *rolC* is related to an increase in cytokinin activity (Nilsson *et al.*, 1996). Hybrid poplar transformed with *rolC* resulted in an increased level of free cytokinins (Nilsson *et al.*, 1996). Transgenic trees had reduced apical dominance with more axial shoots. When the side shoots were removed, the trees showed normal growth and apical dominance with a single shoot (Nilsson *et al.*, 1996). Additional lines exhibited fasciation, enlarged shoot apices and revealed the apical meristem in some cases.

The fasciated apices of the transgenic lines were smaller and more numerous, and the leaves were also smaller and thicker, with larger palisade cells than wild-type leaves. The transgenics also had flattened stems as a result of fasciation (Nilsson *et al.*, 1996). Transgenic lines had less free IAA in the upper leaves and apical meristem, but the conjugated IAA level was not changed, or only showed a slight increase in the apex. Gibberellic acid (GA) activity was markedly lower in the upper regions of the transgenic trees, while cytokinins were unchanged relative to control plants, despite increases in zeatin riboside levels. It appeared that cytokinin levels were regulating conjugation (Nilsson *et al.*, 1996). Additional studies with *rolC* genes have consistently shown changes in hormone levels and growth morphology, as well as changes in timing of dormancy and bud flush (Table 4-4).

Poplar overexpressing *rolC*, characterized by reduced shoot growth and early bud break, have also been examined for changes in wood properties. Wood formation started at the same time as control trees, and there were no changes in wood structure. These observations suggest that the dwarfism was due to a decreased number of cells, as a result of slower cell differentiation rates. However, when compared with controls, cells in transgenics lacked secondary walls and normal lignification, and had discoloured wood and tyloses. In addition, ring borders were not easily identifiable because the transgenics lacked thick-walled fibres associated with latewood (Grunwald *et al.*, 2000). In the control trees, the reactivation of wood formation coincided with bud break, but in the transgenic lines it coincided with full leaf expansion. Given these results, it is possible that

TABLE 4-4  
Summary of genetic modifications in trees, targeting hormone regulation

Gene	Modification	Effects	Reference
<i>ipt</i>	Overexpression in <i>Populus tremula</i> × <i>P. alba</i>	Increased branching shoots, short internodes, unable to root	Von Schwartzberg <i>et al.</i> , 1994
<i>iaaH</i> and <i>iaaM</i>	Overexpression in <i>P. tremula</i> × <i>P. tremuloides</i>	Smaller trees with elevated free and conjugated IAA; decreased axillary bud release following decapitation	Tuominen <i>et al.</i> , 1995
<i>rolC</i>	Overexpression in <i>P. tremula</i> × <i>P. tremuloides</i>	Reduced apical dominance, increased axillary shooting, fasciated apices	Nilsson <i>et al.</i> , 1996
<i>rolC</i>	Overexpression in <i>P. tremula</i> × <i>P. tremuloides</i>	Alterations in hormone levels with a decrease of ABA in pre-dormant buds and during resting; earlier flushing	Fladung, Grossmann and Ahuja, 1997
<i>OSH1</i>	Overexpression in <i>P. nigra</i> L. var. <i>italica</i>	Alterations in phenotypes with three relatively distinct phenotypes identified: I - slender leaves, II - dwarfed, III - multiple shoot apices with tiny leaves	Mohri <i>et al.</i> , 1999
<i>rol</i> genes	Overexpressed in <i>P. tremula</i>	Shorter, but more numerous internodes; axillary shooting; decreased shoot:root ratios; delayed dormancy	Tzfira, Vainstein and Altman, 1999
<i>GA-20 oxidase</i>	Overexpressed in <i>P. tremula</i> × <i>P. tremuloides</i>	Increased height and diameter; increased internode length; longer, broader leaves; increased number of cells; significantly longer xylem cells	Eriksson <i>et al.</i> , 2000
<i>rolC</i>	Overexpressed in <i>P. tremula</i> × <i>P. tremuloides</i>	Reduced shoot growth; early bud break; no change in timing of wood formation; more numerous tyloses formed; lacked thick walled late wood	Grunwald <i>et al.</i> , 2000
<i>iaaM</i>	Overexpressed in <i>P. tremula</i> × <i>P. tremuloides</i>	35% increase in IAA, but no change in radial distribution pattern; increased internode length; decreased occurrence of axillary bud break following decapitation	Tuominen <i>et al.</i> , 2000

the transgenics produced lower amounts of auxin or other factors that are required for cambial division (Grunwald *et al.*, 2000). Discoloration and tyloses may be associated with wounds because, in aspen, tyloses are formed when air enters the wood (Grunwald *et al.*, 2000). Thus, the *rolC* trees may be more susceptible to damage than the controls. The transgenics also did not have fully formed and lignified cell walls, possibly as a result of slower differentiation, causing incomplete development of cells formed at the end of the growing season, or a lack of a signal for latewood maturation (Grunwald *et al.*, 2000).

Gibberellins influence growth and development in plants, including effects on shoot growth, leaf growth and shape, flowering and seed germination. GA also plays a role in cell division and elongation (Kende and Zeevaart, 1997). They are formed through the isoprenoid pathway from mevalonic acid, and are regulated by transcriptional control (Hedden and Proebsting, 1997). GA20 catalyses the production of the immediate precursors to the active gibberellins GA4 and GA1. Overproduction of GA in hybrid poplar resulted in improved growth rates. Transgenics had increased height and diameter growth; increased internode lengths; and longer, broader leaves with longer petioles. The transgenic lines also showed an increase in the number of cells, and the xylem cells were significantly longer than those found in control plants (Eriksson *et al.*, 2000).

Poplar transformed with a gene encoding the rice homeodomain protein *OSH1* also showed morphological abnormalities in the leaves and stems (Mohri *et al.*, 1999). This is similar to results observed previously in rice (Kano-Murakami *et al.*, 1993), *Arabidopsis* (Matsuoka *et al.*, 1993) and tobacco (Kusaba *et al.*, 1998). There were three major phenotypes identified: type 1 had slender leaves; type 2 were dwarf plants with limited life spans; and type 3 had multiple shoot apices and tiny leaves. The expression level was highest in type 3 and lowest in type 1. The phenotype is thought to be the result of a disruption in the balance of plant hormones, as seen in other plants (Mohri *et al.*, 1999).

## WOOD TRAITS

As a consequence of a rapidly growing human population, the world's forests are experiencing increasing pressures to meet demands for wood products, fuel and agricultural land. Moreover, these efforts are being met with more stringent environmental regulations and an increasing interest in sustainability. Clearly, there are huge opportunities for forest and tree biotechnology research, particularly that focused on making wood products available faster, of better quality, and with fewer negative effects on native forests and the environment in general (Boerjan, 2005). The following section highlights traits related to wood fibre chemistry, ultrastructure and growth.

Directed genetic modifications altering the quality and quantity of wood and cell wall components have been pursued by researchers for nearly two decades. Numerous modifications have been reported in model plants such as *Arabidopsis* and tobacco, but they are not useful for the study of wood. *Populus* is not only a valuable model tree, but it is also commercially important, particularly in

the Northern Hemisphere. Because of this and the ease with which it can be manipulated, the vast majority of the tree genetic modification efforts have been with species in this genus. Less frequent, but equally important, are the reports on *Eucalyptus* and industrially important conifers, such as spruce and pine. Genetic modification of genes involved in cell-wall biosynthesis fall into two categories: lignin and non-lignin cellulosic material. The altered expression or regulation of representative genes in both categories has resulted in a range of changes to the tree cell walls, from extreme to no quantifiable difference. The efforts include attempts to up- and down-regulate gene expression, and have employed both endogenes and novel sequences and promoters. Nearly all of the enzymes implicated in the currently accepted lignin biosynthetic pathway have been targeted or modified in some manner (see Table 4-5). Although not as common heretofore, modifying the expression of genes involved in cellulose biosynthesis is a rapidly growing area of interest in tree biotechnology research (see Table 4-6).

TABLE 4-5  
Summary of genetic modifications in plants, targeting lignin biosynthesis

Gene	Modification	Effects	Reference
PAL	Downregulated in tobacco	Reduced lignin content, slightly increased S:G	Sewalt <i>et al.</i> , 1997
	Downregulated in tobacco	Reduced phenylpropanoid compounds in leaves and stems	Blount <i>et al.</i> , 2000
C4H	Downregulated in tobacco	Minor reduction in lignin content, minor changes to S:G	Blee <i>et al.</i> , 2001
	Downregulated in tobacco	Decreased lignin content, decreased S:G	Sewalt <i>et al.</i> , 1997
	Downregulated in tobacco	Altered PAL activity, decreased phenylpropanoid compounds	Blount <i>et al.</i> , 2000
COMT	Downregulated in <i>Populus tremula</i> × <i>alba</i>	4-yr field trial; no dramatic ecological/biological impacts	Halpin <i>et al.</i> , 2007
	Downregulated in <i>P. tremula</i> × <i>alba</i>	Decreased lignin content, decreased S:G	Jouanin <i>et al.</i> , 2000
	Downregulated in <i>P. tremula</i> × <i>alba</i>	Increased G, lower pulping efficiency	Lapierrre <i>et al.</i> , 1999
	Downregulated in <i>P. tremula</i> × <i>alba</i>	4-yr field trial; normal growth	Pilate <i>et al.</i> , 2002
	Downregulated in <i>P. tremuloides</i>	No change in lignin content, S:G decreased, more coniferaldehyde	Tsai <i>et al.</i> , 1998
	Downregulated in <i>P. tremula</i> × <i>alba</i>	No change in lignin content, decreased S, increased G units	Van Doorselaere <i>et al.</i> , 1995
	Downregulated in <i>P. tremula</i> × <i>alba</i>	Lignin contains 5-hydroxyconiferyl alcohol and benzodioxane units	Ralph <i>et al.</i> , 2001
F5H	Overexpressed in <i>P. tremula</i> × <i>alba</i>	Increased S units	Franke <i>et al.</i> , 2000
	Overexpressed in <i>P. tremula</i> × <i>alba</i>	No change in lignin content, increased S:G, decreased kappa	Huntley <i>et al.</i> , 2003
4CL	Downregulated in <i>P. tremuloides</i>	Decreased lignin content, no S:G changes	Hu <i>et al.</i> , 1999
	Downregulated in <i>P. tremuloides</i>	Decreased lignin content, S:G increase,	Li <i>et al.</i> , 2003
	Downregulated in <i>P. tomentosa</i>	Decreased lignin content	Jia <i>et al.</i> , 2004
	Downregulated in <i>P. tremuloides</i>	Decreased lignin content, no S:G changes	Hancock <i>et al.</i> , 2007

TABLE 4-5 (CONTINUED)

Gene	Modification	Effects	Reference
<i>HCT</i>	Downregulated in <i>Pinus radiata</i>	Decreased lignin content, altered monolignol composition	Wagner <i>et al.</i> , 2007
<i>C3H</i>	Downregulated in <i>P. alba</i> × <i>grandidentata</i>	RNAi down regulation, reduced lignin, increased H units, decreased G units	Coleman <i>et al.</i> , 2007
<i>CCR</i>	Downregulated in tobacco	Some reduced lignin, decreased kappa	Chabannes <i>et al.</i> , 2001
	Downregulated in tobacco	Changes in transcriptome and metabolome; decreased phenylpropanoid pathway genes	Dauwe <i>et al.</i> , 2007
	Downregulated in <i>P. tremula</i> × <i>alba</i>	5-year field trial; decreased lignin content, decreased S:G, improved pulping efficiency	Leple <i>et al.</i> , 2007
	Downregulated in tobacco	Decreased lignin content, some increased S:G	O'Connell <i>et al.</i> , 2002
	Downregulated in <i>Picea abies</i>	5-yr field trial; slightly decreased lignin content, decreased H units, slightly decreased kappa	Wadenback <i>et al.</i> , 2008
<i>Cald5H</i>	Downregulated in <i>P. tremuloides</i>	Decreased lignin content, S:G increased	Li <i>et al.</i> , 2003
	Downregulated in <i>P. tremuloides</i>	No change in lignin content, increased S:G	Hancock <i>et al.</i> , 2007
<i>CcoAOMT</i>	Downregulated in <i>P. tremula</i> × <i>alba</i>	Decreased lignin content, slightly increased S:G	Meyermans <i>et al.</i> , 2000
	Downregulated in <i>P. tremula</i> × <i>alba</i>	Decreased lignin content	Zhong <i>et al.</i> , 2000
	Downregulated in <i>P. tremula</i> × <i>alba</i>	Decreased lignin content	Wei <i>et al.</i> , 2001 in Lin <i>et al.</i> , 2006
<i>CAD</i>	Downregulated in <i>P. tremula</i> × <i>alba</i>	Slightly decreased lignin content, increased aldehydes	Baucher <i>et al.</i> , 1996
	Downregulated in tobacco	Some reduced lignin, decreased kappa	Chabannes <i>et al.</i> , 2001
	Downregulated in tobacco	Changes in transcriptome and metabolome; decreased phenylpropanoid pathway genes	Dauwe <i>et al.</i> , 2007
	Overexpressed in tobacco	No change in lignin content; thicker cell walls	Goicoechea <i>et al.</i> , 2005
	Overexpressed in tobacco	Slightly decreased lignin content; more easily extracted	Halpin <i>et al.</i> , 1994
	Downregulated in <i>P. tremula</i> × <i>alba</i>	4-yr field trial; no dramatic ecological or biological impacts	Halpin <i>et al.</i> , 2007
	Downregulated in <i>P. tremula</i> × <i>alba</i>	Decreased lignin content, higher free phenolics, easier pulping	Lapierre <i>et al.</i> , 1999
	Downregulated in tobacco	Increased S, more easily extracted	O'Connell <i>et al.</i> , 2002
	Downregulated in <i>P. tremula</i> × <i>alba</i>	4-yr field trial; normal growth, lower kappa, higher yield, no change in insect interactions	Pilate <i>et al.</i> , 2002
	Downregulated in <i>Eucalyptus camaldulensis</i>	No change in lignin content, quality, composition	Valerio <i>et al.</i> , 2003
	Downregulated in <i>P. taeda</i>	Reduced lignin, brown wood, lignin contains dihydroconiferyl alcohol, increased aldehydes	MacKay <i>et al.</i> , 1997; Ralph <i>et al.</i> , 1997
<i>Laccase</i>	Downregulated in <i>P. trichocarpa</i>	No change to lignin content or composition, deformed xylem, increased phenolics	Ranocha <i>et al.</i> , 2002
<i>Peroxidase</i>	Downregulated in <i>Populus sieboldii</i> × <i>grandidentata</i>	Decreased lignin content, increased S:G ratio	Li <i>et al.</i> , 2003

Notes: Modifications are in trees, and some references review results in tobacco. Tree lignin comprises two main forms, guaiacyl (G) and syringal (S), and their ratio determines many characteristics of the organism. The S:G ratio is therefore an important indicative parameter of suitability for particular uses.

TABLE 4-6  
**Summary of genetic modifications in plants targeting cell wall biosynthesis**

Gene	Modification	Effects	Reference
<i>Xyloglucanase</i>	Expressed in <i>Populus alba</i>	Increased cellulose content, decreased xyloglucans, decreased lignin	Park <i>et al.</i> , 2004
<i>UDP-GD</i>	Downregulated in tobacco	Decreased xylose-containing polymers, increased glucose/xylose content in cell walls	Bindschedler <i>et al.</i> , 2007
<i>4CL</i>	Downregulated in <i>P. tremuloides</i>	Compensatory increase in cellulose content due to lignin reduction	Hu <i>et al.</i> , 1999
<i>Invertase</i>	Expressed in tobacco	Increased cellulose content, reduced growth, increased biomass	Canam <i>et al.</i> , 2006
<i>SuSY/UGPase</i>	Overexpressed in tobacco	No change in cellulose content, increased biomass	Coleman <i>et al.</i> , 2006
<i>UGPase</i>	Expressed in <i>P. alba</i> × <i>grandidentata</i>	Increased cellulose, decreased lignin, increased s units in lignin, reduced growth	Coleman <i>et al.</i> , 2007
<i>CCR</i>	Downregulated in <i>P. tremula</i> × <i>alba</i>	Proportional increase in cellulose due to lignin reduction, reduced hemicellulose content	Leple <i>et al.</i> , 2007
<i>4CL/CAld5H</i>	<i>4CL</i> -downregulated, <i>CAld5H</i> overexpressed in <i>P. tremuloides</i>	Increase in cellulose content, reduced lignin content	Li <i>et al.</i> , 2003
<i>AtCelA1</i>	Overexpressed in <i>P. tremula</i>	Increased cellulose, increased hemicellulose	Shani <i>et al.</i> , 2004

### Lignin content

Hu *et al.* (1999) were among the first to demonstrate the potential of genetic engineering for modifying lignin in trees for industrial applications. *Populus tremuloides* was transformed with antisense 4-coumarate:coenzyme ligase (4CL) constructs that resulted in a 45% reduction in lignin content. This dramatic decrease in total lignin, with no concurrent changes to lignin monomer composition, is advantageous to several industries, including the manufacture of pulp and paper, because lignin removal consumes large amounts of energy and reagents. Pilate *et al.* (2002) conducted a four-year field trial with hybrid poplars (*P. tremula* × *P. alba*) engineered for lower caffeic acid O-methyltransferase (COMT) and cinnamoyl alcohol dehydrogenase (CAD) activity. CAD-reduced trees demonstrated greater ease of delignification and superior yield, whereas COMT-altered trees required more energy for lignin removal. By contrast, in similar efforts with transgenic *Eucalyptus*, reduced CAD expression (antisense) resulted in no change in lignin quality and composition, or pulp yields (Tournier *et al.*, 2003). Importantly, the lignin composition changes seen in the field-grown transgenics studied by Pilate *et al.* (2002) were maintained over the four-year trial. More recently, it was shown, using these same trees, that there does not appear to be dramatic changes in the local insect and soil microbe communities surrounding the transgenic plots. This suggests that, depending on what modifications are done, there may be little or no negative ecological impacts of growing transgenic trees (Halpin *et al.*, 2007). However, it must be emphasized that ecological studies are complicated and critically needed in order to assess more fully the impact of transgenic trees. Furthermore, longer-term trials will be required to fully appreciate the potential for unexpected changes and effects.



Measuring changes in the transcriptome and metabolome of CAD- and cinnamoyl CoA reductase (CCR)-modified tobacco plants has revealed that altering one gene in the lignin biosynthetic pathway affects the expression of other genes within the same pathway, as well as genes involved in detoxification, carbohydrate metabolism, and photorespiration (Dauwe *et al.*, 2007). Although the effect of genetic modifications to forest trees will surely differ from those of tobacco, this research provides an example of the changes that can occur. In fact, more recently, the metabolism and transcript changes in response to CCR down-regulation in *P. tremula* × *P. alba*, which had decreases in total lignin and an increase in G monomer units, suggest that a stress response was elicited (Leple *et al.*, 2007). Also, a general decrease of transcripts related to non-cellulosic cell-wall polymers was observed. Although pulping efficiency was increased for wood from these transgenics, the trees in this five-year-old field trial were stunted. These reports emphasize the importance of long-term field trials and the need to assess non-target effects. Although over 200 field trials exist throughout the world (FAO, 2004; Boerjan, 2005), no published reports have evaluated transgenic trees modified for wood traits over the normal rotation of a forest plantation. This is a critical hole in bridging the gap between tree biotechnology and practical uses, and in gaining public acceptance.

Although the technology for genetic modification of conifers has existed for several years (Ellis *et al.*, 1993), the production and growth of genetically modified conifers is slow and lags behind similar work in angiosperms. Recently, however, Wadenback *et al.* (2008) reported a slight reduction in lignin content (8%) in five-year-old antisense CCR Norway spruce (*Picea abies*). In comparison, down-regulation of CCR activity in tobacco and poplar has demonstrated as much as a 50% reduction in lignin content (Chabannes *et al.*, 2001; Leple *et al.*, 2007). As was reported with the CCR-altered tobacco and poplar, the down-regulation of CCR in spruce led to narrower stems: a form of stunting. Although the reduction in lignin content of the modified spruce is at the lower limit of biological variation (Wadenback *et al.*, 2008), it is important to note that these findings are in an economically important conifer. These results also suggest that it may be necessary to target multiple genes to achieve the desired lignin modifications in trees.

### Lignin composition

Equally important to reducing lignin content for downstream processing is altering the composition of lignin monomers to improve the overall delignification process. An increase in the lignin S:G monomer ratio has been clearly shown to improve the manner and the efficacy of pulping wood (Chang and Sarkanen, 1973; Stewart, Kadla and Mansfield, 2006; Mansfield and Weiniesen, 2007). Over the last decade, substantial effort has been devoted to altering monomer composition (Table 4-5). A significant reduction in total lignin content and a concurrent decrease in S monomers has been achieved by sense-suppression of COMT under the regulation of the 35S promoter (Jouanin *et al.*, 2000). Alternatively, the

overexpression of COMT under the regulation of the *Eucalyptus* CAD promoter resulted in only slight increases in COMT activity in some lines, but the increased COMT activity did not result in altered S:G ratios (Jouanin *et al.*, 2000). This result is interesting because, based on the currently accepted lignin biosynthetic pathway, one would expect that increasing COMT would drive biosynthesis toward S units. The results achieved to date point to the need for careful promoter and gene selection when making targeted genetic modifications.

By down-regulating caffeoyl-coenzyme A O-methyltransferase (*CCoAOMT*), Meyermans *et al.* (2000) were able to generate transgenic lines of hybrid poplar displaying an 11% increase in the S:G ratio, along with a 12% decrease in lignin content. Alternatively, the successful expression of the *Arabidopsis* ferulate-5-hydroxylase (*F5H*) gene under the regulation of the cinnamate 4-hydroxylase (C4H) promoter, employed by Franke *et al.* (2000), led to a substantial increase in the composition of syringyl monomers in *P. tremula* × *P. alba*, at greater than 90% mol S lignin (Huntley *et al.*, 2003). The pulping efficiency of these trees clearly showed dramatic decreases in the energy required for chemical pulping (delignification), with a 23 kappa unit decrease compared with control trees. In addition, pulps had a higher ISO brightness value. In combination, these transgenic trees provide an excellent opportunity to decrease the energy and chemicals required for extracting lignin and obtaining a high-quality pulp.

In an effort to decrease total lignin content while also altering the lignin monomer ratio, Li *et al.* (2003) used a combinatorial approach by simultaneously decreasing 4CL activity and increasing coniferaldehyde 5-hydroxylase (CAlD5H) expression in *P. tremuloides*. The authors reported lignin content reductions as high as 52% and concomitant increases in the S:G ratio. This work highlights the potential benefit from concurrent augmentation and reduction of different gene products within the lignin biosynthesis pathway.

### Efforts to modify cell wall polysaccharides

Genetic modification in trees has resulted in increased cellulose content, both directly and indirectly (Table 4-6). For example, efforts to engineer trees with altered lignin composition has demonstrated the added advantage of indirectly improved cellulosic quantities per unit mass, as demonstrated by Hu *et al.* (1999) and Li *et al.* (2003). Park *et al.* (2004) have successfully increased cellulose and decreased xyloglucan contents in *P. alba* by expressing a fungal xyloglucanase gene. Similarly, *P. tremula* transformed with an *Arabidopsis* endoglucanase (*cel1*) were shown to have a 10% increase in cellulose content (Shani *et al.*, 2004). More recently, Coleman *et al.* (2007) have shown that transgenic *P. alba* × *P. grandidentata* trees expressing a bacterial UDP-glucose pyrophosphorylase (UGPase) gene have substantially increased cellulose content, and decreased lignin. However, these trees grew more slowly than the controls. Alternatively, in tobacco, the expression of yeast-derived invertases has been shown to result in decreased growth rates, but some plants accumulated more biomass and up to 36% more cellulose (Canam *et al.*, 2006). These genetic modifications indicate

that altering carbon allocation is possible, but optimizing cellulose production and growth requires further work.

The identification of genes and enzymes involved in cellulose and hemicellulose biosynthesis is ongoing. For example, Suzuki *et al.* (2006) identified a xylem-specific mannan synthases from *P. trichocarpa* by comparing orthologous genes from *Arabidopsis* and subsequently measuring mannan synthase activity *in vitro*. Looking globally at the genes and proteins involved in cellulose-rich G layer production in poplar, Andersson-Gunneras *et al.* (2006) also revealed some potential targets to alter cellulose production, or perhaps cellulose extractability. The identification of functioning xyloglucan transglycosylases (XETs) in developing secondary xylem in aspen (Bourquin *et al.*, 2002) and its involvement in the formation of the G layer in tension wood (Nishikubo *et al.*, 2007) suggests that XET may be a potential target to modify cellulose properties. If XET is involved in establishing cross-linkages between cellulose microfibrils, altering its expression could greatly affect fibre structure.

## IMPLICATIONS AND FUTURE DIRECTIONS

Although not all efforts have led to an improvement in industrial processes, they have contributed significantly to our understanding of the fundamental mechanisms of cell wall synthesis and formation. For example, a 90% reduction in *CCoAOMT* activity in transgenic poplar only led to an 11% decrease in lignin content (Meyermans *et al.*, 2000), suggesting that *CCoAOMT* has minimal control over the flow of carbon through the lignin pathway (Anterola and Lewis, 2002). Very recently, Wagner *et al.* (2007) revealed a functional hydroxycinnamoyl-CoA:shikimate hydroxycinnamoyltransferase (HCT) in *Pinus radiata* tracheary elements. This gene had not previously been implicated in lignin biosynthesis in gymnosperms, and may be a new target to genetically modify lignin for forestry and biofuels.

Understanding the role of genes involved in cellulose biosynthesis lags behind the progress that has been made with phenylpropanoid biosynthesis. Along with the *CesA* complex itself, the roles of genes such as sucrose synthase (*SuSy*), sucrose phosphate synthase (*SPS*), invertase, *UGPase* and korrigan (*KOR*), to name a few, are examples of genes or gene families that are currently under investigation (for a review, see Joshi and Mansfield, 2007). As information from these studies becomes available, it is very likely that new opportunities for modifying cellulose production, cell wall architecture and growth in trees will become apparent.

The identification of other genetic elements, such as transcription factors, is also an area of tree biotechnology research that affects wood traits. Legay *et al.* (2007) identified a MYB transcription factor, EgMYB1, in Eucalyptus xylem cDNA that represses lignin biosynthetic genes. This discovery could lead to new approaches for modifying wood. For example, Goicoechea *et al.* (2005) overexpressed EgMYB2 in tobacco and showed that the plants had slightly higher S:G, and thicker cell walls. Although no change in total lignin was reported, there could have been an increase that was offset by the additional cell-wall mass (perhaps because of

other cell-wall components). This requires further investigation. Tobacco was also used as a model to study the function of a pine MYB, PtMYB4 (Patzlaff *et al.*, 2003). The authors found that PtMYB4 expression in tobacco caused an increase in overall lignin. Also, identification of transcription factors that affect secondary cell wall deposition in *Arabidopsis* (e.g. Ko *et al.*, 2007) may provide yet another avenue to genetically modify wood traits. Undoubtedly, understanding the role that transcription factors play in lignin and cellulose biosynthesis will be necessary for downstream modifications to be effective.

Other biotechnology applications, such as activation tagging (Weigel *et al.*, 2000) in poplar may also reveal useful and novel modifications that affect cell-wall quality and quantity. Phenotypes from activation-tagged trees have been reported (Busov *et al.*, 2003) and will continue as thousands of activation-tagging transgenic lines are evaluated in the greenhouse and in the field (Harrison *et al.*, 2007; Arborea project, [www.arborea.ulaval.ca/](http://www.arborea.ulaval.ca/)). Finally, the identification of microRNAs involved in tree-specific mechanisms, such as tension wood formation (Lu *et al.*, 2005), is yet another area of research that will soon contribute greatly to our understanding of cell-wall biosynthesis and regulatory mechanisms of wood formation, all of which will be important for identifying future targets of tree biotechnology.

## GROWTH

One of the major goals of plant research is to increase yield, primarily in the form of increased stem biomass through an increase in height and diameter. Numerous, indirect approaches have been used (Table 4-7), some of which were mentioned in previous sections of this chapter. Here we focus on transformations that have resulted in increased growth.

One approach has been to transform poplar with the gene encoding uridine diphosphoglycosyl-transferase, *ugt*, which catalyses the conjugation of IAA with

TABLE 4-7  
Summary of genetic modifications in trees affecting growth

Gene	Modification	Effects	Reference
<i>4CL</i>	Antisense inhibition in <i>Populus</i>	Increased plant growth; structural integrity maintained	Hu <i>et al.</i> , 1999
<i>GS1</i>	Overexpressed in <i>Populus</i>	Increased node and leaf number, larger leaves; increased growth; enhanced nitrogen assimilation and increased growth under both high and low nitrogen conditions	Fu <i>et al.</i> , 2003, Man <i>et al.</i> , 2005
<i>Xylo-glucanase</i>	Overexpression in <i>Populus</i>	Increased stem length and internode length	Park <i>et al.</i> , 2004
<i>cel1</i>	Overexpression in <i>Populus tremula</i>	Increased growth, larger leaves; increased stem diameter	Shani <i>et al.</i> , 2004
<i>ugt</i> and <i>acb</i>	Overexpression in <i>Populus</i> ; sense and antisense expression of <i>acb</i>	<i>ugt</i> plants show increased growth; <i>ugt</i> and <i>acb</i> lower growth than <i>ugt</i> alone; sense <i>acb</i> show increased growth; antisense <i>acb</i> show decreased growth	Salyaev <i>et al.</i> , 2006
<i>vhb</i>	Overexpression in <i>Populus</i>	Increased height and stem diameter	Zhang <i>et al.</i> , 2006
<i>PttEXPA1</i>	Overexpression in <i>Populus</i>	No change in height; increased internode length, fibre diameter and vessel element length; increased leaf expansion	Gray-Mitsumune <i>et al.</i> , 2007

glucose, allowing for a larger pool of IAA for transport. A second gene, *acb*, encoding acyl-CoA binding protein was also used in both sense and antisense directions. Its function is not known, but it is thought to help in the stabilization of membranes (Salyaev *et al.*, 2006). Transgenics overexpressing *ugt* and *acb* showed faster growth, along with elevated IAA concentrations. Transgenics also showed rapid bud and branch development. The height growth of the resultant *ugt* transgenics was about three times that of the control plants, and root elongation was greatly enhanced. Transgenics containing both *ugt* and *acb* had lower height growth than those with *ugt* alone. In contrast, poplar with sense *acb* grew faster than controls and those with *acb* antisense. The increased growth caused by *ugt* was either reduced or cancelled by the effects of *acb* mis-regulation (Salyaev *et al.*, 2006).

Poplars overexpressing an expansin gene, *PttEXPA1*, were recently shown to have increased stem internode length, increased leaf expansion, and larger cells in its leaf epidermis. Fibre diameter growth was increased, as was vessel element length (Gray-Mitsumune *et al.*, 2007). Additionally, poplar overexpressing the *Vitreoscilla* haemoglobin gene showed no significant morphological differences, but three lines had noticeably higher height and diameter growth rates (Zhang *et al.*, 2006). Although gains have been made in the area of increased biomass, more work is required as the trade-off between increased growth and fibre quality in trees is paramount to commercial end uses.

### Flowering control

Before genetically engineered trees can be commercialized, governing bodies will probably require a solid strategy to mitigate the risk of transgene spread and persistence in the environment. One way to satisfy this need is to control flowering (Meilan *et al.*, 2001). The manipulation of flowering can provide many benefits, such as development of a strategy to genetically engineer reproductive sterility. This may help alleviate, or at least reduce, public and regulatory concerns over the commercialization of transgenic trees. Sterility can also reduce genetic pollution from non-transgenic plantations, promote vegetative growth, and eliminate nuisance tissues (e.g. pollen, seed pods). In addition, flowering control may lead to shorter juvenile periods, resulting in shorter breeding cycles.

It is assumed that trees engineered for flowering control will re-direct photosynthate to harvestable products while, at the same time, minimizing gene flow to wild populations. Different types and degrees of sterility may be obtained via polyploidy (e.g. triploids or aneuploids), by genes specifically controlling male or female floral development, or genes controlling the onset of maturation. Ideally, flowering control should be reversible, so that with appropriate stimulus, the tree can be used for conventional breeding.

While each strategy for engineering sterility has advantages, it is unclear which method will work best with trees. Hence, tests are under way that involve a variety of techniques, such as tissue-specific ablation, dominant negative mutations and post-transcriptional gene silencing, including RNA interference. Employing the first approach, Skinner *et al.* (2003) successfully used the promoter from PTD, the

*Populus trichocarpa* homolog of the *Arabidopsis* *APETALA3* gene, to drive the expression of reporter and cytotoxin genes in floral tissues of *Arabidopsis*, tobacco and poplar.

Recently, RNA interference (RNAi) was used to reduce expression of the poplar ortholog of *CENTRORADIALIS* (*PtCENL1*), a gene that plays a key role in maintaining trees in a juvenile state (Mohamed, 2006). When transgenic poplars containing this RNAi vector were grown under field conditions, four of the most strongly silenced lines produced inflorescences or floral buds within two years of planting, which was several years earlier than that observed in wild-type trees. Surprisingly, overexpression of *PtCENL1* also resulted in delayed vegetative budbreak (Mohamed, 2006). Based on this work, it appears that *PtCENL1* is involved in regulating release from winter dormancy and resumption of growth. Hopefully, this work will ultimately lead to the development of methods for shortening breeding cycles, as well as possibly informing further research on flowering control.

Despite indications that one or more of the strategies involving flowering control can be successfully employed to engineer transgene confinement, no single method fulfils the basic requirements for long-term commercial use. Researchers are continuing to determine whether sterility can be complete and stable over several rounds of propagation and growing seasons, successfully identified in juvenile trees, and lack negative growth impacts.

## CONCLUSIONS

The modulation of complex traits such as tree growth, yield, chemical composition, morphology, and health is of vital interest to the plant biotechnology community. These characteristics are influenced by a multitude of environmental and genetic factors. Availability of the full *Populus* genome sequence, along with recent advances in transcript, protein and metabolomic profiling, will continue to lead to a better understanding of genetic modifications and regulation in trees. They will also provide new insights that will be needed to resolve uncertainties concerning the molecular processes that underlie wood formation, growth and plant-environment interactions. These advances, coupled with an improved understanding of the genes and enzymes involved in key metabolic pathways, should enable the genetic manipulation of trees so they will possess the desired properties and produce sufficient volume to satisfy society's ever-increasing demand for forest products.

## REFERENCES

- Akiyoshi, D.E., Klee, H., Amasino, R.M., Nester, E.W. & Gordon, M.P. 1984. T-DNA of *Agrobacterium tumefaciens* encodes an enzyme of cytokinin biosynthesis. *Proceedings of the National Academy of Sciences of the United States of America*, 81: 5994–5998.
- Andersson-Gunneras, S., Mellerowicz, E.J., Love, J., Segerman, B., Ohmiya, Y., Coutinho, P.M., Nilsson, P., Henrissat, B., Moritz, T. & Sundberg, B. 2006. Biosynthesis of cellulose-enriched tension wood in *Populus*: global analysis of transcripts and metabolites identifies biochemical and developmental regulators in secondary wall biosynthesis. *The Plant Journal*, 45: 144–165.



- Anterola, A.M. & Lewis, N.G. 2002. Trends in lignin modification: a comprehensive analysis of the effects of genetic manipulations/mutations on lignification and vascular integrity. *Phytochemistry*, 61: 221–294.
- Arisi, A.C.M., Noctor, G., Foyer, C.H. & Jouanin, L. 1997. Modification of thiol contents in poplars (*Populus tremula* × *alba*) overexpressing enzymes involved in glutathione synthesis. *Planta*, 203: 362–372.
- Arisi, A.C.M., Mocquot, B., Lagriffoul, A., Mench, M., Foyer, C.H. & Jouanin, L. 2000. Responses to cadmium in leaves of transformed poplars overexpressing  $\gamma$ -glutamylcysteine synthetase. *Physiologia Plantarum*, 109: 143–149.
- Baertlein, D.A., Lindow, S.E., Panopoulos, N.J., Lee, S.P., Mindrinos, M.N. & Chen, T.H.H. 1992. Expression of a bacterial ice nucleation gene in plants. *Plant Physiology*, 100: 1730–1736.
- Baker, A. & Brooks, R. 1989. Terrestrial higher plants which hyperaccumulate metallic elements - a review of their distribution, ecology, and phytochemistry. *Biorecovery*, 1: 81–126.
- Barbehenn, R., Jones, C., Yip, L., Tran, L. & Constabel, C. 2007. Limited impact of elevated levels of polyphenol oxidase on tree-feeding caterpillars: assessing individual plant defenses with transgenic poplar. *Oecologia*, 154: 129–140.
- Baucher, M., Chabbert, B., Pilate, G., Van Doorselaere, J., Tollier, M.T., Petit-Conil, M., Cornu, D., Monties, B., Van Montagu, M., Inze, D., Jouanin, L. & Boerjan, W. 1996. Red xylem and higher lignin extractability by down-regulating a cinnamyl alcohol in poplar. *Plant Physiology*, 112: 1479–1490.
- Bindschedler, L.V., Tuerck, J., Maunders, M., Ruel, K., Petit-Conil, M., Danoun, S., Boudet, A.-M., Joseleau, J.-P. & Bolwell, G.P. 2007. Modification of hemicellulose content by antisense down-regulation of UDP-glucuronate decarboxylase in tobacco and its consequences for cellulose extractability. *Phytochemistry*, 68: 2635–2648.
- Bishop-Hurley, S.L., Zabkiewicz, R.J., Grace, L., Gardner, R.C., Wagner, A. & Walter, C. 2001. Conifer genetic engineering: transgenic *Pinus radiata* (D. Don) and *Picea abies* (Karst) plants are resistant to the herbicide Buster. *Plant Cell Reports*, 20: 235–243.
- Bittsanszky, A., Komives, T., Gullner, G., Gyulai, G., Kiss, J., Heszky, L., Radimszky, L. & Rennenberg, H. 2005. Ability of transgenic poplars with elevated glutathione content to tolerate zinc<sup>(2+)</sup> stress. *Environment International, Recent Advances in Bioremediation*, 31: 251–254.
- Bittsanszky, A., Gyulai, G., Humphreys, M., Gullner, G., Csintalan, Z., Kiss, J., Szabo, Z., Lagler, R., Toth, Z., Rennenberg, H., Heszky, L. & Komives, T. 2006. RT-PCR analysis and stress response capacity of transgenic *gshI*-poplar clones (*Populus ×canescens*) in response to paraquat exposure. *Naturforsch*, 61: 699–703.
- Bizily, S.P., Rugh, C.L. & Meagher, R.B. 2000. Phytodetoxification of hazardous organomercurials by genetically engineered plants. *Nature Biotechnology*, 18: 213–217.
- Blee, K., Choi, J.W., O'Connell, A.P., Jupe, S.C., Schuch, W., Lewis, N.G. & Bolwell, G.P. 2001. Antisense and sense expression of cDNA coding for CYP73A15, a class II cinnamate 4-hydroxylase, leads to a delayed and reduced production of lignin in tobacco. *Phytochemistry*, 57: 1159–1166.
- Blennow, K. & Lindkvist, L. 2000. Models of low temperature and high irradiance and their application to explaining the risk of seedling mortality. *Forest Ecology and Management*, 135: 289–301.
- Blount, J.W., Korth, K.L., Masoud, S.A., Rasmussen, S., Lamb, C. & Dixon, R.A. 2000. Altering expression of cinnamic acid 4-hydroxylase in transgenic plants provides evidence for a feedback loop at the entry point into the phenylpropanoid pathway. *Plant Physiology*, 122: 107–116.
- Boerjan, W. 2005. Biotechnology and the domestication of forest trees. *Current Opinion in Biotechnology*, 16: 159–166.

- Bourquin, V., Nishikubo, N., Abe, H., Brumer, H., Denman, S., Eklund, M., Christiernin, M., Teeri, T.T., Sundberg, B. & Mellerowicz, E.J. 2002. Xyloglucan endotransglycosylases have a function during the formation of secondary cell walls of vascular tissues. *Plant Cell*, 14: 3073–3088.
- Brasileiro, A., Tourneur, C., Leple, J.-C., Combes, V.R. & Jouanin, L. 1992. Expression of the mutant *Arabidopsis thaliana* acetolactate synthase gene confers chlorsulfuron resistance to transgenic poplar plants. *Transgenic Research*, 1: 133–141.
- Busov, V.B., Meilan, R., Pearce, D.W., Ma, C., Rood, S.B. & Strauss, S.H. 2003. Activation tagging of a dominant gibberellin catabolism gene (*GA 2-oxidase*) from poplar that regulates tree stature. *Plant Physiology*, 132: 1283–1291.
- Canam, T., Park, J.-Y., Yu, K., Campbell, M., Ellis, D. & Mansfield, S. 2006. Varied growth, biomass and cellulose content in tobacco expressing yeast-derived invertases. *Planta*, 224: 1315–1327.
- Carozzi, N.B. & Koziel, M.G. 1997. *Advances in insect control: the role of transgenic plants*. Taylor and Francis, Inc., New York, USA.
- Chabannes, M., Barakate, A., Lapiere, C., Marita, J.M., Ralph, J., Pean, M., Danoun, S., Halpin, C., Grima-Pettenati, J. & Boudet, A.M. 2001. Strong decrease in lignin content without significant alteration of plant development is induced by simultaneous down-regulation of cinnamoyl CoA reductase (CCR) and cinnamyl alcohol dehydrogenase (CAD) in tobacco plants. *The Plant Journal*, 28: 257–270.
- Chang, H.-M. & Sarkanen, K.V. 1973. Species variation in lignin: effects of species on the rate of Kraft delignification. *TAPPI Journal*, 56: 132–134.
- Che, D.S., Meagher, R.B., Heaton, A.C.P., Lima, A., Rugh, C.L. & Merkle, S.A. 2003. Expression of mercuric ion reductase in eastern cottonwood (*Populus deltoides*) confers mercuric ion reduction and resistance. *Plant Biotechnology Journal*, 1: 311–319.
- Che, D.S., Meagher, R.B., Rugh, C.L., Kim, T., Heaton, A.C.P. & Merkle, S.A. 2006. Expression of organomercurial lyase in eastern cottonwood enhances organomercury resistance. *In vitro Cellular & Developmental Biology-Plant*, 42: 228–234.
- Chen, J., Dai, L.-Y., Wang, X.-P., Tian, Y.-C. & Lu, M.-Z. 2005. The *cry3Aa* gene of *Bacillus thuringiensis* Bt886 encodes a toxin against long-horned beetles. *Applied Microbiology and Biotechnology*, 67: 351–356.
- Chupeau, M.-C., Pautou, V. & Chupeau, Y. 1994. Recovery of transgenic trees after electroporation of poplar protoplasts. *Transgenic Research*, 3: 13–19.
- Cobbett, C.S. 2000. Phytochelatins and their roles in heavy metal detoxification. *Plant Physiology*, 123: 825–832.
- Coleman, H.D., Ellis, D., Gilbert, M. & Mansfield, S.D. 2006. Up-regulation of sucrose synthase and UDP-glucose pyrophosphorylase impacts plant growth and metabolism. *Plant Biotechnology*, 4: 87–101.
- Coleman, H.D., Canam, T., Kang, K.Y., Ellis, D.D. & Mansfield, S.D. 2007. Overexpression of UDP-glucose pyrophosphorylase in hybrid poplar affects carbon allocation. *Journal of Experimental Botany*, 58: 4257–4268.
- Comai, L., Sen, L.C. & Stalker, D.M. 1983. An altered *aroA* gene product confers resistance to the herbicide glyphosate. *Science*, 221: 370–371.
- Confalonieri, M., Allegro, G., Balestrazzi, A., Fogher, C. & Delledonne, M. 1998. Regeneration of *Populus nigra* transgenic plants expressing a Kunitz proteinase inhibitor (*KTi3*) gene. *Molecular Breeding*, 4: 137–145.
- Confalonieri, M., Belenghi, B., Balestrazzi, A., Negri, S., Facciotto, G., Schenone, G. & Delledonne, M. 2000. Transformation of elite white poplar (*Populus alba* L.) cv. 'Villafranca' and evaluation of herbicide resistance. *Plant Cell Reports*, 10: 978–982.
- Costa, A.K., Katz, I.D. & Ivanetich, K.M. 1980. Trichloroethylene: its interaction with hepatic microsomal cytochrome P-450 *in vitro*. *Biochemical Pharmacology*, 29: 433–439.
- Creissen, G., Broadbent, P., Stevens, R., Wellburn, A. & Mullineaux, P. 1996. Manipulation of glutathione metabolism in transgenic plants. *Biochemical Society Transactions*, 24: 465–469.

- Cushman, J.C. & Bohnert, H.J. 2000. Genomic approaches to plant stress tolerance. *Current Opinion in Plant Biology*, 3: 117–124.
- Dauwe, R., Morreel, K., Goeminne, G., Gielen, B., Rohde, A., Van Beeumen, J., Ralph, J., Boudet, A.-M., Kopka, J., Rochange, S.F., Halpin, C., Messens, E. & Boerjan, W. 2007. Molecular phenotyping of lignin-modified tobacco reveals associated changes in cell-wall metabolism, primary metabolism, stress metabolism and photorespiration. *Plant Journal*, 52: 263–285.
- Davis, M.F., Tuskan, G.A., Payne, P., Tschaplinski, T.J. & Meilan, R. 2006. Assessment of *Populus* wood chemistry following the introduction of a Bt toxin gene. *Tree Physiology*, 26: 557–564.
- De Block, M. 1990. Factors influencing the tissue culture and the *Agrobacterium tumefaciens*-mediated transformation of hybrid aspen and poplar clones. *Plant Physiology*, 93: 1110–1116.
- De Kam, M. 1984. *Xanthomonas campestris* pv. *populi*, the causal agent of bark necrosis in poplar. *European Journal of Plant Pathology*, 90: 13–22.
- Delauney, A.J. & Verma, D.P.S. 1993. Proline biosynthesis and osmoregulation in plants. *The Plant Journal*, 4: 215–223.
- Delledonne, M., Allegro, G., Belenghi, B., Balestrazzi, A., Picco, F., Levine, A., Zelasco, S., Calligari, P. & Confalonieri, M. 2001. Transformation of white poplar (*Populus alba* L.) with a novel *Arabidopsis thaliana* cysteine proteinase inhibitor and analysis of insect pest resistance. *Molecular Breeding*, 7: 35–42.
- Dhankher, O.P., Li, Y., Rosen, B. P, Shi, J., Salt, D., Senecoff, J.F., Sashti, N.A. & Meagher, R.B. 2002. Engineering tolerance and hyperaccumulation of arsenic in plants by combining arsenate reductase and gamma-glutamylcysteine synthetase expression. *Nature Biotechnology*, 20: 1140–1145.
- Di Baccio, D., Tognetti, R., Sebastiani, L. & Vitagliano, C. 2003. Responses of *Populus deltoides* × *Populus nigra* (*Populus* × *euramericana*) clone I-214 to high zinc concentrations. *New Phytologist*, 159: 443–452.
- DiCosty, U.R. & Whalon, M.E. 1997. Selection of Colorado potato beetle resistant to *CryIIIa* on transgenic potato plants. *Resistant Pest Management Newsletter*, 9: 33–34.
- Donahue, R.A., Davis, T.D., Michler, C.H., Riemenschneider, D.E., Carter, D.R., Marquardt, P.E., Sankhla, N., Sankhla, D., Haissig, B.E. & Isebrands, J.G. 1994. Growth, photosynthesis, and herbicide tolerance of genetically modified hybrid poplar. *Canadian Journal of Forest Research*, 24: 2377–2383.
- Doty, S.L., Shang, Q.T., Wilson, A.M., Tangen, J., Westergreen, A., Newman, L.A., Strand, S.E. & Gordon, M.P. 2000. Enhanced metabolism of halogenated hydrocarbons in transgenic plants containing mammalian P450 2E1. *Proceedings of the National Academy of Sciences of the United States of America*, 97:6287–6291.
- Doty S.L., James C.A., Moore A.L., Vajzovic A., Singleton G.L., Ma C., Khan Z., Xin G., Kang J.W., Park J.Y., Meilan R., Strauss S.H. and Strand S.E. 2007. Phytoremediation of volatile environmental pollutants with transgenic trees. *Proceedings of the National Academy of Sciences of the United States of America*, 104:16816–16821.
- Edwards, R., Dixon, D.P. & Walbot, V. 2000. Plant glutathione S-transferases: enzymes with multiple functions in sickness and in health. *Trends in Plant Science*, 5: 193–198.
- Ehling, J., Mattheus, N., Aeschliman, D.S., Li, E.Y., Hamberger, B., Cullis, I.F., Zhuang, J., Kaneda, M., Mansfield, S.D., Samuels, L., Ritland, K., Ellis, B.E., Bohlmann, J. & Douglas, C.J. 2005. Global transcript profiling of primary stems from *Arabidopsis thaliana* identifies candidate genes for missing links in lignin biosynthesis and transcriptional regulators of fiber differentiation. *Plant Journal*, 42: 618–640.
- El-Khatib, R.T., Hamerlynck, E.P., Gallardo, F. & Kirby, E.G. 2004. Transgenic poplar characterized by ectopic expression of a pine cytosolic glutamine synthetase gene exhibits enhanced tolerance to water stress. *Tree Physiology*, 24: 729–736.
- Ellis, D.D., McCabe, D.E., McInnis, S., Ramachandran, R., Russell, D.R., Wallace, K.M., Martinell, B.J., Roberts, D.R., Raffa, K.F. & McCown, B.H. 1993. Stable transformation of *Picea glauca* by particle acceleration. *Bio-Technology*, 11: 84–89.

- Eriksson, M.E., Israellsson, M., Olsson, O. & Moritz, T. 2000. Increased gibberellin biosynthesis in transgenic trees promotes growth, biomass production and xylem fiber length. *Nature Biotechnology*, 18: 784–788.
- FAO. 2004. Preliminary review of biotechnology in forestry, including genetic modification. Forest Genetic Resources Working Paper FGR/59E. Forest Resources Development Service, Forest Resources Division. Rome (available at [www.fao.org/docrep/008/ae574e/ae574e00.htm](http://www.fao.org/docrep/008/ae574e/ae574e00.htm)).
- Farago, S. & Brunold, C. 1994. Regulation of thiol contents in maize roots by intermediates and effectors of glutathione synthesis. *Journal of Plant Physiology*, 144: 433–437.
- Fillatti, J.J., Kiser, J., Rose, R. & Comai, L. 1987a. Efficient transfer of a glyphosate tolerance gene into tomato using a binary *Agrobacterium tumefaciens* vector. *Bio-Technology*, 5: 726–730.
- Fillatti, J.J., Sellmer, J., McCown, B., Haissig, B. & Comai, L. 1987b. *Agrobacterium* mediated transformation and regeneration of *Populus*. *Molecular and General Genetics*, 206: 192–199.
- Fladung, M. 1990. Transformation of diploid and tetraploid potato clones with the *rolC* gene of *Agrobacterium rhizogenes* and characterization of transgenic plants. *Plant Breeding*, 104: 295–304.
- Fladung, M., Grossmann, K. & Ahuja, M.R. 1997. Alterations in hormonal and developmental characteristics in transgenic *Populus* conditioned by the *rolC* gene from *Agrobacterium rhizogenes*. *Journal of Plant Physiology*, 150: 420–427.
- Foyer, C.H., Noctor, G., Lelandais, M., Lescure, J.C., Valadier, M.H., Boutin, J.P. & Horton, P. 1994. Short-term effects of nitrate, nitrite and ammonium assimilation on photosynthesis, carbon partitioning and protein-phosphorylation in maize. *Planta*, 192: 211–220.
- Foyer, C.H., Souriau, N., Perret, S., Lelandais, M., Kunert, K.J., Pruvost, C. & Jouanin, L. 1995. Overexpression of glutathione-reductase but not glutathione synthetase leads to increases in antioxidant capacity and resistance to photo-inhibition in poplar trees. *Plant Physiology*, 109: 1047–1057.
- Franke, R., McMichael, C.M., Meyer, K., Shirley, A.M., Cusumano, J.C. & Chapple, C. 2000. Modified lignin in tobacco and poplar plants overexpressing the *Arabidopsis* gene encoding ferulate 5-hydroxylase. *The Plant Journal*, 22: 223–234.
- French, C.E., Rosser, S.J., Davies, G.J., Nicklin, S. & Bruce, N.C. 1999. Biodegradation of explosives by transgenic plants expressing penta-erythritol tetranitrate reductase. *Nature Biotechnology*, 17: 491–494.
- Fu, J., Sampalo, R., Gallardo, F., Cánovas, F.M. & Kirby, E.G. 2003. Assembly of a cytosolic pine glutamine synthetase holoenzyme in leaves of transgenic poplar leads to enhanced vegetative growth in young plants. *Plant, Cell & Environment*, 26: 411–418.
- Fuchs, M. & Gonsalves, D. 2007. Safety of virus-resistant transgenic plants two decades after their introduction: lessons from realistic field risk assessment studies. *Annual Review of Phytopathology*, 45: 173–202.
- Fuentes, S.I., Allen, D.J., Ortiz-Lopez, A. & Hernandez, G. 2001. Overexpression of cytosolic glutamine synthetase increases photosynthesis and growth at low nitrogen concentrations. *Journal of Experimental Botany*, 52: 1071–1081.
- Gallardo, F., Fu, J., Cantón, F.R., García-Gutiérrez, A., Cánovas, F.M. & Kirby, E.G. 1999. Expression of a conifer glutamine synthetase gene in transgenic poplar. *Planta*, 210: 19–26.
- Genissel, A., Leple, J.-C., Millet, N., Augustin, S., Jouanin, L. & Pilate, G. 2003. High tolerance against *Chrysomela tremulae* of transgenic poplar plants expressing a synthetic *cry3Aa* gene from *Bacillus thuringiensis* ssp. *tenebrionis*. *Molecular Breeding*, 11: 103–110.
- Georges, F., Saleem, M. & Cutler, A.J. 1990. Design and cloning of a synthetic gene for the flounder antifreeze protein and its expression in plant cells. *Gene*, 91: 159–165.
- Giorcelli, A., Sparvoli, F., Mattivi, F., Tava, A., Balestrazzi, A., Vrhovsek, U., Calligari, P., Bollini, R. & Confalonieri, M. 2004. Expression of the stilbene synthase (StSy) gene from grapevine in transgenic white poplar results in high accumulation of the antioxidant resveratrol glucosides. *Transgenic Research*, 13: 203–214.

- Gleeson, D., Lelu-Walter, M.-A. & Parkinson, M. 2004. Influence of exogenous L-proline on embryonic cultures of larch (*Larix leptoeuropaea*), Sitka spruce (*Picea sitchensis* (Bong.) Carr.) and oak (*Quercus robur* L.) subjected to cold and salt stress. *Annals of Forest Science*, 61: 125–128.
- Gleeson, D., Lelu-Walter, M.A. and Parkinson M. 2005. Overproduction of proline in transgenic hybrid larch (*Larix × leptoeuropaea* (Dengler)) cultures renders them tolerant to cold, salt and frost. *Molecular Breeding*, 15: 21–29.
- Goicoechea, M., Lacombe, E., Legay, S., Mihaljevic, S., Rech, P., Jauneau, A., Lapierre, C., Pollet, B., Verhaegen, D., Chaubet-Gigot, N. & Grima-Pettenati, J. 2005. EgMYB2, a new transcriptional activator from Eucalyptus xylem, regulates secondary cell wall formation and lignin biosynthesis. *The Plant Journal*, 43: 553–567.
- Gould, F. 1998. Sustainability of transgenic insecticidal cultivars: integrating pest genetics and ecology. *Annual Review of Entomology*, 43: 701–726.
- Grace, L.J., Charity, J.A., Gresham, B., Kay, N. & Walter, C. 2005. Insect-resistant transgenic *Pinus radiata*. *Plant Cell Reports*, 24: 103–111.
- Gray-Mitsumune, M., Blomquist, K., McQueen-Mason, S., Teeri, T.T., Sundberg, B. & Mellerowicz, E.J. 2007. Ectopic expression of a wood-abundant expansin *PttEXPA1* promotes cell expansion in primary and secondary tissues in aspen. *Plant Biotechnology Journal*, 6: 62–72.
- Grunwald, C., Deutsch, F., Eckstein, D. & Fladung, M. 2000. Wood formation in *rolC* transgenic aspen trees. *Trees*, 14: 297–304.
- Gullner, G., Komives, T. & Rennenberg, H. 2001. Enhanced tolerance of transgenic poplar plants overexpressing gamma-glutamylcysteine synthetase towards chloroacetanilide herbicides. *Journal of Experimental Botany*, 52: 971–979.
- Halpin, C., Knight, M.E., Foxon, G.A., Campbell, M.M., Boudet, A.M., Boon, J.J., Chabbert, B., Tollier, M.-T. & Schuch, W. 1994. Manipulation of lignin quality by downregulation of cinnamyl alcohol dehydrogenase. *The Plant Journal*, 6: 339–350.
- Halpin, C., Thain, S., Tilston, E., Guiney, E., Lapierre, C. & Hopkins, D. 2007. Ecological impacts of trees with modified lignin. *Tree Genetics & Genomes*, 3: 101–110.
- Han, K.H., Meilan, R., Ma, C. & Strauss, S.H. 2000. An *Agrobacterium* transformation protocol effective in a variety of cottonwood hybrids (genus *Populus*). *Plant Cell Reports*, 19: 315–320.
- Hancock, J.E., Loya, W.M., Giardina, C.P., Li, L., Chiang, V.L. & Pregitzer, K.S. 2007. Plant growth, biomass partitioning and soil carbon formation in response to altered lignin biosynthesis in *Populus tremuloides*. *New Phytologist*, 173: 732–742.
- Hannink, N., Rosser, S.J., French, C.E., Basran, A., Murray, J.A.H., Nicklin, S. & Bruce, N.C. 2001. Phytodetoxification of TNT by transgenic plants expressing a bacterial nitroreductase. *Nature Biotechnology*, 19: 1168–1172.
- Harcourt, R.L., Kyozuka, J., Floyd, R.B., Bateman, K.S., Tanaka, H., Decroocq, V., Llewellyn, D.J., Zhu, X., Peacock, W.J. & Dennis, E.S. 2000. Insect- and herbicide-resistant transgenic eucalypts. *Molecular Breeding*, 6: 307–315.
- Harrison, E.J., Bush, M., Plett, J.M., McPhee, D.P., Vitez, R., O'Malley, B., Sharma, V., Bosnich, W., Séguin, A., MacKay, J. & Regan, S. 2007. Diverse developmental mutants revealed in an activation-tagged population of poplar. *Canadian Journal of Botany*, 85: 1071–1081.
- Hart, E.R., James, R.R., Nebeker, T.E., Robison, D.J., Raffa, K.F. & Wagner, M.A. 1996. Entomological research in North American *Populus* and *Salix*: an overview. In: Proceedings of the International Poplar Commission, Budapest, Hungary, 1–4 October 1996. 14 p.
- Haworth, R.H. & Spiers, A.G. 1988. Characterisation of bacteria from poplars and willows exhibiting leaf spotting and stem cankering in New Zealand. *Forest Pathology*, 18: 426–436.
- Hedden, P. & Proebsting, W. 1997. Gibberellin biosynthesis: enzymes, genes and their regulation. *Annual Review of Plant Physiology and Plant Molecular Biology*, 48: 431–460.
- Herschbach, C., Jouanin, L. & Rennenberg, H. 1998. Overexpression of  $\gamma$ -glutamylcysteine synthetase, but not of glutathione synthetase, elevates glutathione allocation in the phloem of transgenic poplar trees. *Plant Cell Physiology*, 39: 447–451.



- Herschbach, C. & Kopriva, S. 2002. Transgenic trees as tools in tree and plant physiology. *Trees - Structure and Function*, 16: 250–261.
- Hirel, B., Bertin, P., Quillere, I., Bourdoncle, W., Attagnant, C., Dellay, C., Gouy, A., Cadiou, S., Retailiau, C., Falque, M. & Gallais, A. 2001. Towards a better understanding of the genetic and physiological basis for nitrogen use efficiency in maize. *Plant Physiology*, 125: 1258–1270.
- Hiriga, S., Sasaki, K., Ito, H., Ohashi, T. & Matsu, H. 2001. A large family of Class III plant peroxidases. *Plant Cell Physiology*, 42: 462–468.
- Hu, W.-J., Harding, S.A., Lung, J., Popko, J.L., Ralph, J., Stokke, D.D., Tsai, C.-J. & Chiang, V.L. 1999. Repression of lignin biosynthesis promotes cellulose accumulation and growth in transgenic trees. *Nature*, 17: 808–812.
- Hu, J.J., Tian, Y.C., Han, Y.F., Li, L. & Zhang, B.E. 2001. Field evaluation of insect-resistant transgenic *Populus nigra* trees. *Euphytica*, 121: 123–127.
- Hu, L., Lu, H., Liu, Q., Chen, X. & Jiang, X. 2005. Overexpression of *mtlD* gene in transgenic *Populus tomentosa* improves salt tolerance through accumulation of mannitol. *Tree Physiology*, 25: 1273–1281.
- Huntley, S.K., Ellis, D., Gilbert, M., Chapple, C. & Mansfield, S.D. 2003. Significant increases in pulping efficiency in C4H-F5H transformed poplars: improved chemical savings and reduced environmental toxins. *Journal of Agricultural and Food Chemistry*, 51: 6178–6183.
- Jaglo-Ottosen, K.R., Gilmour, S.J., Zarka, D.G., Schabenberger, O. & Thomashow, M.F. 1998. *Arabidopsis* CBF1 overexpression induces *COR* genes and enhances freezing tolerance. *Science*, 280: 104–106.
- James, R.R. 1997. Utilizing a social ethics toward the environment in assessing genetically engineered insect-resistance in trees. *Agriculture Human Values*, 14: 237–249.
- James, R.R., Croft, B.A. & Strauss, S.H. 1999. Susceptibility of the cottonwood leaf beetle (*Coleoptera: Chrysomelidae*) to different strains and transgenic toxins of *Bacillus thuringiensis*. *Environmental Entomology*, 28: 108–109.
- Jia, C.H., Zhao, H.Y., Wang, H.Z., Xing, Z.F., Du, K.J. & Song, Y.R. 2004. Repressive expression of *4CL* gene and decrease in lignin content in transgenic *Populus tomentosa*. (in Chinese with an English abstract) *China Science Bulletin*, 49: 662–666.
- Jing, Z.P., Gallardo, F., Pascual, M.B., Sampalo, R., Romero, J., de Navarra, A.T. & Canovas, F.M. 2004. Improved growth in a field trial of transgenic hybrid poplar overexpressing glutamine synthetase. *New Phytologist*, 164: 137–145.
- Joshi, C.P. & Mansfield, S.D. 2007. The cellulose paradox—simple molecule, complex biosynthesis. *Current Opinion in Plant Biology*, 10: 220–226.
- Jouanin, L., Goujon, T., de Nadai, V., Martin, M.-T., Mila, I., Vallet, C., Pollet, B., Yoshinaga, A., Chabbert, B., Petit-Conil, M. & Lapierre, C. 2000. Lignification in transgenic poplars with extremely reduced caffeic acid O-methyltransferase activity. *Plant Physiology*, 123: 1363–1374.
- Kano-Murakami, Y., Yanai, T., Tagiri, A. & Matsuoka, M. 1993. A rice homeotic gene, *OSH1*, causes unusual phenotypes in transgenic tobacco. *FEBS Letters*, 334: 365–368.
- Karnosky, D.F., Podila, G.K., Shin, D. & Riemenschneider, D.E. 1997. Differential expression of *aroA* gene in transgenic poplar: influence of promoter and ozone stress. pp. 70–73, in: N.B. Klopfenstein, Y.-W. Chun, M.-S. Kim and M.R. Ahuja (editors). *Micropropagation, genetic engineering, and molecular biology of Populus*. USDA, Forest Service, Rocky Mountain Forest and Range Experimental Station, Fort Collins, Colorado, USA
- Kasuga, M., Liu, Q., Miura, S., Yamaguchi-Shinozaki, K. & Shinozaki, K. 1999. Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor. *Nature Biotechnology*, 17: 287–291.
- Kavi Kishor, P.B., Hong, Z., Miao, G.-H., Hu, C. & Verma, D.P.S. 1995. Overexpression of  $\Delta$ -pyrroline-5-carboxylate synthetase increases proline production and confers osmotolerance in transgenic plants. *Plant Physiology*, 108: 1387–1394.



- Kawaoka, A., Matsunaga, E., Endo, S., Kondo, S., Yoshida, K., Shinmyo, A. & Ebinuma, H. 2003. Ectopic expression of a horseradish peroxidase enhances growth rate and increases oxidative stress resistance in hybrid aspen. *Plant Physiology*, 132: 1177–1185.
- Kawazu, T. 2004. Development of environmental-stress-tolerant eucalyptus and forest plantations. *Japan TAPPI Journal*, 58: 55–61.
- Kelley, S.S., Rials, T.G., Snell, R., Groom, L.H. & Sluiter, A. 2004. Use of near-infrared spectroscopy to measure the chemical and mechanical properties of solid wood. *Wood Science and Technology*, 38: 257–276.
- Kende, H. & Zeevaart, J. 1997. The five “classical” plant hormones. *Plant Cell*, 9: 1197–1210.
- Kent, S.M., Leichti, R.J., Morrell, J.J., Rosowsky, D.V. & Kelley, S.S. 2006. Analytical tools to predict changes in properties of oriented strandboard exposed to the fungus *Postia placenta*. *Holzforschung*, 60: 332–338.
- Kleiner, K.W., Ellis, D.D., McCown, B.H. & Raffa, K.F. 2003. Leaf ontogeny influences leaf phenolics and the efficacy of genetically expressed *Bacillus thuringiensis cry1A(a)* d-endotoxin in hybrid poplar against gypsy moth. *Journal of Chemical Ecology*, 29: 2585–2602.
- Knight, H. & Knight, M.R. 2001. Abiotic stress signalling pathways: specificity and cross-talk. *Trends in Plant Science*, 6: 262–267.
- Knowles, B.H. & Dow, J.A.T. 1993. The crystal  $\beta$ -endotoxins of *Bacillus thuringiensis*: Models for their mechanism of action on the insect gut. *BioEssays*, 15: 469–476.
- Ko, J.-H., Yang, S.H., Park, A.H., Lerouxel, O. & Han, K.-H. 2007. ANAC012, a member of the plant-specific NAC transcription factor family, negatively regulates xylary fiber development in *Arabidopsis thaliana*. *The Plant Journal*, 50: 1035–1048.
- Koprivova, A., Kopriva, S., Jager, D., Will, B., Jouanin, L. & Rennenberg, H. 2002. Evaluation of transgenic poplars overexpressing enzymes of glutathione synthesis for phytoremediation of cadmium. *Plant Biology*, 4: 664–670.
- Kress, W. & Skelly, J.M. 1982. Response of several eastern forest tree species to doses of ozone and nitrogen dioxide. *Plant Disease*, 66: 1149–1152.
- Kusaba, S., Kano-Murakami, Y., Matsuoka, M., Tamaoki, M., Sakamoto, T., Yamaguchi, I. & Fukumoto, M. 1998. Alteration of hormone levels in transgenic tobacco plants overexpressing the rice homeobox gene *OSH1*. *Plant Physiology*, 116: 471–476.
- Labbe, N., Rials, T.G., Kelley, S.S., Cheng, Z.M., Kim, J.Y. & Li, Y. 2005. FT-IR imaging and pyrolysis-molecular beam mass spectrometry: new tools to investigate wood tissues. *Wood Science and Technology*, 39(1): 61–76.
- Lachance, D., Hamel, L.-P., Pelletier, F., Valero, J., Bernier-Cardou, M., Chapman, K., van Frankenhuyzen, K. & Seguin, A. 2007. Expression of a *Bacillus thuringiensis cry1Ab* gene in transgenic white spruce and its efficacy against the spruce budworm (*Choristoneura fumiferana*). *Tree Genetics & Genomes*, 3: 153–167.
- Lam, H.-M., Coschigano, K.T., Oliveira, I.C., Melo-Oliveira, R. & Coruzzi, G.M. 1996. The molecular genetics of nitrogen assimilation into amino acids in higher plants. *Annual Review of Plant Physiology and Plant Molecular Biology*, 47: 569–593.
- Lapierre, C., Pollet, B., Petit-Conil, M., Toval, G., Romero, J., Pilate, G., Leple, J.-C., Boerjan, W., Ferret, V., De Nadai, V. & Jouanin, L. 1999. Structural alterations of lignins in transgenic poplars with depressed cinnamyl alcohol dehydrogenase or caffeic acid O-methyltransferase activity have an opposite impact on the efficiency of industrial Kraft pulping. *Plant Physiology*, 119: 153–164.
- LeDuc, D.L., Abdel Samie, M., Montes-Bayon, M., Wu, C.P., Reisinger, S.J. & Terry, N. 2006. Overexpressing both ATP sulfurylase and selenocysteine methyltransferase enhances selenium phytoremediation traits in Indian mustard. *Environmental Pollutants*, 144: 70–76.
- Lee, K.H., Isenhardt, T.M. & Schultz, R.C. 2003. Sediment and nutrient removal in an established multi-species riparian buffer. *Journal of Soil and Water Conservation*, 58: 1–8.
- Legay, S., Lacombe, E., Goicoechea, M., Briere, C., Seguin, A., Mackay, J. & Grima-Pettenati, J. 2007. Molecular characterization of EgMYB1, a putative transcriptional repressor of the lignin biosynthetic pathway. *Plant Science*, 173: 542–549.

- Lelieveld, J. & Crutzen, P.J. 1990. Influences of cloud photochemical processes on tropospheric ozone. *Nature*, 343: 227–232.
- Leple, J.-C., Brasileiro, A.C.M., Michel, M.F., Delmotte, F. & Jouanin, L. 1992. Transgenic poplars: expression of chimeric genes using four different constructs. *Plant Cell Reports*, 11: 137–141.
- Leple, J.-C., Dauwe, R., Morreel, K., Strome, V., Lapierre, C., Pollet, B., Naumann, A., Kang, K.-Y., Kim, H., Ruel, K., Lefebvre, A., Joseleau, J.-P., Grima-Pettenati, J., De Rycke, R., Andersson-Gunnerase, S., Erban, A., Fehrle, I., Petit-Conil, M., Kopka, J., Polle, A., Messens, E., Sundberg, B., Mansfield, S.D., Ralph, J., Pilate, G. & Boerjan, W. 2007. Downregulation of cinnamoyl-coenzyme A reductase in poplar: multiple-level phenotyping reveals effects on cell wall polymer metabolism and structure. *The Plant Cell*, 19: 3669–3691.
- Li, L., Zhou, Y., Cheng, X., Sun, J., Marita, J.M., Ralph, J. & Chiang, V.L. 2003. Combinatorial modification of multiple lignin traits in trees through multigene co-transformation. *Proceedings of the National Academy of Sciences of the United States of America*, 100: 4939–4944.
- Li, J., Meilan, R., Ma, C., Barish, M. & Strauss, S.H. 2008. Stability of herbicide resistance over 8 years of coppice in field-grown, genetically engineered poplars. *Western Journal of Applied Forestry*, 23(2): 89–93.
- Liang, H., Maynard, C.A., Allen, R.D. & Powell, W.A. 2001. Increased *Septoria musiva* resistance in transgenic hybrid poplar leaves expressing a wheat oxalate oxidase gene. *Plant Molecular Biology*, 45: 619–629.
- Liang, H., Catranis, C.M., Maynard, C.A. & Powell, W.A. 2002. Enhanced resistance to the poplar fungal pathogen, *Septoria musiva*, in hybrid poplar clones transformed with genes encoding antimicrobial peptides. *Biotechnology Letters*, 24: 383–389.
- Lin, S.Z. & Zhang, Z.Y. 2004. *Studies on antifreeze mechanism and molecular biology in poplar* (in Chinese). China Environment and Science Press, Beijing, China.
- Lin, S.-Z., Zhang, Z.-Y., Zhang, Q. & Lin, Y.-Z. 2006. Progress in the study of molecular genetic improvements of poplar in China. *Journal of Integrative Plant Biology*, 48: 1001–1007.
- Little, C.H.A. & Savidge, R. 1987. The role of plant growth regulators in forest tree cambial growth. *Plant Growth Regulators*, 6: 137–169.
- Liu, F.H., Sun, Z.X., Cui, D.C., Du, B.X., Wang, C.R. & Chen, S.Y. 2000. Cloning of *E. coli mtl-D* genes and its expression in transgenic Balizhuangyang (*Populus*) (in Chinese with an English abstract). *Acta Genetica Sinica*, 27: 428–433.
- Liu, B., Li, H.S., Wang, Q.H. & Cui, D.C. 2002. Transformation of *Populus tomentosa* with anti-PLD gene (in Chinese with an English abstract). *Heredity*, 24: 40–44.
- Lius, S., Manshardt, R.M., Fitch, M.M.M., Slightom, J.L., Sanford, J.C. & Gonsalves, D. 1997. Pathogen-derived resistance provides papaya with effective protection against papaya ringspot virus. *Molecular Breeding*, 3: 161–168.
- Lu, S., Sun, Y.-H., Shi, R., Clark, C., Li, L. & Chiang, V.L. 2005. Novel and mechanical stress-responsive microRNAs in *Populus trichocarpa* that are absent from *Arabidopsis*. *Plant Cell*, 17: 2186–2203.
- Luttrell, R.G. & Caprio, M. 1996. Implementing resistance management. pp. 161–163 (vol. 1), in: *Proceedings of Beltwide Cotton Production Research Conference*, Nashville, Tennessee, USA, 9–12 January 1996. National Cotton Council, Memphis, Tennessee, USA.
- Lyyra, S., Meagher, R.B., Kim, T., Heaton, A., Montello, P., Balish, R.S. & Merkle, S.A. 2007. Coupling two mercury resistance genes in Eastern cottonwood enhances the processing of organomercury. *Plant Biotechnology Journal*, 5: 254–262.
- MacKay, J.J., O'Malley, D.M., Presnell, T., Booker, F.L., Campbell, M.M., Whetten, R.W. & Sederoff, R.R. 1997. Inheritance, gene expression, and lignin characterization in a mutant pine deficient in cinnamyl alcohol dehydrogenase. *Proceedings of the National Academy of Sciences of the United States of America*, 94: 8255–8260.
- Man, H., Boriel, R., El-Khatib, R. & Kirby, E.G. 2005. Characterization of transgenic poplar with ectopic expression of pine cytosolic glutamine synthetase under conditions of varying nitrogen availability *New Phytologist*, 167: 31–39.

- Mansfield, S.D. & Weineisen, H. 2007. Wood fibre quality and Kraft pulping efficiencies of trembling Aspen (*Populus tremuloides* Michx) clones. *Journal of Wood Chemistry and Technology*, 27: 35–151.
- Maredia, K.M. & Mihm, J.A. 1997. Sustaining host plant resistance derived through conventional and biotechnological means. pp. 175–179, in: *Insect resistant maize: recent advances and utilization*. Proceedings of the International Symposium, CIMMYT, 27 November–3 December 1994. CIMMYT, Mexico City, Mexico.
- Marrs, K.A. 1996. The functions and regulation of glutathione S-transferases in plants. *Annual Review of Plant Physiology and Plant Molecular Biology*, 47: 127–158.
- Matsuoka, M., Ichikawa, H., Saito, A., Tada, A., Fujimura, T. & Kano-Murakami, Y. 1993. Expression of a rice homeobox gene causes altered morphology of transgenic plants. *Plant Cell*, 5: 1039–1048.
- McCown, B.H., McCabe, D.E., Russell, D.R., Robison, D.J., Barton, K.A. & Raffa, K.F. 1991. Stable transformation of *Populus* and incorporation of pest resistance by electric discharge particle acceleration. *Plant Cell Reports*, 9: 590–594.
- McGaughey, W.H., Gould, F. & Gelernter, W. 1998. Bt resistance management. *Nature Biotechnology*, 16: 144–146.
- Meagher, R.B. 2000. Phytoremediation of toxic elemental and organic pollutants. *Current Opinions in Plant Biology*, 3: 153–162.
- Meagher, R.B., Rugh, C.L., Kandasamy, M.K., Gragson, G. & Wang, N.J. 1998. Engineered phytoremediation of mercury pollution in soil and water using bacterial genes. pp. 203–221, in: *Fourth International Conference on the Biogeochemistry of Trace Elements*. Ann Arbor Press Inc. Berkeley, California, USA.
- Meilan, R., Ma, C., Cheng, S., Eaton, J.A., Miller, L.K., Crockett, R.P., DiFazio, S.P. & Strauss, S.H. 2000. High levels of Roundup® and leaf-beetle resistance in genetically engineered hybrid cottonwoods. pp. 29–38, in: K.A. Blatner, J.D. Johnson & D.M. Baumgartner (editors). *Hybrid poplars in the Pacific Northwest: culture, commerce and capability*. Washington State University Cooperative Extension Bulletin MISC0272, Pullman, Washington, USA.
- Meilan, R., Brunner, A., Skinner, J. & Strauss, S. 2001. Modification of flowering in transgenic trees. pp. 247–256, in: A. Komamine & N. Morohoshi (editors). *Molecular breeding of woody plants*. Elsevier Science BV, Amsterdam, Netherlands.
- Meilan, R., Auerbach, D.J., Ma, C., DiFazio, S.P. & Strauss, S.H. 2002a. Stability of herbicide resistance and GUS expression in transgenic hybrid poplars (*Populus* sp.) during four years of field trials and vegetative propagation. *Hortscience*, 37: 277–280.
- Meilan, R., Han, K.H., Ma, C., DiFazio, S.P., Eaton, J.A., Hoiem, E.A., Stanton, B.J., Crockett, R.P., Taylor, M.L., James, R.R., Skinner, J.S., Jouanin, L., Pilate, G. & Strauss, S.H. 2002b. The CP4 transgene provides high levels of tolerance to Roundup® herbicide in field-grown hybrid poplars. *Canadian Journal of Forest Research*, 32: 967–976.
- Meng, L., Li, H.S., Jin, D.M., Cui, D.C. & Wang, B. 2004. Transformation of *Populus deltoides* with CH5B gene. *Biotechnology Bulletin*, 3: 48–51.
- Mentag, R., Luckevich, M., Morency, M.-J. & Seguin, A. 2003. Bacterial disease resistance of transgenic hybrid poplar expressing the synthetic antimicrobial peptide D4E1. *Tree Physiology*, 23: 405–411.
- Meyermans, H., Morreel, K., Lapierre, C., Pollet, B., De Bruyn, A., Busson, R., Herdewijn, P., Devreese, B., Van Beeumen, J., Marita, J.M., Ralph, J., Chen, C., Burggraeve, B., Van Montagu, M., Messens, E. & Boerjan, W. 2000. Modifications in lignin and accumulation of phenolic glucosides in poplar xylem upon down-regulation of caffeoyl-coenzyme A O-methyltransferase, an enzyme involved in lignin biosynthesis. *Journal of Biological Chemistry*, 275: 36899–36909.
- Mifflin, B.J. & Lea, P.J. 1980. Ammonia assimilation. pp. 169–202, in: B.J. Mifflin (editor). *The biochemistry of plants*. Vol. 5. Academic Press, San Diego, California, USA.
- Mittler, R., Shulaev, V. & Lam, E. 1995. Coordinated activation of programmed cell death and defense mechanisms in transgenic tobacco plants expressing a bacterial proton pump. *Plant Cell*, 7: 29–42.

- Mohamed, R., Meilan, R., Ostry, M.E., Michler, C.S. & Strauss, S.H. 2001. Bacterio-opsin gene overexpression fails to elevate fungal disease resistance in transgenic poplar (*Populus*). *Canadian Journal of Forest Research*, 31: 1–8.
- Mohamed, R. 2006. Expression and function of *Populus* homologs to *TERMINAL FLOWER 1* genes: Roles in onset of flowering and shoot phenology. Ph.D. Thesis, Oregon State University, Corvallis, Oregon, USA. 136 p.
- Mohri, T., Igasaki, T., Futamura, N. & Shinohara, K. 1999. Morphological changes in transgenic poplar induced by expression of the rice homeobox gene *OSH1*. *Plant Cell Reports*, 18: 816–819.
- Murata, N., Ishizaki-Nishizawa, O., Higashi, S., Hayashi, H., Tasaka, Y. & Nishida, I. 1992. Genetically engineered alteration in the chilling sensitivity of plants. *Nature*, 365: 710–713.
- NASS [National Agricultural Statistics Service]. 2005. Papaya Acreage Survey 2005 Results. USDA, Washington, DC, USA.
- Nilsson, O., Moritz, T., Imbault, N., Sandberg, G. & Olsson, O. 1993. Hormonal characterization of transgenic tobacco plants expressing the *rolC* gene of *Agrobacterium rhizogenes* T-DNA. *Plant Physiology*, 102: 363–371.
- Nilsson, O., Moritz, T., Sundberg, B., Sandberg, G. & Olsson, O. 1996. Expression of the *Agrobacterium rhizogenes rolC* gene in a deciduous forest tree alters growth and development and leads to stem fasciation. *Plant Physiology*, 112: 493–502.
- Nishikubo, N., Awano, T., Banasiak, A., Bourquin, V., Ibatullin, F., Funada, R., Brumer, H., Teeri, T.T., Hayashi, T., Sundberg, B. & Mellerowicz, E.J. 2007. Xyloglucan endotransglycosylase (XET) functions in gelatinous layers of tension wood fibers in poplar – a glimpse into the mechanism of the balancing act of trees. *Plant Cell Physiology*, 48: 843–855.
- Noctor, G., Arisi, A.C.M., Jouanin, L. & Foyer, C.H. 1998. Manipulation of glutathione and amino acid biosynthesis in the chloroplast. *Plant Physiology*, 118: 471–482.
- Noctor, G., Arisi, A., Jouanin, L. & Foyer, C. 1999. Photorespiratory glycine enhances glutathione accumulation in both the chloroplastic and cytosolic compartments. *Journal of Experimental Botany*, 50: 1157–1167.
- Noctor, G., Strohm, M., Jouanin, L., Kunert, K.J., Foyer, C.H. & Rennenberg, H. 1996. Synthesis of glutathione in leaves of transgenic poplar overexpressing  $\gamma$ -glutamylcysteine synthetase. *Plant Physiology*, 112: 1071–1078.
- Nwanze, K.F., Seetharama, N., Sharma, H.C., Stenhouse, J.W., Frederiksen, R., Shantharam, S. & Raman, K.V. 1995. Biotechnology in pest management: improving resistance in sorghum to insect pests. Environmental impact and biosafety: issues of genetically engineered sorghum, Nairobi, Kenya. *African Crop Science Journal*, 3: 209–215.
- Obara, M., Kajiura, M., Fukuta, Y., Yano, M., Hayashi, M., Yamaya, T. & Sato, T. 2001. Mapping of QTLs associated with cytosolic glutamine synthetase and NADH-glutamate synthase in rice (*Oryza sativa* L.). *Journal of Experimental Botany*, 52: 1209–1217.
- O’Connell, A., Holt, K., Piquemal, J., Grima-Pettenati, J., Boudet, A., Pollet, B., Lapierre, C., Petit-Conil, M., Schuch, W. & Halpin, C. 2002. Improved paper pulp from plants with suppressed cinnamoyl-CoA reductase or cinnamyl alcohol dehydrogenase. *Transgenic Research*, 11: 495–503.
- Park, Y.W., Baba, K., Furuta, Y., Iida, I., Sameshima, K., Arai, M. & Hayashi, T. 2004. Enhancement of growth and cellulose accumulation by overexpression of xyloglucanase in poplar. *FEBS Letters*, 564: 183–187.
- Pascual, M.B., Jing, Z.P., Kirby, E.G., Canovas, F.M. & Gallardo, F. 2008. Response of transgenic poplar overexpressing cytosolic glutamine synthetase to phosphinothricin. *Phytochemistry*, 69(2): 382–389.
- Pasonen, H.-L., Seppenen, S.-K., Degefu, Y., Rytkenen, A., von Weissenberg, K. & Pappinen, A. 2004. Field performance of chitinase transgenic silver birches (*Betula pendula*): resistance to fungal diseases. *Theoretical and Applied Genetics*, 109: 562–570.

- Patzlaff, A., McInnis, S., Courtenay, A., Surman, C., Newman, L.J., Smith, C., Bevan, M.W., Mansfield, S., Whetten, R.W., Sederoff, R.R. & Campbell, M.M. 2003. Characterisation of a pine MYB that regulates lignification. *The Plant Journal*, 36: 743–754.
- Pilate, G., Guiney, E., Holt, K., Petit-Conil, M., Lapierre, C., Lep le, J.-C., Pollet, B., Mila, I., Webster, E.A., Marstorp, H.G., Hopkins, D.W., Jouanin, L., Boerjan, W., Schuch, W., Cornu, D. & Halpin, C. 2002. Field and pulping performances of transgenic trees with altered lignification. *Nature Biotechnology*, 20: 607–612.
- Pileggi, M., Pereiara, A.A.M., Silva, J.D., Pileggi, S.A.V. & Verma, D.P.S. 2001. An improved method for transformation of lettuce by *Agrobacterium tumefaciens* with a gene that confers freezing resistance. *Brazilian Archives of Biology and Technology*, 24: 191–196.
- Powell, W., Maynard, C., Boyle, B. & Seguin, A. 2006. Fungal and bacterial resistance in transgenic trees. pp. 235–252, in: M. Fladung & D. Ewald (editors). *Tree transgenesis: recent developments*. Springer-Verlag, Berlin, Germany.
- Ralph, J., MacKay, J.J., Hatfield, R.D., O'Malley, D.M., Whetten, R.W. & Sederoff, R.R. 1997. Abnormal lignin in a loblolly pine mutant. *Science*, 277: 235–239.
- Ralph, J., Lapierre, C., Marita, J.M., Kim, H., Lu, F., Hatfield, R.D., Ralph, S., Chapple, C., Franke, R., Hemm, M.R., Van Doorselaere, J., Sederoff, R.R., O'Malley, D.M., Scott, J.T., MacKay, J.J., Yahiaoui, N., Boudet, A.-M., Pean, M., Pilate, G., Jouanin, L. & Boerjan, W. 2001. Elucidation of new structures in lignins of CAD- and COMT-deficient plants by NMR. *Phytochemistry*, 57: 993–1003.
- Ralph, S.G., Yuen, H., Friedmann, M., Aeschliman, D., Zeznik, J.A., Nelson, C.C., Butterfired, Y.S.N., Kirkpatrick, R., Liu, J., Jones, S.J.M., Marra, M.A., Douglas, C.J., Ritland, K. & Bohlmann, J. 2006. Conifer defence against insects: micro-array gene expression profiling of Sitka spruce (*Picea sitchensis*) induced by mechanical wounding or feeding by spruce budworms (*Choristoneura occidentalis*) or white pine weevils (*Pissodes strobi*) reveals large-scale changes of the host transcriptome. *Plant, Cell and Environment*, 29: 1545–1570.
- Ranocha, P., Chabannes, M., Chamayou, S., Danoun, S., Jauneau, A., Boudet, A.M. & Goffner, D. 2002. Laccase down-regulation causes alterations in phenolic metabolism and cell wall structure in poplar. *Plant Physiology*, 129: 145–155.
- Ray, T.B. 1984. Site of action of chlorsulfuron: inhibition of valine and isoleucine biosynthesis in plants. *Plant Physiology*, 75: 827–831.
- Riemenschneider, D.E. & Haissig, B.E. 1991. Producing herbicide-tolerant *Populus* using genetic transformation mediated by *Agrobacterium tumefaciens* C58: a summary of recent research. pp. 247–263, in: M.R. Ahuja (editor). *Woody plant biotechnology*. Plenum Press, New York, USA.
- Riemenschneider, D.E., Haissig, B.E., Sellmer, J. & Fillatti, J.J. 1988. Expression of an herbicide tolerance gene in young plants of a transgenic hybrid poplar clone. pp. 73–80, in: M.R. Ahuja (editor). *Somatic cell genetics of woody plants*. Kluwer Academic Publishers, Dordrecht, Netherlands.
- Rockwood, D.L., Naidu, C.V., Carter, D.R., Rahmani, M., Spriggs, T.A., Lin, C., Alker, G.R., Isebrands, J.G. & Segrest, S.A. 2004. Short-rotation woody crops and phytoremediation: Opportunities for agroforestry? *Agroforest Systems*, 61: 51–63.
- Roush, R.T. 1997. Bt-transgenic crops: just another pretty insecticide or a chance for a new start in resistance management? *Pesticide Science*, 51: 328–334.
- Roush, R.T. & Shelton, A.M. 1997. Assessing the odds: The emergence of resistance to Bt transgenic plants. *Nature Biotechnology*, 15: 816–817.
- Rugh, C.L., Senecoff, J.F., Meagher, R.B. & Merkle, S.A. 1998. Development of transgenic yellow poplar for mercury phytoremediation. *Nature*, 16: 925–928.
- Ruiz, O.N., Hussein, S.H., Terry, N. & Daniell, H. 2003. Phytoremediation of organomercurial compounds via chloroplast genetic engineering. *Plant Physiology*, 132: 1344–1352.
- Salyaev, R., Rekoslavskaya, N., Chepinoga, A., Mapelli, S. & Pacovsky, R. 2006. Transgenic poplar with enhanced growth by introduction of the *ugt* and *acb* genes. *New Forests*, 32: 211–229.



- Sandermann, H. 1996. Ozone and plant health. *Annual Review of Phytopathology*, 34: 347–366.
- Sawahel, W.A. & Hassan, A.H. 2002. Generation of transgenic wheat plants producing high levels of the osmoprotectant proline. *Biotechnology Letters*, 4: 721–725.
- Schneider, S. & Bergmann, L. 1995. Regulation of glutathione synthesis in suspension cultures of parsley and tobacco. *Botanica Acta*, 108: 34–40.
- Schnoor, J.L., Licht, L.A., McCutcheon, S.C., Wolfe, N.L. & Carreira, L.H. 1995. Phytoremediation of contaminated soils and sediments. *Environmental Science Technology*, 29: 318–323.
- Serrano, R., Mulet, J.M., Rios, G., Marquez, J.A., de Larrinoa, I.F., Leube, M.P., Mendizabal, I., Pascual-Ahuir, A., Proft, M., Ros, R. & Montesinos, C. 1999. A glimpse of the mechanisms of ion homeostasis during salt stress. *Journal of Experimental Botany*, 50: 1023–1036.
- Sewalt, V.N., Blount, J.W., Jung, H.G., Masoud, S.A., Howles, P.A., Lamb, C. & Dixon, R.A. 1997. Reduced lignin content and altered lignin composition in transgenic tobacco down-regulated in expression of L-phenylalanine ammonia-lyase or cinnamate 4-hydroxylase. *Plant Physiology*, 115: 41–50.
- Shani, Z., Dekel, M., Tsabary, G., Goren, R. & Shoseyov, O. 2004. Growth enhancement of transgenic poplar plants by overexpression of *Arabidopsis thaliana* endo-1,4-b-glucanase (*cel1*). *Molecular Breeding*, 14: 321–330.
- Shin, D.-I., Podila, G.K., Huang, Y. & Karnosky, D.F. 1994. Transgenic larch expressing genes for herbicide and insect resistance. *Canadian Journal of Forest Research*, 24: 2059–2067.
- Sitbon, F., Sundberg, B., Olsson, O. & Sandberg, G. 1991. Free and conjugated indoleacetic acid (IAA) contents in transgenic tobacco plants expressing the *iaaM* and *iaaH* IAA biosynthesis genes from *Agrobacterium tumefaciens*. *Plant Physiology*, 95: 480–485.
- Sitbon, F., Hennion, S., Sundberg, B., Little, C.H.A., Olsson, O. & Sandberg, G. 1992. Transgenic tobacco plants co-expressing the *Agrobacterium tumefaciens iaaM* and *iaaH* genes display altered growth and indoleacetic acid metabolism. *Plant Physiology*, 99: 1062–1069.
- Skinner, J.S., Meilan, R., Ma, C.P. & Strauss, S.H. 2003. The *Populus* PTD promoter imparts floral-predominant expression and enables high levels of floral-organ ablation in *Populus*, *Nicotiana* and *Arabidopsis*. *Molecular Breeding*, 12: 119–132.
- Stewart, J.J., Kadla, J.F. & Mansfield, S.D. 2006. The influence of lignin chemistry and ultrastructure on the pulping efficiency of clonal aspen (*Populus tremuloides* Michx.). *Holzforschung*, 60: 111–122.
- Strand, S.E., Dossett, M., Harris, C., Wang, X.P. & Doty, S.L. 2005. Mass balance studies of volatile chlorinated hydrocarbon phytoremediation. *Zeitschrift für Naturforschung C-A Journal of Biosciences*, 60: 325–330.
- Strohm, M., Jouanin, L., Kunert, K.J., Pruvost, C., Polle, A., Foyer, C.H. & Rennenberg, H. 1995. Regulation of glutathione synthesis in leaves of transgenic poplar (*Populus tremula* × *alba*) overexpressing glutathione synthetase. *Plant Journal*, 7: 141–145.
- Suárez, M.F., Avila, C., Gallardo, F., Canton, F.R., Garcia-Gutierrez, A., Claros, M.G. & Canovas, F.M. 2002. Molecular and enzymatic analysis of ammonium assimilation in woody plants. *Journal of Experimental Botany*, 53: 891–904.
- Summers, A.O. 1986. Organization, expression, and evolution of genes for mercury resistance. *Annual Review of Microbiology*, 40: 607–634.
- Suzuki, S., Li, L., Sun, Y.-H. & Chiang, V.L. 2006. The cellulose synthase gene superfamily and biochemical functions of xylem-specific cellulose synthase-like genes in *Populus trichocarpa*. *Plant Physiology*, 142: 1233–1245.
- Tabashnik, B.E., Carrière, Y., Dennehy, T.J., Morin, S., Sisterson, M.S., Roush, R.T., Shelton, A.M. & Zhao, J.-Z. 2003. Insect resistance to transgenic Bt crops: lessons from the laboratory and field. *Journal of Economic Entomology*, 96: 1031–1038.
- Tang, W. & Tian, Y. 2003. Transgenic loblolly pine (*Pinus taeda* L.) plants expressing a modified  $\delta$ -endotoxin gene of *Bacillus thuringiensis* with enhanced resistance to *Dendrolimus punctatus* Walker and *Crypytothelea formosicola* Staud. *Journal of Experimental Botany*, 54: 835–844.



- Tang W., Charles T.M. and Newton R.J. 2005. Overexpression of the pepper transcription factor CaPF1 in transgenic Virginia pine (*Pinus virginiana* Mill.) confers multiple stress tolerance and enhances organ growth. *Plant Molecular Biology*, 59: 603–617.
- Tang, W., Peng, X.X. & Newton, R.J. 2005. Enhanced tolerance to salt stress in transgenic loblolly pine simultaneously expressing two genes encoding mannitol-1-phosphate dehydrogenase and glucitol-6-phosphate dehydrogenase. *Plant Physiology and Biochemistry*, 43: 139–146.
- Tang, W., Newton, R.J. & Weidner, D.A. 2007. Genetic transformation and gene silencing mediated by multiple copies of a transgene in eastern white pine. *Journal of Experimental Botany*, 58: 545–554.
- Tang, W., Newton, R.J., Li, C. & Charles, T.M. 2007. Enhanced stress tolerance in transgenic pine expressing the pepper *CaPF1* gene is associated with the polyamine biosynthesis. *Plant Cell Reports*, 26: 115–124.
- Thompson, M.A., Schnepf, H.E. & Feitelson, J.S. 1995. Structure, function and engineering of *Bacillus thuringiensis* toxins. pp. 99–117, in: J.K. Setlow (editor). *Genetic Engineering* Vol. 17. Plenum Press, New York, USA.
- Thompson, C.J., Movva, N.R., Tizard, R., Crameri, R., Davies, J.E., Lauwereys, M. & Botterman, J. 1987. Characterization of the herbicide-resistance gene *bar* from *Streptomyces hygroscopicus*. *European Molecular Biology Organization Journal*, 6: 2519–2523.
- Tiimonen, H., Aronen, T., Laakso, T., Saranpää, P., Chiang, V., Ylioja, T., Roininen, H. & Häggman, H. 2005. Does lignin modification affect feeding preference or growth performance of insect herbivores in transgenic silver birch (*Betula pendula* Roth)? *Planta*, 222: 699–708.
- Tournier, V., Grat, S., Marque, C., El Kayal, W., Penchel, R., de Andrade, G., Boudet, A.-M. & Teullières, C. 2003. An efficient procedure to stably introduce genes into an economically important pulp tree (*Eucalyptus grandis* × *Eucalyptus urophylla*). *Transgenic Research*, 12: 403–411.
- Tsai, C.-J., Popko, J.L., Mielke, M.R., Hu, W.-J., Podila, G.K. & Chiang, V.L. 1998. Suppression of O-methyltransferase gene by homologous sense transgene in quaking aspen causes red-brown wood phenotypes. *Plant Physiology*, 117: 101–112.
- Tsuchikawa, S. 2007. A review of recent near infrared research for wood and paper. *Applied Spectroscopy Reviews*, 42: 43–71.
- Tuominen, H., Sitbon, F., Jacobsson, C., Sandberg, G., Olsson, O. & Sundberg, B. 1995. Altered growth and wood characteristics in transgenic hybrid aspen expressing *Agrobacterium tumefaciens* T-DNA indoleacetic acid-biosynthesis genes. *Plant Physiology*, 109: 1179–1189.
- Tuominen, H., Puech, L., Regan, S., Fink, S., Olsson, O. & Sundberg, B. 2000. Cambial-region-specific expression of the *Agrobacterium iaa* genes in transgenic aspen visualized by a linked *uidA* reporter gene. *Plant Physiology*, 123: 531–541.
- Tuskan, G., West, D., Bradshaw, H.D., Neale, D., Sewell, M., Wheeler, N., Megraw, B., Jech, K., Wiselogle, A., Evans, R., Elam, C., Davis, M. & Dinus, R. 1999. Two high-throughput techniques for determining wood properties as part of a molecular genetics analysis of hybrid poplar and loblolly pine. *Applied Biochemistry and Biotechnology*, 77: 55–65.
- Tuskan, G.A., DiFazio, S., Jansson, S. *et al.* 2006.: The genome of black cottonwood, *Populus trichocarpa* (Torr & Gray). *Science*, 313(5793): 1596–1604.
- Tyystjärvi, E., Riikonen, M., Arisi, A.C.M., Kettunen, R., Jouanin, L. & Foyer, C.H. 1999. Photoinhibition of photosystem II in tobacco plants overexpressing glutathione reductase and poplars overexpressing superoxide dismutase. *Physiologia Plantarum*, 105: 409–416.
- Tzfira, T., Vainstein, A. & Altman, A. 1999. *rol*-gene expression in transgenic aspen (*Populus tremula*) plants results in accelerated growth and improved stem production index. *Trees*, 14: 49–54.
- Uggla, C., Moritz, T., Sandberg, G. & Sundberg, B. 1996. Auxin as a positional signal in pattern formation in plants. *Proceedings of the National Academy of Sciences of the United States of America*, 93: 9282–9286.

- Valerio, L., Carter, D., Rodrigues, J.C., Tournier, V., Gominho, J., Marque, C., Boudet, A.-M., Maunders, M., Pereira, H. & Teulieres, C. 2003. Down regulation of cinnamyl alcohol dehydrogenase, a lignification enzyme, in *Eucalyptus camaldulensis*. *Molecular Breeding*, 12: 157–167.
- Van Doorselaere, J., Baucher, M., Chognot, E., Chabbert, B., Tollier, M.-T., Petit-Conil, M., Leple, J.-C., Pilate, G., Cornu, D., Monties, B., Van Montagu, M., Inze, D., Boerjan, W. & Jouanin, L. 1995. A novel lignin in poplar trees with a reduced caffeic acid/5-hydroxyferulic acid O-methyltransferase activity. *The Plant Journal*, 8: 855–864.
- Von Schwartzberg, K., Doumas, P., Jouanin, L. & Pilate, G. 1994. Enhancement of the endogenous cytokinin concentration in poplar by transformation with *Agrobacterium* T-DNA gene *ipt*. *Tree Physiology*, 14: 27–35.
- Wadenback, J., von Arnold, S., Egertsdotter, U., Walter, M., Grima-Pettenati, J., Goffner, D., Gellerstedt, G., Gullion, T. & Clapham, D. 2008. Lignin biosynthesis in transgenic Norway spruce plants harboring an antisense construct for cinnamoyl CoA reductase (CCR). *Transgenic Research*, 17(3): 379–392.
- Wagner, A., Ralph, J., Akiyama, T., Flint, H., Phillips, L., Torr K., Nanayakkara, B. & Te Kiri, L. 2007. Exploring lignification in conifers by silencing hydroxycinnamoyl-CoA:shikimate hydroxycinnamoyltransferase in *Pinus radiata*. *Proceedings of the National Academy of Sciences of the United States of America*, 104: 11856–11861.
- Wang, W., Vinocur, B. & Altman, A. 2003. Plant responses to drought, salinity and extreme temperatures: towards genetic engineering for stress tolerance. *Planta*, 218: 1–14.
- Wang, G., Castiglione, S., Chen, Y., Li, L., Han, Y., Tian, Y., Gabriel, D.W., Han, Y., Mang, K. & Sala, F. 1996. Poplar (*Populus nigra* L.) plants transformed with a *Bacillus thuringiensis* toxin gene: insecticidal activity and genomic analysis. *Transgenic Research*, 5: 289–301.
- Weigel, D., Ahn, J.H., Blazquez, J., Borevitz, J.O., Christensen, S.K., Fankhauser, C., Ferrandiz, C., Kardailsky, I., Neff, M.M., Nguyen, J.T., Sato, S., Wang, Z., Xia, Y., Dixon, R.A., Harrison, M.J., Lab, C., Yanofsky, M.F. & Chory, J. 2000. Activation tagging in *Arabidopsis*. *Plant Physiology*, 122: 1003–1013.
- Wiklund, S., Karlsson, M., Antti, H., Johnels, D., Sjöstrom, M., Wingsle, G. & Edlund, U. 2005. A new metabonomic strategy for analysing the growth process of the poplar tree. *Plant Biotechnology Journal*, 3: 353–362.
- Wingate, V.P.M., Lawton, M.A. & Lamb, C.J. 1988. Glutathione causes a massive and selective induction of plant defense genes. *Plant Physiology*, 87: 206–210.
- Wu, N.F., Sun, Q., Yao, B., Fan, Y.L., Rao, H.Y., Huang, M.R. & Wang, M.M. 2000. Insect resistant transgenic poplar expressing AaIT gene. *Chinese Journal of Biotechnology*, 16(2): 129–133.
- Yamada-Watanabe, K., Kawaoka, A. & Matsunaga, E. 2003. Molecular breeding of *Eucalyptus*: analysis of salt tolerance of transgenic *Eucalyptus camaldulensis* that overexpress choline oxidase gene (*codA*). In: IUFRO Tree Biotechnology. 7–12 June 2003, Umea, Sweden.
- Yang, C.P., Liu, G.F., Liang, H.W. & Zhang, H. 2001. Study on the transformation of *Populus simonii* × *P. nigra* with salt resistance gene *Bet-A*. (in Chinese with an English abstract) *Scientia Silvae Sinicae*. 37: 32–35.
- Yi, S.Y., Kim, J.H., Joung, Y.H., Lee, S., Kim, W.T., Yu, S.H. & Choi, D. 2004. The pepper transcription factor CaPF1 confers pathogen and freezing tolerance in *Arabidopsis*. *Plant Physiology*, 136: 2862–2874.
- Zelasco, S., Reggi, S., Calligari, P., Balestrazzi, A., Bongiorno, C., Quattrini, E., Delia, G., Bisoffi, S., Fogher, C. & Confalonieri, M. 2006. Expression of the *Vitreoscilla* Hemoglobin (VHb)-encoding gene in transgenic white poplar: plant growth and biomass production, biochemical characterization and cell survival under submergence, oxidative and nitrosative stress conditions. *Molecular Breeding*, 17: 201–216.
- Zhang, B.-Y., Su, X.-H., Li, Y.-L., Huang, Q.-J., Zhang, X.-H. & Zhang, L. 2006. Regeneration of *vgn*-transgenic poplar (*Populus alba* × *P. glandulosa*) and the primary observation of growth. *Chinese Journal of Agricultural Biotechnology*, 3: 59–64.

- Zhao, J.Z., Fan, X.L., Shi, X.P., Zhao, R.M. & Fan, Y.L. 1997. Gene pyramiding: an effective strategy of resistance management for *Helicoverpa armigera* and *Bacillus thuringiensis*. *Resistance and Pest Management*, 9: 19–21.
- Zhao S.M., Zu, G.C., Liu, G.Q., Huang, M.R., Xu, J.X. & Sun, Y.R. 1999. Introduction of rabbit defensin NP-1 gene into poplar (*P. tomentosa*) by Agrobacterium-mediated transformation. *Acta Genetica Sinica*, 26: 711–714.
- Zhong, R.Q., Morrison, W.H., Himmelsbach, D.S., Poole, F.L. & Ye, Z.H. 2000. Essential role of caffeoyl coenzyme A O-methyltransferase in lignin biosynthesis in woody poplar plants. *Plant Physiology*, 124(2): 563–577.
- Zhu, B., Su, J., Chang, M., Verma, D.P.S., Fan, Y. & Wu, R. 1998. Overexpression of a *D-5CS* gene and analysis of tolerance to water and salt-stress in transgenic rice. *Plant Science*, 139: 41–48.
- Zhu, J.K. 2001. Plant salt tolerance. *Trends in Plant Science*, 6: 66–71.
- Zhu, J.K. 2002. Salt and drought stress signal transduction in plants. *Annual Review of Plant Biology*, 53: 247–273.

## 5. Integrating genetically modified traits into tree improvement programmes

*R.D. Burdon and M. Lstibůrek*

Forest tree breeding programmes have typically been based on the progressive population improvement that accrues in breeding populations (White, Adams and Neale, 2007). Seed orchards or clonal stool beds serve as the means of delivering genetic gain using current top-ranked selections from within breeding populations, rather than representing the actual breeding process. A breeding population is usually initiated by selecting superior phenotypes that are both intermated and subjected to evaluation (Namkoong, Kang and Brouard, 1988). After the first generation, the breeding population undergoes recurring cycles of selection, intermating, evaluation, selection and so on, to build up frequencies of favourable alleles of additive effects that will confer cumulative genetic gain. For intermating and evaluation, various crossing designs and field testing schemes may be used (Namkoong 1979; White, Adams and Neale, 2007). Of central importance here, however, is that selection has almost always been based on phenotypic data from individual candidates or relatives, or both.

There are many variations of this process of tree breeding based on cumulative population improvement. For instance, it can be implemented within a single base population, or it can involve hybridization of populations (Fins, Friedman and Brotschol, 1992; White, Adams and Neale, 2007). But, whatever the detailed form that it takes, this process is termed conventional breeding. It depends on naturally occurring genetic variation, unless supplemented by deliberate mutagenesis. And it has the limitation that, with the sexual reproduction that it entails, half of any parent's genes get passed on to offspring. That means that, along with the parent's desirable genes, many unwanted ones will almost certainly be passed on. Eliminating the undesirable genes, by further crossing and selection, is a slow and cumbersome process that will typically take many generations even to approach completion.

Genetic modification, often termed genetic engineering, removes both these limitations. It can be used to introduce DNA sequences (not necessarily complete genes) that are not available in populations of sexually compatible material, or even in nature, and the sequences can be introduced without introducing the unwanted 'genetic baggage' that sexual reproduction brings (El-Kassaby and Mansfield, 2007).

Despite these additional capabilities, genetic modification is seen not as a replacement for conventional breeding but more as a complement to it. To address

the complementarity, the technical options for genetic modification are first reviewed briefly, followed by a review of what has been entailed in conventional breeding, and an outline of how the process can be enhanced by an array of new gene technologies. The appropriate nature of the complementarity must be governed by the fact that conventional breeding entails not only the breeding proper, but also the systems for mass production of either seed from seed orchards, or of planting stock through vegetative propagation as rooted cuttings or by means of tissue culture techniques, such as organogenesis or somatic embryogenesis (El-Kassaby and Krakowski, 2004; Sutton *et al.*, 2004). These aspects are involved in a complex interplay, which must be examined.

### GENETIC MODIFICATION OPTIONS

Genetic modification relates in this context to asexual technologies for generating new genetic combinations. This may take several forms:

- Genetic transformation, involving the insertion of short and specific sequences of DNA into chromosomes, to incorporate new, functioning structural genes or to modify activity levels of 'resident' genes that are already present. Collectively, the various classes of inserted DNA sequences are all referred to here as transgenes. Genetic transformation is now very much the preferred approach for genetic modification.
- Somatic hybridization by fusing cells containing different genomes (each containing  $n$  or  $2n$  chromosomes), and regenerating whole plants from such fusion events (Ma *et al.*, 1998).
- Production of 'cybrids', containing the nuclear genome of one species or population with either or both of the organelle genomes of another (Pelletier *et al.*, 1983).

Further discussion in this chapter will focus on the case of genetic transformation, given its overwhelmingly preferred status.

### CONVENTIONAL BREEDING

Choice of appropriate base populations will remain an essential platform for any genetic improvement. It can be based on general adaptation or basic economic worth, or on special attributes that can be incorporated into domesticated stocks. The choice, once made, sets the base for the recurrent cycles, already referred to, of selection, intermating, evaluation, selection, and so forth.

The phenotypic information used in conventional breeding can be extended beyond expression of field performance. It can include performance under special screening conditions (e.g. response to frosting in growth rooms (Aitken, 2004), or response to inoculation by pathogens in laboratory conditions (Kinloch and Libby 1997). Such specific phenotypic information is effectively a more direct expression of genes of interest. However, there are various ways of screening on the basis of more immediate gene expression, such as using tissue conductance as a measure of frost resistance (Benowicz, L'Hirondelle and El-Kassaby, 2001), or assays for metabolites that are crucial steps towards phenotypic expression (Robinson *et al.*,

2007). The latter approach, however, can be bedevilled by strongly non-linear relationships, or the fact that metabolite fluxes can be more important than the concentrations that are easier to measure, or a combination.

An ideal has long been to locate, and better still identify, genes of desired effects (Neale and Ingvarsson, 2008). Locating polymorphic chromosome regions (termed quantitative trait loci (QTLs)) that exercise detectable phenotypic effects can be exploited by marker-assisted or marker-based selection (collectively, MAS). The actual genes concerned, if they can be characterized, are identified as quantitative trait nucleotides (QTNs), but this again depends on the individual genes having appreciable phenotypic effects (González-Martínez *et al.*, 2006). Past tree breeding has proceeded on the assumption that genetic control of traits of interest in forest trees is strictly polygenic, with allelic differences at individual loci exerting infinitesimal phenotypic effects (Fisher, 1941). With imperfect heritability, however, genes of quite large phenotypic effects may be present without being evident from purely phenotypic data, yet experience with various crop species indicates that such genes of quite large effect are often lurking (Thoday, 1961, 1976), undetected until recently developed genomic analysis has become possible. Such genes, however, may have been mutations favoured in a very long history of domestication (e.g. in maize; Szabó and Burr, 1996), or have been major points of differentiation among populations or even species that have been fused in the course of domestication. For forest trees, detecting such genes of substantial phenotypic effects has often been problematic, with many reported QTLs proving to have been false positives due to the hidden genetic admixture of the studied population (Pritchard, Stephens and Donnelly, 2000; Pritchard, Wen and Falush, 2007). Disease resistance, however, is a case where some major-gene effects can evidently operate in essentially wild, undomesticated populations (Wilcox *et al.*, 1996; Amerson *et al.*, 2005).

Even if new gene technologies are used to identify QTLs or QTNs that thence can serve as selection criteria, this will remain fundamentally an enhancement of classical breeding, rather than an actual application of genetic modification.

Exploiting interactions between genes, in the form of non-additive gene effects (allelic dominance or epistasis between loci, or a combination), generally falls outside the framework of long-term tree breeding. A possible exception is use of reciprocal recurrent selection (or variations thereof), which may have applications in forest trees if different species or populations are bred to produce  $F_1$  hybrids for commercially deployed stocks. Possible examples include interspecific hybrids of poplar (e.g. White, Adams and Neale, 2007), or an intervarietal hybrid of *Pinus radiata* (Low and Smith, 1997). Nevertheless, exploiting non-additive gene effects has been part of the classical breeding tradition, most notably in intercrossing distinct populations (which may even represent different species, e.g. poplars), or in hybrid maize breeding (Simmonds, 1979), which has depended on exhaustive searches for strong heterotic effects in crosses between highly inbred lines. For forest trees, only part of the total non-additive gene effects can be captured by making specific pair-crosses. Identifying the appropriate crossing combinations



can be very difficult and costly but, if this is achieved, vegetative multiplication of offspring may greatly reduce the effective costs of controlled crossing. Clonal forestry in principle offers full capture of non-additive gene effects, but they must be captured afresh each generation rather than contributing over time to cumulative genetic gain.

Where disease resistance is sought, one needs durability against pathogen-strain shifts, through either fresh mutations or changes in frequencies of existing strains. Since durability evidently depends on the presence of multiple resistance mechanisms, we have a situation where epistatic gene effects can be extremely important, yet only targeted readily if resistance-gene effects are quite large.

Population improvement, along the lines of conventional breeding, remains very much the method of choice for exploiting polygenic variation, which will probably be large for many traits in forest trees and yet virtually impossible to characterize as either QTLs or QTNs. Moreover, large populations might be needed to prevent loss of valuable alleles, especially for traits like disease or pest resistance. While genetic modification may have potential for incorporating pathogen resistance, it can face the practical difficulty that many biotic risks may not be identifiable in advance (e.g. pitch canker in Monterey pine).

In many crops, notably cereals, phenotypic uniformity can be crucial to commercial success. This has doubtless led, to varying degrees, to crops shifting to an inbreeding system with pollination being predominantly self-fertilization. Because the inbreeding leads to major purging of genetic load, there is often no obvious inbreeding depression, quite unlike the typical situation with forest trees.

### COMBINING GENETIC MODIFICATION WITH CONVENTIONAL BREEDING

Genetic modification can be combined with conventional breeding in two ways:

- Confining the application of genetic modification to clones produced by the breeding programme; in other words, confining the application to a context of clonal forestry. This basically superimposes genetic modification on the population improvement that is at the core of properly constituted tree-improvement programmes.
- Applying genetic modification to members of the breeding population, such that the inserted DNA sequences are included in the genes that are subject to the genetic recombination that naturally occurs during sexual reproduction. This would be true integration of genetic modification with conventional breeding.

Prevailing opinion (e.g. Burdon and Libby, 2006) is that, for the time being, any pursuit of operational use of genetic modification in forest trees would be based on the former option. In it, transformation would be done afresh on clones selected from untransformed and largely traditional breeding populations.

By contrast, the second option is the way in which genetic modification has been widely applied for annual crops, notably cotton, soybean, canola and maize. In it, transgenes are incorporated into what are effectively the breeding populations, and are delivered into commercial crops by sexual reproduction. It is

appropriate to examine critically the reasons for this contrast, and several reasons stand out:

- The very short generation time for annuals makes it much easier to select effectively for stable expression without adverse side effects on field fitness (although generation intervals can be greatly reduced in some tree species), and to fix the transgenes.
- With the relatively small plants of annuals it is also easier to achieve high intensities for such selection (which could be partly achievable with those forest trees in which generation intervals can be greatly foreshortened).
- Seed propagation is effectively obligate in the annuals, in contrast to many forest trees.
- The only way to achieve crop uniformity with such annuals is by use of intensive inbreeding, in contrast to clonal propagation of forest trees.
- With such inbreeding, the purging of genetic load can be a help in breeding.
- With clonal propagation there is no need to fix transgenes.
- In some of the annual crops, polyploidy would confer some protection against the genetic damage associated with insertional events, whereas the domesticated forest tree species are mostly diploid.
- In annual crops the requirements for field fitness may be less stringent than they are for forest trees.

### **SUPERIMPOSING GENETIC MODIFICATION UPON POPULATION IMPROVEMENT**

Operational use of genetic modification on preselected clones, in conjunction with appropriate risk-management protocols (below), is seen as the precautionary approach. If mass clonal propagation is possible for a species, there will be no need to have any transgenes in a homozygous state, provided they behave as dominant alleles whose presence and activity in the heterozygous state should be easy to confirm. When once identified as containing the desired transgene(s) candidate clones from the breeding population can then be subjected to the usual testing, which should confirm transgene function and lack of adverse side effects of the insertional event(s) and/or the transgene(s), as well as evaluating general performance.

If use of transformation is confined to deployed clones, the requirements for control of flowering may be limited to sterility conferred by inserted transgenes. While the feasibility of conferring complete sterility is debatable, sterility may be a regulatory requirement to assure transgene containment. If conferred successfully, it would have the bonus of avoiding diversion of important resources from producing wood, with the expectation of an increase in wood production (Burdon and Libby, 2006). Yet with genetic modification confined to commercially deployed clones, there would be no need to reverse sterility for flowering on command.

The model described above implies a tandem process of testing candidate clones for basic merit and then testing them after transformation for effect of

transformation. Simultaneous genetic transformation beginning at the seed-embryo stage or soon after it, on multiple candidate clones, to be tested jointly for both successful transformation and overall genetic merit, would pose logistical and technical challenges. There should be scope for prompt initiation of a multistage culling process, eliminating many candidates before even beginning field tests. However, testing both transformed and untransformed ramets of the same clones would seem indicated.

### FULL INTEGRATION OF GENETIC MODIFICATION INTO POPULATION IMPROVEMENT

To apply genetic modification in a context of mass propagation by seed would certainly entail full integration of genetic modification into population improvement. This, however, appears problematic, at the least. To achieve fixation of transgenes without expression of genetic damage from gene insertion, yet with stable vertical transmission through generations assured, could be very slow and difficult – much more so than achieving stable integration without adverse side effects in the heterozygous state in deployed clones. Problems of transgene containment would loom very large, given that the breeding population depends on sexual fertility, creating a strict requirement for flowering ‘on command and command only’ (Brunner *et al.*, 2007).

A context of clonal forestry would be more consistent with the higher level of domestication entailed in use of genetic modification. In this situation, requirements for integrating genetic modification with population management would seem less, but still stringent. It would not be essential to fix transgenes, provided individual offspring can be screened cheaply and early for successful vertical transmission.

An advantage of using genetic modification within a breeding population is that if a transgene is known to show stable vertical transmission, without adverse side-effects, it can be used indefinitely without risks attendant upon re-creating the transformation. At the same time, availability of new transgenes may render it obsolete.

With clonal forestry for deployed crops, only a subsample of a total breeding population would have to be subjected to genetic modification. Choice of subsample, however, might not be straightforward. On present knowledge, choice of subsample may be guided by a need to provide for some failures of vertical transmission through seed. Also, transformation will be needed in sufficient parental clones to assure an appropriate genetic base in the deployed population. The requisite number will depend, as with transformation on deployed clones, on the risk-spread needs for the operation. The needs will be governed, *inter alia*, by the status of the species concerned in any broader frame of risk spread. However, even for a large, stand-alone operation, further risk-spread protection becomes limited as numbers of deployed clones exceed 15 to 20 – the standard error of performance varies according to the inverse of the square root of the effective number of clones, but the confidence limits about expected performance would

decrease more rapidly than that. However, additional provision should be made for transformed parents not contributing to the pool of selected clones. If a breeding population is structured so as to contain specialized breeds, representing different breeding goals (cf. Jayawickrama and Carson, 1990), the particular transformations may be confined to particular breeds, to serve the specific breeding goals.

Note that with multiple transgenes, integration with the breeding population is likely to be less straightforward than with single or very few transgenes.

### **RISK-MANAGEMENT ISSUES**

Accepting that use of genetic modification for forest trees will be based on transformation of pre-selected clones, the existing risk-management protocols for clonal deployment, which are based largely on risk spread (Burdon and Aimers-Halliday, 2006) would remain essentially in place. However, these protocols would need to be supplemented by protocols for managing risks specifically associated with genetic modification (e.g. Burdon, 1999; Burdon and Walter, 2004). A lesson learned from the Southern corn blight epidemic that occurred in the United States of America in 1970 is seen as being relevant to use of genetic modification with forest trees. A massive reliance on a single organelle gene, the Texas male sterile cytoplasmic factor, which was used to simplify the production of 'hybrid maize', contributed greatly to the severity of the epidemic. This was because the gene concerned destroyed the resistance to a mutant strain of the Southern corn blight pathogen (Levings, 1990). The side-effect took many years to become manifest, but was not totally disastrous because the epidemic did not spread quickly enough to destroy crops over the entire United States of America Corn Belt, while there were sufficient stocks of unaffected varieties to meet sowing needs in the following year. The long delay in the side-effect becoming evident, if it involved forest trees, could be disastrous, as it could involve many years' plantings, instead of just one year's sowings. Admittedly, the parallel with genetic modification is by no means exact (Burdon and Walter, 2004). Whereas the cytoplasmic modification involved an organelle genome, genetic transformation is directed at the nuclear genome, although the male sterility was similar to what is often sought now through genetic modification. And, while pathways of gene action can now be traced much more readily, thus improving the chances of predicting unwanted side-effects of transgene action, massive reliance on any single transgene and any single insertion event may still pose some risks. While such risks may involve low probability events, they can be significant because they could involve extreme consequences. Achieving risk spread not only among clones, but also among transgenes to achieve a given objective and among insertion events, would pose challenges, challenges that could be much greater in the context of practising genetic modification within a breeding population.

### **NUMBERS OF GENES INVOLVED IN GENETIC MODIFICATION**

For various reasons the breeder may want to insert multiple transgenes, a measure termed stacking. Engineering sterility (if possible) has already been mentioned as

both a potential regulatory need and a likely boon to commercial productivity. There might also be cases where several other transgenic attributes are sought. However, intensive stacking has its own potential risks. Even if it can be achieved without increasing genetic damage through multiple insertion sites, the number of potential interactions between different transgenes multiplies dramatically as transgene numbers increase (Burdon, 1999). It would only take occasional significant adverse interactions to be disastrous. Speculatively, incorporating multiple transgenes into single constructs, while it may reduce insertion sites, could accentuate interactions among the transgenes.

A specific case where stacking may eventually be strongly indicated is with disease resistance, for which multiple resistance mechanisms may be needed in order to ensure durability of resistance in the face of genetic shifts on the part of a pathogen. This would achieve the same result as ‘pyramiding’ multiple resistance factors by classical breeding, which can be difficult without gene discovery because some resistance genes can mask the expression of others. Indeed, a ‘hybrid’ approach could be used, supplementing naturally occurring resistance factors with transgenic ones in order to achieve effective pyramiding (Burdon and Wilcox, 2007; Wilcox, Echt and Burdon, 2007).

Of special interest is a move to incorporate resistance to chestnut blight in American chestnut by genetic modification, with the eventual aim of restoring the species to its former ecological status (Burdon and Libby, 2006). This is a notable exception to the usual rule of genetic modification being reserved for a context of intensive domestication.

Different classes of transgenes, which include up-regulating, down-regulating and gene-silencing sequences, as well as complete alien genes, will doubtless incur different risks of adverse side-effects. A provisional categorization of risk levels for different transgene categories was given by Burdon and Walter (2004), but better knowledge of such risks will surely come with time.

A precautionary approach based on avoiding intensive stacking of transgenes should not greatly impede the application of genetic modification, because transformation is used to incorporate DNA sequences of large phenotypic effect, which is likely to mean a focus on small numbers of transgenes in any one breeding programme. Capturing polygenic variation conditioned by factors that are widely dispersed through the genome is clearly not a realistic option for genetic modification.

## DEPENDENCE ON PROPAGATION TECHNOLOGIES

Successful use of genetic modification is enormously dependent on adequate propagation technologies. Transformation itself depends on a suitable platform of *in vitro* propagation technology. In many species, major technical challenges remain in achieving transformation with recipient genotypes of choice. And even if successful transformation is achieved, the genotypes for commercial deployment need to be amenable to mass multiplication from *in vitro* culture or more conventional vegetative propagation, or both (e.g. nursery cuttings). If

amenability to *in vitro* culture, transformability and mass-multiplication from *in vitro* culture have to become selection criteria, gain in breeding-goal traits will tend to be eroded or the genetic base of deployed material will be compromised.

### THE SIGNIFICANCE OF GENE DISCOVERY

Detection and use of QTL is a vexed issue for forest trees (e.g. Wilcox, Echt and Burdon, 2007), because the typical outbreeding behaviour means that there is very limited general linkage disequilibrium (Brown *et al.*, 2004), which means in turn that QTL-marker associations are mostly pedigree-specific. Gene discovery has promise for making pervasive contributions to intensive, long-term genetic improvement of forest trees (Burdon and Wilcox, 2007; Wilcox, Echt and Burdon, 2007). Admittedly, it is likely to be relatively slow to achieve with forest trees, especially with the very large genomes of conifers. The use of association genetics with trees can be slow and expensive, since it will involve either causal DNA sequences or linkages over very limited segments of chromosomes (Neale and Savolainen, 2004). Thus, for gene discovery, it will almost certainly need to be heavily supplemented (if not supplanted) by identifying candidate genes from other plants and then confirming their roles in the trees. As indicated earlier, gene discovery, if achieved, can allow selection on the basis of genes of large effect (if they are present) within the framework of classical breeding. Alternatively, such genes when once identified can be cloned and then used for transformation. Moreover, genetic modification followed by observation of phenotype can serve to verify gene discovery.

### POTENTIAL IMPACTS OF FUTURE ADVANCES IN GENE TECHNOLOGY

Engineering sterility, preferably as complete suppression of reproductive structures, has been mentioned as an ideal. Its feasibility, though, is still controversial, but will surely be intensively researched. Gene discovery, apart from the inherent difficulties with forest trees, especially conifers, could pose major challenges (Burdon and Wilcox, 2007; Wilcox, Echt and Burdon, 2007). Some of the challenges arise from the fact that large phenotypic effects can be governed by gene regulatory sequences and possibly sequences that code for RNA with poorly understood roles in developmental processes (see references in Burdon and Wilcox, 2007).

Luo *et al.* (2007) report a technology that has promise for excising transgenes during sexual reproduction, to avoid transmission in seed or pollen. Its potential for forest trees, however, is very problematic. It would not prevent unwanted diversion of resources into reproduction, nor would it favour full integration of transgenes into breeding populations without adding a process to the actual breeding operation.

### ALLOCATION OF RESOURCES AMONG TECHNOLOGIES

In any genetic improvement we now have an array of different technologies that include genetic modification. While different technologies can be used



complementarily, and even synergistically, they can often make strongly competing demands on available resources for research and development. This will create new organizational challenges for optimizing allocation of resources among the various technologies, new and old. Whatever the potential for gene discovery, phenotypic expression will remain a necessary benchmark, if only for calibrating new screening technologies. Achieving the requisite phenotypic expression with typical forest trees will remain slow and expensive, such that intensive genetic improvement cannot in the foreseeable future be undertaken lightly.

## CONCLUSIONS

The topic of this chapter has so far been addressed only very tentatively, if at all, in operational tree breeding programmes. How the integration will actually occur will doubtless depend on many technical developments that are still very uncertain. Accordingly, much of the emphasis has been on reviewing the issues and the possibilities that will need to be explored. Much of the requisite technical knowledge is likely to advance rapidly, but a precautionary approach remains indicated on both technical and political grounds. Achieving such a precautionary approach, however, is a challenge in itself.

## REFERENCES

- Aitken, S.N. 2004. Genecology and adaptation of forest trees. Article 86, pp. 197–204, *in*: J. Evans, J. Burley and J. Youngquist (editors). *Encyclopedia of Forest Sciences*. Elsevier, Amsterdam, Netherlands.
- Amerson, H.V., Kubisiak, T.L., Garcia, S.A., Kuhlman, G.C., Nelson, C.D., McKeand, S.E., Mullin, T.J. & Li, B. 2005. Interacting genes in the pine-fusiform rust forest pathosystem. (abstract) p. 60, *in*: Proceedings of the 28th Biennial Southern Forest Tree Improvement Conference. USDA Forest Service, Raleigh, North Carolina, USA.
- Benowicz, A., L'Hirondelle, S. & El-Kassaby, Y.A. 2001. Patterns of genetic variation in mountain hemlock (*Tsuga mertensiana* (Bong.) Carr.) with respect to height growth and frost hardiness. *Forest Ecology and Management*, 154: 23–33.
- Brown, G.R., Gill, G.P., Kuntz, R.J., Langley, C.H. & Neale, D.B. 2004. Nucleotide diversity and linkage disequilibrium in loblolly pine. *Proceedings of the National Academy of Sciences of the United States of America*, 101: 15255–15260.
- Brunner, A.M., Li, J., Di Fazio, S.P., Shevchenko, O., Montgomery, B.E., Mohamed, R., Wei, H., Ma, C., Elias, A.A., Van Wormer, K. & Strauss, S.H. 2007. Genetic containment of forest plantations. *Tree Genetics and Genomes*, 3: 75–100.
- Burdon, R.D. 1999. Risk-management issues for genetically engineered forest trees. *New Zealand Journal of Forestry Science*, 29: 375–390.
- Burdon, R.D. & Aimers-Halliday, J. 2006. Managing risk in clonal forestry. *CABI Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources*, 1(035): 9. (online journal)
- Burdon, R.D. & Libby, W.J. 2006. *Genetically modified forests: from stone age to modern biotechnology*. Forest History Society, Durham, North Carolina, USA.
- Burdon, R.D. & Walter, C. 2004. Exotic pines and eucalypts: Perspectives on risks of transgenic plantations. pp. 52–75, *in*: S.H. Strauss and H.D. Bradshaw (editors). *The Bioengineered Forest: Challenges for science and society*. RFF Press, Washington, DC, USA.
- Burdon, R.D. & Wilcox, P.L. 2007. Population management: potential impacts of advances in genomics. *New Forests*, 34: 187–206.

- El-Kassaby Y.A. & Krakowski J. 2004. Methods of cloning forest trees. pp. 327–332, in: B. Li and S. McKeand (editors). *IUFRO Joint Conference of Division 2 – Forest genetics and tree breeding in the age of genomics: progress and future*. Charleston, South Carolina, USA, 1–5 November 2004 (available at [www.iufro.org/download/file/4751/4507/20200-charleston04.pdf/](http://www.iufro.org/download/file/4751/4507/20200-charleston04.pdf/)).
- El-Kassaby, Y.A. & Mansfield, S.D. 2007. The use of genetically modified trees in forests: opportunities and challenges. pp. 2–13, in: *Challenges and opportunities of forest research in the policy-making process*. The Chinese Academy of Forestry (CAF) and the International Union of Forest Research Organizations (IUFRO) Symposium, Beijing, China, 29 May 2007.
- Fins, L., Friedman, S.F. & Brotschol, J.V. (editors). 1992. *Handbook of quantitative forest genetics*. Kluwer Academic Publishers, Dordrecht, Netherlands.
- Fisher, R.A. 1941. Average excess and average effect of a gene substitution. *Annals of Eugenics (London)*, 11: 53–63.
- González-Martínez, S.C., Ersoz, E., Brown, G.R., Wheeler, N.C. & Neale, D.B. 2006. DNA sequence variation and selection of tag single-nucleotide polymorphisms at candidate genes for drought-stress response in *Pinus taeda* L. *Genetics*, 172: 1915–1926.
- Jayawickrama, K.J.S. & Carson, M.J. 2000. A breeding strategy for the New Zealand Radiata Pine Breeding Cooperative. *Silvae Genetica*, 49: 82–90.
- Kinloch, B.B. & Libby, W.J. 1997. Variation in susceptibility to western gall rust in *Pinus radiata* and *P. muricata*. pp. 108–112, in: R.D. Burdon & J.M. Moore (editors). *IUFRO '97 Genetics of radiata pine*. Proceedings of conference, 1–4 December 1997, Rotorua, New Zealand. NZ Forest Research Institute, FRI Bulletin, No. 203.
- Levings, C.S. III. 1990. The Texas cytoplasm of maize: Cytoplasmic male sterility and disease susceptibility. *Science*, 250: 942–947.
- Low, C.B. & Smith, T. 1997. Use of the Guadalupe provenance in *Pinus radiata* improvement in New Zealand. pp. 57–61, in: R.D. Burdon & J.M. Moore (editors). *IUFRO '97 Genetics of radiata pine*. Proceedings of conference, 1–4 December 1997, Rotorua, New Zealand. NZ Forest Research Institute, FRI Bulletin, No. 203.
- Luo, K., Duan, H., Zhao, D.; Zheng, X, Deng, W., Chen, Y, Stewart, C.N. Jr, McAvoy, R, Jiang, X., Wu, Y, He, A, Pei, Y. & Li, Y. 2007. 'GM-gene-deletor': fused loxP-FRT recognition sequences dramatically improve the efficiency of FLP or CRE recombinase on transgene excision from pollen and seed of tobacco plants. *Plant Biotechnology Journal*, 5: 263–274.
- Ma, Y., Weber, M., Dumont-BéBoux, N., Webber, J. & von Aderkas, P. 1998. Megagametophytes of Douglas fir (*Pseudotsuga menziesii*) and hybrid larch (*Larix × eurolepis*) in culture: multiplication of neck cells and the formation of binucleate cells. *Protoplasma*, 204(3-4): 219–225.
- Namkoong, G. 1979. *Introduction to quantitative genetics in forestry*. USDA Forestry Service Technical Bulletin No. 1588. USDA, Washington, DC, USA.
- Namkoong, G., Kang, H.C. & Brouard, J.S. 1988. *Tree breeding: principles and strategies*. Monographs in Theoretical and Applied Genetics, No. 11. Springer-Verlag, New York, USA.
- Neale, D.B. & Ingvarsson, P.K. 2008. Population, quantitative and comparative genomics of adaptation in forest trees. *Current Opinion in Plant Biology*, 11: 1–7.
- Neale, D.B. & Savolainen, O. 2004. Association genetics of complex traits in conifers. *Trends in Plant Science*, 9: 325–330.
- Pelletier, G., Primard, C., Vedel, F., Chetrit, P., Remy, R. & Rousselle, R.M. 1983. Intergeneric cytoplasmic hybridization in Cruciferae by protoplast fusion. *Molecular Genomics and Genetics*, 191: 244–250.
- Pritchard, J.K., Stephens, M. & Donnelly, P. 2000. Inference of population structure using multilocus genotype data. *Genetics*, 155: 945–959.
- Pritchard, J.K., Wen, W. & Falush, D. 2007. Documentation for STRUCTURE software: version 2.2 (available at [pritch.bsd.uchicago.edu/software/](http://pritch.bsd.uchicago.edu/software/)).
- Robinson, A.R., Ukrainetz, N.K., Kang, K.-Y. & Mansfield, S.D. 2007. A comprehensive metabolomics analysis of two Douglas fir (*Pseudotsuga menziesii*) progeny test trials. *New Phytologist*, 174: 763–773.

- Simmonds, N.W. 1979. *Principles of crop breeding*. Longman, London, UK, and New York, USA.
- Sutton, S.C.S., Attree, S.M., El-Kassaby, Y.A., Grossnickle, S.C. & Polonenko, D.R. 2004. Commercialisation of somatic embryogenesis for plantation forestry. pp. 275–301, in: C. Walter & M. Carson (editors). *Plantation forest biotechnology for the 21st century*. Research Signpost, Trivandrum, India.
- Szabó V.M. and Burr B. 1996. Simple inheritance of key traits distinguishing maize and teosinte. *Molecular and General Genetics* 252(1-2): 33–41.
- Thoday, J.M. 1961 The location of polygenes. *Nature*, 191: 368–370.
- Thoday, J.M. 1976. Effects of specific genes. pp. 141–159, in: *Proc. of the International Conference on Quantitative Genetics*. The Iowa State University Press, Ames, Iowa, USA.
- White, T.L., Adams, W.T. & Neale, D.B. 2007. *Forest Genetics*. CAB International, Wallingford, UK, and Cambridge, Massachusetts, USA.
- Wilcox, P.L., Amerson, H.V., Kuhlman, G.E., Liu, B.-H., O'Malley, D.M. & Sederoff, R.R. 1996. Detection of a major gene for resistance to fusiform rust disease in loblolly pine by genomic mapping. *Proceedings of the National Academy of Sciences of the United States of America*, 93: 3859–3864.
- Wilcox, P.L., Echt, C.E. & Burdon, R.D. 2007. Gene-assisted selection – Applications of association genetics in forest tree breeding. pp. 213–250, in: N. Oraguzie, E.H.A. Rikkerink, S.E. Gardiner & H.N. De Silva (editors). *Association mapping in plants*. Springer, New York, USA.

## 6. Research, deployment and safety management of genetically modified poplars in China

*Y. Zheng*

China's population is the world's largest, and under demographic pressure from unprecedented expansion coupled with rapid social and economic development. This is exerting severe pressure on its forest resources. This scarcity of forest products is mainly caused by past and contemporary extensive deforestation and forestland degradation. While great efforts are being dedicated to reforestation and afforestation to increase forest resources, the gap between timber supply and demand remains large. To alleviate this shortage, the development of fast-growing, high-yielding short-rotation plantations has been identified as a national top priority.

During the past decade, the country has embarked on and implemented six major national forest programmes, one of which was the development of short-rotation industrial timber plantations. The fast growing nature of poplars and their responsiveness to cultivation made them an ideal species for the establishment of industrial, high-yield short-rotation plantations. Particularly in northern China, poplar is being widely used in large-scale plantations and currently plays an irreplaceable role in both commercial and ecological plantations.

In China, the total area covered by poplar trees is approximately seven million hectares, exceeding the total of other poplar plantations globally (Lu and Hu, 2006). However, the major poplar species in China are often subject to attacks by pests and diseases, which become the major barriers to the development of large-scale poplar plantations (Zhang and Li, 2003; Lu and Hu, 2006; Wang and Lu, 2002). The same problem was also identified for other crops of great economic potential, such as fruit trees and ornamental plants (Cheng *et al.*, 1999; Fang and Wang, 2000; Han *et al.*, 2000, 2006; He *et al.*, 2004). Moreover, China has large areas of saline land currently unsuitable for growing trees, but which might be utilized if trees are bred for salinity tolerance. Thus, the development of trees with improved pest and disease resistance as well as tolerance to stress conditions is paramount for the successful deployment of poplars as high-yield fibre plantations.

Compared with traditional breeding methods that often require a long time, and the inherent difficulty of meeting predetermined breeding objectives, such as improved pest and disease resistance and stress adaptabilities, modern technologies such as genetic engineering, if successful, appear to be very efficient (Lu and Hu, 2006; Su *et al.*, 2003a, b). Additionally, the rapid technological development and the progress accomplished further encourages the application of genetic

engineering in breeding poplar trees, and significant progress has been achieved in genetically modified trees, particularly poplar, in China.

This chapter summarizes the status of genetically modified poplars in China, the research associated with their development, release and commercialization, as well as associated biosafety issues.

## CURRENT STATUS AND PROGRESS OF GENETICALLY MODIFIED TREE APPLICATIONS IN CHINA

### Tree species in genetic modification studies

Poplars have been the most successful trees for genetic transformation studies due to their predisposition to clonal propagation and high transformation efficiency, with the added benefits of poplar being the model tree species for genomic research (a relatively small and completely sequenced genome) and the vast number of molecular and biotechnological studies coupled with its fast growth rate and short rotation period making it amenable to genetic engineering.

The first genetically transformed poplar produced in China was obtained in 1989, where the Bt insect resistance gene was introduced into the *P. nigra* genome (Tian *et al.*, 1993). Following this introduction, several genetically modified trees were placed in field trials and displayed significant improvement in insect resistance (Lu and Hu, 2006). The success of these trials prompted the approval of genetically modified tree release and the establishment of the first genetically modified commercial tree plantations in 2004 in China. Following this, several successful transformations have been achieved in poplar species: *P. tomentosa*, *P. alba*, *P. nigra*, *P. deltoides*, *P. euramericana*, *P. xinjiangensis*, *P. tremuloides*, *P. xiaozhannica* and hybrid poplars of *P. deltoides* × *P. simonii*, *P. alba* × *P. glandulosa*, *P. deltoides* × *P. cathayana*, *P. euramericana* × *yunnanensis*, *P. simonii* × *P. nigra* (Table 6-1).

Other trees used for genetic modification studies were mostly fruit trees of economic importance, such as apple, walnut, orange, date and cherry (Cheng *et al.*, 1999; Fang and Wang, 2000; Han *et al.*, 2000, 2006; He *et al.*, 2004). These activities focused on improving insect resistance, nutritional condition, and fruit taste (Han *et al.*, 1999, 2000, 2006; He and Han, 2000; He *et al.*, 2004; Shi *et al.*, 2000; Zhan *et al.*, 2001; Zhou *et al.*, 2005). Of interest was the development of infertility (i.e. inhibition of the development of reproductive organs) through transformation of *Liquidambar formosana* trees to maximize vegetative growth (Qiao *et al.*, 2007) and the use of the ornamental tree Bergamot (*Citrus medica* var. *sarcodactylis*) to develop a model for a genetic transformation system.

### Target traits for improvement

The main drive for genetic transformation of trees in China was to improve resistance to diseases and pests and the acquisition of pesticide and herbicide resistance. Tolerance to salt and reduction of lignin content were also of interest and have been receiving increased attention as major target traits for improvement (Table 6-1). However, variable extents of improvement were obtained in various target traits under different genetic controls. So far, pest resistance seems to be the

TABLE 6-1  
Summary information on genetically modified poplars in China

Species used	Method	Transferred genes	Target traits	Receptor	Test stage	Reference
<i>P. deltoids</i> × <i>P. simonii</i> 'NL-80106'	Agro-bacterium	<i>Bt</i>	Insect resistance	Stem disc, leaf	Field test	Rao <i>et al.</i> , 2000
<i>P. deltoids</i> × <i>P. simonii</i>	Agro-bacterium	<i>Aa1T</i>	Insect resistance	Leaf, petiole	Lab	Wu <i>et al.</i> , 2000
<i>P. xinjiangensis</i>	Agro-bacterium	<i>cpti</i>	Insect resistance	Leaf	Lab	Zhu <i>et al.</i> , 2003
<i>P. alba</i> × <i>P. glandulosa</i>	Agro-bacterium	<i>SacB</i>	Drought resistance	Leaf	Lab, Greenhouse	Zhang <i>et al.</i> , 2005; Li <i>et al.</i> , 2007a, b, c
<i>P. alba</i> × <i>P. glandulosa</i> '84K'	Agro-bacterium	<i>BtCry3A/OC-I</i>	Resistance to coleopteron	Leaf	Lab	Zhang <i>et al.</i> , 2005
<i>P. deltoides</i> × <i>P. cathayana</i>	Agro-bacterium	<i>mt1D/gutD</i>	Salt resistance	Leaf	Lab	Fan, 2002
<i>P. deltoids</i>	Agro-bacterium	<i>Bt</i>	Insect resistance	Leaf	Lab	Chen <i>et al.</i> , 1995
<i>P. deltoids</i> , <i>P. euramericana</i> , <i>P. nigra</i>	Agro-bacterium	Antibacterial peptide <i>Lcl</i>	Insect resistance	Leaf	Field test	Li <i>et al.</i> , 1996
<i>P. euramericana</i>	Agro-bacterium	<i>cpti</i>	Insect resistance	Leaf	Lab	Zhao <i>et al.</i> , 2005
<i>P. euramericana</i>	Agro-bacterium	<i>CryIA(C)</i>	Insect resistance	Stem, leaf	Lab	Wang <i>et al.</i> , 1997
<i>P. euramericana</i> 'Guariento'	Gene gun	<i>SacB/vgbl/BtCry3A, OC-I/IERF36/NPT II</i>	Insect, drought and salinity resistance	Leaf	Field test	Wang <i>et al.</i> , 2006
<i>P. euramericana</i> , <i>P. euramericana</i> × <i>yunnanensis</i>	Agro-bacterium	<i>CryIA(C)</i>	Insect resistance	Stem disc, leaf	Lab	Wang <i>et al.</i> , 1997
<i>P. nigra</i>	Agro-bacterium	<i>CryIA(c)</i>	Insect resistance	Leaf	Commercial production	Tian <i>et al.</i> , 1993
<i>P. nigra</i>	Gene gun/leaf disc	<i>TA29-BARNASE</i>	Male infertility	Leaf	Lab	Li <i>et al.</i> , 2000a, b
<i>P. simonii</i> × <i>P. nigra</i>	Agro-bacterium	<i>Bet-A</i>	Saline resistance	Leaf	Lab	Yang <i>et al.</i> , 2001, Liu <i>et al.</i> , 2006
<i>P. tomentosa</i>	Polyethylene glycol (PEG)	<i>GUS</i>	Transient expression	leaf	Lab	Wang <i>et al.</i> , 1991
<i>P. tomentosa</i>	Agro-bacterium	Antisense <i>CCoAOMT</i>	Reduction of lignin content	Leaf with petiole	Field test	Zhao, Wei and Lu, 2004
<i>P. tomentosa</i>	Agro-bacterium	<i>cpti</i>	Insect resistance	Leaf	Field test	Hao and Zhu, 2000
<i>P. tomentosa</i>	Agro-bacterium	<i>Bt</i>	Insect resistance	Leaf	Lab	Zheng, Liang and Sun, 1996
<i>P. tomentosa</i>	Agro-bacterium	<i>NP-1</i>	Anti bacteria	Leaf	Lab	Zhao <i>et al.</i> , 1999
<i>P. tomentosa</i> '741'	Agro-bacterium	<i>BtCryIAc/Api</i>	Insect resistance	Leaf	Commercial production	Zheng <i>et al.</i> , 2000
<i>P. tomentosa</i> '741'	Agro-bacterium	<i>Bt</i>	Insect resistance	Leaf	Lab	Zheng <i>et al.</i> , 2000
<i>P. tomentosa</i> (Triploid)	Agro-bacterium	<i>Dγ</i>	Saline resistance	Leaf	Lab	Liu <i>et al.</i> , 2002
<i>P. tremula</i> × <i>P. alba</i>	Agro-bacterium	Antisense <i>CCoAOMT</i>	Reduction of lignin content	Leaf	Field test	Wei <i>et al.</i> , 2001
<i>P. tremuloides</i>	Agro-bacterium	Antisense <i>4CL</i>	Reduction of lignin content	Leaf	Lab	Jia <i>et al.</i> , 2004
<i>P. xiaozhannica</i> 'Balizhuangyang'	Agro-bacterium	<i>mt1D</i>	Saline resistance	Leaf	Field test	Liu <i>et al.</i> , 2000
<i>Populus euramericana</i>	Agro-bacterium	<i>BG2</i>	Resistance to bacteria	Leaf	Lab	Han <i>et al.</i> , 2004



most successful trait attained through genetic transformation, probably attributed to its simple genetic control (i.e. single or major gene resistance) in comparison with other target traits that are likely to be controlled by multiple genes. Salt tolerance is another trait that has shown evidence of improvement in trees using salt-tolerance genes (Sun *et al.*, 2002; Yin *et al.*, 2004; Yang *et al.*, 2001).

### Performance of genetically modified trees

By far, pest resistance in poplars has proven to be the most successful genetically modified trait in China. Transformation of Bt genes significantly improved poplar tree pest resistance. Due to the excellent improvement in pest resistance, two genetically modified trees *P. nigra* and hybrid poplar clone 741 [*Populus alba* × (*P. davidiana* + *P. simonii*) × *P. tomentosa*] have been approved for environmental release and *P. xiaozhannica* ‘Balizhuangyang’ was placed in field tests. The pest resistance of these genetically modified trees was assessed and verified by feeding experiments.

Test feeding of gypsy moths with leaves from genetically modified *P. nigra* trees showed 70–95% mortality on 15% of the trees, and over 50% mortality on 50% of the trees. The genetically modified trees with higher insect mortality rate also suppressed development of the surviving insects. Seven days after start of feeding, the leaves of genetically modified trees were showing evidence of resistance to the insects, with almost no damage, while most leaves of the control trees were eaten.

The genetically modified hybrid poplar clone known as ‘741’ was tested with *Clostera anachoreta*, and insect mortality rate was monitored during the entire development period (23–25 days corresponding to 1–2 days after hatching to cocooning). The results showed that the genetically modified trees induced significantly higher mortality than non-transgenic trees with 91.8% mortality on 14.3% of the genetically modified trees, 40–70% on 23.8 of genetically modified trees, and <20% mortality on 61.9% of the genetically modified trees, while the non-genetically modified ‘741’ clone induced very low mortality (3.4%). Additionally, the insect-resistant genetically modified trees also delayed the development of living insects by 2–8 days from instars I to II, 2–6 days from instars II to III, and 2–8 days from instars IV to V. The average weight of individual insects fed on genetically modified trees was reduced by 41–49% compared with those fed on non-genetically modified trees, providing further evidence of the impact of genetically modified trees on the target insect’s development.

The *P. xiaozhannica* ‘Balizhuangyang’ transformed with the *mtl-D* gene has shown significantly increased saline resistance. Field tests in Shandong and Tianjin indicated both a 70% increase in survival rate of genetically modified trees in soils with 0.3–0.4% salt content and increased growth rate compared with control trees. It should be noted that the upper limit for soil salt content reached 0.43% for the genetically modified trees in the trial.

Although resistances of the genetically modified poplar trees were significantly improved, their growth performance was found to show no notable improvement compared with non-genetically modified trees. However, some trees showed



To date, eight poplar lines of seven different crosses have been transformed (Table 6-1), but only *P. nigra* and the hybrid clone '741' of the native *P. tomentosa* have been approved for commercial production, while seven genetically modified poplar species are in the field testing stage. The first genetically modified poplar trees (*Populus nigra*) were obtained in 1989 and entered field testing in 1994, and finally used in commercial plantation in 2002 (Tian *et al.*, 1993). Plantations of this genetically modified poplar have been established on eight sites in seven provinces or municipalities (Beijing, Jilin, Henan, Shandong, Xinjiang, Shaanxi and Jiangsu) (Su *et al.*, 2003a, b). The hybrid clone '741' (*Populus alba* × [*P. davidiana* + *P. simonii*] × *P. tomentosa*) that was transformed with two genes was approved for commercial planting in 2002 (Tian *et al.*, 2000; Zheng *et al.*, 2000; Su *et al.*, 2003b). The genetically modified poplar clones were granted plant variety rights by the National Plant Variety Rights Protection Agency. The approvals for commercial production of the two genetically modified poplars were given after the completion of all biosafety assessment requirements organized by the National Forestry Biosafety Management Authority in the State Forestry Administration. Seven additional genetically modified poplars, including *Populus xiaozhannica* 'Balizhaungyang' genetically modified with salt-tolerant genes have been approved for field testing (Sun *et al.*, 2002; Yin *et al.*, 2004). Many other genetically modified trees are still restricted to the laboratory testing stage (Table 6-1). However, every year an increasing number of applications for field testing and environment release are being submitted for safety assessment.

## RISK ASSESSMENT AND SAFETY MANAGEMENT

### Concerns with biosafety issues of genetically modified trees

The increasing research and deployment of genetically modified trees has heightened public concerns regarding the safety of genetically modified trees. Worries aroused concern aspects such as pollen contamination of genetically modified trees of other, related species; impacts on non-target insects and other organisms in the soil; and other possible impacts of genetically modified trees on the biotic and abiotic aspects of the ecosystem.

### Biosafety of genetically modified trees

With the rapid development of genetic engineering technologies and the increasing number of released and planted genetically modified trees, increasing attention has been given to biosafety, with the deployment of a number of regulations. In 1993, "Rules for Safety Management of Genetic Engineering" were formulated by the Ministry of Science and Technology (MST, 1993), followed by publication of "Rules for Implementation of Safety Management of Genetic Engineering in Agriculture" in 1996 by the Ministry of Agriculture (MOA, 1996), and the issuance of "Regulations for Biosafety Management of Agricultural GMOs" (State Council, 2001). More recently, regulations were developed to cope with more specific biosafety issues associated with forest trees, including "Rules on Administration of Examination and Approval for Genetic Engineering Activities

of Forest Trees” issued in 2006, followed by “Technical Codes for Biosafety Assessment of Transgenic Forest Plants and Products” in 2007 (State Forestry Administration, 2007). All these regulations and rules have paved the way for managing the biosafety of genetically modified trees.

According to the rules and technical codes, genetically modified trees are classified into three classes based on an evaluation of the risks, and they must pass through a series of tests (laboratory, field, environment release and productivity). Only when these tests are completed and the results are satisfactory can the genetically modified trees be deployed for commercial production. A biosafety assessment must be carried out for each of these tests before granting approval for next step.

### Technical standards for safety assessment and long-term monitoring of genetically modified trees

As we enter the twenty-first century, biosafety of genetically modified trees should be given increasing attention. Studies have been initiated to investigate the potential impacts of genetically modified trees on elements of the ecosystem, such as impacts on soil micro-organisms, target and non-target insects, and gene flow to non-genetically modified trees (Hu *et al.*, 2004; Gao, Li and Liu, 2003; Zhang *et al.*, 2006). Early results from these studies have shown that genetically modified products did not produce any significant changes to the natural ecosystem. However, it may not be feasible to assess many impacts of genetically modified trees during the early stages of tree growth and development due to their long life cycle, so long-term continued monitoring of these impacts is needed to obtain reliable information and reach reliable conclusions. A number of monitoring studies on the long-term impacts of genetically modified trees on soil micro-organisms have been established (Hou, Zhang and Su, 2006) and more information on genetically modified tree impacts on the ecosystem at large will become available in future years.

### FUTURE PROSPECTS

Since the first genetically modified *P. nigra* development, a large number of studies have been carried out with the aim of developing genetically modified trees for different purposes. Although two of the genetically modified poplars are being commercialized and a couple of the genetically modified poplars are at the field testing stage, most genetic modification studies are still at the laboratory stage (Lu and Hu, 2006). Further information on the performance of genetically modified trees of other poplar species still needs to be collected and analysed to develop reliable conclusions.

In addition to pest-resistant genetically modified poplars, progress has been made in saline tolerance in genetically modified trees of several poplar species. Given the large areas of saline land in China, saline-tolerant genetically modified poplar trees appear to be of great potential in afforestation in such areas. Moreover, genetically modified trees of *P. tomentosa* transformed with *4CL* and *CCoAOMT* genes showed great potential for significantly reducing lignin content (Wei *et al.*, 2001; Zhao, Wei and Lu, 2004).

With rapid development of poplar-based industries in China, the plantation area under poplar will be further expanded. Breeding of poplar trees with high resistance to pests and diseases as well as tolerance to saline soils will greatly facilitate the expansion of poplar plantation. As already shown by some genetically modified poplar trees, genetic modification appeared to be a promising tool to achieve this goal.

## REFERENCES<sup>1</sup>

- Chen, Y., Han, Y.F., Tian, Y.C., Li, L. & Xie, S.J. 1995. Study on the plant regeneration from *Populus deltoides* explants transformed with Bt-Toxin gene. *Scientia Silvae Sinicae*, 31(2): 97–103.
- Cheng, J.S., Tian, Y.C., Meng, X.M., Li, W.G., Liu, X.F., Zhang, J.T. & Mang, K.Q. 1999. Transfer Bt-gene to apple. *Acta Agriculturae Boreali-Occidentalis Sinica*, 8(1): 78–81.
- Fan, J.F. 2002. Studies on transformation of salt-resistant *MtLD/gutD* gene to poplar and kiwifruit. Ph.D. thesis. Northwest Agriculture and Forestry University, Yangling, Xi'an, China.
- Fang, H.J. & Wang, G.L. 2000. Somatic embryogenesis of *Juglans nigra* L. and establishment of gene transformation system of walnut. *Acta Horticulturae Sinica*, 27(6): 406–411.
- Gao, B.J., Li, C. & Liu, J.X. 2003. Roles of insect resistant GM trees in pest control and approaches to their appropriate utilization. *Journal of the Agricultural University of Hebei*, 26(4): 25–28.
- Han, M.L., Lu, R.S., Zhu, J.Y., Qiu, X.J. & Huang, H.Y. 1999. Studies on factors affecting gene-transformation by *Agrobacterium*-mediated in *Citrus*. *Acta Botanica Yunnanica*, 21(4): 491–496.
- Han, M.L., Li, G.G., He, H., Zhang, Y.D., Zhang, L.Y. & Lu, R.S. 2000. Preliminary study on establishment of transformation system in *Poncirus trifoliata* Raf. *Guibaia*, 20(1): 47–51.
- Han, L.N., Zhou, C.J., Cui, D.C. & Wang, B. 2004. Transformation of *Populus* with *BG2* gene. *Journal of Shandong Agricultural University*, 35(3): 375–378.
- Han, M.L., Lu, R.S., Wu, Y.J. & Du, X.L. 2006. Studies on factors influencing ShivaA gene transformation mediated by *Agrobacterium* in *Citrus grandis* Osbeck 'Shatianyu'. *Guibaia*, 26(5): 479–482.
- Hao, G.X. & Zhu, Z. 2000. Obtaining of cowpea proteinase inhibitor transgenic *Populus tomentosa*. *Scientia Silvae Sinicae*, 36, Sp(1): 116–119.
- He, H. & Han, M.L. 2000. A Study on the *Agrobacterium*-mediated GUS gene transfer to trifoliolate orange (*Poncirus trifoliata*). *Chinese Traditional and Herbal Drugs*, 31(4): 295–297.
- He, Y.H., Xiong, X.H., Lin, S.Q., Wu, H.T., Lin, L.B., He, G. & Chen, J.H. 2004. Transformation of *Ziziphus jujuba* with antisense ACC synthase gene. *J. Hunan Agricultural University (Natural Sciences)*, 30(1): 33–36.
- Hou, Y.J., Zhang, B.Y. & Su, X.H. 2006. Review of the ecological risks of genetically modified trees. *China Biotechnology*, 26(12): 117–121.
- Hu, J.J., Zhang, Y.Z., Lu, M.Z., Zhang, J.G. & Zhang S.G. 2004. Transgene stability of transgenic *Populus nigra* and its effects on soil microorganism. *Scientia Silvae Sinicae*, 40(5): 105–109.
- Jia, C.H., Wang, H.Z., Du, K.J., Song, Y.R. & Wei, J.H. 2004. Relationship of lignin content with stem colour in transgenic poplar with depressed expression of *4CL* gene. *Journal of Agricultural Biotechnology*, 12(6): 621–624.
- Li, Y., Chen, Y., Li, L. & Han, Y.F. 1996. Transformation of anti-bacterial gene *Lel* into poplar species. *Forest Research*, 9(6): 646–649.
- Li, L., Qi, L.W., Han, Y.F., Wang, Y.C. & Li, W.B. 2000a. A study on the introduction of male sterility of anti-insect transgenic *Populus nigra* by *Ta29-Barnase* gene. *Scientia Silvae Sinicae*, 36(1): 28–32.
- Li, M.L., Zhang, H., Hu, J.J., Han, Y.F. & Tian, Y.C. 2000b. Study on insect resistant transgenic poplar plants containing both Bt and PI genes. *Scientia Silvae Sinicae*, 36(2): 93–97.

<sup>1</sup> All references were published in Chinese with English summary.

- Li, K.Y., Fan, J.F., Zhao, Z., Li, L. & Zhu, H.L. 2007a. Transformation of *CryIAc* and *API* – two insect-resistance genes – to poplar (*Populus alba* × *P. glandulosa*). *Forest Research*, 20(5): 699–704.
- Li, K.Y., Fan, J.F., Zhao, Z., Li, L. & Jia, X.M. 2007b. Resistance to insects of transgenic *Populus tomentosa* Clone-85 plants with two insect-resistance genes. *Acta Botanica Boreali-occidentalia Sinica*, 27(8): 1537–1543.
- Li, Y.L., Su, X.H., Zhang, B.Y. & Zhang, Z.Y. 2007c. Molecular detection and drought tolerance of SacB-transgenic *Populus alba* × *P. glandulosa*. *Journal of Beijing Forestry University*, 29(2): 1–6.
- Liu, B., Li, H.S., Wang, Q.H. & Cui, D.C. 2002. Transformation of *Populus tomentosa* with anti-PLD gene. *Hereditas (Beijing)*, 24(1): 40–44.
- Liu, F.H., Sun, Z.X., Cui, D.C., Du, B.X., Wang, C.R. & Chen, S.Y. 2000. Cloning of *E. coli mtl-D* gene and its expression in transgenic Balizhuangyang (*Populus*). *Acta Genetica Sinica*, 27(5): 428–433.
- Liu, G.F., Cheng, G.L., Jiang, J., Bai, S., Yu, Y., Cai, Z.J., Dong, J.X. & Li, S.J. 2006. The transformation of *BetA* gene into the pollen plantlets of *Populus simonii* × *P. nigra*. *Acta Genetica Sinica*, 32(2): 163–168.
- Lu, M.Z. & Hu, J.J. 2006. Current status of research and development of transgenic poplars in China. *Forestry Science and Technology Development*, 20(6): 1–4.
- MOA [Ministry of Agriculture]. 1996. Rules for implementation of safety management of genetic engineering in agriculture.
- MST [Ministry of Science and Technology]. 1993. Rules for safety management of genetic engineering.
- Qiao, G.R., Luan, W.J., Pan, H.W. & Zhuo, R.Y. 2007. The *LEAFY* gene in RNA interference (RNAi) transgenic *Liquidambar formosana* mediated by *Agrobacterium tumefaciens*. *Journal of Zhejiang Forestry College*, 24(2): 140–144
- Rao, H.Y., Chen, Y., Huang, M.R., Wang, M.M., Wu, N.F. & Fan, Y.L. 2000. Genetic transformation of poplar NL-80106 transferred by Bt gene and its insect-resistance. *Journal of Plant Resources and Environment*, 9(2): 1–5.
- Shi, X.X., Wang, B., Du, G.Q., Wang, M.L. & Gao, Y. 2000. Studies on *Agrobacterium*-mediated *CpTI* g gene transfer in commercial apple cultivars. *Acta Horticulturae Sinica*, 27(4): 282–284.
- State Council. 2001. Regulations for biosafety management of agricultural GMOs.
- State Forestry Administration. 2007. Technical codes for biosafety assessment of transgenic forest plants and products. China Standards Publishing House.
- Su, X.H., Zhang, B.Y., Huang, L.J., Huang, Q.J. & Zhang, X.H. 2003a. Genetic engineering of forest trees. *Forest Research*, 16(1): 95–103.
- Su, X.H., Zhang, B.Y., Huang, L.J., Huang, Q.J. & Zhang, X.H. 2003b. Advances and key research fields in the genetic engineering in forest trees in China. *Scientia Silvae Sinicae*, 39(5): 111–118.
- Sun, Z.X., Yang, H.H., Cui, D.C., Zhao, C.Z. & Zhao, S.P. 2002. Analysis of salt resistance of poplar trees transferred with salt tolerance gene. *Chinese Journal of Biotechnology*, 18(4): 481–485.
- Tian, Y.C., Li, T.Y., Mang, K.Q., Han, Y.F., Li, L., Wang, X.P., Lu, M.Z., Dai, L.Y., Hang, Y.N., Yan, J.J., & Dean, W.G. 1993. Insect tolerance of transgenic *Populus nigra* plants transformed with *Bacillus thuringiensis* toxin gene. *Chinese Journal of Biotechnology*, 9(4): 291–297.
- Tian, Y.C., Zheng, J.B., Yu, H.M., Li, C.Q. & Wang, J.M. 2000. Studies of transgenic hybrid poplar 741 carrying two insect-resistant genes. *Acta Botanica Sinica*, 42(3): 263–268.
- Wang, M.J. & Lu, M.Z. 2002. Current status and development trends in tree breeding by genetic engineering. *World Forestry Research*, 15(3): 7–13.
- Wang, S.P., Xu, N., Xu, Z.H., Huang, M.R. & Wei, Z.M. 1991. Studies on transgenic expression of GUS gene in protoplasts of three *Populus* species by PEG method. *Acta Biologiae Experimentalis Sinica*, 24(1): 71–74.
- Wang, X.P., Han, Y.F., Dai, L.Y. & Li, L. 1997. Studies on insect-resistant transgenic plants (*P. × euramericana*). *Scientia Silvae Sinicae*, 33(1): 69–74.



- Wang, J.G., Su, X.H., Ji, L.L., Zhang, B.Y., Hu, Z.M., Huang, R.F. & Tian, Y.C. 2006. Multiple gene transformation by gene gun of *Populus × euramericana* 'Guariento'. *Science Bulletin*, 51(23): 2755–2760.
- Wang, S.H., Yang, M.Y., Gu, M., Qu, S.C., Yao, Q.H. & Zhang, Z. 2007. *Agrobacterium*-mediated transformation of *Malus micromalus* with trivalent genes *Rirol*. *Journal of Fruit Science*, 24(6): 731–736.
- Wei, J.H., Zhao, H.Y., Zhang, J.Y., Liu, H.R. & Song, Y.R. 2001. Cloning of cDNA encoding CCoAOMT from *Populus tomentosa* and down-regulation of lignin content in transgenic plant expressing antisense gene. *Journal of Integrative Plant Biology*, 43(11): 1179–1183.
- Wu, N.F., Sun, Q., Yao, B., Fan, Y.L., Rao, H.Y. Huang, M.R. & Wang, M.M. 2000. Insect resistant transgenic poplar expressing AaIT gene. *Chinese Journal of Biotechnology*, 16(2): 129–133.
- Yang, C.P., Liu, G.F., Liang, H.M. & Zhang, H. 2001. Study on the transformation of *Populus simonii × P. nigra* with salt resistance gene Bet-A. *Scientia Silvae Sinicae*, 37(6): 34–38.
- Yin, J.D., Sun, Z.X., Wang, Y.X., Li, D.S., Liu, G.J., Feng, X. & Wang, S.Y. 2004. Field test of saline resistant transgenic *Populus × xiaozhannica* 'Balizhuangyang'. *Journal of Northeast Forestry University*, 32(3): 23–25.
- Zhan, Y.G., Liu, Z.H., Wang, Y.H., Wang, Z.Y., Yang, C.P. & Liu, G.F. 2001. Transformation of insect resistance gene into birch. *Journal of the Northeast Forestry University*, 29(6): 4–6.
- Zhang, B.Y., Su, X.H., Huang, Q.J., Zhang, X.H. & Hu, Z.M. 2005. Regeneration of transgenic Poplar (*Populus alba × P. glandulosa*) expressing levansucrase from *Bacillus subtilis*. *Scientia Silvae Sinicae*, 41(3): 48–53.
- Zhang, B.Y., Su, X.H., Li, Y.L., Zhang, Y.A., Qu, L.J., Wang, Y.Z. & Tian, Y.C. 2006. Production of *Populus alba × P. glandulosa* with a coleopterous insect resistant gene and analysis of insect resistance. *Journal of Beijing Forestry University*, 28(2): 102–110.
- Zhang, Q.W. & Li, J.H. 2003. *New poplar varieties for industrial timber production*. China Forestry Publishing House. Beijing, China.
- Zhao, H.Y., Wei, J.H. & Lu, J. 2004. Using antisense CCoAOMT gene to reduce lignin content of the timber of *Populus tomentosa*. *Progress in Natural Sciences*, 14(9): 1067–1071.
- Zhao, S.M., Zu, G.C., Liu, G.Q., Huang, M.R., Xu, J.X. & Sun, Y.R. 1999. Introduction of rabbit defensin NP-1 gene into poplar (*P. tomentosa*) by *Agrobacterium*-mediated transformation. *Acta Genetica Sinica*, 26(6): 711–714.
- Zhao, Q., Zhao, Z.W., Zhang, T.T., Cui, D.C. & Wang, B. 2005. Transformation of *Populus euramericana* with *CpTI* gene. *Biotechnology Bulletin*, (4): 54–58.
- Zheng, J.B., Liang, H.Y., Gao, B.J., Wang, Y.F. & Tian, Y.C. 2000. Selection and insect resistance of transgenic hybrid poplar 741 carrying two insect resistant genes. *Scientia Silvae Sinicae*, 36(2): 13–19.
- Zheng, J.B., Liang, H.Y. & Sun, K.N. 1996. Regeneration of explants of Chinese poplar leaves and its transformation with insect-resistant gene. *Journal of the Hebei Forestry College*, 11(2): 97–101.
- Zhou, C.L., Guo, W.D., Wang, D.J., Yu, C.L., Lu, M. & Li, Y.P. 2005. Exploration of the transformation parameters of the leaf discs of bergamot by particle bombardment in virtue of transient expression of GUS gene. *Acta Botanica Boreali-Occidentalia Sinica*, 25(11): 2145–2150.
- Zhu, G.Q., Wang, J.C., Chen, Y., Huang, M.R. & Wang, M.X. 2003. Establishment of system with high frequency for genetic transformation of *Populus alba* var. *pyramidalis*. *Journal of Plant Resources and Environment*, 12(4): 6–10.