

Demand and supply of feed ingredients for farmed fish and crustaceans

Trends and prospects



Cover photograph:

Drying of farm-made aquafeed for Nile tilapia, Jamalpur, Bangladesh (courtesy of FAO/Mohammad R. Hasan).

Demand and supply of feed ingredients for farmed fish and crustaceans

Trends and prospects

Albert G.J. Tacon
FAO Consultant
Hawaii, United States of America

Mohammad R. Hasan
Aquaculture Officer
Aquaculture Service
FAO Fisheries and Aquaculture Department
Rome, Italy

and

Marc Metian
Littoral Environment and Societies
University of La Rochelle
La Rochelle, France

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views of FAO.

ISBN 978-92-5-106933-2

All rights reserved. FAO encourages reproduction and dissemination of material in this information product. Non-commercial uses will be authorized free of charge, upon request. Reproduction for resale or other commercial purposes, including educational purposes, may incur fees. Applications for permission to reproduce or disseminate FAO copyright materials, and all queries concerning rights and licences, should be addressed by e-mail to copyright@fao.org or to the Chief, Publishing Policy and Support Branch, Office of Knowledge Exchange, Research and Extension, FAO, Viale delle Terme di Caracalla, 00153 Rome, Italy.

Preparation of this document

This document was prepared as part of the Food and Agriculture Organization of the United Nations (FAO) Aquaculture Service's (FIRA) ongoing regular work programme on "Study and analysis on use of feed and nutrient in sustainable aquaculture development", under strategic objective "Sustainable management and use of fisheries and aquaculture resources", and was partially funded by the FAO Multidonor Partnership Programme.

This technical paper was reviewed for technical content by Dr Methil N. Kutty. Pedro Bueno is acknowledged for his comments on the manuscript. For consistency and conformity, scientific and English common names of fish species were used from FishBase (www.fishbase.org/search.php).

The authors would also like to acknowledge the following colleagues (commercial feed manufacturers and/or nutritionists) for their kind assistance in providing information during an electronic survey conducted for this study: Syed Ahamad Ali, Niels Alsted, Michel Autin, Christian Berger, Aliro Borquez, Hans Erik Bylling, Ian Carr, Rodrigo Carvalho, Silvio Coelho, José Duarte, Carlos Mastrokalo Durand, Fabricio Vargas Elias, Abdel-Fattah El-Sayed, Kevin Fitzsimmons, Brett Glencross, Ozlem Guzel, Brad Hicks, Santiago Panné Huidobro, Patricia Infante, Dave Jackson, Jeongdae Kim, Claudio Larraín, Daniel Lemos, Menghe Li, Christian Lückstädt, Umberto Luzzana, Cesar Molina, Jessica Montaña, Sigve Nordum, Dave Robb, Cheryl Shew, Shi-Yen Shiau, Supis Thongrod, Francisco Velasquez, Miao Weimin, Iannis Zarkadis and Jesus Zendejas.

The authors acknowledge Maria Giannini for her assistance in quality control and FAO house style and Koen Ivens for layout design. The publishing and distribution of the document were undertaken by FAO, Rome. Finally, Jiansan Jia, Chief of the Aquaculture Service of the FAO Fisheries and Aquaculture Department is acknowledged for providing the necessary support to initiate the study and to complete the publication.

Abstract

The rise into global prominence and rapid growth of finfish and crustacean aquaculture has been due, in part, to the availability and on-farm provision of feed inputs within the major producing countries. More than 46 percent of the total global aquaculture production in 2008 was dependent upon the supply of external feed inputs. For the aquaculture sector to maintain its current average growth rate of 8 to 10 percent per year to 2025, the supply of nutrient and feed inputs will have to grow at a similar rate. This had been readily attainable when the industry was young. It may not be the case anymore as the sector has grown into a major consumer of and competitor for feed resources. This paper reviews the dietary feeding practices employed for the production of the major cultured fed species, the total global production and market availability of the major feed ingredient sources used and the major constraints to feed ingredient usage, and recommends approaches to feed ingredient selection and usage for the major species of cultivated fish and crustacean. Emphasis is placed on the need for major producing countries to maximize the use of locally available feed-grade ingredient sources, and, in particular, to select and use those nutritionally sound and safe feed ingredient sources whose production and growth can keep pace with the 8 to 10 percent annual average growth of the fed finfish and crustacean aquaculture sector.

Tacon, A.G.J.; Hasan, M.R.; Metian, M.

Demand and supply of feed ingredients for farmed fish and crustaceans: trends and prospects.

FAO Fisheries and Aquaculture Technical Paper No. 564. FAO, 2011. 87 pp.

Contents

Preparation of this document	iii
Abstract	iv
List of tables and figures	vi
Contributors	viii
Abbreviations and acronyms	ix
Executive summary	x
1. Introduction	1
2. Current feeds and feeding practices	3
2.1 Major fed fish and crustacean species	3
2.2 In-country fed species production and feeding practices	7
2.3 Global aquaculture feed production by major species group and country	16
3. Feed ingredient production and availability	21
3.1 Animal nutrient sources	21
3.2 Plant nutrient sources	32
3.3 Microbial ingredient sources	50
4. Current levels of feed ingredient usage and constraints	51
4.1 Continued use of fishmeal and fish oil as major dietary animal protein and lipid sources	51
4.2 Increased use of terrestrial animal protein meals and oils as dietary nutrient sources	54
4.3 Continued and increased use of plant protein meals and oils as dietary nutrient sources	61
4.4 Ingredient competition with other users	62
4.5 Growing importance of feed and food safety	65
5. Recommended approaches to feed ingredient selection and use	67
5.1 Reduce country dependence upon imported feed ingredient sources	67
5.2 Select feed ingredients that can be sustainably produced and grow with the sector	68
5.3 Minimize environmental and ecosystem impact of feeds and feeding regimes	68
5.4 Give special attention to small-scale farmers using farm-made/ semi-commercial aquafeeds	69
References	71
Annexes	
1. Global production of finfish and crustaceans	79
2. Fed cultured species production by country	87

TABLES

1. Top 20 country producers of fed fish and crustacean species in 2008	7
2. Top 53 major fed cultured fish and crustacean species/species groups by main country producers in 2008	10
3. Estimated global aquaculture production and use of commercial aquafeeds, 1995–2020	14
4. Major country producers of commercial aquaculture feeds, 2007–2010	17
5. Total reported production of aquatic animal protein meals and lipids	22
6. IFFO estimate of global fishmeal production derived from fisheries by-products, 2007	27
7. Total reported production of terrestrial animal protein meals and lipids, 2008	29
8. Estimated global production and trade of major agricultural food/feed commodities, 2008/09	33
9. Global production and growth of major oilseed crops, 1995–2009	43
10. Country responses regarding feed ingredient usage for major cultivated species groups	55
11. Estimated global use and demand for fishmeal and fish oil, 1995–2020	59
12. Biofuel production by country, 2007	65

FIGURES

1. Total global production of fed fish and crustacean species by major FAO species grouping, 1980–2008	3
2. Growth of fed freshwater fish species, 1980–2008	4
3. Growth of fed crustacean species, 1980–2008	5
4. Growth of fed diadromous fish species, 1980–2008	6
5. Growth of fed marine fish species, 1980–2008	6
6. Estimated global production of commercial aquaculture feeds by major species grouping, 2008	16
7. Historical development of feeds in the Norwegian salmon industry in relation to prevailing technologies and typical inclusion levels of crude proteins, digestible energy and digestible protein levels, 1960–2000	18
8. Total capture fisheries and aquaculture production and volume of the catch destined for reduction and other non-food uses, 1970–2008	23
9. Major country producers of fishmeal and fish oil, 2007	23
10. Historical production trend of fishmeal and fish oil, 1962–2008	24
11. Major exporters of fishmeal and fish oil, 2007	25
12. Major importers of fishmeal and fish oil, 2007	26
13. Composition of rendered animal protein meal by meal type in Australia, 2008	30
14. Annual production of animal protein meals in the United States, 2003–2008	31
15. Annual production of animal fats and grease in the United States, 2003–2008	31
16. Total global production of cereals by commodity, 1995–2009	32
17. Composition of commodities in global production of cereals, 2009	38
18. Top cereal producer countries by commodity, 2009	38
19. Total global production of cereals by country, 1995–2009	39
20. Top cereal importers and exporters, 2008/09	40
21. Production of corn feed by-products from alcohol biorefineries in the United States, 1990/91 to 2008/09	41
22. Global production of major oilseed crops, 1995–2009	42
23. Composition of commodities in global production of oilseed crops, 2009	42
24. Global production of soybean oilseed crop by country, 1995–2009	44
25. Global production of soybean oilseed crop by country, 2009	44
26. Global production of major oilseed meals, 2005/06 to 2008/09	45

27.	Global production of major oilseed meals by commodity, 2008/09	45
28.	Global production of major oilseed oils, 2005/06 to 2008/09	46
29.	Global production of palm oil by country, 2008/09	46
30.	Global production of soybean oil by country, 2008/09	47
31.	Top agricultural exports by quantity from the United States, 2007	47
32.	Top agricultural imports by quantity in China, 2007	48
33.	Top agricultural imports by quantity in the European Union, 2007	48
34.	Global production of dry peas by country, 2009	49
35.	Global production of lupins by country, 2009	49
36.	Estimated global consumption of fishmeal by major aquaculture species group, 2008	51
37.	Estimated global consumption of fish oil by major aquaculture species group, 2008	52
38.	Estimated total global consumption of fishmeal and fish oil by major aquaculture species group, 2008	53
39.	Variation in price of fair average quality Peru fishmeal, January 2006 to October 2009	53
40.	Variation in price of FAQ fishmeal and soybean meal, 2006 to 2010	63
41.	United States and European Union production of biodiesel, 2001–2008	64
42.	United States corn destined for ethanol production, 1988/89 to 2009/10	65

Contributors

Mohammad R. Hasan, Aquaculture Officer, Aquaculture Service
FAO Fisheries and Aquaculture Department, Rome, Italy
E-mail: mohammad.hasan@fao.org

Marc Metian, Littoral Environment and Societies
University of La Rochelle, La Rochelle, France
E-mail: marc.metian@univ-lr.fr

Albert G.J. Tacon, FAO Consultant, Aquatic Farms Ltd
Hawaii, United States of America
E-mail: agjtacon@aol.com

Abbreviations and acronyms

ABP	animal by-products
BM	blood meal
CGM	corn gluten meal
CP	crude protein
CPC	canola protein concentrate
CSM	cottonseed meal
DE	digestible energy
DHA	docosahexaenoic acid
DP	digestible protein
EFCR	economic feed conversion ratio
EPA	eicosapentaenoic acid
EU	European Union
FAQ	fair average quality
FBM	faba bean meal
FM	fishmeal
FO	fish oil
FOB	free on board
FPM	field pea meal
G/PM	groundnut/peanut meal
HFM	hydrolysed feather meal
IFFO	International Fishmeal and Fish Oil Organisation
KM	krill meal
LKM	lupin kernel meal
MA	maize/corn
MBM	meat and bone meal
MC	mustard seed cake
MM	meat meal
Mtoe	million tonnes of oil equivalent
nei	not elsewhere included
PBM	poultry by-product meal
PO	poultry oil
RB	rice bran
R/CM	rapeseed/canola meal
R/CO	rapeseed/canola oil
SBM	soybean meal
SCP	single cell protein
SL	soy lecithin
SM	squid meal
SO	soybean oil
SSM	sunflower seed meal
US\$	US dollar
USDA	United States Department of Agriculture
WB	wheat bran
WF	wheat flour
WGM	wheat gluten meal
WH	wheat
WM	wheat middlings

Executive summary

The rapid rise and growth of finfish and crustacean aquaculture has been due, in part, to the availability and on-farm provision of feed inputs within the major producing countries. If the aquaculture sector is to maintain its current average growth rate of 8 to 10 percent per year to 2025, the supply of nutrient and feed inputs will need to grow at a comparable rate. While this may have been readily attainable when the industry was still in its infancy, this may not be the case in the future as the sector matures and grows into a major consumer and competitor for feed resources.

It is estimated that about 31.5 million tonnes of farmed fish and crustaceans (46.1 percent of the total global aquaculture production in 2008) is dependent upon the supply of external nutrient inputs provided in the form of fresh feed items, farm-made feeds or commercially manufactured feeds. Total industrial compound aquafeed production increased more than threefold, from 7.6 million tonnes in 1995 to 29.2 million tonnes in 2008, with production growing at an average rate of 11.0 percent per year. Aquafeed production is expected to continue growing at a similar rate to 71.0 million tonnes by 2020. Although current estimates for industrially produced aquafeed for the period 2007–2010 vary between 24.4 and 28.9 million tonnes, aquafeed volume represents only 4 percent of the total global animal feed production of the over 708 million tonnes in 2009. In contrast to compound aquaculture feeds, there is no comprehensive information on the global production of farm-made aquafeeds (estimated at between 18.7 and 30.7 million tonnes in 2006) and/or on the use of low-value fish/trash fish as feed, with 2008 estimates for China at 6 to 8 million tonnes.

Fed aquaculture production, in particular, of higher trophic level finfish and crustaceans (includes marine shrimps, salmonids, marine finfishes, eels) are largely dependent upon capture fisheries for the supply of their major dietary source of protein and lipids. For example, on a global basis, it is estimated that the aquaculture sector consumed 3.72 million tonnes of fishmeal (60.8 percent of global fishmeal production) and 0.78 million tonnes of fish oil (73.8 percent of global fish oil production) in 2008; it was 3.84 million tonnes of fishmeal (or 68.4 percent of global production) and 0.82 million tonnes of fish oil (or 81.3 percent of global production) in 2007. Despite this continued dependence of aquaculture production on fishmeal and fish oil, there remains a wide variation in fishmeal and fish oil usage between major producing countries for individual farmed species. This variation mainly reflects differences between countries concerning the selection and use of fishmeal and fish oil replacers from plant sources or by the use of land animal proteins and fats in feeds for high trophic-level fish and crustacean species.

The total use of fishmeal by the aquaculture sector is expected to decrease in the long term. It has gone down from 4.23 million tonnes in 2005 to 3.72 million tonnes in 2008 (or 12.8 percent of total aquafeeds by weight), and is expected to decrease to 3.49 million tonnes by 2020 (or 4.9 percent of total aquafeeds). The reasons for this are the diminishing amount of fishmeal and fish oil supplies owing to tighter quota setting and additional controls on unregulated fishing and the increased use of more cost-effective dietary fishmeal replacers. On the contrary, the use of fish oil by the aquaculture sector will probably increase in the long run albeit slowly; total usage will increase by more than 16 percent, from 782 000 tonnes (2.7 percent of total feeds by weight) in 2008 to the estimated 908 000 tonnes (1.3 percent of total aquafeeds for that year) by 2020. Increased usage will shift from salmonids to marine finfishes and crustaceans because of the current absence of cost-effective alternative lipid sources that are rich in long-chain polyunsaturated fatty acids. Increasing volumes of fishmeal and fish oil are likely to come from fisheries by-products, extracted from both wild capture and farmed fish. Estimates have been made that around 25 percent of fishmeal production in 2007 came from by-products. This will grow as it becomes increasingly viable to process this material.

It is estimated that the total usage of terrestrial animal by-product meals and oils within compound aquafeeds ranges between 0.15 and 0.30 million tonnes, or less than 1 percent of total global compound aquafeed production – clearly, there is considerable room for increased usage. In addition to meat meal, or, to a lesser extent meat and bone meal, ingredients such as blood meal, poultry by-product meal and poultry oil have all been very effective in feeds for a number of aquatic species.

Soybean meal is the most common source of plant proteins used in compound aquafeeds, with feeds for herbivorous and omnivorous fish species and crustaceans usually containing from 15 to 30 percent soybean meal, with a mean of 25 percent in 2008. In global usage terms, and based on a total compound aquafeed production of 27.1 million tonnes in 2007, it is estimated that the aquaculture feed sector consumed about 6.8 million tonnes of soybean meal (25.1 percent of total compound aquafeeds by weight). Other plant proteins being increasingly used include corn products, pulses, oilseed meals and protein from other cereals products.

Alternative lipid sources to fish oil are being used in greater amounts. Key alternatives include vegetable oils, preferably those with high omega-3 contents, and poultry oil. The use of oil from farmed fish offal is also a potential omega-3 source for other farmed fish. The production of marine microalgae or bacteria with very high contents of highly unsaturated fatty acids is currently expensive for use in most aquaculture feeds, but more cost efficient production methods will change this.

Prices for food and feed ingredients have been rising and are likely to continue to rise owing to the increasing demands from an increasing population, the diversion of some grains for use in biofuels, the increasing costs of production and transport, and the changes in global trade owing to the demand of food and raw materials from China and other emerging economies. The focus on carbohydrate-rich fractions for production of biofuels may indeed provide an opportunity to use protein fractions for feed ingredients.

Although current discussion on the use of marine products as aquafeed ingredients focuses on fishmeal and fish oil resources, the sustainability of the aquaculture sector is more likely to be linked with the sustained supply of terrestrial animal and plant proteins, oils and carbohydrate sources for aquafeeds. This is because a significant proportion of aquaculture production is of the non-carnivorous species. Therefore, aquaculture producing countries should place more emphasis to maximize the use of locally available feed-grade ingredient sources and use nutritionally sound and safe feed ingredients that can be sustainably produced and grow with the sector.