

Food and Agriculture Organization of the United Nations



Viale delle Terme di Caracalla, 00153 Rome, Italy - Tel: (+39) 06 57051 - E-mail: codex@fao.org - www.**codex**alimentarius.org

Agenda Item 7

CX/CF 19/13/7 March 2019

# JOINT FAO/WHO FOOD STANDARDS PROGRAMME CODEX COMMITTEE ON CONTAMINANTS IN FOODS

13<sup>th</sup> Session

# Yogyakarta, Indonesia, 29 April – 3 May 2019

## DRAFT CODE OF PRACTICE FOR THE REDUCTION OF 3-MONOCHLOROPROPANE-1,2- DIOL ESTERS (3-MCPDE) AND GLYCIDYL ESTERS (GE) IN REFINED OILS AND FOOD PRODUCTS MADE WITH REFINED OILS

(Prepared by the electronic working group led by the United States of America, Malaysia and European Union)

Codex members and Observers wishing to submit comments at Step 6 on this draft should do so as instructed in CL 2019/09-CF available on the Codex webpage/Circular Letters: <u>http://www.fao.org/fao-who-codexalimentarius/resources/circular-letters/en/</u>.

## BACKGROUND

1. CCCF11 (2017) agreed to endorse the proposal for new work for adoption by CAC40 (2017) on a Code of Practice for the Reduction of 3-monochloropropane-1,2-diol esters and glycidyl esters in refined oils and products made with refined oils, especially infant formula and to establish an Electronic Working Group (EWG), chaired by the United States of America and co-chaired by the European Union and Malaysia, working in English only, to follow-up on this new work<sup>1</sup>. CAC40 approved the new work<sup>2</sup>.

2. CCCF12 (2018) discussed the proposed COP and noted the following issues:

- Removal of the term "vegetable" from the title would broaden the scope of the COP and allow for the inclusion of non-vegetable oils, e.g. fish oils, since these are also refined oils used in food (including infant formula) and prone to formation of these contaminants;
- Inclusion of a new paragraph to the introduction of the COP explaining that the text could also be applicable to fish oils and subsequent additional references to fish oils could also be added elsewhere in the text as appropriate; and
- Proposed revisions to the text regarding specific practices on matters such as: low lipase activity, irrigation water, polar solvents, degumming, bleaching clay, or the inclusion of specific references to fish oils would be kept in square brackets and deferred to a re-established EWG for further discussion.

3. In order to cover all edible oils in the COP, Codex members and observers interested in this matter should provide additional examples of practices or information that were missing from the proposed COP to the Electronic Working Group for the further revision of the COP.

4. CCCF12 agreed<sup>3</sup>:

- i. that the scope of the COP covers refined oils and food products made with refined oils;
- ii. to forward the COP (with the sections in square brackets related to the points raised in 2<sup>nd</sup> and 3<sup>rd</sup> bullets of paragraph 2) to CAC41 for adoption at Step 5; and
- iii. to establish an EWG chaired by the United States of America, co-chaired by Malaysia and the European Union to revise the COP based on the comments and information submitted by Codex members and observers and to resolve all outstanding issues in order to submit a new draft for consideration by CCCF13.

5. CAC41 (2018) approved the proposed draft COP at Step 5 and advanced it to Step 6 for comments and further consideration by CCCF13 (2019).<sup>4</sup>

<sup>&</sup>lt;sup>1</sup> REP17/CF, para. 151 and Appendix V

<sup>&</sup>lt;sup>2</sup> REP17/CAC, Appendix VI

<sup>&</sup>lt;sup>3</sup> REP18/CF, paras. 99 – 102 and Appendix VI

<sup>&</sup>lt;sup>4</sup> REP18/CAC, para. 63, Appendix IV

## DISCUSSION

6. In developing this draft COP, the EWG considered changes proposed during CCCF12 and input provided through the electronic platform. These included:

- Broadening the scope of the COP to include fish oils;
- Modifications to mitigation measures; and
- Editorial changes.

## Fish oils

7. The EWG expanded the scope to include fish oils, including modifying the title to cover refined oils rather than vegetable oils, inserting a paragraph on the steps of fish oil production, and adding fish oils to relevant mitigation measures.

## Mitigation measures

8. The EWG made changes to draft mitigation practices, e.g. in paragraphs on the use of plant varieties with low lipase activity, the use of irrigation water during cultivation of palm fruits, the temperature used to inactivate lipases, techniques for washing crude vegetable oils, the type of acid used for degumming, and the oils most appropriate for post refining treatment. Modifications were made to reflect requested changes where technical information was provided to support the changes.

9. One outstanding issue identified in brackets in paragraph 28 is the use of water/alcohol (ethanol) mixture to remove chlorine compounds when washing crude vegetable oil.

10. Additional information was added on agricultural practices, specifically oilseeds. A statement on washing fruits prior to sterilization was also added.

## Editorial changes

11. The EWG made editorial changes, including removing examples of food products that contain 3-MCPDE and GE, removing extraneous references to infant formula, and removing unnecessary comparisons between palm oil and other oils.

## Recommendations

12. CCCF:

- note the revisions to the draft COP based on the discussion held and comments submitted at CCCF12 as well as the submissions made to the EWG as summarized in paragraphs 6 to 11; and
- consider the draft COP as set out in Appendix I together with comments submitted in reply to CL 2019/09-CF.

## APPENDIX I

## DRAFT CODE OF PRACTICE FOR THE REDUCTION OF 3-MONOCHLOROPROPANE-1,2- DIOL ESTERS (3-MCPDE) AND GLYCIDYL ESTERS (GE) IN REFINED OILS AND FOOD PRODUCTS MADE WITH REFINED OILS

## (COMMENTS REQUESTED THROUGH CL 2019/09-CF)

### INTRODUCTION

- 1. Edible oils are produced from various commodities, including fruits, seeds, nuts, and fish. Refining of edible oils (at temperatures of about 200°C or higher) can produce 3-monochloropropane-1,2-diol (MCPD) esters (3-MCPDE) and glycidyl esters (GE).
- Exposure to 3-MCPDE and GE can occur through consumption of refined oils and food products containing refined oils, including infant formula, dietary supplements, fried potato products, and fine bakery wares.
- 3. 3-MCPDE and 3-MCPD have toxic effects on the kidney and male reproductive organs, and 3-MCPD is a non-genotoxic carcinogen. GE and glycidol are genotoxic carcinogens.<sup>5</sup>
- 4. The 83rd JECFA session evaluated 3-MCPDE and GE and recommended that efforts to reduce 3-MCPDE and 3-MCPD in infant formula be implemented and that measures to reduce GE and glycidol in fats and oils continue, particularly when used in infant formula.
- Different types of unrefined oils have different capacities to form 3-MCPDE and GE during deodorization (part of the refining process). For example, refined palm oil has historically been reported to have higher concentrations of these esters than other refined edible oils.
- 6. Processing conditions during refining have an important effect on formation of 3-MCPDE and GE for all oil types. Most unrefined oils do not contain detectable levels of 3-MCPDE or GE.
- For vegetable oils, factors that contribute to capacity to form 3-MCPDE and GE during refining include climate, soil and growth conditions of source plants, plant genotype, and harvesting techniques. These factors all affect the levels of precursors of 3-MCPDE and GE (e.g. acylglycerols, chlorine-containing compounds).
- 8. 3-MCPDE forms primarily from the reaction between chlorine containing-compounds and acylglycerols like triacylglycerols (TAGs), diacylglycerols (DAGs), and monoacylglycerols (MAGs). GE forms primarily from DAGs or MAGs.
- 9. Some chlorinated compounds are precursors for 3-MCPDE formation. Oil palm fruits absorb chloride ions (in the form of chlorinated compounds) during tree growth from soil (including from fertilizers and pesticides) and from water, and these chloride ions are converted into reactive chlorinated compounds such as hydrochloric acid during oil refining, leading to formation of 3-MCPDE.
- 10. Oil fruits and seeds contain the enzyme lipase; lipase activity increases with fruit maturation, while the lipase activity in seeds remains stable. Lipase interacts with oil from mature fruits to rapidly degrade TAGs into free fatty acids (FFAs), DAGs, and MAGs, while the effect of lipase in seeds that are appropriately stored is negligible.
- 11. GE formation begins at about 200°C. When DAGs exceed 3-4% of total lipids, GE formation increases exponentially with increasing temperature. Formation of 3-MCPDE occurs at temperatures as low as 160-200°C, and formation does not increase with higher temperatures.
- 12. Because 3-MCPDE and GE are formed via different mechanisms, different mitigation strategies are needed to control their formation. Due to the different formation mechanisms, there generally is no relationship between relative levels of 3-MCPDE and GE in individual oil samples.
- 13. GE is generally easier to mitigate than 3-MCPDE, because its formation is directly associated with elevated temperatures (with formation beginning at about 200°C and becoming more significant at temperatures >230°C). GE is formed primarily from DAGs and does not require the presence of chlorinated compounds. Oils can be deodorized at temperatures below 230°C to avoid significant GE formation. However, it is not practical to decrease deodorization temperatures below the threshold that would lead to 3-MCPDE formation, as that could affect the quality and safety of the oil.

<sup>&</sup>lt;sup>5</sup> 3-MCPDE and GE, following consumption, are broken down in the body to 3-MCPD and glycidol, respectively.

- 14. Although 3-MCPDE and GE are primarily produced during deodorization, mitigation measures can be applied across the edible oil production chain, from agricultural practices for vegetable oils (e.g. cultivation, harvesting, transporting, and storing of fruits and seeds), to oil milling and refining (e.g. crude oil production and treatment, degumming/bleaching, and deodorization), as well as to post-refining measures (e.g. additional bleaching and deodorization and use of activated bleaching earth). Where possible, it may be best to remove precursors at the earlier stages of processing, to minimize the formation of 3-MCPDE and GE.
- 15. There are a wide range of methods to mitigate 3-MCPDE and GE, and the applicable methods used will vary depending on different conditions (including the oil source, the refining process, and the type of equipment in use). In addition, multiple methods may need to be combined to reduce 3-MCPDE and GE in oils. Manufacturers should select and apply those techniques that are appropriate to their own processes and products.
- 16. In concert with mitigation of 3-MCPDE and GE, it is important to also consider the overall impacts on the quality of refined oils and oil-based products, including product properties such as smell and taste, FFA profiles, stability attributes, levels of nutrients, and removal of contaminants such as pesticides and mycotoxins. In addition, environmental impacts of the recommended mitigation practices should be considered.
- 17. Although most work on mitigation of 3-MCPDE and GE in refined oils has focused on palm oil, some of the information and experience on mitigation of 3-MCPDE and GE in palm oil may be applicable to mitigation of 3-MCPDE and GE in other refined oils. Therefore, where data are available, this document specifies when the mitigation approach is specific to palm oil, and when it may be more widely applicable to other refined oils, including fish oils.

## SCOPE

- 18. This Code of Practice intends to provide national and local authorities, producers, manufacturers, and other relevant bodies with guidance to prevent and reduce formation of 3-MCPDE and GE in refined oils and food products made with refined oils. This guidance covers three strategies (where information is available) for reducing 3-MCPDE and GE formation:
  - (i) Good agricultural practices,
  - (ii) Good manufacturing practices, and
  - (iii) Selection and uses of refined oils in food products made from these oils.

# RECOMMENDED PRACTICES BASED ON GOOD AGRICULTURAL PRACTICES (GAP) AND GOOD MANUFACTURING PRACTICES (GMP)

- 19. Producing edible vegetable oils involves several major steps: cultivating, harvesting, transporting, and storing the fruits and seeds for further processing; palm oil milling where fruit is sterilized and crude oil is extracted; oilseed crushing where oilseeds are cleaned, ground, and steamed and crude oil is extracted; and refining of the crude oils.
- 20. Producing edible fish oils involves several major steps: harvesting the fish, steam cooking, dewatering/wet reduction (which involves pressing the liquor, separating the oil and water, and optionally, water washing the oil), and refining.
- 21. Refining edible oils consists of two main types; chemical or physical refining. Chemical refining consists of degumming (removal of phospholipids); neutralization (addition of hydroxide solution to remove FFAs through formation of soaps); bleaching (using clays) to reduce colors and remove remaining soaps and gums, trace metals, and degradation products; and deodorization (i.e. a steam-distillation process carried out at low pressures, 1.5-6.0 mbar, and elevated temperatures, 180 270°C) to remove FFA, colors, and volatile compounds. Physical refining involves degumming, bleaching, and deodorization, but does not have a neutralization step. While several factors influence the selection of physical refining, it is typically conducted on oils containing low levels of phospholipids.

## AGRICULTURAL PRACTICES FOR VEGETABLE OILS

- 22. When planting new trees, consider selecting oil palm plant varieties with low lipase activity in oil fruits, as low lipase activity is one factor that can reduce formation of FFAs and acylglycerol precursors.
- 23. Consider storing oil seeds at cool temperatures (e.g. < 25°C) and dry conditions (optimally <7% moisture content) to help ensure low levels of lipase.
- 24. During cultivation of palm fruits, minimize use of substances such as fertilizers, pesticides, and water that have excessive amounts of chlorine-containing compounds, in order to reduce chlorine uptake by the fruits.

- 25. Harvest oil palm fruits when they are at optimal ripeness. Minimize handling of the fruits to reduce bruising and prevent formation of FFAs. Avoid using damaged or overripe fruits, which may be associated with higher 3-MCPDE and GE formation.
- 26. Transport oil palm fruits to oil mills as soon as possible.

## **OIL MILLING AND REFINING**

## Crude Oil Production and Treatment

- 27. Following receipt of oil palm fruits at the mill, sterilize the fruits immediately (preferably within less than 2 days of harvesting) at temperatures at or below 140°C to inactivate lipases (with temperatures varying depending on the sterilization method). (Fruits may be washed prior to sterilization to remove chlorine precursors.) For oilseeds, clean, grind, and steam to inactivate lipases.
- 28. Wash crude vegetable oil with polar solvents like chlorine-free water [or water/alcohol (ethanol)mixtures] to remove chlorine-containing compounds.
- 29. Avoid using residual vegetable oil recovered from solvents or additional extractions, as this oil tends to have higher levels of precursors (e.g. DAGs, chlorine-containing compounds).
- 30. Assess precursors in batches of crude vegetable oils or fish oils (e.g. DAGs, chlorine-containing compounds) to adjust refining parameters and target appropriate mitigation strategies depending on the type of vegetable oil or fish oil being processed and processing conditions.
- 31. Preferentially refining crude vegetable oil or fish oil with low concentrations of precursors can produce finished oils with lower levels of 3-MCPDE and GE.

## Degumming

- 32. Use milder and less acidic conditions (e.g. either degumming with a low concentration of phosphoric, citric, or other acids or water degumming) to decrease 3-MCPDE in vegetable oils or fish oils. The concentration of acid needed depends on the quality of the crude vegetable oil or fish oil. Care should be taken to remove sufficient concentrations of phospholipids and acid to ensure quality.
- 33. Lowering the degumming temperature may help to reduce formation of 3-MCPDE precursors in vegetable oils; however, the degumming temperature will depend on numerous factors including the type of vegetable oil.

## Neutralization

34. Using chemical refining (i.e., neutralization) in place of physical refining can help remove precursors (e.g. chloride) and reduce FFAs, which may allow for lower deodorization temperatures in vegetable oils or fish oils. However, chemical refining can lead to excessive oil loss (especially for palm oil due to higher FFA levels) and may have a greater environmental impact than physical refining.

## Bleaching

- 35. Use of greater amounts of bleaching clay may reduce formation of 3-MCPDE and GE in all vegetable oils and fish oils. However, bleaching clays that contain significant amounts of chlorine-containing compounds should be avoided.
- 36. Use of more pH-neutral clays reduces the acidity and potential to form 3-MCPDE in palm oil, some seed oils, and fish oil.

## Deodorization

- 37. Consider conducting deodorization of vegetable oils and fish oils at reduced temperatures to decrease formation of GE. For example, it has been suggested to conduct deodorization at 190-230°C for vegetable oils and less than 190°C for fish oils. The temperature will vary depending on the residence time of oil.
- 38. As an alternative to traditional deodorization, conduct dual deodorization of vegetable oils and fish oils (2-stage deodorization) to reduce thermal load in oil. This includes both a shorter deodorization period at a higher temperature and a longer deodorization period at a lower temperature. Consideration needs to be given to parameters such as temperature, vacuum pressure, and time, and variations in equipment design and capability. Also, additional post processing may be required to reduce levels of GE.
- 39. Use of a stronger vacuum facilitates evaporation of volatile compounds due to the increased steam volume and rate of stripping, contributing to decreased deodorization temperatures and reduced formation of GE, and to a lesser extent 3-MCPDE, in vegetable and fish oils.

40. Short-path distillation<sup>6</sup> (in place of deodorization) has been shown to reduce the thermal load and formation of esters in fish oil, contributing to lower amounts of 3-MCPDE and GE in comparison to conventional deodorization. However, additional post processing using mild deodorization (e.g. 160-180°C) is needed to address sensory considerations.

## TREATMENT POST REFINING

- 41. The following recommended practices can be used for reducing levels of 3-MCPDE and GE in refined oils. These practices may be most appropriate for oils with 3-MCPDE and GE levels that are higher than desired for their intended use.
- 42. Additional bleaching and deodorization following initial bleaching and deodorization has been shown to achieve lower levels of GE in refined palm oil. (The second deodorization should occur at a lower temperature than the first deodorization.)
- 43. Application of activated bleaching earth during post refining has been shown to reduce GE in refined vegetable oils.
- 44. Use of short-path distillation (pressure: <1 mbar and temperature: 120 to 270°C) on bleached and deodorized vegetable oil can reduce acylglycerol components and levels of 3-MCPDE and GE.
- 45. Treatment of refined MCT (medium-chain triacylglycerols) oil with one or more bases (including carbonate, bicarbonate, hydroxide, oxide, alkoxide, amine bases, hydrides, and phosphines) converts 3-MCPDE and GE to TAGs.

## SELECTION AND USES OF REFINED OILS IN FOOD PRODUCTS MADE FROM THESE OILS

## Oil selection

46. Selecting refined vegetable oils and fish oils with low levels of 3-MCPDE and GE (e.g. either through natural occurrence or through application of mitigation measures) results in lower levels of 3-MCPDE and GE in finished products containing these oils. For example, variation in levels of 3-MCPDE and GE in infant formula has been observed, which may be due to the use of oils with different levels of 3-MCPDE and GE; therefore, selection of oils low in 3-MCPDE and GE can result in infant formulas with lower 3-MCPDE and GE levels. However, manufacturers also may have to consider quality or compositional factors. For example, for infant formula, refined oils are selected by manufacturers to ensure these products meet compositional criteria, e.g. national criteria or those established in the *Standard for Infant Formula and Formulas for Special Medical Purposes Intended for Infants* (CXS 72-1981).

## **Processing modifications**

- 47. Reducing the amount of refined vegetable oils and fish oils used in finished products is expected to reduce the levels of 3-MCPDE and GE in the finished product. However, this could impact the organoleptic or nutritional qualities of the finished products.
- 48. Use of refined vegetable oils themselves during frying does not contribute to formation of additional 3-MCPDE and GE, but rather the formation of additional 3-MCPDE during frying may result from the type of food that is fried (e.g. meat and fish products).

<sup>&</sup>lt;sup>6</sup> Short-path distillation enables gentle removal of volatile compounds at relatively low temperatures. This is accomplished through reduced pressure, where the boiling point of the compound to be separated is lowered and there is increased efficiency due to the short distance between the evaporator and the condenser surface.

7

# POTENTIAL MITIGATION MEASURES FOR REDUCING 3-MCPDE AND GE The mitigation measures are not listed in order of priority. It is recommended that reduction measures be tested to identify the most successful for your own product.

Productior Stage	Mitigation measures
AGRICULTURAL PRACTICES FOR VEGEGETABLE OILS	<ul> <li>Select oil palm plant varieties with low lipase activity.</li> <li>Store oil seeds at cool temperatures and dry conditions.</li> <li>Minimize use of substances such as fertilizers, pesticides, and irrigation water that contain excessive amounts of chlorine-containing compounds during oil palm cultivation.</li> <li>Harvest oil palm fruits when they are at optimal ripeness. Minimize handling of the fruit. Avoid using damaged or overripe fruit.</li> <li>Transport oil palm fruits to oil mills as soon as possible.</li> </ul>
OIL MILLING AND REFINING	<ul> <li>Crude Oil Production and Treatment</li> <li>Sterilize oil palm fruit at temperatures at or below 140°C.</li> <li>Wash crude vegetable oil with polar solvents (e.g., chlorine-free water [or water/alcohol mixtures]).</li> <li>Avoid using residual vegetable oil recovered from solvents or extractions.</li> <li>Assess precursors (e.g., DAGs and chlorine compounds) in batches of crude vegetable oil or fish oil to adjust refining parameters.</li> <li>Preferentially refine crude vegetable oil or fish oil with low</li> </ul>
	<ul> <li>Degumming</li> <li>Use milder and less acidic conditions (e.g., either degumming with a low concentration of acid or water degumming) in vegetable oils or fish oils.</li> <li>Lower the degumming temperature in vegetable oils.</li> </ul>
OIL M	<ul> <li>Neutralization</li> <li>Use chemical refining (i.e., neutralization) in place of physical refining in vegetable oils or fish oils.</li> </ul>
	<ul> <li>Bleaching</li> <li>Use greater amounts of bleaching clay in vegetable oils and fish oils.</li> <li>Use more pH-neutral clays to reduce acidity in palm oils, some seed oils, and fish oils.</li> </ul>

# POTENTIAL MITIGATION MEASURES FOR REDUCING 3-MCPDE AND GE

The mitigation measures are not listed in order of priority.

It is recommended that reduction measures be tested to identify the most successful for your own product.

Production Stage	Mitigation measures
OIL MILLING AND REFINING	<ul> <li>Deodorization</li> <li>Conduct deodorization of vegetable oils or fish oils at reduced temperatures. The temperatures will vary depending on residence time of oil.</li> <li>Conduct dual deodorization of vegetable oils and fish oils (2-stage deodorization) as an alternative to traditional deodorization.</li> <li>Use a stronger vacuum to facilitate evaporation of volatile compounds and to contribute to decreased deodorization temperatures in vegetable oils and fish oils.</li> <li>Use short-path distillation (in place of deodorization) to reduce the thermal load in fish oil.</li> </ul>
TREATMENT POST REFINING	<ul> <li>Conduct additional bleaching and deodorization following initial bleaching and deodorization of refined palm oil.</li> <li>Apply activated bleaching clay to refined vegetable oils.</li> <li>Use short-path distillation on bleached and deodorized vegetable oils.</li> <li>Treat refined MCT (medium-chain triglyceride) oil with bases to convert 3-MCPDE and GE to triglycerides.</li> </ul>
SELECTION AND USES OF REFINED OILS	<ul> <li>OIL SELECTION <ul> <li>Select refined vegetable oils or fish oils with lower levels of 3-MCPDE and GE.</li> </ul> </li> <li>PROCESS MODIFICATIONS <ul> <li>Reduce the amount of refined vegetable oils or fish oils in finished products.</li> </ul> </li> </ul>

## **APPENDIX II**

#### LIST OF PARTICIPANTS

CHAIR United States Eileen Abt U.S. Food and Drug Administration Center for Food Safety and Applied Nutrition eileen.abt@fda.hhs.gov

CO-CHAIR European Union Frans Verstraete European Commission frans.verstraete@ec.europa.eu

CO-CHAIR Malaysia Rosidah Radzian Malaysian Palm Oil Board rosidah@mpob.gov.my

#### Argentina

Lic. Silvana Ruarte Jefe de Servicio Analítica de Alimentos Departamento Control y Desarrollo Dirección de Fiscalización, Vigilancia y Gestión de Riesgo Instituto Nacional de Alimentos sruarte@anmat.gov.ar; codex@magyp.gob.ar

#### Australia

Matthew O'Mullane Section Manager Food Standards Australia New Zealand Matthew.O'Mullane@foodstandards.gov.au

Glenn Paul Stanley Deputy Section Manager Food Standards Australia New Zealand <u>Glenn.Stanley@foodstandards.gov.au</u>

Luisa Trevisan Food Standards Australia New Zealand

#### Belgium

Korati Safia Regulatory Expert Federal Public Service Health Food Safety Safia.korati@sante.belgique.be

#### Brazil

Lígia Lindner Schreiner Risk Assessment Manager Brazilian Health Regulatory Agency - ANVISA ligia.schreiner@anvisa.gov.br

Larissa Bertollo Gomes Porto Health Regulation Specialist Brazilian Health Regulatory Agency - ANVISA Jarissa.porto@anvisa.gov.br

#### Canada

Luc Pelletier Scientific Evaluator, Food Contaminants Section Bureau of Chemical Safety Health Canada <u>luc.pelletier@hc-sc.gc.ca</u> Elizabeth Elliott Head, Food Contaminants Section Bureau of Chemical Safety Health Canada Elizabeth.Elliott@hc-sc.gc.ca

Mark Feeley Associate Director Bureau of Chemical Safety Food Directorate Health Canada Mark.Feeley@canada.ca

#### China

Yongning Wu Professor, Chief Scientist China National Center of Food Safety Risk Assessment (CFSA) wuyongning@cfsa.net.cn

Stephen Chung Wai-Cheung Senior Chemist Center for Food Safety

Ms. Yi Shao Research Associate China National Center of Food Safety Risk Assessment (CFSA) <u>shaoyi@cfsa.net.cn</u>

Mr. Jingguang Li Researcher China National Center for Food Safety Risk Assessment (CFSA) Iijg@cfsa.net.cn

#### Colombia

Wilmer Humberto Fajardo Jimenez Coordinator Grupo del Sistema de Analisis de Riesgos Quimicos en Alimentos y Bebidas Invima wfajardo@invima.gov.co

Sonia Buitrago Contratista Minsterio de Salud y Proteccion Social <u>sbuitrago@minsalud.gov.co</u> **Dominican Republic** 

Fatima del Rosario Cabrera General Directorate of Medicines, Food and Health Products Ministry of Public Health and Social Assistance <u>codex.pccdor@msp.gob.do</u>

Elsa Maritza Piantini

## **European Union**

Frans Verstraete European Commission Director General for Health and Consumers <u>frans.verstraete@ec.europa.eu</u>

Germany Annette Rexroth Senior Officer Federal Ministry for Nutrition and Agriculture Annette.Rexroth@bmel.bund.de

#### India

National Codex Contact Point Food Safety Standards and Authority of India Ministry of Health and Family Welfare Codex-india@nic.in

#### Indonesia

Dyah Setyowati Head of Section of Food Standard Harmonization Directorate of Processed Food Standardization National Agency of Drug and Food Control codexbpom@yahoo.com

#### Japan

Yukiko Yamada Advisor to Vice-Minister Ministry of Agriculture, Forestry and Fisheries of Japan yukiko\_yamada530@maff.go.jp

Yoshiyuki TAKAGISHI Associate Director Food Safety Policy Division, Food Safety and Consumer Affairs Bureau, Ministry of Agriculture, Forestry and Fisheries of Japan yoshiyuki takagis500@maff.go.jp

## Kazakhstan

Zhanar Tolysbayeva Technical Expert Ministry of Healthcare tolyzhan@gmail.com

Korea Min Yoo Researcher Ministry of Food and Drug Safety <u>minyoo83@korea.kr</u> Codexkorea@korea.kr

Lee Yeonkyi Ministry of Food and Drug Safety Malaysia Rosidah Radzian Director, Product Development & Advisory Services Division Malaysian Palm Oil Board rosidah@mpob.gov.my

Raznim Arni Abd. Razak Senior Research Officer Malaysian Palm Oil Board raznim@mpob.gov.my

Ainie Kuntom Senior Research Fellow Malaysian Palm Oil Board <u>ainie@mpob.gov.my</u>

Raizawanis Abdul Rahman Principal Assistant Director Malaysian Palm Oil Board raizawanis@moh.gov.my

#### Mexico

Tania Daniela Fosado Soriano Economic Secretary

## New Zealand

Andrew Pearson Manager, Food Risk Assessment (Acting) Ministry for Primary Industries Andrew.pearson@mpi.govt.nz

Jeane Nicolas Senior Advisor Toxicology Ministry for Primary Industries Jeane.nicolas@mpi.govt.nz

#### Norway

Codex Contact Point Norwegian Food Safety Authority

Oda Walle Almeland Mattilsynet

**Peru** Javier Aguilar Zapata Ministry of Agriculture

Jorge Pastor Miranda Ministry of Agriculture

#### Singapore Yat Yun Wei Senior Analytical Scientist Yat yun\_wei@hsa.gov.sg

Spain David Merino Fernandez Risk Manager Agencia Española de Consumo, Seguridad Alimentaria y Nutrición dmerrino@msssi.es

#### Switzerland

Lucia Klauser Scientific Officer Federal Food Safety and Veterinary Office Iucia.klauser@blv.admin.ch

#### Thailand

Korwadee Phonkliang Standards Officer, Officer of Standard Development National Bureau of Agricultural Commodity and Food Standards

codex@acfs.go.th; korwadeep@hotmail.com

#### United Kingdom Izaak Fryer-Kanssen Chemical Contaminants and Residues Office Food Standards Agency Izaak.fryerkanssen@food.gov.uk

## **United States of America**

Eileen Abt U.S. Food and Drug Administration Center for Food Safety and Applied Nutrition <u>eileen.abt@fda.hhs.gov</u>

Henry Kim U.S. Food and Drug Administration Center for Food Safety and Applied Nutrition henry.kim@fda.hhs.gov

Shaun MacMahon U.S. Food and Drug Administration Center for Food Safety and Applied Nutrition shaun.macmahon@fda.hhs.gov

Lauren Posnick Robin U.S. Delegate U.S. Food and Drug Administration Center for Food Safety and Applied Nutrition lauren.robin@fda.hhs.gov

Paul South U.S. Food and Drug Administration Center for Food Safety and Applied Nutrition paul.south@fda.hhs.gov

#### AOCS (American Oil Chemists' Society)

Scott Bloomer Director, Technical Services Scott.bloomer@aocs.org

# FAO (JECFA)

Markus Lipp Senior Officer Agriculture and Consumer Protection Department Food and Agriculture Organization of the UN Viale delle Terme di Caracalla markus.lipp@fao.org Vittorio Fattori Food Safety Officer Agriculture and Consumer Protection Department Food and Agriculture Organization of the UN Viale delle Terme di Caracalla vittorio.fattori@fao.org

# FEDIOL

Kalila Hajjar Senior Manager, Scientific and Regulatory Affairs <u>khajjar@fediol.eu</u>

## FoodDrinkEurope

Eoin Keane Manager Food Policy, Science and R&D <u>e.keane@fooddrinkeurope.eu</u>

Alejandro Rodarte

GOED (Global Organization for EPA and DHA Omega-3s) Gerard Bannenberg Director of Compliance and Scientific Outreach gerard@goedomega3.com

IFFO (The Marine Ingredients Organisation) Alejandra Aguilar Solis Regulatory Research Associate aaguilar@iffo.net

IFT (Institute of Food Technology) James R. Coughlin Coughlin & Associates jrcoughlin@cox.net

ICGMA (International Council of Grocery Manufacturers Associations/) Nichole Mitchell Lead Delegate International Council of Grocery Manufacturers Associations (ICGMA) nmitchell@gmaonline.org

ISDI (International Special Dietary Foods Industries) Jean Christophe Kremer Secretary General secretariat@isdi.org

WHO (JECFA) Angelika Tritscher Coordinator, Risk Assessment and Management tritschera@who.int