

Molecular breeding: Challenges and perspectives

Roberto Tuberosa

Dept. of Agroenvironmental Sciences & Technology
University of Bologna, Italy



The ABDC Congress
1-4 March 2010
Gudalajara, Mexico



Definition

In molecular or marker-assisted breeding (**MB**), DNA markers are used as a substitute for phenotypic selection and to accelerate the release of improved cultivars.

Marker-assisted selection (**MAS**). Selection of individuals with specific alleles for traits controlled by a limited number of loci (up to 6-8).

Marker-assisted backcrossing (**MABC**). Transfer of a limited number of loci (e.g. transgene, disease resistance loci, etc.) from one genetic background to another.

Marker-assisted recurrent selection (**MARS**). The identification and selection of several genomic regions (up to 20 or even more) for complex traits within a single population.

Critical factors for the success of molecular breeding

Germplasm

- Elite vs. non-elite
- Wild relatives
- Structure

Phenotyping

- Identify key traits
- Stress management
- Accuracy


MAS and conventional breeding are not mutually exclusive; instead, they are complementary under most breeding schemes (Ribaut et al., COPB 2010).

- High-throughput level
- Cost-effectiveness
- Informatics
- Data management

Breeding

- Choice of MB strategy
- Breeders fluent with MB
- Stakeholders involvement
- Seed industry for delivery

From gene/QTL discovery to cultivar release



Issues for using molecular breeding in crop improvement


- Cost of each single data point
- Availability and complexity of molecular platforms
- Reliability of marker profiling and scoring (SSR vs. SNP)
- Degree of polymorphism
- Availability of equipment
- Availability of technical expertise
- Integration with breeding programs

MAS is preferable to conventional breeding when:

- Phenotypic screening is particularly expensive or laborious (e.g. root traits, osmotic adjustment, disease resistance, etc.)
- Phenotyping is possible only after flowering
- Linked marker is co-dominant and target gene is dominant
- Strong influence of environment (low heritability)
- Pyramiding multiple resistance genes
- Marker profiles can be obtained from seeds


Marker-assisted backcrossing (MABC)

Susceptible line




X


Resistant line



2-3 cycles of MABC



Gene/QTL



Los Angeles Times April 13, 2008

QTLs

An aerial photograph of a vast agricultural field filled with numerous combine harvesters. The machines are arranged in a grid-like pattern, moving across the field and kicking up dust or chaff. The overall scene is one of intense agricultural activity. Overlaid on the left side of the image is the text 'QTLs' in large, bold, red letters with a black outline.

It follows that a given QTL can have positive, null, or negative effects depending on the environment. This complication has slowed considerably the utilization of QTL data for breeding.

Collins et al. (2008). Plant Physiol. 147: 469-486.

Nested Association Mapping (NAM)



large

large

small

small

Additive effect

Additive effect

Genome-wide selection

Genome wide selection (**GWS**) was first suggested in animal breeding (Meuwissen, 2001)

GWS focuses exclusively on prediction of performance based on as many loci as possible (unlimited number)

GWS avoids QTL mapping altogether

- In GWS, the joint effects of all markers are fitted as random effects in a linear model
- Trait values are predicted from a weighed index calculated for each marker

Genome-wide selection

- Simulation studies have shown that across different numbers of QTLs (from 20 up to 100) and levels of h^2 , responses to GWS were 18 to 43% larger than MARS (Bernardo and Yu, 2007)
- GWS was found more useful with complex traits and low h^2
- GWS focuses on the genetic improvement of quantitative traits rather than on understanding their genetic basis
- GWS and QTL discovery are not mutually exclusive.

...looking ahead...



Future challenges in molecular breeding

- Developing high-density SNP platforms in target crops
- Platforms to further reduce the cost of SNP profiling
- Multiparental populations for a “breeding-by-design” approach
- Optimize MARS and GWS procedures
- Identifying genes/QTLs for yield potential and stability
- Identifying high-yielding, stable genotypes (low G x E x M)
- Integration of modeling (es. QTL effects)
- Closer collaboration with breeders in DCs
- Training of breeders to integrate MB in their programs

Tying it all together

- On a case-by-case basis, develop appropriate breeding strategies for the improvement of multiple traits
 - Genetic dissection of the developmental and functional correlation between multiple traits, including yield and yield stability
 - Understanding of genetic networks for correlated traits
 - Construction of selection indices across traits
- Delivering improved germplasm and new cultivars will require a close collaboration and an effective coordination among all the relevant stakeholders.
- Only an appropriate engagement of the breeders and seed companies in target areas will allow the above to materialize and tangibly impact crops yield and yield quality.

A vibrant illustration of a cornfield. The corn plants are green with long, pointed leaves and yellow tassels. The background is a clear blue sky. The text "Thanks for your attention!" is written in a bold, 3D, orange-to-yellow gradient font, slanted across the middle of the image. At the bottom, there is a decorative border with a repeating geometric pattern in red and black.

Thanks for your attention!