

FAO Specifications for Plant Protection Products

OXADIXYL (AGP:CP/358)



FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS
Rome, 1998

DISCLAIMER ¹

FAO specifications are developed with the basic objective of promoting, as far as practicable, the manufacture, distribution and use of pesticides that meet basic quality requirements.

Compliance with the specifications does not constitute an endorsement or warranty of the fitness of a particular pesticide for a particular purpose, including its suitability for the control of any given pest, or its suitability for use in a particular area. Owing to the complexity of the problems involved, the suitability of pesticides for a particular purpose and the content of the labelling instructions must be decided at the national or provincial level.

Furthermore, pesticides which are manufactured to comply with these specifications are not exempted from any safety regulation or other legal or administrative provision applicable to their manufacture, sale, transportation, storage, handling, preparation and/or use.

FAO disclaims any and all liability for any injury, death, loss, damage or other prejudice of any kind that may arise as a result of, or in connection with, the manufacture, sale, transportation, storage, handling, preparation and/or use of pesticides which are found, or are claimed, to have been manufactured to comply with these specifications.

Additionally, FAO wishes to alert users to the fact that improper storage, handling, preparation and/or use of pesticides can result in either a lowering or complete loss of safety and/or efficacy.

FAO is not responsible, and does not accept any liability, for the testing of pesticides for compliance with the specifications, nor for any methods recommended and/or used for testing compliance. As a result, FAO does not in any way warrant or represent that any pesticide claimed to comply with a FAO specification actually does so.

¹ This disclaimer applies to all specifications published by FAO.

INTRODUCTION TO FAO SPECIFICATIONS DEVELOPED UNDER THE OLD PROCEDURE

Between 1975 and 2000, FAO published booklets of specifications for technical materials and related formulations of plant protection products. Revisions of, and additions to, already published specifications will be issued when necessary. However, all changes and revisions of FAO specifications are now subject to the new procedure described in the *Manual on the development and use of FAO and WHO Specifications for Plant Protection Products*, FAO Plant Production and Protection Paper No. 173, Rome 2002 (*Revised First Edition* available only on the FAO home page of the Internet at: <http://www.fao.org/ag/agp/agpp/pesticid/>)

FAO specifications developed under the old procedure are based on the requirements defined in the Fourth Edition of the *Manual on the development and use of FAO specifications for plant protection products*, Plant Production and Protection Paper No. 128, Rome 1995.

This manual contained detailed definitions and other essential background information on basic procedures and technical principles adopted by the group on Pesticide Specifications of the FAO Panel of Experts on Pesticide Specifications, Registration Requirements, Application Standards and Prior Informed Consent, such as:

1. Categories of Specifications (Section 3.1 of the Manual)

FAO Tentative Specifications (Code 'S/T', formerly 'TS') are those which have been recommended by FAO as preliminary specifications and which are based on minimum requirements. The methods of analysis cited are normally supplied by the manufacturer or may already have been published or be the subject of collaborative work.

FAO Provisional Specifications [Code 'S/P', formerly ('S')] are those for which more evidence of the necessary parameters is available and where some collaborative study of the methods of analysis has been carried out.

FAO (full) Specifications (Code 'S/F', formerly 'S').

Specifications that have all necessary requirements together with CIPAC (full) methods, or other collaboratively studied (proven) methods.^{2,3}

Wherever possible, standards for apparatus and common names for pesticides are those approved by the International Organization for Standardization (ISO).

2. Expression of active ingredient content (Section 4.2.5 of the Manual)

- for solids, liquid technical materials, volatile liquids (of maximum boiling point 50°C) and viscous liquids (with minimum kinematic viscosity of $1 \times 10^3 \text{ m}^2/\text{s}$ at 20°C) the FAO Specification shall be based on expression of the content as g/kg;

- for all other liquids the active ingredient content of the product shall be declared in terms of g/kg *or* g/l at 20°C. If the customer requires both g/kg *and* g/l at 20°C, then in case of dispute the analytical results shall be calculated as g/kg.

3. Tolerance on content (Section 4.2.7 of the Manual)

A declared content of active ingredient must be included in all specifications, and one of the problems immediately arising is the level of tolerance acceptable about the nominal figure. The tolerance is influenced by (a) the reproducibility of the method of analysis, (b) the sampling error and (c) the manufacturing variance.

Allowable variations in analytical results (i.e. tolerances in content of active ingredient) with respect to specific pesticide consignments are intended to cover reasonable variations in the contents of active ingredients. For examples of such tolerances, see the table in Section 4.2.7 of the Manual.

4. Containers/packaging

FAO guidelines are in preparation.

Containers shall comply with pertinent national and international transport and safety regulations.

Technical materials, dustable powders and granules

Containers shall be suitable, clean, dry and as specified, and shall not adversely affect, or be affected by, the contents, but shall adequately protect them against external conditions.

Wettable powders

The product shall be packed in suitable, clean, dry containers as specified in the order. The container shall provide all necessary protection against compaction, atmospheric moisture, loss by vaporization and/or contamination to ensure that the product suffers no deterioration under normal transit and storage conditions.

The product shall be protected by an adequate moisture barrier. This may be a suitable bag of polyethylene or alternative means of giving equal or better protection.

Solutions and emulsifiable concentrates

Containers shall be lined, where necessary, with a suitable material, or the interior surfaces shall be treated to prevent corrosion and/or deterioration of the contents.

Additional information should be given in all specifications where particular pesticides present problems in packaging.

5. Biological information

Phytotoxicity

No test can be specified to cover the possible phytotoxicity of a formulation to all crops. When a crop is not mentioned in the instructions for use, purchasers should check with the supplier that the material is suitable, always provided that such a use is not restricted or legally forbidden.

Wetting of crops

The dilute spray should satisfactorily wet the leaves of the specified crops when used in accordance with the instructions. Test method MT 53.2, CIPAC F, p.162, may be useful.

¹ *Should national pesticide specifications developed from these approved FAO specifications deviate from them, the National Authority responsible for making such changes is requested to inform the FAO Plant Protection Service of the nature of, and the reasons for, the modifications.*

² *Methods of analysis and miscellaneous techniques referred to in these specifications have been developed and adopted by CIPAC (Collaborative International Pesticides Analytical Council Ltd.). See CIPAC Handbooks 1 (1970), 1A (1980), 1B (1983), 1C (1985), D (1988), E (1993), F (1995), G (1995), CIPAC Proceedings 1980 and 1981, obtainable from Black Bear Press Limited, King's Hedges Road, Cambridge CB4 2PQ, England. The page numbers of specific methods are given in parentheses in the specifications. Copies of methods not yet published can be obtained from the FAO Plant Protection Service.*

³ *Information on standard waters for laboratory evaluation of pesticidal formulations will be found in CIPAC Monograph 1, Standard Waters and an FAO Survey on Naturally Occurring Waters (1972), Black Bear Press Limited, King's Hedges Road, Cambridge CB4 2PQ, England.*

SUBMISSION OF DRAFT SPECIFICATIONS TO FAO

Any organization, commercial firm or interested individual is encouraged to submit relevant specifications, or proposals for revision of existing specifications, for pesticide products for consideration and possible adoption by FAO. Correspondence should be addressed to the Pesticide Management Group, Plant Production and Protection Division, FAO, Viale delle Terme di Caracalla, 00153 Rome, Italy.

General guidelines on preparing draft specifications are given in the *Manual on the development and use of FAO and WHO Specifications for Plant Protection Products*, FAO Plant Production and Protection Paper No. 173, Rome 2002 (Revised First Edition available only on the FAO home page of the Internet at: <http://www.fao.org/ag/agp/agpp/pesticid/>).

Specifications which are considered suitable for further processing are assigned priorities and circulated to appropriate organizations and specialists to comment. Comments, together with other relevant information, are then reviewed in detail by the Group on Specifications of the FAO Panel of Experts on Pesticide Specifications, Registration Requirements, Application Standards and Prior Informed Consent. The drafts are converted into FAO Provisional Specifications, or full FAO Specifications.


OXADIXYL

2-methoxy-N-(2-oxo-1,3-oxazolidin-3-yl)acet-2',6'-xylidide

INFORMATION

COMMON NAME: oxadixyl (ISO)

STRUCTURAL FORMULA:



EMPIRICAL FORMULA: C₁₄H₁₈ N₂ O₄

RMM: 278.3

CAS REGISTRY NUMBER: 77732-09-3

CIPAC CODE NUMBER: 397

CHEMICAL NAMES:

2-methoxy-N-(2-oxo-1,3-oxazolidin-3-yl)acet-2',6'-xylidide (IUPAC)

N-(2,6-dimethylphenyl)-2-methoxy-N-(2-oxo-3 oxazolidinyl)acetamide (CA)

OXADIXYL TECHNICAL

FAO Specification 397/TC/S/F (1997)

1. DESCRIPTION

The material shall consist of oxadixyl together with related manufacturing impurities and shall be in the form of a light beige to brownish crystalline powder, free from visible extraneous matter and added modifying agents.

2. ACTIVE INGREDIENT

2.1. Identity tests (397/TC/M/2, CIPAC G, p.122)

An identity test is required if the identity of the active ingredient is in doubt. Then it shall comply with at least one additional test, e.g. HPLC or IR spectrum.

2.2. Oxadixyl (397/TC/M/3, CIPAC G, p.122)

The oxadixyl content shall be declared (not less than 960 g/kg) and, when determined, the content obtained shall not differ from that declared by more than ± 25 g/kg.

3. IMPURITIES

No relevant impurities.

OXADIXYL WETTABLE POWDERS

FAO Specification 397/WP/S/F (1997)

1. DESCRIPTION

The material shall consist of a homogeneous mixture of technical oxadixyl, complying with the requirements of FAO specification 397/TC/S/F (1997) together with filler(s) and any other necessary formulants. It shall be in the form of a fine powder free from visible extraneous matter and hard lumps.

2. ACTIVE INGREDIENT

2.1. Identity tests (397/WP/M/2, CIPAC G, p.125)

An identity test is required if the identity of the active ingredient is in doubt,. Then the isolated active ingredient shall comply with at least one additional test, e.g. HPLC or IR spectrum.

2.2. Oxadixyl (397/WP/M/3, CIPAC G, p. 125)

The oxadixyl content shall be declared (g/kg) and, when determined, the content obtained shall not differ from that declared by more than the following amounts:

<u>Declared content</u>	<u>Permitted tolerance</u>
Above 25 up to 100 g/kg	± 10 % of the declared content
Above 100 up to 250 g/kg	± 6 % of the declared content
Above 250 up to 500 g/kg	± 5 % of the declared content

3. IMPURITIES

No relevant.

4. PHYSICAL PROPERTIES

4.1. pH range (MT 75, CIPAC F, p.205)

pH range: 5.0 to 10.0.

4.2. Wet sieve test (MT 59.3, CIPAC F, p.179)

Maximum: 1 % retained on a 75 µm test sieve.

4.3. Suspensibility (MT 15.1, CIPAC F, p.45. Notes 1 & 2)

A minimum of 60 % of the oxadixyl content found under 2.2. shall be in suspension after 30 min. at $30 \pm 2^\circ\text{C}$ in CIPAC standard water D.

Alternatively, if the buyer requires other CIPAC standard waters or temperature to be used, then this shall be specified when ordering.

4.4. Persistent foam (MT 47.2, CIPAC F, p.152. Note 3)

Maximum: 25 ml after 1 min.

4.5. Wetting of the material (MT 53.3.1, CIPAC F, p.164)

It shall be completely wetted in 1 min. without swirling.

5. STORAGE STABILITY

5.1. Stability at 54 °C (MT 46.1.1, CIPAC F, p.149)

After storage at $54 \pm 2^\circ\text{C}$ for 14 days, the determined average active ingredient content must not be lower than 97 % relative to the determined average content found under 2.2. (Note 4) and the product shall continue to comply with 4.1., 4.2., 4.3. and 4.5.

Note 1 The product should be tested at the highest and lowest rates of use recommended by the supplier, provided this does not exceed the conditions given in Method MT 15.1.

Note 2 Chemical assay is the only fully reliable method to measure the mass of active ingredient still in suspension. However, simpler methods such as gravimetric and solvent extraction determination may be used on a routine basis provided that these methods have been shown to give equal results to those of the chemical assay method. In case of dispute, the chemical method shall be the “Referee method”.

Note 3 The mass of the sample to be used in the test should be specified at the highest rate of use recommended by the supplier.

Note 4 Samples of the product taken before and after the storage stability test should be analyzed together after the test, to reduce the analytical error.

OXADIXYL AQUEOUS SUSPENSION CONCENTRATES

FAO Specification 397/SC/S/F (1997)

1. DESCRIPTION

The material shall consist of a suspension of fine particles of technical oxadixyl, complying with the requirements of FAO specification 397/TC/S/F (1997) in an aqueous phase together with suitable formulants. After gentle agitation the material shall be homogeneous (Note 1) and suitable for further dilution in water.

2. ACTIVE INGREDIENT

2.1. Identity tests (397/WP/M/2, CIPAC G, p.125. Note 2)

An identity test is required if the identity of the active ingredient is in doubt,. Then the isolated active ingredient shall comply with at least one additional test, e.g. HPLC or IR spectrum.

2.2. Oxadixyl (397/WP/M/3, CIPAC G, p. 122. Note 2)

The oxadixyl content shall be declared (g/kg or g/l at $20 \pm 0.5^{\circ}\text{C}$. Note 3) and, when determined, the content obtained shall not differ from that declared by more than the following amounts:

<u>Declared content</u>	<u>Permitted tolerance</u>
Above 25 up to 100 g/kg or g/l	$\pm 10\%$ of the declared content
Above 100 up to 250 g/kg or g/l	$\pm 6\%$ of the declared content
Above 250 up to 500 g/kg or g/l	$\pm 5\%$ of the declared content

3. IMPURITIES

Not relevant.

4. PHYSICAL PROPERTIES

4.1. Mass per millilitre at 20°C (MT 3.3, CIPAC F, p.18)

If required, the range of the mass per millilitre (g/ml) at $20 \pm 0.5^{\circ}\text{C}$ shall be declared.

4.2. pH range (MT 75, CIPAC F, p.205)

pH range: 5.0 to 10.0.

4.3. Rinsability (MT 148.1, CIPAC H, to be published)

Maximum non-volatile residue: 0.01 %.

4.4. Spontaneity of dispersion (MT 160, CIPAC F, p.391. Note 4)

A minimum of 90 % of the oxadixyl content found under 2.2. shall be in suspension after 5 min. in CIPAC Standard Water D at $30 \pm 2^\circ\text{C}$ (Note 5).

4.5. Suspensibility (MT 161, CIPAC F, p.394. Note 4)

A minimum of 70 % of the oxadixyl content found under 2.2. shall be in suspension after 30 min. in CIPAC Standard Water D at $30 \pm 2^\circ\text{C}$ (Note 5).

4.6. Wet sieve test (MT 59.3, CIPAC F, p.179)

Maximum: 1 % retained on a 75 μm sieve.

4.7. Persistent foam (MT 47.2, CIPAC F, p.152. Note 6)

Maximum: 25 ml after 1 min.

5. STORAGE STABILITY

5.1. Stability at 0 °C (MT 39.1, CIPAC F, p.128)

After storage at $0 \pm 1^\circ\text{C}$ for 7 days, the product shall continue to comply with 4.4., 4.5. and 4.6.

5.2. Stability at 54 °C (MT 46.1.2, CIPAC F, p.149)

After storage at $54 \pm 2^\circ\text{C}$ for 14 days, the determined average active ingredient content must not be lower than 97 % relative to the determined average content found under 2.2. (Note 7) and the product shall continue to comply with 4.2., 4.3., 4.4., 4.5. and 4.6.

Note 1 Before sampling to verify the product quality, inspect the commercial container carefully. On standing, suspension concentrates usually develop a concentration gradient from the top to the bottom of the container. This may even result in the appearance of a clear liquid on the top and/or of sediment at the bottom. Therefore, before sampling, the product must be homogenized according to the

instructions given by the manufacturer or, in the absence of such instructions, by gentle shaking of the commercial container (for example by inverting the closed container several times). Large containers must be opened and stirred adequately. After this procedure the container should not contain a sticky layer of non-dispersed product at the bottom. A suitable and simple method for checking for a non-dispersed sticky layer “cake” is by probing with a glass rod or similar device adapted to the size of the container. All the chemical and physical tests must be carried out on laboratory samples taken after the recommended homogenization procedure.

- Note 2 Method for wettable powders is applicable to aqueous suspension concentrates.
- Note 3 If the buyer requires both g/kg and g/l at 20°C then, in case of dispute, the analytical results shall be calculated as g/kg.
- Note 4 Chemical assay is the only fully reliable method to measure the mass of active ingredient still in suspension. However, simpler methods such as gravimetric determination or solvent extraction may be used on a routine basis provided that these methods have been shown to give equal results to those of the chemical assay method. In case of dispute, the chemical method shall be the “Referee method”.
- Note 5 Unless other CIPAC waters, temperatures, or times are specified.
- Note 6 The mass of the sample to be used in the test should be specified at the highest rate of use recommended by the supplier.
- Note 7 Samples of the product taken before and after the storage stability test should be analyzed together after the test to reduce the analytical error.

OXADIXYL WATER DISPERSIBLE GRANULES

FAO Specification 397/WG/S/F (1997)

1. DESCRIPTION

The material shall consist of a homogeneous mixture of technical oxadixyl, complying with the requirement of FAO specification 397/TC/S/F (1997), together with suitable carriers and any other necessary formulants. It shall be in the form of granules (Note 1) for application after disintegration and dispersion in water. The product shall be dry, free-flowing and free from visible extraneous matter and hard lumps.

2. ACTIVE INGREDIENT

2.1. Identity tests (397/WP/M/2, CIPAC G, p. 125. Note 2)

An identity test is required when the identity of the active ingredient is in doubt,. Then the isolated active ingredient shall comply with at least one additional test, e.g. HPLC or IR spectrum.

2.2. Oxadixyl (397/WP/M/3, CIPAC G, p. 125. Note 2)

The oxadixyl content shall be declared (g/kg) and, when determined, the content obtained shall not differ from that declared by more than the following amounts:

<u>Declared content</u>	<u>Permitted tolerance</u>
Above 25 up to 100 g/kg	± 10 % of the declared content
Above 100 up to 250 g/kg	± 6 % of the declared content
Above 250 up to 500 g/kg	± 5 % of the declared content

3. IMPURITIES

Not relevant.

4. PHYSICAL PROPERTIES

4.1. pH range (MT 75, CIPAC F, p.205)

pH range: 5.0 to 10.0.

4.2. Wetting of the material (MT 53.3. Note 3)

The product shall be completely wetted in 1 min.

4.3. Wet sieve test (MT 167, CIPAC F, p.416)

Maximum: 2 % retained on a 75 µm sieve.

4.4. Suspensibility (MT 168, CIPAC F, p.417)

A minimum of 60 % of the oxadixyl content found under 2.2. shall be in suspension after 30 minutes in CIPAC standard water D at $30 \pm 2^\circ\text{C}$, Notes 4 & 5).

Alternatively, if the buyer requires other CIPAC standard waters, temperature or time to be used, then this shall be specified when ordering.

4.5. Degree of dispersion (MT 174, CIPAC F, p.152)

Minimum: 70% of the insoluble content shall be dispersed after 1 min. of stirring in standard water D at $30 \pm 2^\circ\text{C}$.

Alternatively, if the buyer requires other CIPAC standard waters, temperature or time to be used, then this shall be specified when ordering.

4.6. Persistent foam (MT 47.2, CIPAC F, p.152. Note 6)

Maximum: 50 ml after 1 min.

4.7. Dustiness (MT 171, gravimetric method, CIPAC F, p.425)

Maximum: 30 mg collected dust (If optical method is used: max. 25 mg).

4.8. Flowability (MT 172, CIPAC F, p.430)

After storage under a pressure of 25 g/cm^2 at $54 \pm 2^\circ\text{C}$ for 14 days the granules shall drop through the test sieve of 5 mm mesh size (Note 7) spontaneously or after 5 liftings, leaving no residue on the sieve.

5. STORAGE STABILITY

5.1. Stability at 54°C (MT 46.1.1, CIPAC F, p.148. Note 8)

After storage at $54 \pm 2^\circ\text{C}$ for 14 days, the determined average active ingredient content must not be lower than 97 % relative to the determined average content found under 2.2. and the product shall continue to comply with 4.1., 4.2., 4.3., 4.4., 4.5. 4.7. and 4.8.

- Note 1 To describe a specific products, it is recommended to add information about the form (e.g. irregular shape, nearly spherical, cylindrical, ...) and to state the nominal size range.
- Note 2 If necessary, the sample of the granule must be further homogenized. Method for wettable powders is applicable to water dispersible granules, when 3 ml of water are added to the sample after the addition of the internal standard..
- Note 3 The method to be used shall be stated, either with or without swirling.
- Note 4 The product should be tested at the highest and lowest rates of use recommended by the supplier, provided this does not exceed the conditions given in method MT 168.
- Note 5 Chemical assay is the only fully reliable method to measure the mass of active ingredient still in suspension. However, simpler methods such as gravimetric determination or solvent extraction determination may be used on a routine basis provided, that these methods have been shown to give equal results to those of the chemical assay method. In case of dispute, the chemical method shall be the “Referee method”.
- Note 6 The mass of the sample to be used in the test should be specified at the highest rate of use recommended by the supplier.
- Note 7 If the granules do not drop through the sieve before the test, adapt the test sieve.
- Note 8 Samples of the product taken before and after the storage stability test should be analyzed together after the test to reduce the analytical error.

OXADIXYL WATER DISPERSIBLE POWDERS

FOR SLURRY SEED TREATMENT

FAO Specification 397/WS/S/F.(1997)

1. DESCRIPTION

The material shall consist of a homogeneous mixture of technical oxadixyl, complying with the requirements of FAO specification 397/TC/S/F (1997), together with carrier(s) and any other necessary formulants, including colouring matter (Note 1). It shall be in a form of a fine powder free from visible extraneous matter and hard lumps.

2. ACTIVE INGREDIENT

2.1. Identity tests (397/WP/M/2, CIPAC G, p.125. Note 2)

An identity test is required if the identity of the active ingredient is in doubt,. Then the isolated active ingredient shall comply with at least one additional test, e.g. HPLC or IR spectrum.

2.2. Oxadixyl (397/WP/M/3, CIPAC G, p.125. Note 2)

The content of oxadixyl shall be declared (g/kg) and, when determined, the content obtained shall not differ from that declared by more than the following amounts:

<u>Declared content</u>	<u>Permitted tolerance</u>
Above 25 up to 100 g/kg	± 10 % of the declared content
Above 100 up to 250 g/kg	± 6 % of the declared content
Above 250 up to 500 g/kg	± 5 % of the declared content

3. IMPURITIES

Not relevant.

4. PHYSICAL PROPERTIES

4.1. pH range (MT 75, CIPAC F, p.205)

pH range: 5.0 to 10.0.

4.2. Wet sieve test (MT 59.3, CIPAC F, p.416. Note 3)

Maximum: 2 % retained on a 75 µm test sieve.

4.3. Persistent foam (MT 47.2, CIPAC F, p.152. Note 3)

Maximum: 25 ml after 1 min.

4.4. Wetting of the material (MT 53.3.1, CIPAC F, p.164)

The material shall be completely wetted in 1 min. without swirling.

5. STORAGE STABILITY

5.1. Stability at 54 °C (MT 46.1.1, CIPAC F, p.148. Note 4)

After storage at 54 ± 2 °C for 14 days, the determined average active ingredient content must not be lower than 97 % relative to determined average content found under 2.2. and the product shall continue to comply with 4.1., 4.2. and 4.4.

Note 1 The product shall contain a dye that permanently colours the seed after treatment (red is recommended) and cannot be removed by washing with water. In some countries, there may be a legal requirement that a specific colour shall be used. The same colour should not be used for denaturing seeds to be used as livestock feeding stuffs.

Note 2 Method for wettable powders is applicable.

Note 3 Test to be carried out at the application concentration.

Note 4 Samples of the product taken before and after the storage stability test should be analyzed together after the test to reduce the analytical error.

OXADIXYL SUSPENSION CONCENTRATES

FOR SEED TREATMENT

FAO Specification 397/FS/S/F.(1997)

1. DESCRIPTION

The material shall consist of a suspension of fine particles of technical oxadixyl, complying with the requirements of FAO specification 397/TC/S/F (1997), in an aqueous phase together with any suitable formulants including colouring matter (Note 1) for seed treatment.

After gentle stirring or shaking on the commercial container, the material shall be homogeneous (Note 2) and suitable for dilution with water, if necessary.

2. ACTIVE INGREDIENT

2.1. Identity tests (397/WP/M/2, CIPAC G, p. 125. Note 3)

An identity test is required if the identity of the active ingredient is in doubt,. Then the isolated active ingredient shall comply with at least one additional test, e.g. HPLC or IR spectrum.

2.2. Oxadixyl (397/WP/M/3, CIPAC G, p. 125. Note 3)

The oxadixyl content shall be declared (g/kg or g/l at $20 \pm 0.5^{\circ}\text{C}$. Note 4) and, when determined, the content obtained shall not differ from that declared by more than the following amounts:

<u>Declared content</u>	<u>Permitted tolerance</u>
Above 25 up to 100 g/kg or g/l	$\pm 10\%$ of the declared content
Above 100 up to 250 g/kg or g/l	$\pm 6\%$ of the declared content
Above 250 up to 500 g/kg or g/l	$\pm 5\%$ of the declared content

3. IMPURITIES

Not relevant.

4. PHYSICAL PROPERTIES

4.1. Mass per millilitre at 20 °C (MT 3.3, CIPAC F, p.18)

If required, the range of the mass per millilitre (g/ml) at $20 \pm 0.5^\circ\text{C}$ shall be declared.

4.2. pH range (MT 75, CIPAC F, p.205)

pH range: 5.0 to 10.0.

4.3. Rinsability (MT 148.1, CIPAC H, to be published)

Maximum non-volatile residue: 0.01 %.

4.4. Wet sieve test (MT 59.3, CIPAC F, p.179. Note 5)

Maximum: 2 % retained on a 75 μm sieve.

4.5. Persistent foam (MT 47.2, CIPAC F, p.152)

Maximum: 25 ml after 1 min.

4.6. Flash point (MT 12, CIPAC F, 31)

If relevant, the flash point shall not be lower than the minimum declared flash point. The procedure used shall be stated (Note 6).

5. STORAGE STABILITY

5.1. Stability at 0 °C (MT 39.1, CIPAC F, p.128)

After storage at $0 \pm 1^\circ\text{C}$ for 7 days, the product shall continue to comply with 4.4.

5.2. Stability at 54 °C (MT 46.1.2, CIPAC F, p.149. Note 7)

After storage at $54 \pm 2^\circ\text{C}$ for 14 days, the determined average active ingredient content must not be lower than 97 % relative to the determined average content found under 2.2. and the product shall continue to comply with 4.2., 4.3. and 4.4.

Note 1 The product shall contain a dye that permanently colours the seed after treatment (red is recommended). In some countries, there may be a legal requirement that a specific colour shall be used. The same colour should not be used for denaturing seeds to be used as livestock feeding stuffs.

- Note 2 Before sampling to verify the product quality, inspect the commercial container carefully. On standing, suspension concentrates usually develop a concentration gradient from the top to the bottom of the container. This may even result in the appearance of a clear liquid on the top and/or of sediment at the bottom. Therefore, before sampling, the product must be homogenized according to the instructions given by the manufacturer or, in the absence of such instructions, by gentle shaking of the commercial container (for example by inverting the closed container several times). Large containers must be opened and stirred adequately. After this procedure the container should not contain a sticky layer of non-dispersed product at the bottom. A suitable and simple method for checking for a non-dispersed sticky layer “cake” is by probing with a glass rod or similar device adapted to the size of the container. All the chemical and physical tests must be carried out on laboratory samples taken after the recommended homogenization procedure.
- Note 3 Method for wettable powders is applicable.
- Note 4 Unless homogenization is carried out carefully, it is possible for the sample to become aerated. This can lead to errors in the determination of the density and in calculation of the active ingredient content (in g/l) if methods other than MT3.3 are used. If the buyer requires both g/kg and g/l at 20°C, then in case of dispute the analytical result shall be calculated as g/kg.
- Note 5 Test to be carried out at the application concentration.
- Note 6 Attention is drawn to the appropriate national and for international regulations on handling and transport of flammable materials.
- Note 7 Samples of the product taken before and after the storage stability test should be analyzed after the test at the same time to reduce the analytical error.