

Indicators Utilisation	Standard Module(s)	Usual source
<b>III.1. Outcome indicators (anthropometry)</b>		
Low birth weight babies (% of births)	body measurement	MICS/DHS/health facilities
prevalence of wasting (weight for height) (% of children under 5)	body measurement	MICS/DHS
prevalence of stunting (height for age) (% of children under 5)	body measurement	MICS/DHS
prevalence of underweight (weight for age) (% of children under 5)	body measurement	MICS/DHS
Mid-Upper Arm Circumference (mm)	body measurement	MICS/DHS
BMI-for-age (kg/m <sup>2</sup> )	body measurement	MICS/DHS
Chronic Energy Deficiency (CED) (or, Percentage of adults with BMI < 18.5) (%)	body measurement	MICS/DHS
<b>III.2. Outcome indicators (nutritional and health status)</b>		
Life expectancy at birth (years)		MICS/DHS/health surveys
Maternal mortality ratio (per 100,000 live births)		MICS/DHS/health surveys
Infant mortality rate (%)		MICS/DHS/health surveys
Under-five mortality rate (%)		MICS/DHS/health surveys
Exclusive breastfeeding (% of children under 6 months)		MICS/DHS
Vitamin A Deficiency	Blood test	MICS/DHS
Vitamin A supplementation coverage rate (% of children ages 6-59 months)		MICS/DHS
Prevalence of Anemia	Blood test	MICS/DHS
Consumption of iodized salt (% of households)	Salt test	MICS/DHS
Prevalence of Diarrhea		MICS/DHS
ARI treatment (% children)		MICS/DHS
Prevalence of HIV (% of population ages 15-49)	Blood test	MICS/DHS
Access to improved water and sanitation		MICS/DHS