



JOINT FAO/WHO EXPERT COMMITTEE ON FOOD ADDITIVES
Eight-sixth meeting
Geneva, 12–21 June 2018

SUMMARY AND CONCLUSIONS

Issued 3 July 2018

A meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA) was held in Geneva, Switzerland, from 12 to 21 June 2018. The purpose of the meeting was to evaluate certain food additives (including flavouring agents).

Dr A. Mattia, Center for Food Safety and Applied Nutrition, United States Food and Drug Administration, served as Chairperson, and Dr. Richard Cantrill, Canada, served as Vice-Chairperson.

Dr M. Lipp, Office for Food Safety, Food and Agriculture Organization of the United Nations, and Dr A. Tritscher, Department of Food Safety and Zoonoses, World Health Organization, served as Joint Secretaries.

The present meeting was the eighty-sixth in a series of similar meetings. The tasks before the Committee were (a) to undertake safety evaluations of certain food additives (including flavouring agents); and (b) to review and prepare specifications for certain food additives (including flavouring agents).

The Committee evaluated the safety of eight food additives, revised the specifications for 19 other food additives (including 16 modified starches), evaluated 69 flavouring agents according to the revised Procedure for the Safety Evaluation of Flavouring Agents and revised the specifications for three flavouring agents.

The report of the meeting will be published in the WHO Technical Report Series. Its presentation will be similar to that of previous reports – namely, general considerations, comments on specific substances and recommendations for future work. An annex will include detailed tables (similar to the tables in this report) summarizing the main conclusions of the Committee in terms of acceptable daily intakes and other toxicological, dietary exposure and safety recommendations. Information on the specifications for the identity and purity of certain food additives (including flavouring agents) examined by the Committee will also be included.

The participants in the meeting are listed in Annex 1. Items of a general nature that the Committee would like to disseminate quickly are included in Annex 2. Future work and recommendations are listed in Annex 3.

Toxicological and dietary exposure monographs on most of the substances that were considered will be published in WHO Food Additives Series No. 77. New and revised specifications for the identity and purity of the compounds will be published in FAO JECFA Monographs 22.

More information on the work of JECFA is available at:

<http://www.fao.org/food/food-safety-quality/scientific-advice/jecfa/en/>

and

http://www.who.int/foodsafety/areas_work/chemical-risks/jecfa/en/

The issuance of this document does not constitute formal publication. The document may, however, be freely reviewed, abstracted, reproduced or translated, in whole or in part, but not for sale or use in conjunction with commercial purposes.

Toxicological information and information on specifications

Food additives evaluated toxicologically and assessed for dietary exposure

| Food additive | Specifications | Acceptable daily intakes (ADIs) and other toxicological and dietary exposure conclusions |
|--------------------------------------|-------------------|---|
| Anionic methacrylate copolymer (AMC) | N, T ^a | The Committee was unable to complete the evaluation of AMC. While the copolymer itself is not of health concern, genotoxicity concerns remains for the residual monomer methacrylic acid. The specifications were made tentative pending the completion of the safety evaluation of AMC. |
| Basic methacrylate copolymer (BMC) | N | The Committee established an ADI “not specified” for basic methacrylate copolymer. The Committee concluded that the use of BMC that complies with the specifications established at the current meeting is not of safety concern when the food additive is used as a coating or glazing agent for solid food supplements and for foods for special medical purposes and micronutrient encapsulation for food fortification. The NOAELs for BMC ranged from 750-2000 mg/kg bw per day which were the highest doses tested. The Committee evaluated exposure to BMC for the copolymer and its monomers (n-butyl methacrylate, 2-(dimethylamino)ethyl methacrylate and methyl methacrylate). Estimated exposures to BMC range from 3.0 to 135 mg/kg bw per day. The total monomeric content of BMC is less than 0.3%. The Committee concluded that the toxicological data on the residual monomers do not give rise to concerns when taking into account the low dietary exposures. |
| Erythrosine | R ^b | The Committee concluded that the new data that have become available since the previous evaluation of erythrosine do not give reason to revise the ADI and confirmed the previous ADI of 0–0.1 mg/kg bw. The Committee noted that the dietary exposure estimate for erythrosine of 0.09 mg/kg bw per day (95 th percentile for children) was close to the upper bound of the ADI. Given that this estimate of exposure is for children and it is a high percentile for consumers only, such a level is unlikely to occur every day over a lifetime. Therefore, the Committee concluded that dietary exposures to erythrosine for all age groups do not present a health concern. |

| Food additive | Specifications | Acceptable daily intakes (ADIs) and other toxicological and dietary exposure conclusions |
|--------------------------------------|------------------|--|
| Indigotine | R ^b | <p>The Committee considered the new data that had become available since the previous evaluation as well as previously evaluated studies and concluded that there are no reasons to revise the ADI and confirmed the previous ADI of 0–5 mg/kg bw.</p> <p>The Committee noted that the conservative dietary exposure estimate of 0.8 mg/kg bw per day (95th percentile for children and toddlers) is less than the upper bound of the ADI of 0–5 mg/kg bw. The Committee concluded that dietary exposure to indigotine for all age groups does not present a health concern.</p> |
| Lutein | R ^{c,d} | <p>Free lutein, lutein esters and free zeaxanthin including <i>meso</i>-zeaxanthin are biochemically and toxicologically equivalent. At the present meeting the Committee concluded that there were sufficient toxicological data to complete a safety assessment of lutein and lutein esters from <i>Tagetes erecta</i>, synthetic zeaxanthin and <i>meso</i>-zeaxanthin. Free lutein, lutein esters and free zeaxanthin and <i>meso</i>-zeaxanthin are substances of low toxicity for which no adverse effects have been observed in a broad range of toxicological studies in laboratory animals and clinical studies in humans.</p> <p>Based on the absence of toxicity in a wide range of studies, the Committee established a group ADI "not specified" for lutein from <i>Tagetes erecta</i>, lutein esters from <i>Tagetes erecta</i> and zeaxanthin (synthetic).</p> <p>Meso-zeaxanthin was not included in this group ADI, as specifications are not currently available.</p> <p>The group ADI of 0-2 mg/kg bw for lutein from <i>Tagetes erecta</i> and zeaxanthin (synthetic) was withdrawn.</p> |
| Neutral methacrylate copolymer (NMC) | N, T | <p>The Committee established an ADI "not specified" for NMC. The ADI "not specified" was made temporary because the specifications are tentative.</p> <p>The Committee concluded that the use of NMC that complies with the specifications established at the current meeting is not of safety concern when the food additive is used as a coating or glazing agent for solid food supplements and for foods for special medical purposes. The NOAELs for NMC ranged from 454–2000 mg/kg bw per day, and these were the highest doses tested.</p> <p>The Committee evaluated exposure to NMC for the copolymer and its monomers (methyl methacrylate and ethyl acrylate). Estimated exposures to NMC range from 5.8 to 86 mg/kg bw per day. The total monomeric content of NMC is less than 0.01%. Toxicological data on the residual monomers do not give rise to concerns when taking into account the low dietary exposures.</p> |
| Sorbitol syrup | - | <p>Sorbitol syrup (INS 420(ii)) is currently included in the Codex General Standard for Food Additives (GSFA) although it has not been assigned an ADI or determined, on the basis of other criteria, to be safe. The Committee was therefore requested to consider the previous evaluations of sorbitol, hydrogenated glucose syrups and other relevant substances, and advise on the need for a</p> |

| Food additive | Specifications | Acceptable daily intakes (ADIs) and other toxicological and dietary exposure conclusions |
|-------------------|----------------|---|
| | | <p>separate evaluation of sorbitol syrup or if the ADI “not specified” for sorbitol is also applicable for sorbitol syrup.</p> <p>Based on the similarity of the chemical constituents of sorbitol syrup to the previously evaluated sorbitol, maltitol syrup and polyglycitol syrup, the Committee concluded that there is no need for a separate evaluation of sorbitol syrup and established an ADI “not specified” for sorbitol syrup.</p> |
| Spirulina extract | N, T | <p>The Committee established a temporary ADI “not specified” for spirulina extract. The ADI was based on the absence of toxicity in repeated-dose animal studies with spirulina extract and dried spirulina. The ADI “not specified” was made temporary due to the tentative nature of the specifications.</p> <p>Expressed as phycocyanins, estimated dietary exposure from the use of spirulina extract as a food colour based on the Budget method and exposure to spirulina extract and dried spirulina from other dietary sources, including food ingredients, dietary supplements, and coatings of food supplements was 190 mg/kg bw for adults (60 kg/person) and 650 mg/kg bw for a child (15 kg/person). The Committee concluded that this dietary exposure does not present a health concern.</p> |

- : no specifications prepared; N: new specifications; R: existing specifications revised; T: tentative specifications

^a The specifications were made tentative pending the completion of the safety evaluation of AMC.

^b At the current meeting, high-performance liquid chromatographic (HPLC) methods were added for determining subsidiary colouring matters and organic compounds other than colouring matters. The method of assay was changed to visible spectrophotometry, and spectrophotometric data were provided for the colour dissolved in water.

^c The specifications for lutein esters from *Tagetes erecta* and zeaxanthin (synthetic) were maintained.

^d At the current meeting, the identity test for melting range was deleted, the identity tests for carotenoids and spectrophotometry were updated, the test for propylene glycol was incorporated verbatim and the previous reference removed, and the method of assay was updated.

Food additives considered for specifications only

| Food additive | Specifications |
|--|--------------------|
| Cassia gum | R ^a |
| Citric and fatty acid esters of glycerol | R, T ^b |
| Glycerol ester of wood rosin | R ^c |
| Modified starches | R ^d , T |

R: existing specifications revised; T: tentative specifications

^a The Committee, at its current meeting, received analytical methods and included the most suitable validated method in the specifications monograph. However, this method uses chloroform for the extraction of anthraquinones. Extraction with n-hexane and diethyl ether resulted in poor recovery of anthraquinones. The Committee recommends that the JECFA Secretariat be notified if an alternative extraction solvent is identified. The specifications were revised and the tentative status was removed.

^b The Committee did not receive a replacement method for the obsolete packed column gas chromatographic method for the determination of total citric acid, in its specifications monograph. The Committee noted further that the method for total glycerol still uses chloroform. The Committee encouraged the submission of a method for total glycerol that eliminates the use of chloroform. Specifications were revised and made tentative pending the availability of data. Specifications will be withdrawn if suitable information is not provided by December 2019.

^c The Committee received information on the manufacture of GEWR from the rosin obtained from the stumps of two additional species namely *Pinus halepensis* and *Pinus brutia* as source materials. Recognizing the natural variability of the composition of wood rosin, the Committee removed the restriction to certain pine species within the specifications. Since the specifications monograph for GEWR does not contain an assay, the Committee

recommended that the JECFA Secretariat be notified upon the development and validation of an appropriate assay. The existing specifications were revised.

^d The Committee reviewed data on the method of manufacture, identity, and purity of all 16 modified starches. Based on the information received, and available information the Committee noted that:

- All processes are performed under similar manufacturing conditions and result in minor chemical modifications. Given the chemical and physical similarities of modified starches, the Committee at previous meetings considered the application of a read-across approach to be appropriate for the toxicological evaluation of these substances.
- All 16 modified starches had been assigned an ADI of “not specified”.
- All modified starches can be additionally bleached or fragmented; therefore revision in the specifications of bleached or fragmented starches would imply the revision of all 16 monographs;
- Microbiological specifications were not present in the existing specifications for all modified starches.
- Several specifications were common to all modified starches (such as for heavy metals impurities content and microbiological considerations). Revision of those common specifications would affect all 16 monographs;
- As a result of the wide range of products manufactured, the identification tests required to unambiguously chemically characterize each modified starch in individual specifications may be cumbersome, potentially unavailable, and unlikely to reflect market requirements.
- It may not be possible to publish identification tests based on market requirements without unduly revealing proprietary information.
- Based on the points noted above, individual specifications for several modified starches may remain tentative for an indefinite period or may need to be withdrawn.

The Committee therefore recommended that a new approach to the specifications monographs should be introduced to account for the chemical similarity between all modified starches, their functional diversity, the variety of chemicals used in their manufacture, and the corresponding diversity of impurities. The Committee recommended that all modified starches be included in a modular monograph titled ‘Modified Starches’ that contains common requirements [General specifications for modified starches] consisting of specifications that apply to all 16 modified starches (INS 1400, 1401, 1402, 1403, 1404, 1405, 1410, 1412, 1413, 1414, 1420, 1422, 1440, 1442, 1450, 1451), and annexes with specifications applicable to each individual modified starch based on the treatment(s) received. The Committee drafted a new modular specifications monograph titled “Modified starches” consisting of an explanatory introduction, “General specifications for modified starches,” and eight annexes. The new modular specifications monograph for modified starches is printed in FAO Monograph 22, and will replace the 16 existing individual specifications for modified starches (INS 1400, 1401, 1402, 1403, 1404, 1405, 1410, 1412, 1413, 1414, 1420, 1422, 1440, 1442, 1450, 1451).

The specification for lead included in the General specifications be decreased from 2 mg/kg to 0.2 mg/kg. The limit of lead for starch sodium octenylsuccinate for use in infant formula and formula for special medical purposes intended for infants was set to 0.1 mg/kg in the General specifications.

The methods for the determination of free adipic acid and adipate groups, residual vinyl acetate, free octenyl succinic acid and octenyl succinate esters were revised and a method for the determination of propylene chlorohydrins was added.

Flavouring agents evaluated by the revised Procedure for the Safety Evaluation of Flavouring Agents

A. Alicyclic primary alcohols, aldehydes, acids and related esters

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|--|------------|-----------------------|---|
| Structural class I | | | |
| Mixture of 1-Vinyl-3-cyclohexenecarbaldehyde and 4-Vinyl-1-cyclohexenecarbaldehyde | 2253 | N | No safety concern |
| <i>p</i> -Mentha-1,8-dien-7-ol | 974 | N | No safety concern |
| <i>p</i> -Mentha-1,8-dien-7-yl acetate | 975 | N | No safety concern |
| Formyl-6,6-dimethylbicyclo[3.1.1]hept-2-ene | 980 | N | No safety concern |
| Myrtenol | 981 | N | No safety concern |
| Myrtenyl acetate | 982 | M | No safety concern |

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|---|------|----------------|--|
| Structural class II | | | |
| (1-Methyl-2-(1,2,2-trimethylbicyclo[3.1.0]hex-3-ylmethyl)cyclopropyl)methanol | 2254 | N | No safety concern |
| Structural class III | | | |
| (±)-Bicyclo[2.2.1]hept-5-ene-2-carboxylic acid, ethyl ester | 2255 | N | No safety concern |
| Flavouring agent excluded at Step 1 of the Procedure | | | |
| <i>p</i> -Mentha-1,8-dien-7-al (Perillaldehyde) | 973 | M | Genotoxicity data for <i>p</i> -mentha-1,8-dien-7-al raise concerns for potential genotoxicity |

N: new specifications

M: existing specifications maintained;

B. Carvone and structurally related substances

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|---|-------|----------------|--|
| Structural class I | | | |
| Pinocarvyl isobutyrate | 2242 | N | No safety concern |
| Carvyl palmitate | 2243 | N | No safety concern |
| Structural class III | | | |
| 6-Hydroxycarvone | 2244 | N | No safety concern |
| Flavouring agents not evaluated according to the revised Procedure | | | |
| (+)-Carvone | 380.1 | M | The Committee did not re-evaluate (+)-carvone (No. 380.1) according to the revised Procedure given the lack of information on the oral exposure from all sources and the need to review the ADI. A review of the ADI is recommended based on the evaluation of all biochemical and toxicological data. Also, data are needed for an exposure assessment for oral exposure to (+)-carvone from all sources to complete the evaluation for (+)-carvone. |
| (-)-Carvone | 380.2 | M | The Committee did not re-evaluate (-)-carvone (No. 380.2) according to the revised Procedure given the lack of information on the oral exposure from all sources |

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|------------------|-----|----------------|--|
| | | | and the lack of toxicological data. |

M: existing specifications maintained; N: new specifications

C. Furan-substituted aliphatic hydrocarbons, alcohols, aldehydes, ketones, carboxylic acids and related esters, sulfides, disulfides and ethers

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|---|------|----------------|--|
| Structural class III | | | |
| 2-Pentylfuran | 1491 | M ^a | No safety concern |
| 2-Heptylfuran | 1492 | M ^a | No safety concern |
| 2-Decylfuran | 1493 | M ^a | No safety concern |
| 3-Methyl-2-(3-methylbut-2-enyl)-furan | 1494 | M ^a | No safety concern |
| 2,3-Dimethylbenzofuran | 1495 | M ^a | No safety concern |
| 2,4-Difurfurylfuran | 1496 | M ^a | No safety concern |
| 3-(2-Furyl)acrolein | 1497 | M ^a | No safety concern |
| 2-Methyl-3(2-furyl)acrolein | 1498 | M ^a | No safety concern |
| 3-(5-Methyl-2-furyl)prop-2-enal | 1499 | M ^a | No safety concern |
| 3-(5-Methyl-2-furyl)butanal | 1500 | M ^a | No safety concern |
| 2-Furfurylidene-butyraldehyde | 1501 | M ^a | No safety concern |
| 2-Phenyl-3-(2-furyl)prop-2-enal | 1502 | M ^a | No safety concern |
| 2-Furyl methyl ketone | 1503 | M ^a | No safety concern |
| 2-Acetyl-5-methylfuran | 1504 | M ^a | No safety concern |
| 2-Acetyl-3,5-dimethylfuran | 1505 | M ^a | No safety concern |
| 3-Acetyl-2,5-dimethylfuran | 1506 | M ^a | No safety concern |
| 2-Butyrylfuran | 1507 | M ^a | No safety concern |
| (2-Furyl)-2-propanone | 1508 | M ^a | No safety concern |
| 2-Pentanoylfuran | 1509 | M ^a | No safety concern |
| 1-(2-Furyl)butan-3-one | 1510 | M ^a | No safety concern |
| 4-(2-Furyl)-3-buten-2-one | 1511 | M ^a | No safety concern |
| Pentyl 2-furyl ketone | 1512 | M ^a | No safety concern |
| Ethyl 3-(2-furyl)propanoate | 1513 | M ^a | No safety concern |
| Isobutyl 3-(2-furan)propionate | 1514 | M ^a | No safety concern |
| Isoamyl 3-(2-furan)propionate | 1515 | M ^a | No safety concern |
| Isoamyl 3-(2-furan)butyrate | 1516 | M ^a | No safety concern |
| Phenethyl 2-furoate | 1517 | M ^a | No safety concern |
| Propyl 2-furanacrylate | 1518 | M ^a | No safety concern |
| 2,5-Dimethyl-3-oxo-(2H)-fur-4-yl butyrate | 1519 | M ^a | No safety concern |
| Furfuryl methyl ether | 1520 | M ^a | No safety concern |
| Ethyl furfuryl ether | 1521 | M ^a | No safety concern |

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|--|------|----------------|--|
| Difurfuryl ether | 1522 | M ^a | No safety concern |
| 2,5-Dimethyl-3-furanthiol acetate | 1523 | M ^a | No safety concern |
| Furfuryl 2-methyl-3-furyl disulfide | 1524 | M ^a | No safety concern |
| 3-[(2-Methyl-3-furyl)thio]-2-butanone | 1525 | M ^a | No safety concern |
| O-Ethyl S-(2-furylmethyl)thiocarbonate | 1526 | M ^a | No safety concern |
| (E)-Ethyl 3-(2-furyl)acrylate | 2103 | M ^a | No safety concern |
| di-2-Furylmethane | 2104 | M ^a | No safety concern |
| 2-Methylbenzofuran | 2105 | M ^a | No safety concern |

M: existing specifications maintained

^a The text indicating that the safety evaluation for these flavouring agents had not been completed was removed from the specifications and the specifications were maintained as full

D. Linear and branched-chain aliphatic, unsaturated, unconjugated alcohols, aldehydes, acids and related esters

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|---------------------------|------|----------------|--|
| Structural class I | | | |
| <i>trans</i> -6-Octenal | 2240 | N | No safety concern |
| 2,6-Dimethyl-5-heptenol | 2241 | N | No safety concern |

N: new specifications

E. Maltol and related substances

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|-----------------------------|------|----------------|--|
| Structural class II | | | |
| Maltol | 1480 | M | No safety concern ^a |
| Structural class III | | | |
| Ethyl maltol isobutyrate | 2252 | N | No safety concern |

M: existing specifications maintained

N: new specifications

^a The previously established ADI for maltol was withdrawn by the Committee.

F. Menthol and structurally related substances

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|-------------------------------|------|----------------|--|
| Structural class I | | | |
| Menthyl formate | 2246 | N | No safety concern |
| Menthyl propionate | 2247 | N | No safety concern |
| <i>l</i> -Menthyl butyrate | 2248 | N | No safety concern |
| <i>d</i> <i>l</i> -Isomenthol | 2249 | N | No safety concern |

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|--|------|----------------|--|
| Dimethyl glutarate | 2250 | N | No safety concern |
| Menthol | 427 | M | No safety concern ^a |
| Structural class III | | | |
| (±)-2-[(2- <i>p</i> -Menthoxy)ethoxy]ethanol | 2251 | N | No safety concern |

M: existing specifications maintained

N: new specifications

^a The ADI of menthol of 0–4 mg/kg bw established at the fifty-first meeting was maintained.

G. Miscellaneous nitrogen-containing substances

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|---|------|----------------|--|
| Structural class III | | | |
| 2-(((3-(2,3-Dimethoxyphenyl)-1 <i>H</i> -1,2,4-triazol-5-yl)thio)methyl)pyridine | 2235 | N | No safety concern |
| S)-1-(3-(((4-Amino-2,2-dioxido-1 <i>H</i> -benzo[<i>c</i>][1,2,6]thiadiazin-5-yl)oxy)methyl)piperidin-1-yl)-3-methylbutan-1-one | 2236 | N | No safety concern |
| 2-(4-Methylphenoxy)- <i>N</i> -(1 <i>H</i> -pyrazol-3-yl)- <i>N</i> -(thiophen-2-ylmethyl)acetamide | 2237 | N | No safety concern |

N: new specifications

H. Saturated aliphatic acyclic branched-chain primary alcohols, aldehydes, and acids

| Flavouring agent | No. | Specifications | Conclusion based on current estimated dietary exposure |
|---------------------------|------|----------------|--|
| Structural class I | | | |
| 8-Methyldecanal | 2238 | N | No safety concern |
| 8-Methylnonanal | 2239 | N | No safety concern |

N: new specifications

Flavouring agents considered for specifications only

| Flavouring agent | No. | Specifications | |
|------------------------|------|----------------|----------------|
| L-menthyl lactate | 433 | | R ^a |
| L-malic acid | 619 | | R ^b |
| Glutamyl-valyl-glycine | 2123 | | R ^c |

^a The CAS number was changed from 59259-38-0 to 61597-98-6 and the name to L-menthyl L-lactate.

^b The specification for specific rotation were removed

^c The melting point range was revised.

Annex 1**Eight-sixth meeting of the
Joint FAO/WHO Expert Committee on Food Additives**
Geneva, 12–21 June 2018**Members**

Dr S. Barlow, Brighton, East Sussex, England, United Kingdom
Dr J. Bend, Department of Pathology and Laboratory Medicine, Schulich Medicine & Dentistry, Western University, London, Ontario, Canada
Dr D. Benford, Cheddington, London, England, United Kingdom
Dr R. Cantrill, Halifax, Nova Scotia, Canada (Vice-Chairperson)
Dr E. Dessipri, General Chemical State Laboratory, Athens, Greece, currently European Directorate for the Quality of Medicines and Health Care, Strasbourg, France
Dr D. Folmer, Office of Food Additive Safety, Center for Food Safety and Applied Nutrition, United States Food and Drug Administration, College Park, Maryland, USA (Joint Rapporteur)
Ms T. Hambridge, Food Standards Australia New Zealand (FSANZ), Kingston, ACT, Australia
Dr A. Mattia, Office of Food Additive Safety, Center for Food Safety and Applied Nutrition, United States Food and Drug Administration, College Park, Maryland, USA (Chairperson)
Dr U. Mueller, Australian Pesticides and Veterinary Medicines Authority (APVMA), Symonston, Australian Capital Territory (ACT), Australia (Joint Rapporteur)
Dr O.E. Orisakwe, University of Port Harcourt, Choba, Port Harcourt, Rivers State, Nigeria
Dr J. Schlatter, Zurich, Switzerland
Dr J. Smith, Bio|Food|Tech, Charlottetown, Prince Edward Island, Canada
Dr J.R. Srinivasan, Division of Biotech and GRAS Notice Review, Office of Food Additive Safety, Center for Food Safety and Applied Nutrition, United States Food and Drug Administration, College Park, Maryland, USA
Dr M. Veerabhadra Rao, Hyderabad, India

Secretariat

Dr J.H. Andersen, National Food Institute, Technical University of Denmark, Lyngby, Denmark (FAO Expert)
Dr J.N. Barrows, Office of Cosmetics and Colors, Center for Food Safety and Applied Nutrition, United States Food and Drug Administration, College Park, Maryland, USA (FAO Expert)
Dr C. Béchaux, ANSES, Maisons-Alfort, France (WHO Temporary Adviser)
Dr M. DiNovi, Office of Food Additive Safety, Center for Food Safety and Applied Nutrition, United States Food and Drug Administration, College Park, Maryland, USA (WHO Temporary Adviser)
Dr B. Fallico, Food Science and Technology Unit, University of Catania, Catania, Italy (FAO Expert)
Mr Y. Fan, China National Center for Food Safety Risk Assessment, Beijing, China (CCFA Chairperson)
Dr M.J.F. Fernandez, Universidad Miguel Hernández, Alicante, Spain (FAO Expert)
Dr B. Fields, Food Standards Australia New Zealand, Barton, Australian Capital Territory (ACT), Australia (WHO Temporary Adviser)
Ms F. Hill, Food Standards Agency, London, United Kingdom (WHO Temporary Adviser)
Dr S.M.F. Jeurissen, Centre for Nutrition, Prevention and Health Services, National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands (WHO Temporary Adviser)
Dr X. Jia, Laboratory of Toxicology, China National Center for Food Safety Risk Assessment, Beijing, China (WHO Temporary Adviser)
Dr S. Kim, Department of Food Safety and Zoonoses, World Health Organization, Geneva, Switzerland (WHO Secretariat)
Dr K. Laurvick, United States Pharmacopeial Convention, Rockville, Maryland, USA (FAO Expert)
Dr M. Lipp, Agriculture and Consumer Protection Department, Food and Agriculture Organization of the United Nations, Rome, Italy (FAO Joint Secretary)

- Dr P. Mosesso, Department of Ecological and Biological Sciences, Università degli Studi della Tuscia, Viterbo, Italy (WHO Temporary Adviser)
- Ms J. Odrowaz, Toronto, Ontario, Canada (WHO Technical Editor)
- Mr K. Petersen, Department of Food Safety and Zoonoses, World Health Organization, Geneva, Switzerland (WHO Secretariat)
- Dr L. Rosenfeld, Division of Petition Review, Office of Food Additive Safety, Center for Food Safety and Applied Nutrition, United States Food and Drug Administration, College Park, Maryland, USA (WHO Temporary Adviser)
- Dr J. Rotstein, Pre-Market Toxicology Assessment Section, Chemical Health Hazard Assessment Division, Bureau Chemical Safety, Food Directorate, Health Products and Food Branch, Health Canada, Ottawa, Ontario, Canada (WHO Temporary Adviser)
- Dr N. Sugimoto, Division of Food Additives, National Institute of Health Sciences (NIHS), Kanagawa, Japan (FAO Expert)
- Dr S. Takasu, Division of Pathology, Biological Safety Research Center, National Institute of Health Sciences, Kawasaki-shi, Kanagawa, Japan (WHO Temporary Adviser)
- Dr A. Tritscher, Department of Food Safety and Zoonoses, World Health Organization, Geneva, Switzerland (WHO Joint Secretary)
- Dr T. Umemura, Faculty of Animal Science Technology, Yamazaki Gakuen University, Tokyo, Japan (WHO Temporary Adviser)
- Dr X. Yang, Food Safety and Health Research Center, Southern Medical University, Guangzhou, Guangdong Province, China (WHO Temporary Adviser)
- Ms L. Zhang, Joint FAO/WHO Food Standards Programme, Food and Agriculture Organization of the United Nations, Rome, Italy (Codex Secretariat)

Annex 2

General considerations

An edited version of this section will appear in the report of the eighty-sixth meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA). It is reproduced here so that the information can be disseminated quickly. This draft will be subject to editing.


.....

Corrigenda for specifications monographs

The following requests for corrections in JECFA Food Additives Specifications Monographs were received by the JECFA Secretariat. The Committee at the current meeting evaluated the information provided and made the following corrections. These corrections will be published in the electronic versions and in the online database of JECFA Food Additives Specifications Monographs. The information is provided here to make interested parties aware of these changes.

| Food additive | Original text | New text | Additional information |
|--|---|---|---|
| Calcium disodium ethylenediaminetetraacetate (INS 385) Monograph 1 (2006) | CAS No. 662-33-9 | CAS No. 62-33-9 | Transcription error |
| Chlorophyllins, copper complexes sodium and potassium salts (INS 141(ii)) Monograph 5 (2008) Test for "Free ionisable copper" | Accurately weigh about 1 g of the sample and dissolve in 20 ml of arachid oil.... | Accurately weigh about 1 g of the sample and mix in 20 ml of arachid oil.... | Correction |
| Curcumin (INS: 100(i)) Monograph 1 (2006) | The criteria for several residual solvents are listed under the heading "Residual solvents" (see Fig. 1). | Acetone: Not more than 30 mg/kg Hexane: Not more than 25 mg/kg Methanol: Not more than 50 mg/kg Ethanol: Not more than 50 mg/kg Isopropanol: Not more than 50 mg/kg | Improves readability It was unclear whether the criterion "Not more than 50 mg/kg" extended to methanol, ethanol, isopropanol and ethyl acetate. |

| Food additive | Original text | New text | Additional information |
|---|--|---|------------------------|
| | | Ethyl acetate: Not more than 50 mg/kg | |
| Ethyl acetoacetate ethyleneglycol ketal JECFA No: 1969 JECFA 73 (2010) | CAS No. 1648615 | CAS No. 6413-10-1 | Transcription error |
| Ethyl 2-methyl pentanoate JECFA No: 214 JECFA 55 (2000) | CAS No. 28959-02-6 | CAS No. 39255-32-8 | Wrong CAS number |
| cis-3-Hexen-1-ol JECFA No.: 315 JECFA 51 (1998) | 98.0% (sum of (<i>Z</i>) and (<i>E</i>) isomers, =<92.0% (<i>Z</i>)) | 98.0% (sum of (<i>Z</i>) and (<i>E</i>) isomers, =>92.0% (<i>Z</i>)) | Transcription error |
| Monosodium L-glutamate (INS: 621) Monograph 1 (2006) | CAS No. 142-47-2 | CAS No. 6106-04-3 | Wrong CAS number |
| Myrcene JECFA No.: 1327 JECFA 63 (2004) | Specific gravity: 0.789–1.793 | Specific gravity: 0.789–0.793 | Transcription error |
| Polyoxyethylene (20) sorbitan monostearat (Polysorbate 60) (INS 435) Monograph 16 (2014) | CAS No. 9005-07-6 | CAS No. 9005-67-8 | Wrong CAS number |
| Sodium aluminium silicate (INS 554) Monograph 20 (2017) | Within the assay, the limits for silicon dioxide, aluminium oxide and sodium oxide are expressed “on dried basis”. | Within the assay, the limits for silicon dioxide, aluminium oxide and sodium oxide are expressed “on ignited basis”. | Transcription error |
| Silicon dioxide, amorphous (INS 551) Monograph 20 (2017) | CAS No. 112696-00-8 (hydrated silica) | CAS No. 112926-00-8 (hydrated silica) | Transcription error |
| | Pyrogenic silica is produced in an essentially anhydrous state, whereas the wet process products are obtained as hydrates or contain surface absorbed water. | Pyrogenic silica is produced in an essentially anhydrous state, whereas the wet process products are obtained as hydrates or contain surface adsorbed water. | Transcription error |

| Food additive | Original text | New text | Additional information |
|---|---|--|---|
| Sodium thiosulfate (INS 539) Monograph 1 (2006) | CAS No. 7772-98-7 | CAS No. 10102-17-7 | CAS No. 7772-98-7 refers to the anhydrous form. The specifications in the monograph refer to the pentahydrate form. |
| Brown HT and its aluminium lake (FAO JECFA Monographs 19, 82nd meeting, 2016) | Text in the Table 1 “Values for synthetic colours for use in performing tests for colouring matters content by spectrophotometry” | See Table 1, below | |
| Fast Green FCF (FAO JECFA Monographs 19, 82nd meeting, 2016) | Chemical structure in Table 1 “Values for synthetic colours for use in performing tests for colouring matters content by spectrophotometry” |  | |

CAS: Chemical Abstracts Service; INS: International Numbering System for Food Additives; No.: number

Bolding and underlining for clarity only. This formatting will not be shown in the online database.

The criteria for several residual solvents are listed under the heading “Residual solvents” (see Fig. 1)

Figure1: Residual solvent criteria for curcumin as displayed in Monograph 1, 2006


| <u>Residual solvents</u> (Vol. 4) | | |
|-----------------------------------|--|------------------------|
| Acetone: | | Not more than 30 mg/kg |
| Hexane: | | Not more than 25 mg/kg |
| Methanol: | | } |
| Ethanol: | | |
| Isopropanol: | | Not more than 50 mg/kg |
| Ethyl acetate: | | |

Table 1

Replacement of the text for the spectrophotometric data for Brown HT and its aluminium lake originally published in “Table 1. Values for synthetic colours for use in performing tests for Colouring Matters Content by Spectrophotometry” (FAO JECFA Monographs 19, 82nd meeting, 2016)


| JECFA name | Sample weight | Structure | Spectral data | Visible absorption spectrum |
|------------|---------------|-----------|---------------|-----------------------------|
| | | | | |

Brown 245.6
HT mg



Water, pH 7

$\lambda_{max} = 464$
A = 0.9957
Spec abs = 403
a = 40.3



Water

$\lambda_{max} = 464$
A = 0.9804
Spec abs = 397
a = 39.7

0.04 N AmAc

$\lambda_{max} = 461$
A = 0.9206
Spec abs = 373
a = 37.3


Brown 53.3
HT mg
Aluminium
m Lake

Straight colour
(blue)

0.04 N AmAc
 $\lambda_{max} = 461$
A = 0.9206

Lake (red)

0.04 N AmAc
 $\lambda_{max} = 458$
A = 1.0451



Annex 3

Future work and recommendations

Specific food additives (other than flavouring agents)

Anionic Methacrylate Copolymer

The Committee noted that there were insufficient data to reach a conclusion on the genotoxic potential of methacrylic acid. Further studies to clarify the in vivo carcinogenic potential are required.

Citric and fatty acid esters of glycerol

The specifications of CITREM were made tentative, requiring a suitable validated method for the determination of total citric acid content, along with performance characteristics of the method and data on the total citric acid content in at least five batches of products currently available in commerce, determined using that method.

The Committee noted that the method for total glycerol still uses chloroform. The Committee encouraged the submission of a method for total glycerol that eliminates the use of chloroform. Specifications were revised and made tentative. Specifications will be withdrawn if suitable information is not provided by **December 2019**.

Neutral Methacrylate Copolymer

The Committee noted that there was no data submitted for a suitable method of assay. Tentative specifications for NMC were prepared and made tentative requiring a suitable validated method of assay.

Spirulina extract

The Committee received limited analytical data on *spirulina* extract. In order to remove the tentative designation from the specifications, the following information on the products of commerce is requested by **December 2019**:

- Full compositional characterization of commercial products in both liquid and powder forms.
- Full compositional characterization of the aqueous extract before formulation/standardization.
- Validated analytical methods for identification of the substance with a suitable specificity (including validation data and representative batch data).
- Validated analytical methods for the determination of the purity of the substance with a suitable specificity (including validation data and representative batch data).

Modified starches

The Committee requested additional data and a suitable method for the determination of propylene chlorohydrins in Hydroxypropyl starch (INS 1440) and Hydroxypropyl distarch phosphate (INS 1442) in order to consider lowering this limit.

The Committee requests suitable microbiological acceptance criteria and supporting data for all modified starches.

Table 1. The annexes and the modified starches to which they apply along with required information:

| ANNEX | Modification | Starches to which it applies | Information required |
|-------|---|---|--|
| 1 | Minor fragmentation | INS 1400: Dextrin roasted starch; INS 1401: Acid treated starch; INS 1402: Alkaline treated starch; INS 1405: Enzyme-treated starch All modified starches that are additionally fragmented. | A suitable method for dispersion and a method for reducing sugars and data on at least 5 representative batches using the method(s) from each of the fragmentation processes. |
| 2 | Bleaching | INS 1403: Bleached starch All modified starches if additionally bleached. | Suitable method(s) for the determination of residual reagents and data on at least 5 representative batches using the method(s). |
| 3 | Esterification and/or crosslinking with phosphorus containing compounds | INS 1410: Monostarch phosphate; INS 1412: Distarch phosphate; INS 1413: Phosphated distarch phosphate; INS 1414: Acetylated distarch phosphate; INS 1442: Hydroxypropyl distarch phosphate | A suitable method for identification of crosslinking and data on at least 5 representative batches of crosslinked and non-crosslinked starches. |
| 4 | Acetylation | INS 1420: Starch acetate; INS 1414: Acetylated distarch phosphate; INS 1422: Acetylated distarch adipate; INS 1451: Acetylated oxidized starch | Currently no additional information required. |
| 5 | Oxidation | INS 1404: Oxidized starch; INS 1451: Acetylated oxidized starch | A suitable method for determination of residual hypochlorite and data on at least 5 representative batches using the method. |
| 6 | Esterification with octenyl succinic anhydride | INS 1450: Starch sodium octenyl succinate | Currently no additional information required. |
| 7 | Etherification with propylene epoxide | INS 1440: Hydroxypropyl starch; INS 1442: Hydroxypropyl distarch phosphate | A suitable method for the determination of propylene chlorohydrin with detection limit lower than 0.1 mg/kg and data on at least 5 representative batches of Hydroxypropyl starch using the method |
| 8 | Crosslinking with adipic anhydride | INS 1422: Acetylated distarch adipate | A suitable method for identification of crosslinking and data on at least 5 representative batches of crosslinked and non-crosslinked starches. Levels of free adipic acid in at least 5 representative batches |

Flavouring agents

Carvone and structurally related substances

For (+)-carvone (No. 380.1), the Committee concluded that a review of the ADI is recommended based on the evaluation of all biochemical and toxicological data. Also, data are needed for an exposure assessment for the oral exposure to (+)-carvone from all sources.

The ADI for (+)-carvone is maintained pending review of the ADI at a future meeting. The Committee recommends that the re-evaluation is completed within three years.

For (-)-carvone (No. 380.2), the Committee concluded that toxicological data on (-)-carvone are necessary. Also, data are needed for an exposure assessment for the oral exposure to (-)-carvone from all sources

Maltol and related substances

The Committee could not verify the NOEL of 100 mg/kg bw in rats that was used to derive the ADI of 0–1 mg/kg bw for maltol (No. 1480) during its twenty-fifth meeting because of uncertainties in the administered dose levels and the effects observed in several studies described in the monograph of that meeting.

The Committee withdrew the ADI for maltol. The Committee concluded that access to either the original studies or submission of new data would be needed to reaffirm or amend the ADI.

The ADI for ethyl maltol was maintained.