

**STREET FOODS IN NAIROBI, KENYA: THEIR ROLE AS A SOURCE  
OF MICRONUTRIENTS IN LOW INCOME GROUPS**

**Oyunga-Ogubi MA<sup>1\*</sup>, Waudu NJ<sup>2</sup>, Afullo A<sup>3</sup> and SO Oiye<sup>4</sup>**



**Mary Oyunga- Ogubi**

\*Corresponding author Email: [ogubim2003@yahoo.com](mailto:ogubim2003@yahoo.com)

<sup>1</sup>KARI-National Potato Research Center, Tigoni, P.O Box 338, 00217, Limuru, Kenya

<sup>2</sup>Department of Foods, Nutrition and Dietetics, Kenyatta University, P.O Box 43844-00100 Nairobi, Kenya

<sup>3</sup>Public Health Department, Kenyatta University, P.O Box 43844-00100, Nairobi, Kenya

<sup>4</sup>Food and Nutrition Consultant, P.O Box 17094, 00510, Makongeni, Nairobi, Kenya

## ABSTRACT

Vitamin A (VA), iron and zinc deficiencies are of public health concern in Kenya. Although more than 40% of Nairobi's lower-income groups consume street foods (SFs), inadequate information exists on the contribution or potential contribution to micronutrient intake. The study objective was to determine household intake of VA, iron and zinc, and the proportion of the intakes from SFs. Cross-sectional surveys and a non-repetitive 24hr-dietary recall were employed. The study sites were Kangemi and Dandora in Nairobi, Kenya (low and middle-low income groups) respectively. Student's t-test was used to compare the means of independent data while  $\chi^2$  test was used to compare categorical data. The level of significance was set at  $p=0.05$ . Results indicated that there was no difference in the consumption pattern of SFs between the two sites. Street foods were less important VA source in Kangemi (4.4% of total intake and 2.64% of the RDA) than in Dandora (26% of total intake and 9.46% of the RDA) and there was a significant difference. The proportion of those who obtained at least 50% VA from SFs was significantly higher in Dandora than Kangemi. Differences in VA intake from SFs in the two sites were attributed to quantities consumed because there was no difference in the pattern and in the levels of VA in SFs from the two sites. Iron intake contributed by SFs significantly differed with Dandora being higher (30%) than Kangemi (21%). Although not significantly different, those who obtained at least 50% iron from SFs was higher in Dandora than Kangemi. Iron intake from SFs in both sites (15mg/AE in Kangemi and 25mg/AE in Dandora) was sufficient to meet the RDA for iron for adults (5-28mg/AE). From the quantities consumed, the difference in iron intakes was partly attributed to the consumption pattern since the consumption pattern of some iron-rich foods was statistically different between the two sites. Zinc intake contributed by SFs foods was higher in Dandora (25.2%) than Kangemi (16.7%). Those who obtained at least 50% on their zinc intake from the SFs was also higher in Dandora (12%) than Kangemi (7%) although neither of these differences was significant.

**Key words:** urban-poor, zinc, iron, vitamin A

## INTRODUCTION

Street foods (SFs) have become an indispensable component of the socio-economic environment of many cities in developing countries. Their nutritional significance to urban populations is considerable [1]. In Nairobi, the vending of street foods increased substantially during the 1980s and 1990s, particularly in the low income and in the industrial areas [2]. More recently, due to the increase in urban populations, the number of street food vendors/hawkers has risen greatly in response to the need to feed larger populations. This has been heightened by the daily movement from residential areas to working places that has created need among many working people to eat outside the home more often [3]. Although more than 40% of households in Nairobi's lower-income areas consume SFs, there is paucity of knowledge on the contribution or potential contribution of these foods to micronutrient intake [4]. A study by Mwaniki *et al.* [5] showed that sixty percent of pre-school children in Kenya are estimated to have iron deficiency anemia while seventy percent are estimated to be vitamin A deficiency (VAD), one of the highest rates in sub-Saharan Africa. Further, the study found a mean serum s-zinc concentration (non-fasting <65µg/dl) of 65.07 µg/dL with a prevalence of 50.8% among children.

Micronutrients function as co-factors that enable the body to produce enzymes, hormones and other substances essential for proper growth and development. Although only small amounts are required, the consequences of their absence are severe [6]. Diets poor in micronutrients are associated with increased risk of infection and are known to cause blindness, premature death, reduced productivity, and impaired mental development, particularly among women and children in developing countries [7]. Known as "hidden hunger," micronutrient malnutrition can exist even when poor people have enough to eat, but lack the resources to afford a complete diet of fruits, vegetables, meat, and other foods rich in vitamins and minerals [8]. Given the devastating and increasing malnutrition rates among Kenyan urban low-income populations [9, 10], understanding the micronutrient accessibility through the major food systems of the poor is important [11].

In Kenya, fortification of food is not yet an important source of most micronutrients especially for the poor [12]. The micronutrient composition of the SFs sold and consumed is not known although it has been noted by Mwangi *et al.* that these sources of are inadequate [3]. Consequently, the potential role of SFs in terms of micronutrient intake and scope for increasing micronutrient contribution needs to be explored and incorporated into Kenyan policy programs. These are policies that aim at eradicating poverty and inequality and improving physical and economic access by all, at all times, to sufficient, nutritionally adequate and safe food and its effective utilization. Because SFs vary so much in type from country to country, for specific country recommendations, it is impossible to draw any general conclusions from the few analyses that have been done in some countries [13]. Therefore, this study was undertaken and the results are expected to inform the various stakeholders such as local authorities, Ministries of Health (Public Health officials), Agriculture (Bio-fortification programs), relevant local and international NGOs such as FAO and financial institutions on the way forward for street foods in Kenya.

## MATERIALS AND METHODS

### Study site selection

Street foods are found in other large and small cities in Kenya, but Nairobi was selected because, being the capital city, it houses people from all ethnic groups and races in Kenya. In addition, differences in history, intensity and types of food vending in Nairobi compared to other large (metropolitan) cities such as Mombasa, Nakuru and Kisumu were not expected to be significant. The study compared two different socio-economic sites, the low and middle-low income levels in Nairobi [14]. Non-probability sampling was used to cluster Nairobi into two areas; West and East based on the opposite regions, and taking the Central Business District as the reference point. Thus, to the West of Nairobi, areas were categorized into 4 socio-economic levels outlined by the Government of Kenya [10] as; high, middle, middle-low and low income areas. Purposively, a list was made of all settlement sites belonging to the middle-low and low-income areas (where SFs are commonly sold and consumed). One area was randomly selected - Kangemi (low income area). To the East of Nairobi a similar procedure was followed except that low-income areas were avoided, a site that falls into this category had already been selected. Dandora a middle-low income site was randomly selected. Dandora is about 30km from Kangemi.

### Household selection and sample determination

The study areas were stratified based on villages (Kangemi) and phases (Dandora). From each stratum, villages and phases were selected by the use of random numbers. The populations in Kangemi and Dandora are 19,298 and 36,691, respectively [14]. This formed the sampling frame from where the sample size was drawn. In obtaining a representative sample size, the Fisher *et al.* equation recommended by FAO, and Mugenda and Mugenda [15-17], was used to calculate the number of caregivers to be interviewed per area. From the selected villages/phases, households were selected by the use of random numbers (chits). A total of 600 households were selected with 300 households per area. If the selected household had no respondent, the survey proceeded to the next household until the sample size was obtained. For household dietary surveys, the respondent was usually the caregiver (95%) or in few cases (5%) were adults or older children (15years old).

To estimate the quantities of ingredients used per pot, an interactive 24-hour dietary recall developed by Gibson *et al.* [18] was used to obtain the sample size. The calculations for sample size were repeated for each of the 3 micronutrients where the largest sample size, considered as the worst case was applied. All values used for calculation of sample size are as reported by Mwangi *et al.* [3]. The respondent was usually the female involved in food preparation. Sixty households per site were surveyed. It was assumed that the ingredients used per pot were evenly distributed among the servings. Nutrients provided per serving from home prepared foods (HPFs) (VA and iron) were calculated using the Kenya Food Composition Tables and compared to the Recommended Daily Allowances (RDA) of an average Kenyan adult male referred to as Adult Equivalent (AE) [19]. Food and Agriculture Organization (FAO) data were used to calculate zinc values since these were absent in the Kenyan database [20]. The results of laboratory analysis on nutrient content of SFs in this

study were used to calculate the nutrient intake from SFs. The pro-vitamin A carotenoids were converted to (Retinol Equivalent) RE using the 6:1 ratio [19] and not the conversion ratio of 12:1 currently proposed by the Institute of Medicine [21].

### Vendor selection

From initial visits, it was found that there were approximately 50 vendors/processors of SFs within each study site. They were all considered for the vendor/processor survey as long as they sold SFs as defined in this study. Street foods in this study have been defined as 'ready-to-eat' foods and beverages, processed or fresh, which are sold on the streets as opposed to stores and licensed establishments, and which are sold at stationary locations or by mobile vendors and are not only restricted to snack like items [4]. Thus, 38 and 53 vendors were interviewed in Kangemi and Dandora, respectively.

### Data Analysis

Data were analyzed using SPSS version 12 for windows. Student's t-test was used to compare the means of independent data from the two study sites. Proportions were compared using  $\chi^2$  as well as  $\phi$  and Cramer's V statistics. The level of significance was set at  $p=0.05$ .

### RESULTS


Most household heads (48%) in both sites were permanently employed with a very small proportion (3%) being unemployed. Among the occupation status classification groups, only the proportion of temporarily employed was found to significantly differ between the two sites; 26.5% in Kangemi and 18.1% in Dandora. The average household income was higher in Dandora at Ksh.11,604.4  $\pm$  8653.1/month but was not significantly different from Kangemi with Ksh.10,518.7  $\pm$  9444.7/month (Table.1). There was high deviation from the mean income, meaning that the incomes of the two areas were very much spread – that is, the differences between the lowest and highest income in the respective study sites was huge. It is important to note that the two sites had marked differences in socio-economic status and this was more evident in the type of housing (building material used) and access to electricity than the occupation and household income of the dwellers. Dandora in this regard had better housing and other facilities than Kangemi.

The distribution and consumption pattern of SFs were similar in both sites. Most people (33.3%) on average consumed SFs 3-5 days a week and about one-fifth consumed SFs on a daily basis. In both areas about 17.4% did not consume SFs throughout the year (Table 2). Half of Kangemi dwellers (49.8%) gave lack of sufficient time to cook as the reason for consuming SFs and this was different from Dandora where only 2.7% gave such a reason. Further, more people in Kangemi (24.7%) than Dandora (17.4%) gave lack of money as a reason for not relying on SFs.

There were more female vendors in Dandora (51.9%) than in Kangemi (28.9%). Most vendors (50%) had only operated for 1 year or less at the time of the survey (Table. 3). A majority of the vendors in both sites obtained cooking ingredients from retailers (52%) and wholesalers (34%). Charcoal was the main cooking fuel and was more

important in Kangemi (81.6%) than in Dandora (40.7%). Wood was also an important source of energy and was used more by vendors in Kangemi than in Dandora. Electric power was the least used source of cooking fuel in both sites. Most vendors (57.9%) reported that consumers were of all ages and gender but a considerable proportion (23.6%) from both sites said that foods were mostly sold to adult males.

Street food vendors in both sites sell a wide range of foods but mostly those that are protein- and carbohydrate- rich. When these foods were classified in the major food groups on average a vendor in Dandora prepared and sold at least one SF food rich in protein (animal and vegetable). This was higher when compared to Kangemi who supplied less than one SF (0.7) rich in protein. All vendors in the two sites prepared about two SFs rich in carbohydrates and were comparable (*chapatti, ugali and mandazi*) (Table 4). The proportion of vendors selling carbohydrate foods was also the highest among the different food classes- 79% in Kangemi and 76.9% in Dandora. Street foods rich in micronutrients were least sold in the two sites; only 0.3 vendors sold these foods in Kangemi while 0.5 sold the same in Dandora (these were orange fleshed sweetpotato (OFSP), fruits, and vegetables). The proportion of vendors selling SFs rich in mixed nutrients (*githeri, mokimo and Kienyeji (irio)*) was not as high as those selling SFs rich in protein and carbohydrates alone, which was 39% and 48% in Kangemi and Dandora, respectively. There were more households closer to the SFs vending points (<10 metres) in Dandora (43%) than in Kangemi (27%) (Fig.1).



**Figure 1: Distance to nearest street food selling point in Kangemi and Dandora**

In both sites 50% of households were able to access SFs which were between 10-50 meters away. About 6% in both sites had SFs located more than 100 meters away which made it difficult for these households to access these foods. Eighty percent of respondents said they prefer easy access to the SFs.

### **Household intake of vitamin A, iron and zinc from street foods**

Vitamin A intake from HPFs was higher in Kangemi (428 $\mu$ g RE/AE) than in Dandora (208 $\mu$ g RE/AE), while VA intake from SFs was higher in Dandora (71 $\mu$ g RE/AE) than Kangemi (20 $\mu$ g RE/AE). In this regard total household VA intake was higher in Kangemi (447 $\mu$ g RE/AE) than in Dandora (279 $\mu$ g RE/AE). This translated to about 60 and 40 percent of the RDA in Kangemi and Dandora, respectively (Table 5). Street foods thus were a less important VA source in Kangemi (4.4% of total intake and 2.64 of the RDA) than in Dandora (25.4% of total intake and 9.46% of the RDA) and there was significant difference between the two sites with regard to the contribution to the daily household intake. The proportion of those who obtained at least 50% VA from SFs was also significantly higher in Dandora (23%) than in Kangemi (8.6%). On average, SFs contributed about 12% of VA consumed from the two study sites and about 16% sourced at least half of the VA from SFs. Although not consumed as much as the cereals and pulses, the major contributors to VA intake were orange-fleshed sweetpotato (OFSP) (130 $\mu$ g RE/100gdw), African black nightshades (34.7 $\mu$ g RE/100gdw, beef sausages (33.3 $\mu$ g RE/100gdw), *samosas* (vegetable) (26.2 $\mu$ g RE/100gdw), green grams (27.56 $\mu$ g RE/100gdw), kales (25.4 $\mu$ g RE/100gdw), animal offal (14.5 $\mu$ g RE/100gdw), and beef (13.3 $\mu$ g RE/100gdw).

Iron intake from HPFs in the two study sites was comparable, averaging about 57mg/day (Table 6). Intake from SFs was, however, higher in Dandora (26mg/AE) than Kangemi (15mg/AE). Total iron intake was also higher in Dandora (85.24 mg/AE) than in Kangemi (68.73 mg/AE). The proportion of iron intake contributed by SFs differed between the two sites, with the contribution in Dandora (30%) being significantly higher than Kangemi (21%). The proportion of those who obtained at least 50% of iron intake from the SFs was not significantly higher in Dandora (11.7%) than Kangemi (5%). Street foods play an important role as source of iron providing about a quarter of the daily intake but was more important in Dandora than in Kangemi. Iron from SFs alone was adequate to meet 85.94% of RDA in Kangemi and 150.71% of the RDA in Dandora. The intake was way above the RDA of 5-9mg for an adult male and 14-28mg for adult female [19]. Fried cabbages had the highest level of iron 17.9mg/100g dw with 50% bioavailability. Other sources high in iron bioavailability (72-97.5%) were beef sausages (2.1 mg/100g dw), animal offal (10.2 mg/100g dw), stewed beef (9.3 mg/100g dw), potato chips (7.8 mg/100g dw), *mandazi* (doughnut) (5.7 mg/100g dw). Those low in iron bioavailability (0-48%) were African black nightshades (11.1 mg/100g dw), Kales (11.3 mg/100g dw), fried beans (7.1 mg/100g dw), *githeri* (boiled/fried maize and beans) (6.1 mg/100g dw).

Zinc intake from HPFs, SFs and consequently total intake was higher in Dandora than in Kangemi. In Dandora zinc intake from SFs were 2.5mg/AE while in Kangemi it was only 1.2mg/AE. The RDA for zinc was assumed to be 10mg/AE per day as

reported by [22]. In this regard, zinc intake from SFs accounted for only 11.9% of RDA in Kangemi and 24.6% in Dandora (Table .7). The proportion of zinc intake contributed by SFs although not significant was higher in Dandora (25.2%) than in Kangemi (16.2%). Those who obtained at least 50% of zinc intake from the SFs were higher in Dandora (12%) than in Kangemi (7%) but this difference was not significant. Fried cabbages had the highest level of zinc (19.34mg/100g dw) with 63.5% bioavailability. Other sources high in zinc bioavailability (55-90%) were beef sausages (11.9mg/100g dw), animal offal (9.9mg/100g dw), stewed beef (7.8 mg/100g dw), potato chips (3.7mg/100g dw), *mandazi* (4.2mg/100g dw), *githeri* (4.4mg/100g dw), fried beans (5.4mg/100g dw) and kale (5.7mg'100gdw). Those low in zinc bioavailability (0-46%) were, fried African black nightshades (7.4 mg/100g dw), *samosa* (5.6 mg/100g dw), stewed green grams (5.4 mg/100g dw), OFSP (2.7mg/100g dw) and tea (0mg/100g dw).

## DISCUSSION

The study has shown that SFs are equally important in both sites with the only difference occurring in the variety and quantities consumed. This contradicts earlier findings by van't Riet which found that SF consumption in the low-income area was higher than in the low-middle-income of Dandora [4]. Income levels and general socio-economic status are major determinants of the degree of an urban population's access to food and of the nature of their choice of foods [23]. In urban areas distance from markets and low income levels force people to buy small amounts of food, often on a daily basis, which cumulatively tend to be more expensive than purchasing larger amounts at once. Furthermore, these people cannot afford the high cost of cooking fuel and cooking space which makes SFs more convenient [4]. This information is important for targeting food 'vehicles' for micronutrient improvement for different socio-economic groups. Targeting SFs in areas where low income groups live is likely to improve their nutritional status. The study found more female vendors in Dandora than in Kangemi. Female vendors have been found to be better at preparing quality foods especially where the consumers are willing to pay. Mwangi *et al.* [2] in Nairobi found that carbohydrate-rich foods were associated with male vendors while mixed nutrient (*irio*) foods and foods rich in micronutrients were associated with female vendors. This explains why Dandora had more food varieties and, therefore, more nutrient intake as compared to Kangemi where most vendors were male. Retail and wholesale markets are the main sources of cooking ingredients, which means that vendors have no control over nutrient quality of what they purchase. A report by FAO indicates that one of the difficulties in the control of SFs quality is that vendors often operate according to informal rules with their suppliers and clients [1]. Most of the vendors had been in operation for one year or less. This is because they operate in a permanent state of uncertainty, with little or no recourse to justice [4]. This is linked to the lack of official recognition and the absence of any form of policy governing the SFs sector [24]. Although a majority of vendors had attained both primary (39%) and secondary (51%) education, this did not determine the quality of foods made. This confirms findings by FAO that SFs vending does not require specific training other than the domestic experience of preparing food [1]. Given that increasing employment is one of the major strategies of the Kenyan government to boost economic develop-

ment, the contribution of the informal sector has been found to be significant, and any effort to eliminate SFs vendors may be retrogressive [4].

Vendors particularly those in Dandora chose their selling points carefully, concentrating in a variety of strategic spots, which contrary to expectation, were not in exclusively commercial areas. There were more households closer to the SFs vending points (<10 meters) in Dandora compared to Kangemi. This implies more accessibility of SFs in Dandora. From observations Kangemi appeared more crowded and vendors had to seek out- spots that were far from households unlike Dandora where the community was more organized in terms of buildings and access to basic services, thus availing more space for vendors to get close to households. Furthermore Dandora had a larger population (36,691) than Kangemi (19,298) [10]. This supports the suggestion by Tinker that there is a direct association between the number of SF enterprises and increases in city sizes [25]. Although there are a variety of food groups in both sites, the highest food group produced is carbohydrates. This is consistent with findings by Mwangi and van't Riet [2,4], where they found that although the SF sector in Nairobi offered products from all food groups, most vendors are one-food group sellers with cereals as the predominant group. There is a higher consumption of protein, especially meat in Dandora, which favor consumers in Dandora since the consumption of protein especially meat improves the absorption of most micronutrients in the meal [26].

Though it was observed that there was no difference in mean household income of the two study sites, proportion of income spent on food may have been lower in Kangemi than in Dandora thus prompting more of the households in the former to rely on the much cheaper SFs. Generally, the higher the income the lower the proportion of income spent on food, and *vice versa* [27]. Additionally, with less variety of SFs in Kangemi, the consumers here appeared to be purchasing poor quality foods. Overall, SFs were consumed mainly because there was no sufficient time to cook and because they were generally cheaper as compared to home prepared foods as reported by the consumers. These findings are consistent with those of FAO and van't Riet [1, 4].

None of the total household VA intakes of the two areas was close to the recommended adult RDA of 750 $\mu$ g RE as reported by Sehmi [19]. The amount of VA from SFs did not meet the RDA in both sites and this confirms the observation by Mwangi *et al.* [3] that SFs in low income groups lack in variety for quality diet [3]. Fruits which are the major plant sources of VA are considered expensive, contain a lot of water, are not stomach filling and hence not in the interest of the urban poor [2]. Since the pro-vitamin A carotenoids were converted to Retinol Equivalent (RE) using the 6:1 ratio [19] and not the conversion ratio of 12:1 currently proposed by IOM [21], a larger amount of provitamin A carotenoids, may be needed to meet VA requirement in the two sites. This also means that the conversion of VA from provitamin A carotenoids using Kenyan data base has been overestimated. Intake of iron in the two areas was far above the RDA of 5-9 mg for an adult male and 14-28 mg for adult female [19]. These findings are consistent with those of Mwangi *et al.* that SFs in Nairobi are able to provide adequate dietary iron and even far in excess of the RDA, irrespective of the type of the main meal [2]. Although the amount of iron taken in both study sites is adequate, the problem of iron deficiency anemia still persists.

This supports the suggestion that nutritional anemia needs to be effectively prevented and controlled by a combination of both vitamins and minerals in the diet [28]. Further, the common practice (>80% of consumers) of consuming tea alongside most meals has the effect of reducing the absorption of non-heme iron from plant sources. Fruits such as oranges and lemons were hardly consumed during meals. Vitamin C from these fruits enhances non-heme iron absorption in the presence of all known inhibitors of iron absorption including phytates (cereals and legumes), and polyphenols (tea, coffee, cocoa, certain vegetables and grains) [26]. The differences in quantities of iron intake could be partly attributed to the consumption pattern since the consumption of some iron-rich foods was different in the two sites. These foods are meat, *githeri*, French fries, which were found to be high in iron. The high iron level in French fries needs further investigation to ascertain the source of iron. Zinc intake in the two sites performed very poorly. These findings are in agreement with those of Okoth that the most commonly consumed staples in Kenya can hardly meet the RDA of adults for zinc [29]. None of the vendors fermented, soaked, or germinated the cooking ingredients used in SF preparation. These practices are known to improve the bioavailability of minerals including zinc [30].

## CONCLUSION

Although vitamin A, iron and zinc intake from SFs were comparatively lower among the low-income groups as compared to middle low-income groups, these differences were not significant. In terms of consumption frequencies, there were no significant differences in the consumption patterns between these two areas. Of the 3 micronutrients only iron intake from SFs met the RDA in both sites. To combat these deficiencies in urban areas, the local authorities together with the Central government and other stakeholders need to intervene by promoting programmes targeting critical 'vehicle' foods. Street food trade deserves recognition by local and national authorities and the attention of urban policy makers, in order to improve the opportunities of vendors to ensure their livelihood and to ensure the availability of cheap, safe and nutritious food for low-income consumers. This will in turn help to feed the increasing number of working urbanites with quality foods.

## ACKNOWLEDGMENT

Special thanks to Urban Harvest and International Potato Center, Nairobi for funding part of this study. Additional funds from World Bank through the Kenya Agricultural Productivity Project, Clive West Micronutrient Grant of the University of Wageningen and Food Science and Nutrition Network for Africa are highly appreciated.

**Table 1: Occupation and income status in Kangemi and Dandora Nairobi**

Characteristics	% household		
	Kangemi n=300)	Dandora n=300)	Mean of both sites
Occupation status of household head (%)			
Unemployed	3.4	2.6	3.0
Temporarily employed <sup>‡‡</sup>	26.5	18.1	22.3
Permanently employed	47.7	46.7	47.2
Self employed	24.4	32.6	28.5
Average household income per month (mean± SD in Kshs)	10,518.7 ±9444.7	11,604.4 ±8653.1	11,0604.8 ±9044.9
Proportion staying in single rooms (%)	75.9	47.1	60.2
Type of the roof of the house (%)			
Grass thatched	0.3	0.0	0.2
Tiles	2.7	0.7	1.7
Corrugated iron sheets <sup>‡‡</sup>	94.3	60.7	77.5
Cemented <sup>‡‡</sup>	2.7	38.0	20.4
Type of floor of the house (%)			
Cement	96.7	97.7	97.2
Wooden	1.0	0.5	0.5
Linolium	1.3	2.0	1.7
Type of wall for the house (%)			
Unfired mud brick	0.3	0.0	0.2
Fired mud brick	1.7	1.3	1.5
Stones and cement <sup>‡‡</sup>	34.4	96.0	65.3
Wood <sup>‡‡</sup>	13.7	0.3	7.0
Iron sheet <sup>‡‡</sup>	49.8	2.3	26.0
Using electricity <sup>‡‡</sup> (%)	61.9	91.0	76.5

<sup>‡‡</sup> There is statistical difference ( $p < 0.05$ ) using  $\chi^2$ , ( $\phi$  and Cramer's V statistics)

**Table 2: Frequency of street food consumption in Kangemi and Dandora, Nairobi**

Days/week	% households consuming street foods		
	Kangemi n=300)	Dandora (n=300)	Mean of both sites (n=600)
0 days/week	17.4	17.4	17.4
1-2 days/week	29.5	25.8	27.6
3-5 days/week	30.9	35.8	33.3
6-7 days/week	22.1	21.1	21.6
Mean frequency (mean ± SD)	3.1 ±2.4	3.24 ±2.4	3.09±2.4

<sup>‡‡</sup> There is statistical difference ( $p < 0.05$ ) using  $\chi^2$ , ( $\phi$  and Cramer's V statistics)

**Table 3: Selected characteristics of street food vendors in Kangemi and Dandora, Nairobi**

Characteristics	Kangemi (n=38)	Dandora (n=53)	Mean of both sites (n=91)
Age of the vendors (mean±SD)	30.6±8.8	32.6 ±9.7	31.8±9.3
Sex of the vendors (% female) <sup>‡‡</sup>	28.9	51.9	42.2
Level of education (%)			
Primary level <sup>‡‡</sup>	50.0	30.8	38.9
Secondary level	47.4	53.8	51.1
Post-secondary level <sup>‡‡</sup>	26.0	7.7	5.60
Never been to school <sup>‡‡</sup>	0.0	7.7	4.40
Number of years in SF selling (%)			
1 year and below	50.0	50.0	50.0
Between 1-2 years <sup>‡‡</sup>	21.1	13.5	16.7
More than 2 years <sup>‡‡</sup>	28.6	38.5	33.3
Source of cooking ingredients (%)			
Wholesale markets <sup>‡‡</sup>	43.2	27.5	34.1
Retail markets <sup>‡‡</sup>	43.2	58.8	52.3
Own farm	0.0	0.0	0.0
Delivery from agents	8.1	9.8	9.1
Various sources	5.4	3.9	4.5
Main customers (%)			
Adult males	26.3	21.6	23.6
Adult females	10.5	13.7	12.4
Children	5.2	4.0	4.4
All ages and gender	57.9	60.8	59.6
Source of energy for cooking (%)			
Charcoal <sup>‡‡</sup>	81.6	40.7	52.7
Wood <sup>‡‡</sup>	42.1	19.78	26.4
Electricity	2.6	7.7	6.2

<sup>‡‡</sup> There is statistical difference ( $p=0.05$ ) using  $\chi^2$ , ( $\phi$  and Cramer's V statistics)

**Table 4: Classification of street foods sold by vendor in Kangemi and Dandora, Nairobi**

Classification	Mean number of foods produced by the vendors	
	Kangemi (n=38)	Dandora (n=52)
Protein**	0.7 (58.0)	1.1 (71.2)
Carbohydrates	1.8 (79.0)	1.6 (76.9)
Micronutrient	0.3 (29.0)	0.5 (34.6)
Mixed nutrients (e.g. Maize+Beans)	0.4 (38.8)	0.5 (48.1)

In parenthesis are % vendors selling the food group. \*\*There is statistical difference ( $p<0.05$ ) using student's t-test.

**Table 5: Daily household vitamin A intake in  $\mu\text{gRE/AE}$  ( $\pm$  SD), Kangemi and Dandora, Nairobi**

	<b>Kangemi (n=60)</b>	<b>Dandora (n=60)</b>	<b>Mean of both sites</b>	<b>p-value</b>
Vitamin A intake from household prepared food	427.92 $\pm$ 7.51	208.43 $\pm$ 3.62	321.40	-
Vitamin A intake from street food	19.83 $\pm$ 0.77	71.19 $\pm$ 3.81	44.76	-
Total vitamin A intake	447.75 $\pm$ 8.28	279.62 $\pm$ 7.43	366.15	-
Proportion of vitamin A intake from street food (%)	4.43	25.46	12.22	0.03
Percent RDA from streets food	2.64	9.49	6.06	-
Proportion of those who obtain at least 50% of vitamin A from street food (%)	8.60	23.30	15.83	0.03

**Table 6: Daily household iron intake in mg/AE ( $\pm$  SD), Kangemi and Dandora, Nairobi.**

	<b>Kangemi (n=60)</b>	<b>Dandora (n=60)</b>	<b>Mean of both sites</b>	<b>p-value</b>
Iron intake from household prepared food	54.12 $\pm$ 6.88	59.62 $\pm$ 7.61	56.75	-
Iron intake from street food	14.61 $\pm$ 1.92	25.62 $\pm$ 3.32	19.88	-
Total iron intake	68.73 $\pm$ 8.80	85.24 $\pm$ 10.93	76.64	-
Proportion of iron intake from street (%)	21.26	30.06	25.95	0.04
Percent RDA from streets	85.94	150.71	118.32	-
Proportion of those who obtain at least 50% of iron from street (%)	5.00	11.70	8.35	0.19

**Table 7: Daily household zinc intake in mg/AE ( $\pm$  SD), Kangemi and Dandora, Nairobi**

	<b>Kangemi (n=60)</b>	<b>Dandora (n=60)</b>	<b>Mean of both sites</b>	<b>p-value</b>
Zinc intake from household prepared food	5.95 $\pm$ 0.75	7.28 $\pm$ 0.92	6.59	-
Zinc intake from street food	1.19 $\pm$ 0.15	2.46 $\pm$ 0.32	1.80	-
Total zinc intake	7.14 $\pm$ 0.91	9.74 $\pm$ 1.2	8.39	-
Proportion of zinc intake from street (%)	16.72	25.22	21.45	0.20
Percent RDA from streets	11.9	24.6	18.25	-
Proportion of those who obtain 50% of zinc from street (%)	6.67	11.67	9.17	0.34

## REFERENCES

1. **FAO.** The State of Food and Agriculture; FAO, Rome, 1997.
2. **Mwangi AM** Nutritional, hygienic and socio-economic dimensions of street foods in urban areas: the case of Nairobi. PhD-thesis Wageningen University Wageningen, Ponsen en Looijen, 2002.
3. **Mwangi AM, Hartog AP den, Foeken DWJ, Riet H van't, Mwadime RKN, Staveren WA** The ecology of street foods in Nairobi. *Ecol. Food Nutr.* 2001; **40**: 497-523.
4. **H van't Riet, AP den Hartog, Mwangi AM, Mwadime RKN, Foeken DWJ and WA van Staveren** The role of street foods in dietary pattern of two low-income groups in Nairobi. *European Journal of Clinical Nutrition*, 2001; **55**:562-570.
5. **Mwaniki DL, Omwega AM, Munia EM, Mutunga JN, Akelola R, Shako BR, Gotink MH and AM Peter** Anemia and status of iron, vitamin A and zinc in Kenya. The 1999 national survey. Overview of findings, conclusions and recommendations. Policy brief in anemia and status of iron, vitamin A and zinc, 2001.
6. **WHO.** Nutrition, Micronutrients.  
<http://www.who.int/nutrition/topics/micronutrients/en/index.html>.
7. **UNICEF/MI.** Vitamin and mineral deficiency: A global progress report, the United Nations' Children's Fund, 2004.
8. **FAO.** The scourge of hidden hunger: Global dimensions of micronutrient deficiencies, Food, Nutrition and Agriculture, Food and Agriculture Organization of the United Nations, 2003.
9. **GoK/UNICEF.** Situation analysis of women and children in Kenya, United Nations Office, Nairobi, 1998.
10. **GoK/ UNDP.** *Kenya Human Development Report*. Vol.II: Poverty and social indicators, United Nations Office, Nairobi, 1999.
11. **Mwadime RK, Oyunga AM.** A qualitative survey on issues related to VAD in Nyarongi Division, Homabay District, Sight and Life, 1998; **4**: 29-35.
12. **MI.** Solutions for hidden hunger: Accelerating food fortification. A rapid results initiative in Kenya. [www.micronutrient.org](http://www.micronutrient.org). 2007.

13. **Draper A** Street foods in developing countries: The Potential for micronutrient fortification, London School of Hygiene and Tropical Medicine, OMNI/USAID Publication, 1996.
14. **GoK.** Population distribution by administrative areas and urban centers. Volume 1. The 1999 population and housing census. Central Bureau of Statistics, Ministry of Finance and Planning, Nairobi, Kenya. 2001.
15. **Fisher AA, Laing JE, Stoeckel JE and JW Townsend** Handbook for family planning operations research design. 2<sup>nd</sup> ed., Population Council, New York, 1998.
16. **FAO.** Conducting small scale nutrition surveys. A Field Manual. Nutrition in Agriculture, 1990; No. 5.
17. **Mugenda OM and AG Mugenda** **Research Methods.** Quantitative and Qualitative Approaches. Acts Press African Center for Technology studies (ACTS) Nairobi, Kenya, 1999.
18. **Gibson RS and EL Ferguson** An interactive 24-hour recall for assessing the adequacy of iron and zinc intakes in developing countries. ILSI press, Washington DC, USA: 1999; 120-124.
19. **Sehmi JK** National food and composition tables and planning of satisfactory diets in Kenya: Ministry of Health/National Public Health Laboratory Services, Kenya, 1993.
20. **FAO-Infoods** [http://www.fao.org/infoods/software\\_worldfood\\_en.stm](http://www.fao.org/infoods/software_worldfood_en.stm).
21. **Institute of Medicine** Dietary reference intakes for vitamin A, vitamin K, arsenic, boron chromium, copper, iodine iron manganese molybdenum nickel, silicon, vanadium and zinc. Institute of Medicine. National Academy Press, Washington D.C. 2001; pp 4-22.
22. **Williams SR** Nutrition and diet therapy. Mosby-year Book, Inc. St Louis, MO. USA, 1993.
23. **Gopalan C** Micronutrient deficiencies: Public health implications. NFI Bulletin **15(3):1- 6.** 1994.
24. **Boccas F** Codex Alimentarius: Control of street foods. Food, Nutrition and Agriculture, 1996.**17/18;** 70-72.
25. **Tinker I** The case for legalizing street foods. Ceres; 1987; **20 (5):** 26-31.

26. **Jonnalagadda SS, Seshadri S.** In vitro availability of iron from cereal meal with the addition of protein isolates and fenugreek leaves (*Trigonella foenum-graecum*). *Plant Foods Hum. Nutr.* 1994; **45**: 119-125.
27. **Engel.**Engels Law and curves,  
<http://faculty.washington.edu/krumme/resources/engel.html>.
28. **Fisher MS, Christian P and KP West Jr.** The role of vitamins in the prevention and control of Anemia, *Public Health Nutrition*; Johns Hopkins Bloomberg School of Public Health, 2000; **3 (2)**: 125-150.
29. **Okoth JK.** Bioavailability of iron and zinc in selected foods commonly consumed in Kenya. 2004; MSc Thesis. Jomo Kenyatta University of Agricultural and Technology (JKUAT).
30. **Walingo M** Indigenous food processing methods that improve zinc absorption and bioavailability of plant diets consumed by the Kenyan population. *Af. J. Food Agric. Nutr. and Dev.*, 2009; **9 (1)**: 523-535.