

PHYTOEXTRACTION OF METAL POLLUTED SOILS IN LATIN AMERICA

Maria N. Dos Santos Utmazian and Walter W. Wenzel*

Department of Forest- and Soil Sciences. University of Natural Resources and Applied Life Sciences, Vienna. Peter-Jordan-Str. 82, A-1190 Vienna.

Tel.: ++43147654/ 3125. Fax Nr. 3130*

*corresponding author: walter.wenzel@boku.ac.at

Abstract

Phytoextraction is a cost-effective method that could be an alternative to remediate polluted sites in Latin America. In this communication, basic concepts relating to phytoextraction of metal-polluted soils are discussed as well as some limitations to and perspectives for applying this method in the region are considered.

Keywords: phytoremediation, phytoextraction, metals, contamination, Latin America.

Introduction

The long mining history in Latin America was not supported by adequate environmental laws and regulations until recent decades. Therefore, metal mining imposed severe and diverse environmental problems in the region, including wild environments, croplands, and human health (Ginocchio, 2003). Most of the mining sites are in the Andean countries (Table 1) and therefore high natural metal concentrations in soil and metal contamination problems can be expected. On the other hand, some soils in industrialized areas are contaminated either by heavy metals atmospheric deposition or through application of metal-rich sewage sludge, fertilizers, pesticides, etc. Unfortunately, there is no complete inventory showing polluted sites in Latin America, but there are several examples reported in the literature (Table 1). Likewise, site remediation is generally not a priority in these countries and current technologies for soil remediation are time consuming, too expensive, and in some cases create additional risks to the workers and produce secondary waste. Therefore, it is imperative to develop techniques that can treat and stabilize contaminants in situ in an efficient and cost effective manner.

Phytoremediation are technologies that use green or higher terrestrial plants for treating chemically- or radioactively- polluted soils (Wenzel et al., 1999). These technologies permit the decontamination in-situ, preserves the topsoil, reduces the amount of hazardous materials generated during clean-up and have public acceptance. Thereafter, the polluted plants have a proper control of the disposal and could be used to generate energy. Some of the disadvantages are related to the depth of the treatment and time required.

There are different fundamental processes by which plants can be used to remediate metal contaminated soils/ sediments (Wenzel et al., 1999):

- Containment processes: phytostabilisation and phytoimmobilisation; and
- Removal processes: phytoextraction, phytodegradation and phytovolatilization.

Among these technologies cleanup may be achieved with the removal processes to an extent depending on the type of contaminants and soil-chemical properties. The objective of this communication is to discuss basic concepts relating to phytoextraction of metal-polluted soils as well as some limitations to and perspectives for applying this method in Latin America.

Table 1 Mining ores and some polluted sites reported in Latin America.

Country	Mine ores Ginocchio (2003)	Metal contaminated sites	
Argentina	Zn, Pb, Cu, Au		
Bolivia	Au, Sn, Sb.	Muñoz, 2004	Hg
Brazil	Al, Au, Fe, Mn, Sn	Biller, 1994 Da Silva et al., 2003	Hg Cd, Zn
Chile	Cu	Ginocchio, 2000, 2003; Bech, 2002	Cu, As
Colombia	Ni, Au, and Fe-Ni		
Cuba, Puerto Rico Dominican Republic	Fe-Ni, Co		
Ecuador		Bech, 2002 Requelme et al., 2003	As, Cu, Zn, Cd Hg
Mexico	Ag, As, Cu, Pb, Mn, Mo, Zn	Bech et al., 1997; Razo et al., 2004 Valdés & Cabrera, 1999	As Pb
Peru	Cu, Au, Fe, Ag, Pb, Sn, Zn	Tolmos, 2000; Bech, 1997	Cu, As, Zn, Cd
Uruguay		Intendencia Municipal de Montevideo, 2003	Pb
Venezuela	Al, Au, Fe, Co, steel, bauxite, alumina, Fe-Ni		

Basically, a phytoremediation project consist of different phases, the application of which is mainly a function of the available time-costs:

- Sampling and analysis of native species and soils in polluted areas.
- Greenhouse experiments:
 - Hydroponic experiments (1-2 months): Rooted plants grown in pots containing a modified nutrient solution. After an acclimatization period, plants are exposed to different metal treatments. It is recommended in particular that the capacity of the plant for (hyper)accumulation and tolerance to metals be evaluated, phytotoxicity essays, or to carry out a fast screening among many species;
 - Pot experiments (1 vegetation period): the most promising species are planted in naturally polluted or spiked soils;
- Field experiments (minimum 1 vegetation period): the most promising species are planted in the polluted soil. The same species should be also evaluated in other soils and under other environmental conditions to verify the phytoremediation potential of the selected plants, and to confirm whether the property of accumulation, for example, is constitutive in the plant.

In parallel, experiments can be carried out to enhance the phytoremediation potential of the plants, e.g. improving the biomass production or increasing the metal bioavailability. The use of mycorrhiza has also been suggested to enhance the

phytoextraction process or to help establishment of the plants in polluted soils (González-Chávez et al., 2002). Genomic approaches are increasingly being employed in phytoremediation-related research (Krämer, 2005) but there is a limitation related to public acceptance.

Phytoextraction

Phytoextraction processes extract both metallic and organic constituents from soil by direct uptake into plants and translocation to aboveground biomass using metal-(hyper)accumulating plants. The criteria described by Brooks (1998) to define hyperaccumulators are:

- Exceeding the hyperaccumulation threshold: Concentration 100 times higher than in normal plants for each metal of interest, i.e. (mg kg^{-1}): 100 Cd; 1000 As, Ni; 10000 Zn, Mn, etc.;
- Bioconcentration factor > 1 (concentration of the element in the plant $>$ concentration in the soil); and
- Translocation factor > 1 (element concentration in the overground part of the plant $>$ than in roots).

To date, the best studied hyperaccumulators ($< 0.2\%$ angiosperms) are: *Brassica juncea*, *Brassica oleracea*, *Berkeya coddii*, *Allysum bertolonii*, *Thlaspi caerulescens* and *Thlaspi goesingense*. Likewise, numerous plants with hyperaccumulation potential or tolerance to metals have been discovered in Brazil, Chile, Cuba, the Dominican Republic and Venezuela and include Ni (89%), Cu (5%) and As (3%) hyperaccumulators (Ginocchio & Backer, 2004).

Hyperaccumulators have been suggested for phytoextraction because of the large quantities of a specific metal that they are able to accumulate. However, most of these plants achieve low shoot biomass, which consequently considerable time period necessary for the remediation before an acceptable metal level in the soils is reached. Therefore, the use of trees in phytoextraction has been suggested and recently reviewed by Pulford & Dickinson (2005). *Alnus incana*, *Betula pendula*, *Fraxinus excelsior*, *Sorbus mongoeotii*, *Salix* and *Populus* species are some of the trees reported to grow on contaminated lands. Based on screenings, it has been found that *Salix* (willows) and *Populus* (poplars) are capable of accumulating metals, especially Cd and Zn in substantially large amounts in their leaves. Large metal concentrations ($300 \text{ mg Cd kg}^{-1}$) were found in leaves of *Salix smithiana* BOKU 03 CZ-001, *S. purpurea* BOKU 05 CZ-001 and *S. caprea* BOKU 01 AT-004 clones growing on contaminated soils ($\text{Cd}_{\text{total}} 30 \text{ mg kg}^{-1}$) (Dos Santos et al., submitted) and in shoots of *S. viminalis* ($2695 \text{ mg Zn kg}^{-1}$) grown on an acidic soil ($\text{Zn}_{\text{total}} 1158 \text{ mg kg}^{-1}$) (Hammer et al., 2003).

The phytoextraction process can be continuous (natural) using (hyper)accumulators or induced through the addition of chelates to increase the soil bioavailability. However, some authors have reported the persistence of metallic complexes in the water of the soil for several weeks in experiments using EDTA (ethylenediamine tetraacetic acid) to induce the phytoextraction (Lombi et al., 2001; Wenzel et al., 2003). For this reason, the chelates must be handled with care to avoid possible environmental risks due to the leachate enriched with metals.

Perspectives and limitations in Latin America

Currently, public institutions and private companies in Latin America are applying phytoremediation technologies to remediate soils polluted with organic compounds (Heidtmann, 2001, Fernandez et al., 2001; Olguin, 2001). However, there is a scarcity of studies aimed at cleaning up sites contaminated with heavy metals using

phytoremediation technologies. Therefore this innovative and cost-effective alternative may present a potential market in Latin American countries, considering the numerous metallic ores in the region.

Potential species for phytoextraction purposes such as willows and poplars can be profitably used by the farmers i.e. for heat/ electricity production through the burning of harvested biomass. Also the comparative economics of the phytoremediation processes in general will also be a driving force for acceptance and include simple methods which can be applied by farmers.

Research on the following areas related to plant characterization and remediation of metal contaminated sites has been carried out in Latin America:

- Description of metal tolerant and hyperaccumulating plants (Brooks et al., 1992; Barreto and Casale, 2002; Flores-Tavizón et al., 2003; Reeves, 2003; Ginocchio, 2003, Ginocchio & Backer, 2004),
- Determination of soils, sediments and plants metal concentrations of species growing near the polluted sources, smelters, etc. (Jordao et al., 1999; Bech et al., 2002; Arribere et al., 2003; Carillo & González, 2005; Ginocchio, 2000),
- Studies related to metal accumulation and tolerance in seeds (Marques et al., 2000) and plants (Dos Santos et al., submitted);
- Revegetation (PUC-ECU, 1998),
- Use of microorganisms for phytoremediation purposes (Pérez et al., 2002) and
- Phytoremediation studies (Cusato et al., 2002, Pletsch et. al, 1999; Ortiz, 2002; Miretzki et al., 2004, Anderson et al., 2005).

Few metal-tolerant and metal-hyper accumulating plants have been reported for Latin America (172) in comparison with other areas of the world. Due to the scarcity of scientific studies of the native vegetation growing on natural mineralised or metal-contaminated sites (Ginocchio & Baker, 2004).

Plant selection criteria –prior to running a phytoremediation program in Latin America- should include the life cycle of the plant (dormancy period and harvest), plant cultivation, maintenance requirements and germination potential (Merkl et al., 2004), plant tolerance to the metals of interest, metal accumulation potential, easy propagation, availability of the plant material (commercially/ field), favourable root system and use of native (preferable) or introduced species.

Many studies related to soil and water characterization are unpublished, in the property of private companies or are not up to date. Only few sites near mining areas have been characterized due to the metal contamination and/ or recognized environmental and human risks. The scarcity of relevant publications in international journals does not help dissemination and effective transfer of information; this gives rise to unnecessary repetition and/ or loss of data of basic research.

It is very important not only to adapt the available study techniques to the climate, soil and vegetation conditions of the individual regions, but also consider the available technologies (maintenance/ monitoring) before phytoremediation technologies can be applied in the region. Special attention should be given to the metal-enriched biomass that requires a safe disposal. Recycling, incineration and/ or landfill solutions that should not be a problem in developed countries could be of limited availability in Latin America.

The social, economic and political situation in a given country also influences the strategies that governments choose to manage the problems of contamination (limit values), revegetation, etc. (Ginocchio, 2003). Local scientists have not been strongly involved in phytoremediation issues until recent years, maybe as result of the scarcity

of resources (basic research, field experiments, analytical laboratories, etc.) and limited transfer of knowledge in this topic.

Conclusions

Phytoextraction in Latin America and in the developing countries in general, offers a feasible and economic alternative to achieve remediation of contaminated soils. It is hoped that governments will recognize the importance of the issues raised and support research on phytoremediation as a practical and effective technology for soil remediation. It is likely that governments could also draw on collateral funds from international agencies and other donors.

References

- Anderson, C.; Moreno, F.; Meech, J. A field demonstration of gold phytoextraction technology: *Minerals Engineering*. v. 18, n. 4, p. 385-392, 2005.
- Arribere, M.A.; Guevara, S.R.; Sanchez, R.S.; Gil, M.I.; Ross, G.R.; Daurade, L.E.; Fajon, V.; Horvat, M.; Alcalde, R.; Kestelman, A.J. Heavy metals in the vicinity of a chlor-alkali factory in the upper Negro River ecosystem, Northern Patagonia, Argentina. *Science of the Total Environment*, v. 301, n. 1-3, p. 187-203, 2003.
- Barreto, L.; Casale, I. Caracterización de las plantas asociadas a los suelos serpentiniticos de Loma de Hierro. Venezuela. In: Congreso Latinoamericano de Botánica, 8th. Proceedings. Cartagena de Indias. 2002. p. 357.
- Bech, J.; Poschenrieder, C.; Llugany, M., Barceló, J.; Tume, P.; Tobias, F.J.; Barranzuela, J.L.; Vásquez, E.R. Arsenic and heavy metal contamination of soil and vegetation around a copper mine in Northern Peru. *Science Total Environment*, v. 203, p. 83-91, 1997.
- Bech, J.; Poschenrieder, C.; Barceló, J.; Lansac, A. Plants from mine spoils in the South American area as potential source of germplasm for phytoremediation technologies. *Acta Biotechnologica* v. 22, p. 5-11, 2002.
- Billier, A. Informal Gold Mining and Mercury Pollution in Brazil. Paper provided by The World Bank in its series Policy Research Working Paper Series with number 1304, 1994. Accessed on 11 Jan. 2005 Available in:
<http://ideas.repec.org/p/wbk/wbrwps/1304.html> Accessed on 11 May 2006
- Brooks, R.R.; Reeves, R.D. ; Baker, A.J.M. The serpentine vegetation of the Goiás State, Brazil. In: Baker, A.J.M.; Proctor, J.; Reeves, R.D. (Ed.) *The vegetation of ultramafic (serpentine) soils*. Andover, UK: Intercept Ltd, 1992, 509 p.
- Brooks, R.R. *Plants that hyperaccumulate heavy metals. Their role in phytoremediation, microbiology, archaeology, mineral exploration and phytomining*. Wallingford, Oxon : CAB International, 1998. 380 p.
- Carrillo, R.; González M.C.A. Metal Accumulation in Wild Plants Surrounding Mining Wastes. In: International Conference on the Biogeochemistry of Trace Elements, 8. Proceedings. Adelaide, 2005.
- Cusato, M.S.; Valiente, L.; Tortosa, R.D.; Bartoloni, N.; Puelles, M.M.; D'ambrogio, A. Uso de discaria americana en fitorremediación. In: Jornadas de Desarrollo e Innovación, 4. Proceedings. Buenos Aires, 2002. p 1-2.
- Da Silva, A.C.; Mendonca, E.D.; Fontes, R.L.F.; De Marco, P., Ribeiro, F.P. Potential environmental consequences of zinc and cadmium contamination in kaolin minespoil located in Minas Gerais, Brazil. *Communications in Soil Science and Plant Analysis*, v. 34, n. 5-6, p. 671-680, 2003.

- Dos Santos Utmazian, M.N.; Wieshammer, G.; Vega Montoya, R.M. and Wenzel, W.W. Hydroponic screening for metal resistance and accumulation of cadmium and zinc in willows and poplars, submitted.
- Dos Santos Utmazian, M.N. and Wenzel W.W. Screening for cadmium and zinc accumulation by willow and poplar species grown on polluted soils, submitted.
- Fernández, L.; Molina, I.; Quiñónez, E.; Hernández, E ; Ferrera, R. (2001) Potential of phytoremediation in Mexican oil contaminated soil. In: International In-situ and On-site Bioremediation Symposium, 6th. Proceedings. San Diego.
- Flores-Tavizon, E.; Alarcon-Herrera, M.T.; Gonzalez-Elizondo, S.; Olguin, E.J. Arsenic tolerating plants from mine sites and hot springs in the semi-arid region of Chihuahua, Mexico. *Acta Biotechnologica*, vol. 23, n. 2-3, p. 113-119, 2003.
- Ginocchio, R. Effects of a copper smelter on a grassland community in the Puchuncavi Valley, Chile. *Chemosphere*, 41, 1-2, p. 15-23, 2000
- Ginocchio, R. Mining for ores and metal tolerant plants in Latin America. In: International Meeting on Preservation and Ecological Restoration in Tropical Mining Environments. Proceedings. Noumea. 2003.
- Ginocchio, R.; Baker, A.J.M. Metallophytes in Latin America: a remarkable biological and genetic resource scarcely known and studied in the region. *Revista Chilena de Historia Natural*. v. 77, p. 185-194, 2004.
- Gonzalez-Chavez, C.; Harris, P.J.; Dodd, J.; Meharg, A.A. Arbuscular mycorrhizal fungi confer enhanced arsenate resistance on *Holcus lanatus*. *New Phytologist*, v. 155, p. 163-171, 2002.
- Hammer, D.; Kayser, A.; Keller, C. Phytoextraction of Cd and Zn with *Salix viminalis* in field trials. *Soil use and Management*, vol. 19, n. 3, p. 187-192, 2003.
- Heidtmann, M. The Potential of Phytoremediation in Venezuelan Soils contaminated by the Petroleum Industry. Höhenheim, 2001. Thesis (Diplomarbeit) – University of Höhenheim.
- Intendencia Municipal de Montevideo. Informe Ambiental. Contaminación por metales en el suelo, 2003. Available in: <http://www.montevideo.gub.uy/ambiente/documentos/infoamb03c.pdf> Accessed on 11 May 2006
- Jordao C.P.; Pereira J.L., Jham G.N., Bellato C.R. Distribution of heavy metals in environmental samples near smelters and mining areas in Brazil. *Environmental technology*, v 20, n. 5, pp. 489-498, 1999.
- Krämer, U. Phytoremediation: novel approaches to cleaning up polluted soils. *Current Opinion in Biotechnology*, v. 16, p. 133-141, 2005.
- Lombi, E. Zhao, F.J.; Dunham, S.J.; Mcgrath, S.P. Phytoremediation of Heavy Metal-Contaminated Soils: Natural Hyperaccumulation versus Chemically Enhanced Phytoextraction. *Journal of Environmental Quality* v. 30, p. 1919-1926, 2001.
- Marques, T. C. L. L. ; Moreira, F. M.; Siqueira, J. O. Crescimento e teor de metais de mudas de espécies arbóreas cultivadas em solo contaminado com metais pesados. *Pesquisa Agropecuária Brasileira*. vol.35, n.1, p.121-132, 2000.
- Merkel, N.; Schultze-Kraft, R.; Infante, C. Phytoremediation of Petroleum-contaminated soils in the tropics- Pre-selection of plant species from eastern Venezuela. *Journal of Applied Botany and Food Quality*, v. 78, p. 185-192, 2004.

- Miretzky, P.; Saralegui, A.; Cirelli, A.F. Aquatic macrophytes potential for the simultaneous removal of heavy metals (Buenos Aires, Argentina). *Chemosphere*, v. 57, n.8, p. 997-1005, 2004
- Muñoz, A. Gestión Sostenible de la Diversidad Biológica en Bolivia: Factor desaprovechado en el Desarrollo Nacional. *Ambiente Ecológico*, 88, 2004. Available in: <http://www.ambiente-ecologico.com/> Accessed on 11 May 2006
- Olguin, E.J. Application of phytoremediation in Mexico. In: *Phytoremediation. ISEB 2001 Meeting. Proceedings. Leipzig. 2001*, p. 44.
- Ortiz, C.; Li Kao, J.; Orellana, S. (2002). Acumulación y tolerancia a Cu⁺⁺ en una gramínea silvestre: Potencial de uso en fitorremediación. *Biological Research*, v. 35, p. R-93.
- Perez, J.; García, G.; Esparza, F. Papel Ecológico de la flora rizosférica en fitorremediación. *Avance y Perspectiva*, v. 21, p. 297-300, 2002.
- Pletsch, M.; De Araujo, B.S.; Charlwood, B.V. Novel biotechnological approaches in environmental remediation research. *Biotechnology Advances*, v. 17, (8), p. 679-687, 1999.
- PUC-ECUS. Estudio sobre revegetación potencial de sustratos contaminados con metales pesados en Chile: el Valle de Puchuncaví como caso de estudio. Informe Técnico, Ministerio de Minería, 1998.
- Pulford, I.D.; Dickinson, N.M.. *Phytoremediation Technologies Using Trees*. In Prasad, M.N.V.; Sajwan, K.S.; Naidu, R. Trace elements in the environment. Boca Raton, Fla: Lewis. 2005, p. 375-395.
- Razo, I.; Carrizales, L.; Castro, J.; Diaz-Barriga, F.; Monroy, M. Arsenic and heavy metal pollution of soil, water and sediments in a semi-arid climate mining area in Mexico. *Water Air and Soil Pollution*, vol. 152, n. 1-4, p. 129-152, 2004.
- Reeves, R.D. Tropical hyperaccumulators of metals and their potential for phytoextraction. *Plant and Soil*, v. 249, n. 1, p. 57-65, 2003.
- Requelme, M.E.R.; Ramos, J.F.F.; Angelica, R.S.; Brabo, E.S. Assessment of Hg-contamination in soils and stream sediments in the mineral district of Nambija, Ecuadorian Amazon (example of an impacted area affected by artisanal gold mining) *Applied Geochemistry*, v.18, n. 3, p. 371-381, 2003.
- Tolmos, R. (2000) Minería y Medio Ambiente en Perú. *Revista Ambiente y Desarrollo de CIPMA*. Available in: <http://www.cipma.cl/hyperforum/MineriaMedio%20AmbientePeru.doc> Accessed on 11th May 2006.
- Valdés, F.; Cabrera, V. (1999) Contaminación por metales pesados en Torreón, Coahuila, México. Texas Center for Policy Studies. Available in: <http://www.texascenter.org/publications/torreon.pdf> Accessed on 11th May 2006.
- Wenzel, W.W.; Lombi, E.; Adriano, D. Biogeochemical Processes in the Rhizosphere: Role in Phytoremediation of Metal-Polluted Soils. In *Heavy metal stress in plants - from molecules to ecosystems*. Prasad, N.; Hagemeyer, J. (Ed) Heidelberg, Springer Verlag. 1999, pp. 273-303.
- Wenzel, W.W.; Unterbrunner, R.; Sommer, P.; Sacco, P. Chelate assisted phytoextraction using canola (*Brassica napus L.*) in outdoors pot and lysimeter experiments. *Plant and soil* 249: 83-9, 2003.