



Forestry Department

Food and Agriculture Organization of the United Nations

Forest Genetic Resources Working Papers

*State of forest genetic resources
conservation and management in
Bangladesh*

by

Sk. Sirajul Islam

Bangladesh Forest Research Institute

2003

Forest Resources Development Service
Forest Resources Division

Working Paper FGR/68E
FAO, Rome, Italy

Disclaimer

The Forest Genetic Resources Working Papers report on issues and activities in related to the conservation, sustainable use and management of forest genetic resources. The purpose of these papers is to provide early information on on-going activities and programmes and to stimulate discussion.

The designations employed and the presentation of material in this publication do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

Quantitative information regarding the status of forest resources has been compiled according to sources, methodologies and protocols identified and selected by the author, for assessing the diversity and status of genetic resources. For standardized methodologies and assessments on forest resources, please refer to FAO, 2003. *State of the World's Forests 2003*; and to FAO, 2001. *Global Forest Resources Assessment 2000 (FRA 2000)*. FAO Forestry Paper No 140. Official information can also be found at the FAO Internet site (<http://www.fao.org/forestry/Forestry.asp>).

This paper is based on a country report prepared for the Asia Pacific Forest Genetic Resources Programme (APFORGEN) Inception Workshop, held at Forest Research Institute Malaysia (FRIM), Kuala Lumpur, Malaysia, 15-18 July 200. The Workshop was organized by the Asia Pacific Association of Forestry Research Institutions (APAFRI) and the International Plant Genetic Resources Institute (IPGRI), Regional Office for Asia, the Pacific and Oceania, in technical collaboration with FAO. The proceedings of the workshop, including this paper, will be published in "Proceedings of the Asia Pacific Forest Genetic Resources Programme (APFORGEN) Inception Workshop, Kepong, Kuala Lumpur, Malaysia, 15-18 July, 2003" and available on line at: "<http://www.apforgen.org/>".

Comments and feedback are welcome.

For further information please contact:

Pierre Sigaud, Forestry Officer (Forest Genetic Resources)
Forestry Department
FAO, Viale delle Terme di Caracalla
00100 Rome, Italy
Fax: + 39 06 570 55 137
[Email: pierre.sigaud@fao.org](mailto:pierre.sigaud@fao.org)

Sim Heok-Choh, Ph.D.
Executive Director
APAFRI Secretariat
c/o Forest Research Institute Malaysia
Kepong, 52109 Kuala Lumpur
Malaysia
[Email: simhc@frim.gov.my](mailto:simhc@frim.gov.my)

For quotation:

Sirajul Islam, Sk. 2003. *State of forest genetic resources conservation and management in Bangladesh*. Forest Genetic Resources Working Papers, Working Paper FGR/68E. Forest Resources Development Service, Forest Resources Division. FAO, Rome.

TABLE OF CONTENTS

1. INTRODUCTION	1
1.1 FOREST RESOURCES	1
1.2. STATUS OF FOREST GENETIC RESOURCES	2
1.3. IDENTIFICATION OF THREATS	2
2. LINKS BETWEEN FORESTRY SECTOR AND FGR	3
3. LINKS BETWEEN AGRICULTURE AND FGR.....	3
3.1. LINKS BETWEEN AGROFORESTRY AND FGR	4
4. PAST AND PRESENT ACTIVITIES IN CONSERVATION, UTILIZATION AND MANAGEMENT OF FGR	4
4.1. CONSERVATION STRATEGIES	5
4.1.1. <i>In situ</i> conservation	5
4.1.2. <i>Ex situ</i> conservation	8
4.1.3. Conservation of provenances within a species	9
5. INSTITUTIONAL FRAMEWORK	9
5.1. NATIONAL FOREST POLICY	9
6. LIST OF NATIONAL PRIORITY SPECIES	10
7. RESEARCH ON FGR.....	11
8. CONCLUSION AND RECOMMENDATIONS.....	11
REFERENCES.....	13
APPENDICES.....	14
Appendix 1. Value and use of main forest species in Bangladesh	15
Appendix 2. Conservation and management of important FGR by ecogeographic zone in Bangladesh.....	19
Appendix 3. Level and nature of threats to the integrity of species/populations of important tree species	23
Appendix 4. List of priority species for conservation, improvement or seed procurement, their uses and conservation activities needed	25

Tables

TABLE 1. NATIONAL PARKS IN BANGLADESH	6
TABLE 2. WILDLIFE SANCTUARIES IN BANGLADESH.....	6

1. Introduction

Bangladesh is located between latitudes 20°34' to 26°38' north and longitudes 88°01' to 92°41' east. The country consists mostly of flood plains (80%) with some hilly areas (12%), with a sub-tropical monsoon climate. In winter, temperature ranges from a minimum of 7° to 13°C to a maximum of 23° to 32°C. In summer, the temperature varies from 36°C to 41°C. The mean annual rainfall ranges from 143 to 434 cm (BBS 1994). The population stood at 131.6 million in 2001.

1.1 Forest resources

Bangladesh has a total area of 14.39 million hectares, of which 9.12 million ha is cultivated, 2.14 million ha public forests, 0.27 million ha village groves, and 1.64 million ha constantly under water. The remaining land area (1.22 million hectares) is occupied by tea gardens, uncultivable areas, rural and urban houses and ponds (Kibria *et al.* 2000). The area covered by government and village forests is about 16% of the total land area; however only 0.93 million ha (6.5%) is under tree cover, which is about 40% of the forests controlled by the government. The remaining 60% includes denuded lands (grassland, scrubland and encroached areas). About 24 000 ha of forest is lost annually as a result of homestead development, urbanization and deforestation (Anon 1992).

The mangrove forests are composed of two major components, *viz.*, the Sundarbans and the Coastal Afforestation. The Sundarbans forest is spread over 0.58 million ha in the south-western region (Ali 1989), with an annual depletion rate of about 1.7%. A large-scale coastal afforestation programme has been undertaken since 1966, covering about 170 000 ha. The hill forests cover an area of 0.67 million ha. The growing stock has been estimated to be about 28 million m³ or 100 m³ ha⁻¹ (Anon 1992). Annual depletion of the growing stock stands at 1.65%. The estimated average annual forest growth is 2.5 m³ ha⁻¹. The forests cover of 0.12 million ha and to a great degree consists of *Shorea robusta* (sal) located in the central and northern part of the country. The area under tree cover in this region has been estimated to be 32%. A large portion of this forestland has been encroached. About 0.02 million ha of plantations has been established. The unclassified state forests include 0.73 million ha of hilly land located at the southeast corner of the country. The village forest area is 0.27 million ha (TFYP 1985). The growing stock of the woodlots and bamboo resources were estimated to be 54.7 million m³ and 7480 million culms respectively.

The contribution of forestry to the national Gross Domestic Product (GDP) at current prices has been estimated to be 3.28% (BBS 1994). The supply of various forest products such as timber, poles, fuelwood, bamboo, etc. cannot meet the present demand. Village forest areas, being one-tenth of the national forest area, supply 70% of sawlogs, 90% of fuelwood and 90% of bamboo consumption of the country (Douglas 1982). The annual per capita consumption of timber and fuelwood was estimated to be 0.01 m³ and 0.08 m³, respectively, based on a population of 90 million as at 1980 (Byron 1984). The figure would be less if the present forest production and population were considered. The gap between demand and supply has been increasing with the increase in population. This gap may be narrowed through the establishment of plantations of fast growing trees in the denuded forest areas, wastelands as well as homestead areas.

1.2. Status of forest genetic resources

Bangladesh, located in the humid tropical region is rich in species diversity and is unique in the diversity of genetic resources compared to its land area. It has about 5700 species of angiosperms and four species of gymnosperms (Khan 1977; Troup 1975) of which some 2260 species are reported from the Chittagong region (Anon 1992). The major forest tree species of the country are shown in Appendix 1.

Khan (1996) reported that there are about 86 timber species, 130 species yielding fibre and 29 medicinal plant species available in the country. The Bangladesh National Herbarium (BNH) prepared a list of 500 medicinal plants. Bamboo resources of 18 taxa, both wild and planted are available in the country (Alam 1982). There are at least nine species of rattans, including a recently reported one (Alam and Basu 1988), along with 12 other palm species (Khan 1996). However, it must be noted that comprehensive information on forest genetic resources (FGR) is not available in the country. Information on species diversity in terms of (i) chromosome numbers, (ii) morphological variation, (iii) flowering and seed production habits, (iv) flowering time and nature and (v) seed morphology and viability, etc. are available only for a few agricultural species. Forest species are neglected in this respect (Hassan 1995).

1.3. Identification of threats

Forests in Bangladesh are declining at an alarming rate. An estimated 73 000 ha of forests was lost due to encroachment for agriculture and aquaculture by the year 1980. Some species are disappearing fast and are considered as threatened. A total of 19 tree species and nine rattan species need immediate conservation measures (Khan 1996). The on-going loss of germplasm is a threat to FGR. High population pressure on land, clearing of forests, exploitation, draining and filling of wetlands, introduction of exotic species, introduction of improved genotypes, pests, improper silvicultural techniques and management and lack of public awareness are some of the major threats to FGR. Poverty and the attitude of the people towards exploitation of natural habitats of plants as free goods also contribute to the loss of germplasm in the country.

The destruction of forests by shifting cultivation is another problem in Bangladesh. The forests are cut and cultivated for a short time only, and when the residual nutrients in the soil are leached as a result of erosion removing the topsoil, the shifting cultivators move to another location.

The country has four wild gymnosperms, viz. *Cycas pectinata*, *Gnetum scandens*, *G. funiculare* and *Podocarpus nerifolia*. *C. pectinata* is regionally threatened and the two *Gnetum* species have become very rare while the populations of *P. nerifolia* are much depleted. Immediate appropriate conservation measures are needed to protect these species in the country. Khan (1996) reported that the number of plant species threatened in the country is 45. Among mammals, birds, reptiles and amphibians, 15 species have become extinct and 33 species are endangered. According to the Government of Bangladesh (GOB 1992), there are 27 threatened and 39 endangered species of wildlife in Bangladesh at present.

2. Links between forestry sector and FGR

Genetic resources play an important role in improving the quality and quantity of forest products. In this respect, FGR provide the source material for selection of mother trees, provisional plus trees and establishment of seed orchards. It is associated with the conservation of superior genetic materials and their subsequent improvement. Ultimately, the improved materials will be utilized in national programmes to enhance the productivity of forests, and in order to do so the following activities are undertaken by the Forest Department:

- Use of fast-growing genotypes for increasing the production
- Use of best genotype/provenance of proven trees for particular site type
- Use of best genotype to develop interim and alternate sources of improved seeds required for national plantation programme
- Use of best genotypes in tree improvement programmes

3. Links between agriculture and FGR

The Bangladesh Agriculture Research Institute (BARI), Bangladesh Rice Research Institute (BRRI) and Bangladesh Jute Research Institute (BJRI) initiated systematic germplasm conservation activities around the mid-70s. The assistance of Institute of Jute Organization (IJO) added extra emphasis and BJRI now has a global germplasm collection of jute and associated fibre crops. FGR have not received much attention until recently.

Although Bangladesh is a small country it has a diverse ecosystem with hills, plains, coastal areas and wetlands, and a wide variety of FGR as a result of the diverse agroecological conditions [38 agroecological zones and 88 subzones]. The landraces have survived here under the hostile climates over centuries and acquired some characteristics, which are of immense importance to the breeders for development of new cultivars, hybrids, etc, and face challenges of more violent pathogens, pests and a hostile climate (drought, salinity, temperature, etc).

With regard to tree species, collecting, conserving and utilizing fruit germplasm has received some attention. A total of 817 fruit germplasms have been collected so far. Moreover, 53 fruit species have been identified to be occurring mainly in the forests. The conservation of these fruit species is implemented by the BARI, the Department of Agriculture Extension (DAE) and the Bangladesh Agriculture Development Corporation (BADC).

FGR are needed for further incorporation of genotypes into the existing agricultural production systems, reforestation programmes, agroforestry, social forestry and commercial timber estates. Agriculture utilizes FGR in the following ways:

- Selection of superior traits
- Selection breeding
- Identification of high-yielding varieties (HYV)
- Domestication of fruit germplasms

3.1. Links between agroforestry and FGR

Large-scale agroforestry projects have been initiated in heavily degraded *Shorea robusta* forests of the central and northern regions of the country. A community forestry project at Dinajpur is considered a success story of the Forest Department (FD). In this area, plantations have been established in close cooperation with local people on a benefit-sharing basis. At the moment, the people are harvesting the trees. The major tree species are *Eucalyptus camaldulensis*, *Acacia auriculiformis* and *Cassia siamea*. During the first two to three years after planting, people cultivated rice, wheat and vegetables, etc.

The farmers in the northern region are also cultivating tree species along with their agricultural crops. They keep different tree species at a wide distance in their crop fields. The species used include *Acacia nilotica*, *A. catechu*, *Azadirachta indica*, *Eucalyptus spp.*, *Dalbergia sissoo*, *Borassus flabellifer*, *Phoenix sylvestris*, etc. In some areas, they have also planted mulberry for feeding silkworms to produce silk.

There are homestead agroforestry systems all over the country. These homesteads are rich in genetic resources. Ninety tree species and 13 bamboo species are grown in homestead forests. The Bangladesh Forest Research Institute (BFRI) has started research to develop improved rubber agroforestry systems in some rubber estates and trying to find out shade tolerant genotypes of economic crops. Ginger, turmeric and *Colocasia* (Arum) have given satisfactory results when planted at the early stage of rubber plantation.

For agroforestry purposes, farmers use different forest species for soil fertility management, fodder and living fences, as well as timber. Agroforestry utilizes forest genetics resources in the following ways:

- Selection of superior trees
- Selection of MPTs and fast-growing tree species
- Selection of leguminous tree species
- Selection of wild crops

4. Past and present activities in conservation, utilization and management of FGR

Establishment of forest plantations in Bangladesh started in 1871 with teak (*Tectona grandis*) using seeds brought from Myanmar. Since then plantation forestry became a part of the overall clearfelling silvicultural systems. Teak was the main species planted because of its high value. Other species, such as *Gmelina arborea*, *Artocarpus integrifolius*, *Dipterocarpus turbinatus*, *Swietenia mahagoni*, *Lagerstroemia speciosa*, *Toona ciliata*, *Artocarpus chaplasha*, *Xylia kerrii* and *Syzygium grande* were introduced later. Most of the plantations were monocultures established by shifting cultivators through taungya system. Since the plantations were established through clearfelling followed by artificial regeneration, there was a severe loss of native vegetation. Moreover, with the development of mechanized logging for commercial purposes, the shifting cultivators could not cope with the extensive area cleared, which resulted in a rapid loss of FGR. In many cases plantation establishment failed. The above-mentioned species used for plantation establishment were slow-growing and long-rotation species. Plantations with these species were unable to meet the growing needs of the rapidly increasing population.

In 1974, the Forest Department (FD) began to establish plantations with fast growing species such as *Gmelina arborea*, *Paraserianthes falcataria* and *Anthocephalus chinensis*. During this period, plantations of industrial species such as rubber, oil palm, mulberry and cashew were established but the results were disappointing.

Presently, industrial and fuelwood plantations comprise a mixture of species with long rotation (40 years), medium rotation (12-18 years) and short rotation (5-6 years). In addition, agroforestry plantations are being raised in encroached land with the active participation of local people on a benefit-sharing basis. Encroachment is a serious problem both in sal forests and Hill Forests.

Forest plantations cannot be the substitute to natural forests. The most important reason is that plantations are severely degraded in genetic resources compared to natural forests. In fact, the forests of Bangladesh, particularly the plain land sal forests and hill forests, are severely degraded due to indiscriminate exploitation, the sal forest are even on the verge of extinction. Therefore, the GOB has taken initiatives for the conservation of ecosystems and forest gene resources in the remaining natural forests.

4.1. Conservation strategies

Major international efforts to conserve FGR began in the 1960s with the guidance and support of FAO. Conservation efforts of FGR have been implemented with the following strategies.

4.1.1. *In situ* conservation

In situ conservation is carried out in the following areas: nature reserves, national parks, wildlife sanctuaries and world heritage sites.

- **Nature Reserves**

The objectives of a nature reserve are to protect communities and species and to maintain natural processes in order to have ecologically representative examples of the natural environment. However, in Bangladesh there are no nature reserves. There are some wilderness areas in Chittagong Hill Tracts but there are no records of these areas.

- **National Parks**

There are four national parks in Bangladesh. Details of the national parks are presented in Table 1, next page.

- **World Heritage Site**

The Sundarbans has been declared a world heritage site.

TABLE 1. NATIONAL PARKS IN BANGLADESH

Name & location	Area (ha)	Year est.	Purpose/Flora
Himchari National Park (Cox's Bazar)	1729	1980	To preserve the fauna and habitats as well as to provide facilities for research, education and recreation. / Semi-evergreen and moist deciduous forests. <i>Dipterocarpus</i> spp., <i>Albizia procera</i> , <i>Artocarpus chaplasha</i> , <i>Salmalia malabarca</i> , <i>Sterculia alata</i> , <i>Quercus</i> spp., <i>Castanopsis</i> spp., <i>Eugenia</i> spp., <i>Lannea grandis</i> , undergrowth mixed with canes, palms, orchids and ferns and sometimes as pure stands.
Bhawal National Park (Gazipur)	5022	1982	To preserve and develop the habitats and provide facilities for research, education and recreation. About 90 percent of the forest cover is composed of coppice-origin sal (<i>Shorea robusta</i>) in association with <i>Dillenia pentagyna</i> , <i>Lagerstroemia parviflora</i> , <i>Adina cordifolia</i> , <i>Miliusa velutina</i> , <i>Lannea grandis</i> , <i>Albizia</i> spp., <i>Bauhinia variegata</i> , <i>Spondias mangifera</i> , <i>Butea frondosa</i> , <i>Barringtonia acutangula</i> , etc. Among undergrowth, <i>Eupatorium</i> spp., <i>Pennisetum setosum</i> , <i>Asparagus racemosus</i> and <i>Rauvolfia serpentina</i> are common.
Madhupur National Park (Tangail)	8436	1982	To preserve and develop the habitats and provide facilities for research, education and recreation. Flora is the same as that of the Bhawal National Park. About 40% of the forest flora is composed of sal.
Ramsagar National Park (Dinajpur)	52	1974	To preserve and develop the habitats and provide facilities for research, education and recreation. The park does not contain any type of natural forest of the country but is artificially raised. The species planted consist mostly of fruit and ornamental trees.

- **Wildlife Sanctuaries**

There are 14 wildlife sanctuaries and game reserves in Bangladesh and details on these are as given in Table 2 below.

TABLE 2. WILDLIFE SANCTUARIES IN BANGLADESH (SARKER 1989; RAHMAN 1996)

Name & location	Area (ha)	Year est.	Purpose/Flora
Sundarbans World Heritage Site (Khulna)	16 065	1998	To preserve breeding habitats. The Sundarbans mangrove forest is fairly evergreen with low floristic diversity. The two main dominant species are <i>Heritiera fomes</i> and <i>Excoecaria agallocha</i> . Both species are gregarious in nature but the latter is more saline tolerant and therefore, predominates in the saline zones. Other species are <i>Rhizophora</i> spp., <i>Ceriops</i> spp., <i>Kandelia candel</i> , <i>Sonneratia apetala</i> , <i>Avicennia</i> spp., <i>Aegiceras corniculatum</i> , <i>Bruguiera gymnorrhiza</i> , <i>Phoenix paludosa</i> , <i>Nipa fruticans</i> , etc. and among grasses <i>Imperata cylindrica</i> , <i>Typha elephantina</i> and <i>Phragmites karka</i> are prominent.
Sundarbans East (Khulna)	5439	1977	To preserve the Royal Bengal Tiger (<i>Panthera tigris</i>) and its habitats. Flora is same as in the Sundarbans World Heritage Site.
Sundarbans South (Khulna)	17 878	1977	To preserve the Royal Bengal Tiger (<i>Panthera tigris</i>) and its habitats. Flora is same as in the Sundarbans World Heritage Site.
Sundarbans West (Khulna)	9069	1977	To preserve the Royal Bengal Tiger (<i>Panthera tigris</i>) and its habitats. Flora is same as in the Sundarbans World Heritage Site.
Rema-Kalenga (Sylhet)	1095	1981	To preserve the existing fauna and flora in the area. Flora is same as in tropical evergreen and semi-evergreen forests.

Name & location	Area (ha)	Year est.	Purpose/Flora
Char Kukri-Mukri (Bhola)	40	1981	To preserve the existing habitat used by local and migratory birds. The sanctuary area is covered very thickly by a small spiny plant locally called tambulkanta that grows up to 2.5 metres in height. There are patches of hogla (<i>Typha elephantina</i>), hargoja (<i>Acanthus illicifolius</i>) and keora (<i>Sonneratia apetala</i>) scatteredly present throughout the sanctuary area. Khalisha (<i>Aegiceras malus</i>) is also seen.
Pablakhali WS (Chittagong Hill Tracts)	42 087	1983	To preserve fauna and habitat for white-winged wood duck (<i>Cairina scutulata</i>). The following types of forest cover are recognized in the sanctuary area: a) Tropical evergreen; important species are <i>Pterygota alata</i> and <i>Quercus</i> spp. b) Tropical semi-evergreen; important trees are <i>Dipterocarpus</i> spp., <i>Mangifera</i> , <i>Amoora</i> , <i>Cinnamomum</i> , <i>Syzygium</i> , <i>Tetrameles</i> , <i>Artocarpus</i> , <i>Salmalia</i> , <i>Albizia</i> , etc. c) Tropical moist deciduous; the characteristic tree species are <i>Albizia</i> , <i>Salmalia</i> , <i>Terminalia</i> , <i>Ficus</i> spp., etc. Bamboos grow in all the types as undergrowth.
Hail Haor (Sylhet)	1427	1983	To preserve habitats for migratory ducks. At present all the wetland areas are under management control of Haor Development Board. Their activities probably place more emphasis on paddy production and other agricultural crops. It is the most threatened habitat in the country.
Hazarikhil (Chittagong)	2909	1970	To preserve local fauna and habitats. The forest cover of the sanctuary is of evergreen/semi-evergreen type. Main species of trees are <i>Dipterocarpus</i> spp., <i>Artocarpus chaplasha</i> , <i>Tetrameles nudiflora</i> , <i>Cedrela toona</i> , <i>Mesua ferrea</i> , <i>Eugenia</i> spp., <i>Ficus</i> spp., <i>Albizia procera</i> , etc. Undergrowth consists mostly of bamboo and <i>Eupatorium odoratum</i> .
Rampahar-Sitapahar (Rangamati)	3026	1973	To preserve local fauna and habitats. Same as Hazarikhil except bamboos occur as prominent undergrowth.
Chunati (Chittagong)	7764	1986	To preserve habitats. As in the case of mixed evergreen forests, the upper storey of the sanctuary forest consists of deciduous trees such as <i>Dipterocarpus turbinatus</i> with fewer <i>D. pilosus</i> and <i>D. costatus</i> , <i>A. Chaplasha</i> , <i>Salmalia insignis</i> and <i>S. malabarica</i> , <i>Ficus</i> spp., <i>Swintonia floribunda</i> , <i>Bandarholla</i> , etc. The second storey consists of evergreen species, such as <i>Quercus</i> spp., <i>Eugenia</i> spp., <i>Lannea</i> spp., <i>Lagerstroemia</i> spp., <i>Aphanomixis</i> spp., <i>Hargaza</i> , etc. The undergrowth consists mostly of smaller evergreen trees with bamboo and <i>assamlata</i> (<i>Eupatorium odoratum</i>).
Dulhazara Safari Park(Cox's Bazar)	1600	1995	To preserve habitats. Flora is same as in tropical semi-evergreen forests.
Khagrachari (Chittagong Hill Tracts)	n.a.	n.a.	To preserve elephants. Flora is same as in tropical wet evergreen forests.
Teknaf (Cox's Bazar)	11 615	1983	To preserve the Asian elephant (<i>Elphas maximus</i>) and its habitats. Semi-evergreen and moist deciduous forests. <i>Dipterocarpus</i> spp., <i>Albizia procera</i> , <i>Artocarpus chaplasha</i> , <i>Salmalia malabarica</i> , <i>Sterculia alata</i> , <i>Quercus</i> spp., <i>Castanopsis</i> spp. <i>Eugenia</i> spp., <i>Lannea grandis</i> , <i>Lagerstroemia</i> spp., <i>Amoora</i> spp. etc. Bamboo occurs as undergrowth mixed with canes, palms, orchids and ferns and sometimes as pure stands. This category does not fall in any of the ten categories described by the IUCN.

4.1.2. *Ex situ* conservation

In contrast to *in situ* conservation, *ex situ* conservation includes any practices that conserve genetic materials outside the natural habitat of the parent population. *Ex situ* conservation methods and materials include genebanks for seed or pollen as well as clone banks, arboreta, preservation plots, sample plots, etc:

- **Preservation plots**

The BFRI has established five preservation plots at different hill forest areas and 27 at the Sundarbans (mangrove) forest.

- **Clone banks**

The BFRI has established two clonal banks, one at Hayko, Chittagong (4 ha) and another at Ukhia, Cox's Bazar (4 ha). Seven tree species (*Tectona grandis*, *Gmelina arborea*, *Bombax ceiba*, *Dipterocarpus turbinatus*, *Syzygium grande*, *Swietenia mahagoni* and *Paraserianthes falcataria*) are preserved in these two locations.

- **Botanical gardens**

- Mirpur Botanical Garden: area 85 ha, with 255 tree species (total 28 200 plants), 310 shrub species (8400 plants), 385 herb species (10 400 plants). The total number of families of trees, herbs and shrubs is 114 (Ranjit 1997).
- Baldha garden: area 1.15 ha with 18 000 trees, herbs and shrubs from 820 species and 92 families (Ranjit 1997).

- **BFRI Bamboo Arboretum**

The BFRI Bambusetum (1.5 ha) has been established at the BFRI campus. This arboretum contains 27 bamboo species (*Bambusa balcooa*, *B. bambos* var. *spinosa*, *B. burmanica*, *B. cacharensis*, *B. comillensis*, *B. jaintiana*, *B. multiplex*, *B. nutans*, *B. polymorpha*, *B. salarkhanii*, *B. tulda*, *B. vulgaris*, *B. ventricosa*, *Dendrocalamus giganteus*, *D. hamiltonii*, *D. longispathus*, *D. strictus*, *D. brandisii*, *Gigantochloa andamanica*, *G. atroviolacea*, *G. apus*, *Melocalamus compactiflorus*, *Melocanna baccifera*, *Schizostachyum dullooa*, *Thyrsostachys oliveri*, *T. regis* and *T. siamensis*), including six exotic species. One arboretum of medicinal plants (1 ha) has been established at the BFRI campus with a collection of 40 species. One cane arboretum (0.5 ha) of seven species has also been established (Banik 1997). Three arboreta of tree species have been established at the BFRI HQ with 56 species, Keochia Forest Research Station with 56 species and Charaljani Silviculture Research Station with 52 species.

- **Seed storage**

There is a National Forest Seed Center (NFSC) at the BFRI; however, the centre does not have any facility for long time storage of seeds.

- **Tissue culture**

Tissue culture on forest tree species has been done only at the BFRI tissue culture laboratory. The BFRI has so far developed tissue culture techniques for six tree species and seven bamboo species.

Appendices 2-4 provide information on the conservation of important tree species in Bangladesh, their use and threats.

4.1.3. Conservation of provenances within a species

The BFRI has established provenance trials of *Acacia mangium*, *A. auriculiformis*, *Eucalyptus camaldulensis*, *E. brassiana*, *E. tereticornis*, *E. urophylla*, *Tectona grandis*, *Gmelina arborea*, *Pinus caribaea*, *P. oocarpa*, *Paraserianthes falcataria*, *Leucaena leucocephala*, *Melaleuca leucadendra*, *Gliricidia sepium* and *Populus deltoides* from 68 provenances.

5. Institutional framework

Roles of different institutions are crucial in guiding the course of events and ensuring the successful achievement of aims and objectives. The Forestry Master Plan considered five interrelated institutions, i.e. policy, legislation, organizational structure, human resource development, research and extension.

Today the forestry and forest institutions in Bangladesh are judged in much wider context than before. The interrelated and multiple roles of forests are vital for human welfare and sustained socioeconomic development.

Bangladesh signed the convention on the Earth Summit on 5 June 1992 in Rio and subsequently ratified it on 20 March 1994. As a result, the country has certain obligations under the convention.

5.1. National forest policy

In accordance with the National Forest Policy promulgated in October 1994, the following policy objectives are set in order to eliminate any uncertainty regarding the aims of the Government:

- To meet the basic needs of the present and future generations and also to ensure greater contribution of the forestry sector in economic development.
- To create employment opportunities, strengthening the rural and national economy; the scope for poverty alleviation and trees and forest-based rural development sectors will be extended and consolidated.
- Biodiversity of the existing degraded forests will be enriched by conserving the remaining natural habitats of birds and animals.
- Agricultural sector will be strengthened by conserving the land and water resources.
- National responsibilities and commitments will be fulfilled by implementing various international efforts and agreements ratified by the government relating to global warming, desertification and control of trade and commerce of wild birds and animals.
- Through the participation of the local people, illegal occupation of the forestlands, illegal tree felling and hunting of wild animals will be prevented.
- Effective use and utilization of the forest goods at various stages of processing will be encouraged.

- Implementation of afforestation programmes on both public and private lands will be provided with encouragement and assistance.

6. List of national priority species

The following species are the priority species in the Forest Department plantation programme:

Long rotation plantation species

Tectona grandis, *Dipterocarpus turbinatus*, *Syzygium grande*, *Swietenia macrophylla*, *Chukrasia tabularis*, *Michelia champaca*, *Hopea odorata*, *Xylia kerrii*, *Lagerstroemia flos-reginae*, *Shorea robusta* and *Toona ciliata*

Medium rotation species

In addition to the long rotation plantation species *Pinus caribaea*, *Albizia flacatana*, *Bombax ceiba*, *Gmelina arborea*, *Anthocephalus chinensis* and *Eucalyptus camaldulensis*, *E. tereticornis*, *Dalbergia sissoo*, *Azadirachta indica*, *Samanea saman*, *Bombax ceiba*, *Acacia nilotica* and *A. catechu*

Short rotation species

Acacia auriculiformis, *Acacia mangium*, *Eucalyptus camaldulensis*, *Melia azadirachta*, *Albizia chinensis*, *Leucaena leucocephala*, *Trewia nudiflora* and *Casuarina equisetifolia*

Village groves

Artocarpus heterophyllus, *Mangifera indica*, *Aegle mermelos*, *Litchi chinensis*, *Psidium guajava*, *Ziziphus spp.*, *Syzygium*, *Albizia*, *Barringtonia*, *Eucalyptus*, *Erythea*, *Ficus*, *Albizia fuman*, *Anthocephalus*, *Tamarindus indica*, *Bombax ceiba*, *Swietenia macrophylla*, *Alstonia scholaris*, *Cocos nucifera*, palmyra palm and bamboo

On marginal lands such as roadsides

Tectona grandis, *Mangifera indices*, *Artocarpus heterophyllus*, *Dalbergia sissoo*, *Butea frondosa*, *Polyanthina longifolia*, *Eucalyptus camaldulensis*, *Acacia auriculiformis*, *Swietenia*, *Albizia*, *Samanea*, *Syzygium* and *Casuarina equisetifolia*

Multipurpose tree species for different zones:

Hillzones

Albizia lebbeck, *A. procera*, *Phyllanthus emblica*, *Eucalyptus camaldulensis*, *Elaeocarpus robusta*, *Artocarpus heterophyllus*, *Acacia auriculiformis* and rattans

Coastal zone

Casuarina equisetifolia, *Albizia lebbeck*, *Acacia procera*, *S. grandiflora*, *Cocos nucifera*, *Phonek sylvestria* and *Erythrina indica*

Mangrove

Heritiera fomes, *Avicennia sp.*, *Bruguiera gymnorrhiza*, *Ceriops decandra*, *Rhizophora mucronata* and *Sonneratia apetala*

7. Research on FGR

The BFRI conducts research under 12 programme areas. Each year the Institute undertakes a number of priority research studies following the suggestions of the Bangladesh Forest Department, Bangladesh Forest Industrial Development Corporation, Bangladesh Chemical Industries Corporation, Bangladesh Tea Board, Rural Electrification Board, other wood-based industries, private owners and non-government organizations. The suggested studies are scrutinized by a Technical Committee. The selected studies are approved by an Advisory Committee.

The BFRI conducts a number studies on FGR conservation and management under the following programme areas: (i) Biodiversity and its conservation, (ii) Production of quality planting materials, (iii) Plantation techniques and forest management, (iv) Breeding and improvement, (v) Social and non-timber forest products, (v) Social forestry and farming system research and (vi) Pest and diseases. Seed Orchard Division, Silviculture Research Division, Silviculture Genetics Division, Mangrove Silviculture Division, Plantation Trial Unit Division, Minor Forest Products Division, Soil Science Division, Forest Protection Division and Farming System Research Component are involved in conducting these studies.

A total of 44 technologies have been developed and out of these, 28 technologies have been transferred to end-users. A total of 16 technologies on conservation and management of FGR have been developed and transferred to different end-users. Training programmes based on these new technologies are arranged as and when required.

8. Conclusion and recommendations

The natural forests of Bangladesh have been seriously degraded, resulting in serious genetic erosion of FGR. There is a critical need to develop coordinated efforts to conserve and manage FGR. Effective and hopeful efforts have been developed into conservation activities, but national and international financial and technical assistance are needed to bring about success. The following recommendations have been put forward for the conservation and sustainable utilization of FGR in Bangladesh:

- Development of a database on the present status of flora and fauna in different ecosystems of Bangladesh.
- *In situ* and *ex situ* programmes to conserve, manage and use FGR should be significantly expanded.
- Community-based resource conservation needs to be emphasized.
- Improved silvicultural methods should be applied in the management of natural and plantation forests.
- The method of clear-felling followed by burning for plantation establishment must be stopped.
- Silvicultural measures for aided natural regeneration should be followed.
- Enrichment planting should be conducted in the forest gaps with diversified genetic resources collected from natural regeneration in the forest floor.
- Establishment of preservation plots and permanent sample plots in the reserved forest.
- Establishment of a genebank for conservation of FGR.

- Logging in the remaining natural forests must be stopped.
- Creation of diversified job opportunities for the hill people through Farming System approach.
- Improvement of shifting cultivation by improved technologies like SALT, DSA, etc.
- Introduction of a forest certification system for sustainable forest resources management.
- Awareness should be developed among the shifting cultivators about the detrimental effect of shifting cultivation.
- Education and training of professionals and technicians should be given to equip them with the latest knowledge of forest genetic resource survey, management and conservation.
- Strengthening the international cooperation for FGR conservation.

References

- Alam, M.K.** 1982. A guide to eighteen species of bamboo. Bulletin 2. Plant Taxonomy Series, Bangladesh Forest Research Institute, Chittagong, Bangladesh.
- Alam, M.K. and Basu, S.K.** 1988. On the occurrence of *Calamus longisetus* in Bangladesh. Bana Biggyan Patrika, Vol. 17 (1,2). Bangladesh Forest Research Institute, Chittagong, Bangladesh.
- Ali, M.O.** 1989. Forestry Research in Bangladesh: Problems and prospects BARC, Farmgate, Dhaka, Bangladesh.
- Anon.** 1992. Forestry Master Plan (FMP) of Bangladesh. Ministry of Environment and Forest, Government of Bangladesh.
- Banik, R.L.** 1997. Bamboo resources of Bangladesh. Pp. 183-207 in *Agroforestry: Bangladesh perspective* (M.K. Alam, F.U. Abrned and S.M.R. Amin, eds.). APAN-NAWG-BARC. Dhaka.
- BBS.** 1994. Statistical Year Book of Bangladesh. Twelve edition, Bangladesh Statistics Division, Ministry of Planning, Dhaka. 628 pp.
- Byron, N.** 1984. Peoples' forestry - a novel perspective of forestry Bangladesh. ADAB News, Dhaka. 11:31-37.
- Douglas, J.J.** 1982. Consumption and supply of wood and bamboo in Bangladesh. Field document no. 2. UNDP/FAO Project. BGD/78/010, Planning Commission, Dhaka. 162p.
- GOB (Government of Bangladesh)** 1992. The Government of Bangladesh and the International Union for Conservation of Nature and Natural Resources. Towards sustainable development, national conservation strategy of Bangladesh. 214p.
- Hassan, M.M.** 1995. Biodiversity Conservation and Management in Bangladesh: A state of the art, Review. ICIMOD, Kathmandu, Nepal.
- Huq, M.F. and Banik, R.L.** 1992. Country Report for Bangladesh. Pp. 19-48 in *Proceedings of the Regional Workshop on Tree Breeding and Propagation, held in Bangkok, Thailand 10-14 July 1990*. Field document no. 2 (RAS/88/025).
- Khan, M.S.** 1977. Flora of Bangladesh. Report 4. Camelinaceae. Bangladesh National Herbarium, Bangladesh Agriculture Research Council (BARC), Farmgate, Dhaka.
- Khan, M.S.** 1991. National Conservation Strategy of Bangladesh - Towards sustainable development: Genetic Resources in Bangladesh. Bangladesh Agricultural Research Council, IUCN, Dhaka. Pp. 162-172.
- Khan, M.S.** 1996. National Obligations in the Implementation of the Conservation on Biological diversity: Bangladesh perspective BARC, IUCN, the World Conservation Union, Dhaka 1991. Pp. 162-172.
- Kibria, M.G., Sarker, D.C., Hossain, M.A.T, Manna, M.A., Motaleb, M.A. and Islam, S.S.** 2000. Forest Statistics of Bangladesh, Bulletin 4, Forest Economics Division, Bangladesh Forest Research Institute, Chittagong, Bangladesh. 119p.
- Rahman, M.K.** 1996. Bangladesh Protected areas: Wildlife. Tiger paper 23(3):9-13.
- Sarker, N.M.** 1989. Protected areas of Bangladesh - an overview of present management status. Country report. Proceedings of the First International Workshop Cum Seminar for Conservation of Wildlife in Bangladesh.
- TFYP.** 1985. Planning Commission, Ministry of Planning, Dhaka. Pp. 57-65.
- Troup, R.S.** 1975. Silviculture of Indian Trees (Revised edition). Forest Research Institute Press, Dehradun, India.

APPENDICES

Appendix 1. Value and use of main forest species in Bangladesh

Species name	Value code	Present, future or potential use											
		ti	po	wo	nw	pu	fo	fd	sh	ag	so	am	xx**
1. <i>Acacia auriculiformis</i>	1	+		+		+		+	+	+			
2. <i>A. catechu</i>	1	+	+	+	+			+					
3. <i>A. farnesiana</i>	3			+	+			+					
4. <i>A. mangium</i>	1	+		+		+		+		+			pb,v
5. <i>A. nilotica</i>	1	+		+	+			+					cw,agi
6. <i>Acrocarpus fraxinifolius</i>	3	+											pw
7. <i>Aegiceras corniculatum</i>	3			+	+		+						
8. <i>Albizia chinensis</i>	1	+				+		+	+				pw
9. <i>A. lebbeck</i>	1	+		+			+	+					v
10. <i>A. lucidor</i>	3	+		+				+					
11. <i>A. odoratissima</i>	2	+		+				+				+	
12. <i>A. richardiana</i>	2	+			+		+					+	bb
13. <i>A. procera</i>	1	+		+				+					
14. <i>Alstonia scholaris</i>	1	+			+	+						+	pc,n
15. <i>Amoora cucullata</i>	3		+	+									
16. <i>Anacardium occidentale</i>	1	+			+		+						
17. <i>Anogeissus acuminata</i>	3	+		+	+							+	agi
18. <i>Anthocephalus chinensis</i>	1					+	+					+	
19. <i>Antidesma ghaesembilla</i>	3	+					+	+					
20. <i>Aphanamix polystachya</i>	2	+			+								
21. <i>Aporosa dioica</i>	3		+		+		+						
22. <i>Aquilaria agallocha</i>	2												ab
23. <i>Artocarpus chama</i>	1	+				+	+						
24. <i>A. heterophyllus</i>	1	+			+		+	+		+		+	
25. <i>Artocarpus lacucha</i>	2	+			+		+	+					agi
26. <i>Avicennia alba</i>	2			+									pw
27. <i>Azadirachta indica</i>	1				+					+		+	
28. <i>Bambusa sp.</i>	1			+		+	+	+					pb
29. <i>Barringtonia acutangula</i>	2	+		+	+								
30. <i>Bauhinia purpurea</i>	3				+		+	+					agi
31. <i>Bauhinia variegata</i>	3				+		+	+					
32. <i>Beilschmiedia pseudomicrocarpa</i>	3	+											
33. <i>Berrya cordifolia</i>	3	+										+	agi,cw
34. <i>Bischofia javanica</i>	2	+	+		+			+					agi
35. <i>Bombax ceiba</i>	1	+				+	+	+					k
36. <i>Bridelia retusa</i>	3		+		+		+	+					
37. <i>Bouea oppositifolia</i>	3	+					+						agi
38. <i>Bruguiera gymnorhiza</i>	2	+	+		+								
39. <i>B. sexangula</i>	3	+	+		+								
40. <i>Buchanania lanzan</i>	3				+		+						
41. <i>Butea monosperma</i>	2	+			+			+					
42. <i>Callicarpa tomentosa</i>	3		+	+				+					

Species name	Value code	Present, future or potential use											
		ti	po	wo	nw	pu	fo	fd	sh	ag	so	am	xx**
43. <i>Calophyllum inophyllum</i>	2				+								m,bb,rs
44. <i>Canarium resiniferum</i>	3				+								v,pw
45. <i>Cassia fistula</i>	2	+			+			+				+	c
46. <i>C. nodosa</i>	2	+										+	
47. <i>C. siamea</i>	1	+		+				+				+	c
48. <i>Castanopsis tribuloides</i>	3	+						+					agi
49. <i>Casuarina cunninghamiana</i>	3								+			+	
50. <i>C. equisetifolia</i>	1	+		+	+					+		+	bb,m
51. <i>Ceiba pentandra</i>	1	+			+			+					k
52. <i>Ceriops decandra</i>	3	+	+		+								
53. <i>Chukrasia velutina</i>	1	+								+			
54. <i>Cinnamomum iners</i>	3												p
55. <i>Cordia dichotoma</i>	3	+		+	+		+	+					
56. <i>Crateva magna</i>	3	+			+		+						t,n
57. <i>Croton oblongifolius</i>	3	+		+	+								
58. <i>Crypteronia paniculata</i>	3												cw,c,rs
59. <i>Dalbergia sissoo</i>	1	+			+			+					a,v
60. <i>Delonix regia</i>	2	+		+					+			+	
61. <i>Dillenia indica</i>	3		+	+	+		+						bb
62. <i>D. scabrella</i>	3	+					+						
63. <i>Diospyros montana</i>	3	+			+								
64. <i>D. nigricans</i>	3		+										
65. <i>D. peregrina</i>	3	+			+		+						ts
66. <i>D. toposia</i>	3	+					+						
67. <i>Dipterocarpus alatus</i>	2												bb,rs
68. <i>D. turbinatus</i>	1									+			bb,c
69. <i>Dolichandrone spathacea</i>	3											+	
70. <i>Duabanga grandiflora</i>	3	+				+							pw,d
71. <i>Dysoxylum binectariferum</i>	3	+			+								
72. <i>D. hamiltonii</i>	3	+			+		+						
73. <i>Ehretia serrata</i>	3	+					+	+					agi
74. <i>Elaeocarpus floribundus</i>	1	+					+						
75. <i>E. sphaericus</i>	3											+	
76. <i>E. varunna</i>	3									+			tc
77. <i>Engelhardtia spicata</i>	3	+			+								
78. <i>Erioglossum rubiginosum</i>	3	+					+						c
79. <i>Erythrina fusca</i>	3			+				+	+				
80. <i>E. variegata</i>	2			+				+	+				d,v,t
81. <i>E. fusea</i>	3												
82. <i>Eucalyptus alba</i>	3				+							+	
83. <i>E. camaldulensis</i>	1			+			+						c,v,rs,pw

Species name	Value code	Present, future or potential use											
		ti	po	wo	nw	pu	fo	fd	sh	ag	so	am	xx**
84. <i>E. brassina</i>	2			+			+						
85. <i>E. tereticornis</i>	2			+			+						
86. <i>E. urophylla</i>	2			+			+						
87. <i>E. citriodora</i>	2	+			+								c, agi
88. <i>Excoecaria agallocha</i>	1					+							
89. <i>Ficus benghalensis</i>	1	+		+	+			+					
90. <i>F. hispida</i>	2			+			+	+					
91. <i>F. racemosa</i>	2	+		+	+		+	+					
92. <i>F. religiosa</i>	1			+				+				+	pc
93. <i>Flacourtia jangomas</i>	3	+			+		+						
94. <i>Garcinia cowa</i>	2	+			+		+						va
95. <i>Garuga pinnata</i>	2	+			+	+	+	+	+				
96. <i>Gliricidia sepium</i>	2			+				+	+				
97. <i>Gmelina arborea</i>	1	+			+	+		+		+			
98. <i>Grevillea robusta</i>	3	+											v,d
99. <i>Grewia tiliaefolia</i>	3						+						agi,m
99. <i>Heritiera fomes</i>	1		+										bb,c
100. <i>Hevea brasiliensis</i>	1	+				+							v,r
101. <i>Holarrhena pubescence</i>	3	+			+								n
102. <i>Hopea odorata</i>	1	+	+		+					+			
103. <i>Hydnocarpus kurzii</i>	2				+								
104. <i>Hymenodictyon orixensis</i>	2	+			+			+					tc,m
105. <i>Jacaranda mimosifolia</i>	3				+							+	
107. <i>Kandelia candel</i>	3				+								
108. <i>Lagerstroemia macrocarpa</i>	2		+										c,bb
109. <i>L. speciosa</i>	1	+	+									+	bb,c
110. <i>Lannea coromandelica</i>	3	+			+	+		+					
111. <i>Leucaena leucocephala</i>	3			+				+		+			
112. <i>Lithocarpus elegans</i>	2		+	+									
113. <i>L. pachyphylla</i>	3			+									c
114. <i>Litsea glutinosa</i>	3	+		+	+								
115. <i>L. monopetala</i>	3	+			+			+				+	
116. <i>Madhuca indica</i>	3	+			+		+						
117. <i>Mallotus philippensis</i>	3				+								
118. <i>Mangifera indica</i>	1	+			+		+	+		+			
119. <i>Melaleuca leucadendra</i>	3		+	+	+	+	+						rs,bb
120. <i>Michelia champaca</i>	1	+			+		+			+		+	
121. <i>Olea dioica</i>	3			+									
122. <i>Oroxylum indicum</i>	3			+	+		+	+					
123. <i>Paraserianthes falcataria</i>	2					+							pw,m
124. <i>Phyllanthus emblica</i>	1				+		+			+			
125. <i>Pinus caribaea</i>	2					+						+	

Species name	Value code	Present, future or potential use											
		ti	po	wo	nw	pu	fo	fd	sh	ag	so	am	xx**
126. <i>Pithecellobium dulce</i>	2	+		+	+		+	+					
127. <i>Pongamia pinnata</i>	2				+		+						
128. <i>Prosopis juliflora</i>	3	+		+				+					
129. <i>Protium serratum</i>	3						+						c,rs
130. <i>Pterospermum acerifolium</i>	2	+			+			+				+	agi
131. <i>Pterygota alata</i>	2	+					+	+					
132. <i>Rhizophora mucronata</i>	1		+	+	+		+						
133. <i>Samanea saman</i>	1	+		+				+				+	
134. <i>Saraca asoca</i>	2				+							+	
135. <i>Schima wallichii</i>	2			+									c
136. <i>Schleichera oleosa</i>	2				+		+						
137. <i>Sesbania grandiflora</i>	1			+		+	+	+		+			
138. <i>Semecarpus anacardium</i>	3				+		+						
139. <i>Shorea robusta</i>	1	+			+								c
140. <i>Sonneratia apetala</i>	1			+			+	+					
141. <i>Sterculia villosa</i>	2	+			+		+						
142. <i>Stereospermum suaveolens</i>	3	+		+									
143. <i>Swietenia mahagoni</i>	1	+										+	
144. <i>Syzygium grandis</i>	2		+							+			
145. <i>Tamarindus indica</i>	1	+		+	+		+	+					
146. <i>Tamarix dioica</i>	3			+							+		
147. <i>Tectona grandis</i>	1	+			+								c,bb
148. <i>Terminalia bellirica</i>	1	+			+		+	+					
149. <i>T. catappa</i>	2				+		+						
150. <i>T. chebula</i>	1				+		+						
151. <i>Toona ciliata</i>	1	+											bb
152. <i>Trema orientalis</i>	3	+		+				+					
153. <i>Vatica lanceaefolia</i>	3				+								rs
154. <i>Xanthophyllum flavescens</i>	3	+					+						
155. <i>Xylia kerrii</i>	2		+										c
156. <i>Xylocarpus granatum</i>	2				+								
157. <i>Zanthoxylum rhetsa</i>	3	+			+		+					+	
158. <i>Ziziphus mauritiana</i>	1	+		+			+	+					agi

VALUE: 1 = Species of current socioeconomic importance; 2= Species with clear potential of future value; 3 = Species of unknown value given present knowledge and technology

UTILIZATION: ti = timber production; po = posts, poles, roundwood; wo = fuelwood, charcoal; nw = non-wood products (gums, resins, oils, tannins, medicines, dyes, etc.); pu = pulp and paper; fo = food; fd = fodder; sh = shade, shelter; ag = agroforestry systems; so = soil and water conservators; am = amenity, antithetic, ethical values; **xx** other:** v = veneer; pw = plywood; cw = cartwheel; pb = particle board; c = construction work; k = Kapok; irs = railway sleepers; mb = mast of boat; p = planking; t = toys; n = novelties; d = decorative; tc = tea chest; va = varnish; agi = agricultural implements; bb = boat building; pc = packing cages; ab = agar batti; m = match splints and boxes; r = rubber

Appendix 2. Conservation and management of important FGR by ecogeographic zone in Bangladesh

Species in ecogeographic (or genealogical) zones	Nature reserves, protected areas	<i>In situ</i> conservation stands	Managed forests	Unmanaged forests	Plantations	<i>Ex situ</i> conservation stands	Villages, fields, homesteads	Experiment, fields, trials
Hill Forest (Chittagong, Chittagong Hill Tracts, Sylhet):								
<i>Artocarpus chaplasha</i>	+	+	+		+			+
<i>Swintonia floribunda</i>	+	+	+					+
<i>Dipterocarpus turbinatus</i>	+	+	+		+			+
<i>D. pilosus</i>	+							
<i>D. costatus</i>	+							
<i>D. gracilis</i>	+							
<i>Mesua ferrea</i>							+	
<i>Hopea odorata</i>	+	+	+		+		+	+
<i>Syzygium</i> spp.		+	+		+			+
<i>Calophyllum</i> spp.	+							
<i>Palaquium</i> spp.	+							
<i>Chukrasia tabularis</i>	+	+	+		+			+
<i>Ficus</i> spp.	+							
<i>Michelia champaca</i>	+	+			+		+	+
<i>Pterygota alata</i>	+							
<i>Lophopetalum fimbriatum</i>	+							
<i>Amoora</i> spp.	+							
<i>Dysoxylum</i> spp.	+							
<i>Albizia procera</i>	+	+			+		+	+
<i>A. lebbeck</i>	+	+			+		+	+
<i>A. chinensis</i>	+	+			+		+	+
<i>Gmelina arborea</i>	+	+			+		+	+
<i>Alstonia scholaris</i>	+	+						
<i>Toona ciliata</i>	+	+					+	+
<i>Quercus semiserrata</i>	+							
<i>Q. gomeziana</i>	+							
<i>Podocarpus neriloforcis</i>	+							

Species in ecogeographic (or genealogical) zones	Nature reserves, protected areas	<i>In situ</i> conservation stands	Managed forests	Unmanag ed forests	Plantations	<i>Ex situ</i> conservation stands	Villages, fields, homesteads	Experiment, fields, trials
<i>Cassia fistula</i>	+							
<i>Phyllanthus emblica</i>	+						+	+
<i>Tetrameles nudiflora</i>	+							
<i>Bombax insigne</i>	+							
<i>B. ceiba</i>	+						+	
<i>Duabanga grandiflora</i>	+							
<i>Lithocarpus elegans</i>	+							
<i>Castanopsis tribuloides</i>	+							
<i>Calophyllum polyanthum</i>	+							
<i>Macaranga spp.</i>	+							
<i>Terminalia bellirica</i>	+							
<i>Pterospermum acerifolium</i>	+							
<i>Diospyros embryopteris</i>	+							
<i>Sterculia villosa</i>	+							
<i>Garuga pinnata</i>	+							
<i>Meliosma pinnata</i>	+							
<i>Callicarpa macrophylla</i>	+							
<i>Vitex glabrata</i>	+							
<i>Saraca indica</i>	+							
<i>Elaeocarpus robustus</i>	+							
<i>Lagerstroemia spp.</i>	+					+		+
<i>Mitragyna parvifolia</i>	+							
<i>Calamus guruba</i>	+					+		+
<i>C. viminalis</i>	+					+		+
<i>C. latifolius</i>	+					+		+
<i>Daemonorops jenkinsanus</i>	+					+		+
<i>Melocanna baccifera</i>	+					+		+
<i>Dendrocalamus longispathus</i>	+							+
<i>D. hamiltonii</i>	+							+
<i>Neohouzeaua dullooa</i>	+							+

Species in ecogeographic (or genealogical) zones	Nature reserves, protected areas	<i>In situ</i> conservation stands	Managed forests	Unmanaged forests	Plantations	<i>Ex situ</i> conservation stands	Villages, fields, homesteads	Experiment, fields, trials
<i>Bambusa tulda</i>	+							+
<i>B. polymorpha</i>	+							+
<i>Melocalamus compactiflorus</i>	+							+
<i>Oxytenanthera nigrocalata</i>	+							+
<i>B. vulgaris</i>				+			+	+
Plainland Sal Forest (Comilla, Dhaka, Dinajpur):								
<i>Shorea robusta</i>	+							+
<i>Terminalia bellirica</i>	+							
<i>T. chebula</i>	+							
<i>Miliusa velutina</i>	+							
<i>Albizia procera</i>	+				+			+
<i>Dillenia pentagyna</i>	+				+			+
<i>Lagerstroemia spp.</i>	+							
<i>Garuga spp.</i>	+							
<i>Cassia fistula</i>	+							
<i>Phyllanthus emblica</i>	+				+			+
<i>Adina cordifolia</i>	+							
<i>Butea monosperma</i>	+							
<i>Careya arborea</i>	+							
<i>Schleichera oleosa</i>	+							
<i>Sterculia spp.</i>	+							
<i>Semecarpus anacardium</i>	+							
<i>Litsea polyantha</i>	+							
<i>Aphanamixis polystachya</i>	+							
<i>Microcos paniculata</i>	+							

Species in ecogeographic (or genealogical) zones	Nature reserves, protected areas	<i>In situ</i> conservation stands	Managed forests	Unmanag ed forests	Plantations	<i>Ex situ</i> conservation stands	Villages, fields, homesteads	Experiment, fields, trials
Littoral and Swamp Forest								
<i>Casuarina equisetifolia</i>					+			+
<i>Calophyllum inophyllum</i>	+							
<i>Terminalia catappa</i>	+						+	
<i>Erythrina variegata</i>	+							
<i>Barringtonia spp.</i>	+							
<i>Hibiscus tiliaceus</i>	+							
<i>Thespesia populnea</i>	+							
<i>Vitex negundo</i>	+							
<i>Trewia nudiflora</i>	+							
<i>Dolichandrone spathacea</i>	+							
Mangrove Forest (Sundarban and Coastal Forest):								
<i>Heritiera fomes</i>	+				+			+
<i>Excoecaria agallocha</i>	+				+			+
<i>Sonneratia apetala</i>	+				+			+
<i>Avicennia officinalis</i>	+				+			+
<i>Xylocarpus granatum</i>	+				+			+
<i>Nipa fruticans</i>	+				+			+

+ = Available

Appendix 3. Level and nature of threats to the integrity of species/populations of important tree species (Huq and Banik 1992; Khan 1991)

Species in ecogeographic (or genealogical) zones	Nature reserves, prot. areas	<i>In situ</i> conservati on stands	Managed forests	Un- managed forests	Plantations	<i>Ex situ</i> conservation stands	Villages, fields, homesteads	Experiment fields, trials	Degree of threat index
1. <i>Aglaonema clarkei</i>	+								
2. <i>Aldrovanda vesiculosa</i>	+								
3. <i>Aquilaria agallocha</i>	+								
4. <i>Cirrhopetalum roxburghii</i>	+								
5. <i>Cymbopogon osmastonil</i>	+								
6. <i>Debregeasia dentata</i>	+								
7. <i>Elaeocarpus lucidus</i>	+								
8. <i>Hippocratea marcantha</i>	+								
9. <i>Homalium schlichil</i>	+								
10. <i>Justicia orephylla</i>	+								
11. <i>Knema benghalensis</i>	+								
12. <i>Limnophila cana</i> (endemic)	+								
13. <i>Mantisia spathutala</i> (endemic)	+								
14. <i>Marsdenia thyrsoflora</i>	+								
15. <i>Ophiorrhiza villosa</i>	+								
16. <i>Phrynium imbricum</i>	+								
17. <i>Quercus acuminata</i>	+								
18. <i>Rotala simpliciuscula</i> (endemic)	+								
19. <i>Semecarpus</i> <i>sulepanduriformis</i> (endemic)	+								
20. <i>Sonneratia griffithi</i>	+								
21. <i>Spatholobus listeri</i> (endemic)	+								
22. <i>Tournefortia roxburghii</i>	+								

Species in ecogeographic (or genealogical) zones	Nature reserves, prot. areas	<i>In situ</i> conservati on stands	Managed forests	Unmana ged forests	Plantations	<i>Ex situ</i> conservation stands	Villages, fields, homesteads	Experiment fields, trials	Degree of threat index
23. <i>Typhonium listeu</i> (endemic)	+								
24. <i>Vatica scaphula</i> (endemic)	+								
25. <i>Vernonia thomsonii</i>	+								
26. <i>Adina cordifolia</i>	+								
27. <i>Aphanamixis polystachya</i>	+								
28. <i>Bassia latifolia</i>	+								
29. <i>Bauhinia malabarica</i>	+								
30. <i>Castanopsis tribuloides</i>	+								
31. <i>Derris robusta</i>	+								
32. <i>Diospyros cordifolia</i>	+								
33. <i>Hydnocarpus kurzii</i>	+								
34. <i>Lophopetalum fimbriatum</i>	+								
35. <i>Mesua ferrea</i>	+								
36. <i>Mitragyna parvifolia</i>	+								
37. <i>Podocarpus neriifolius</i>	+								
38. <i>Pterospermum acerifolium</i>	+								
39. <i>Pterygota alata</i>	+								
40. <i>Schleichera oleosa</i>	+								
41. <i>Sterculia foetida</i>	+								
42. <i>Swintonia floribunda</i>	+								
43. <i>Tamarindus indica</i>	+								

+ = Available

Appendix 4. List of priority species for conservation, improvement or seed procurement, their uses and conservation activities needed

Species	End use				Operations / activities needed								REMARKS
	W	NW	FW	O	Exploration & collection		Evaluation		Conservation		Germplasm use		
	1	2	3	4	5	6	7	8	9	10	11	12	
1. <i>Acacia auriculiformis</i>	+		+					+		+		PVT	Narrow Genetic Base (NGB)
2. <i>A. catechu</i>		+		+					+	+			
3. <i>A. mangium</i>	+		+					+		+		PVT	NGB
4. <i>A. nilotica</i>	+	+	+	+					+	+			
5. <i>Albizia chinensis</i>	+		+										
6. <i>A. lebeck</i>	+												
7. <i>A. procera</i>	+												
8. <i>Alstonia scholaris</i>				+									
9. <i>Anacardium occidentale</i>				+									
10. <i>Anthocephalus chinensis</i>	+			+									
11. <i>Artocarpus heterophyllus</i>	+		+	+					+				MPTS
12. <i>Azadirachta indica</i>	+	+	+	+					+				MPTS
13. <i>Bombax ceiba</i>				+									
14. <i>Cassia siamea</i>			+										
15. <i>Casuarina equisetifolia</i>			+										
16. <i>Ceiba pentandra</i>				+									
17. <i>Chukrasia velutina</i>	+		+										
18. <i>Dalbergia sissoo</i>	+		+	+									NGB
19. <i>Dipterocarpus turbinatus</i>	+	+											
20. <i>Elaeocarpus floribundus</i>				+									
21. <i>Eucalyptus alba</i>	+		+										

Species	End use				Operations / activities needed								REMARKS
	W	NW	FW	O	Exploration & collection		Evaluation		Conservation		Germplasm use		
	1	2	3	4	5	6	7	8	9	10	11	12	
22. <i>E. camaldulensis</i>	+		+										
23. <i>E. brassina</i>	+		+									PVT	NGB
24. <i>E. tereticornis</i>	+		+									PVT	NGB
25. <i>E. urophylla</i>	+		+									PVT	NGB
26. <i>E. citriodora</i>	+		+									PVT	NGB
27. <i>Excoecaria agallocha</i>	+		+	+									
28. <i>Ficus benghalensis</i>			+	+									
29. <i>F. religiosa</i>			+	+									
30. <i>Gliricidia sepium</i>			+	+									
31. <i>Gmelina arborea</i>	+		+	+									
32. <i>Heritiera fomes</i>	+		+										
33. <i>Hevea brasiliensis</i>	+	+	+										
34. <i>Hopea odorata</i>	+		+										
35. <i>Hydnocarpus kurzii</i>			+	+									
36. <i>Lagerstroemia speciosa</i>	+		+										
37. <i>Leucaena leucocephala</i>			+	+									
38. <i>Madhuca indica</i>	+	+	+	+									
39. <i>Mangifera indica</i>	+		+	+									
40. <i>Michelia champaca</i>	+		+										
41. <i>Paraserianthes falcataria</i>	+		+										

Species	End use				Operations / activities needed								REMARKS
	W	NW	FW	O	Exploration & collection		Evaluation		Conservation		Germplasms use		
	1	2	3	4	5	6	7	8	9	10	11	12	
42. <i>Phyllanthus emblica</i>			+	+									
43. <i>Pinus caribaea</i>	+	+	+										
44. <i>Rhizophora mucronata</i>	+		+										
45. <i>Samanea saman</i>	+		+	+									
46. <i>Saraca asoca</i>			+	+									
47. <i>Sesbania grandiflora</i>			+	+									
48. <i>Shorea robusta</i>	+		+										
49. <i>Sonneratia apetala</i>	+		+										
50. <i>Swietenia mahagoni</i>	+		+										
51. <i>Syzygium grande</i>	+		+										
52. <i>Tamarindus indica</i>			+	+									
53. <i>Tectona grandis</i>	+												
54. <i>Terminalia bellirica</i>	+		+	+									
55. <i>T. chebula</i>	+		+	+									
56. <i>Toona ciliata</i>	+		+										
57. <i>Xylia kerrii</i>	+												
58. <i>Xylocarpus granatum</i>	+		+										
59. <i>Ziziphus mauritiana</i>			+	+									

End uses: **1** = Industrial wood products (logs, sawtimber, construction wood, plywood, chip and particle board, wood pulp etc.); **2** = Industrial non-wood products (gums, resin, oils, tannins); **3** = Fuelwood, posts, poles (firewood, charcoal, roundwood used on-farm, wood for carving); **4** = Other uses, goods and services (food, medicinal use, fodder, land stabilization/amelioration, shade, shelter, environmental values).

Exploration & collection: **5** = Biological information (natural distribution, taxonomy, genecology, phenology etc.); **6** = Collection of germplasm for evaluation

Evaluation: **7** = *In situ* (population studies); **8** = *Ex situ* (provenance and progeny tests)

Conservation: **9** = *In situ*; **10** = *Ex situ*

Reproductive use/germplasm use: **11** = Semi-bulk/bulk seedlots, reproductive materials; **12** = Selection and improvement

Remarks (13): PVT = provenance trials; E = endangered at species or provenance level; PGT = progeny trials; MPTS = multi-purpose tree species; CLT = clonal trials; SO = seed orchard; NGB = narrow genetic base