ISSN 1726-5274



Food and Agriculture Organization of the United Nations



# Measures for the control of *Campylobacter* spp. in chicken meat

### Meeting report





MICROBIOLOGICAL RISK ASSESSMENT SERIES



MICROBIOLOGICAL RISK ASSESSMENT SERIES

# Measures for the control of *Campylobacter* spp. in chicken meat

Meeting report

Food and Agriculture Organization of the United Nations World Health Organization

Rome, 2024

Required citation:

FAO & WHO. 2024. *Measures for the control of* Campylobacter *spp. in chicken meat – Meeting report.* Microbiological Risk Assessment Series, No. 46. Rome. https://doi.org/10.4060/cc9607en

This publication contains the collective views of an international group of experts and does not necessarily represent the decisions or the policies of FAO or WHO. The expert group members alone are responsible for the views expressed in this publication and they do not necessarily represent the views, decisions or policies of the institutions with which they are affiliated.

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) or the World Health Organization (WHO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO or WHO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO or WHO.

ISSN 1726-5274 [Print] ISSN 1728-0605 [Online]

ISBN [FAO] 978-92-5-138584-5 ISBN [WHO] 978-92-4-008808-5 (electronic version) ISBN [WHO] 978-92-4-008809-2 (print version) © FAO and WHO, 2024



Some rights reserved. This work is made available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; https://creativecommons.org/licenses/by-nc-sa/3.0/igo/legalcode).

Under the terms of this licence, this work may be copied, redistributed and adapted for non-commercial purposes, provided that the work is appropriately cited. In any use of this work, there should be no suggestion that FAO or WHO endorses any specific organization, products or services. The use of the FAO or WHO logo is not permitted. If the work is adapted, then it must be licensed under the same or equivalent Creative Commons licence. If a translation of this work is created, it must include the following disclaimer along with the required citation: "This translation was not created by the Food and Agriculture Organization of the United Nations (FAO) or the World Health Organization (WHO). Neither FAO nor WHO is responsible for the content or accuracy of this translation. The original English edition shall be the authoritative edition.

Disputes arising under the licence that cannot be settled amicably will be resolved by mediation and arbitration as described in Article 8 of the licence except as otherwise provided herein. The applicable mediation rules will be the mediation rules of the World Intellectual Property Organization http:// www.wipo.int/amc/en/mediation/rules and any arbitration will be conducted in accordance with the Arbitration Rules of the United Nations Commission on International Trade Law (UNCITRAL).

**Third-party materials.** Users wishing to reuse material from this work that is attributed to a third party, such as tables, figures or images, are responsible for determining whether permission is needed for that reuse and for obtaining permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

**Sales, rights and licensing**. FAO information products are available on the FAO website (www.fao. org/publications) and can be purchased through publications-sales@fao.org. Requests for commercial use should be submitted via: www.fao.org/contact-us/licence-request. Queries regarding rights and licensing should be submitted to: copyright@fao.org.

Cover picture cDennis Kunkel Microscopy, Inc.

# Contents

Contributors											
Acknowledgements											
Abbreviations											
Declaration of interests Executive summary Scope and objectives Conclusions											
						1	Introduction				
							1.1	Reque	est from Codex	1	
							1.2	Consu	mption and production	2	
		1.2.1	Global consumption of chicken meat	2							
		1.2.2	Intensification of broiler chicken production to meet growing	g							
	10	<b>c</b>	needs	3							
	1.3	Scope	of this report	3							
	1.4	Litera	ture survey	4							
2	Con	ntrol measures									
	2.1.	Contro	ols during primary production	5							
		2.1.1.	Biosecurity, management and awareness of <i>Campylobacter</i> prevalence addressed in approaches to control <i>Campylobac</i> spp.	ter 5							
		2.1.2.	Vaccine- and bacteriophage-based approaches for the control of <i>Campylobacter</i> spp.	9							
		2.1.3.	Feed and drinking water additives approaches for the contro <i>Campylobacter</i> spp.	ol of 12							
	2.2.	Contro	ol measures during processing	16							
		2.2.1	Effectiveness of specific processing procedures	18							
	2.3.	. Controls during post-processing, retail and consumer levels									
		2.3.2	Effectiveness of specific retail/post-processing measures	22							
3	Review of the Code of Practice			25							

4	Concluding remarks								
5	Intervention summary synopsis								
6	Refere	nces							
7	Annex				55				
	Annex 1	sed for searching the control measures of spp. in poultry	57						
	Annex 2	The questions for the two-step relevance screening and confirmation							
		A2.1. F	Relevance	escreening	59				
		A2.2 Relevance confirmation							
	Annex 3	Annex 3 Recommended revisions to the <i>Guidelines for the Control of</i> Campylobacter <i>and</i> Salmonella <i>in Chicken Meat (GXG 78-20</i> as they relate specifically to the control of <i>Campylobacter</i> sp							
	Annex 4	Bibliography used for scoping review							
		A 4.1	Primar	ý	71				
			A4.1.1	Biosecurity and management approaches Review of the Codex guidelines	71				
			A4.1.2	Vaccine and phage based approaches	77				
			A4.1.3	Feed and water additives approaches	80				
		A4.2	Process	sing	87				
		A4.3	Post-pi	rocessing	105				

## Contributors

### Experts

**Marianne Chemaly**, French Agency for Food, Environmental and Occupational Health & Safety, France

**Frances Colles**, University of Oxford, the United Kingdom of Great Britain and Northern Ireland

Alessandra De Cesare, Department of Veterinary Medical Sciences, University of Bologna, Italy

Moses Gathura Gichia, Food Safety Consultant, Nairobi, Kenya

**Ihab Habib**, College of Agriculture and Veterinary Medicine, United Arab Emirates University, the United Arab Emirates

**Nicol Janecko**, Quadram Institute Bioscience, the United Kingdom of Great Britain and Northern Ireland (served as Rapporteur)

**Catherine M Logue**, Department of Population Health, College of Veterinary Medicine, University of Georgia, the United States of America

Marcos Sanchez-Plata, Texas Tech University, the United States of America

Elina Tast-Lahti, National Veterinary Institute, Sweden

**Jaap A. Wagenaar**, Faculty of Veterinary Medicine, Utrecht University, the Kingdom of the Netherlands (served as Chair)

Bing Wang, University of Nebraska-Lincoln, the United States of America

#### **Resource persons**

Jose Emilio Esteban, Codex Committee on Food Hygiene, the United States of America

**Evelyne Mbandi**, Codex Committee on Food Hygiene, the United States of America

Sarah Cahill, Joint FAO/WHO Food Standards Programme, Italy

**Jorge Pinto Ferreira**, Food Systems and Food Safety Division, Food and Agriculture Organization of the United Nations, Italy

#### Secretariat

**Akio Hasegawa**, Department of Nutrition and Food Safety, World Health Organization, Switzerland

**Christine Kopko**, Food Systems and Food Safety Division, Food and Agriculture Organization of the United Nations, Italy

**Jeffrey LeJeune**, Food Systems and Food Safety Division, Food and Agriculture Organization of the United Nations, Italy

**Juliana de Oliveira Mota**, Department of Nutrition and Food Safety, World Health Organization, Switzerland

**Moez Sanaa**, Department of Nutrition and Food Safety, World Health Organization, Switzerland

**Kang Zhou**, Food Systems and Food Safety Division, Food and Agriculture Organization of the United Nations, Italy

## Acknowledgements

The Food and Agriculture Organization of the United Nations (FAO) and the World Health Organization (WHO) would like to express their appreciation to all those who contributed to the preparation of this report through the provision of their time and expertise, data and other relevant information before, during and after the meeting. Special appreciation is extended to all the members of the expert committee for their dedication to this project, to Jaap Wagenaar for his expert chairing of the committee, and to Nicol Janecko for her excellent support as rapporteur. All contributors are listed in the following page. Appreciation is also extended to all those who responded to the calls for data that were issued by FAO and WHO.

The preparatory work and expert meeting convened to prepare this report was coordinated by the Secretariat of the Joint FAO/WHO Expert Meetings on Microbiological Risk Assessment (JEMRA).

## Abbreviations

CCFH	Codex Committee on Food Hygiene	
CFU	Colony forming unit	
DNA	deoxyribonucleic acid	
FAO	Food and Agriculture Organization of the United Nations	
GHP	good hygiene practices	
НАССР	Hazard Analysis and Critical Control Points	
JEMRA	Joint FAO/WHO Expert Meetings on Microbiological Risk Assessment	
QMRA	quantitative microbial risk assessment	
WHO	World Health Organization	
WOAH	World Organisation for Animal Health	

## Declaration of interests

All participants completed a Declaration of Interests form in advance of the meeting. All experts were not considered by FAO and WHO to have declared any interest that may be perceived as a potential conflict with regard to the objectives of the meeting.

All the declarations, together with any updates, were made known and available to all the participants at the beginning of the meeting.

All the experts participated in their individual capacities and not as representatives of their countries, governments, or organizations.

## Executive summary

### **Scope and objectives**

In response to a request from the 52nd Session of the Codex Committee on Food Hygiene (CCFH), the Joint FAO/WHO Expert Meeting on Microbiological Risk Assessment (JEMRA) convened a meeting in Rome, Italy from 6 to 10 February 2023, to collate and assess the most recent scientific information relevant to the control of thermotolerant *Campylobacter* species *C. jejuni* and *C. coli* (hereafter *Campylobacter*) in broiler production and chicken meat, including a review of the Codex *Guidelines for the Control of Campylobacter and Salmonella in Chicken Meat* (CXG 78-2011).<sup>1</sup>

The scope was focused on aspects of broiler primary production from the point of chick placement into production establishments to consumer handling.

The objectives were to identify and assess control measures for *Campylobacter* in the broiler production chain. The expert committee reviewed the available data on *Campylobacter* control including scientific literature published from 2008 to October 2022 and data submitted in response to a call for data for this meeting. The experts: 1) determined the quality and quantity of evidence of control measures for *Campylobacter*, 2) evaluated the impact of measures to control *Campylobacter* in the broiler production chain, 3) determined which hazard-based interventions pertained specifically to *Campylobacter* and which were general to the control of foodborne pathogens in the pre- and post-harvest broiler production chain, and 4) reviewed and recommended revisions to the *Guidelines for the Control of Campylobacter and Salmonella in Chicken Meat* (CXG 78-2011), paragraphs 1 to 115, based on the currently available scientific evidence (Annex 3).

Control measure evaluation began at the time of chick placement since there is currently no evidence that parent flocks or hatchery practices contribute to the colonization of broiler chicks. The available literature on interventions was predominantly based on laboratory and pilot studies, with few commercial scale applications; therefore, limited conclusions could be reached. The experts recommend the use of a combination of multiple interventions (multihurdle

<sup>1</sup> FAO and WHO. 2011. Codex Guidelines for the Control of Campylobacter and Salmonella in Chicken Meat (CXG 78-2011). Rome. https://www.fao.org/fao-who-codexalimentarius/shproxy/de/?lnk=1&url=https%253A%252F%252Fworkspace.fao.org%252Fsites%252Fcodex %252FStandards%252FCXG%2B78-2011%252FCXG\_078e.pdf

approach) suitable to production and processing stages to lower *Campylobacter* contamination on chicken meat.

### Conclusions

Several interventions were identified through primary production to post-processing stages. The expert consultation concluded the following in each stage of production:

#### Assessment of primary production interventions for the control of Campylobacter

#### Biosecurity and management approaches

- Using strict biosecurity measures (hygienic practices and sanitation) can enhance the control of *Campylobacter* in broiler flocks.
- Risk factors for *Campylobacter* contamination at primary production establishments, such as partial depopulation, litter management, down period length, proximity to other livestock and slaughter age can help guide intervention strategies.

#### Vaccination-based approaches

- Currently, there are no commercial vaccines for *Campylobacter* readily available for any stage of primary production, and vaccination studies were limited to *C. jejuni* only.
- Several potential vaccine candidates are in the proof-of-concept phase but cannot be considered yet as an intervention.
- Some vaccines induced a cellular or humoral response in the chicken host, but this did not always translate to reduced caecal colonization by *Campylobacter* in pilot studies.

#### Bacteriophage-based approaches

- There are currently no commercial products available for use in primary production.
- The effects of phage therapy may be transitory and prone to resistance.

#### Feed and water additives

#### Organic acids

• In feed, short- and medium-chain fatty acids, and in particular, caprylic acid, show promise as feed additives in reducing *Campylobacter* in pilot studies.

• In drinking water, organic acids reduced *Campylobacter* in caecal/ faecal specimens at the end of the primary production period; however, the effects were not sustained to the end of production in pilot studies.

#### Probiotics

• In feed, there is inconsistent evidence on the efficacy of probiotics as an intervention for reducing *Campylobacter* in broilers at primary production level.

#### Plant-based additives

• In feed, the efficacy of some plant-based molecules in *in vivo* pilot studies showed limited reduction of *Campylobacter* in caecal/faecal specimens at the end of the primary production period.

#### Assessment of processing interventions for the control of Campylobacter

- Good hygienic practices (GHP) during processing are important in minimizing *Campylobacter* contamination on meat.
- The effectiveness of interventions during processing is dependent upon the incoming flock prevalence and concentration of *Campylobacter* in the gastrointestinal tract and on the bird.
- The impact of processing practices can be enhanced by a combination of a multihurdle approach, processing effects, physical and/or chemical interventions.

#### **Processing effects**

- Logistic slaughter scheduling can reduce Campylobacter cross-contamination.
- Qualitative and quantitative targets for *Campylobacter* may be used to optimize process control.
- Scalding reduces the carcass surface concentration and prevalence of *Campylobacter*, The result depends on the temperature, and dilution effect.
- Defeathering and evisceration may increase both prevalence and concentration of *Campylobacter* on carcasses.
- Immersion chilling can reduce (dilute) the carcass concentration of *Campylobacter*; however, this is dependent on the initial *Campylobacter* load on the incoming birds.
- In combination with processing aids, immersion chilling may reduce the carcass prevalence of *Campylobacter*.

• Air chilling may reduce concentration of *Campylobacter*, but the efficacy of air chilling in reducing prevalence of *Campylobacter* when used without other processing aids is inconclusive.

#### Physical

- Irradiation is effective at eliminating Campylobacter on meat.
- Freezing meat reduces the concentration of *Campylobacter*.
- Steam, ultrasonication, high-intensity light pulse, visible light, UV-C and other technologies have shown promise either at the laboratory or pilot scale, but their impact is unknown at commercial scale.

#### Chemical

- Processing aids such as chlorine derivatives, peroxyacetic acids, and organic acids added to water used for washing and/or dipping may reduce *Campylobacter* on carcasses.
- Some marination ingredients have shown reductions in *Campylobacter* on meat.

#### Post-processing interventions for the control of Campylobacter

- Thorough cooking is effective at eliminating *Campylobacter* on meat.
- The application of GHPs is important in reducing *Campylobacter* on meat.
- Freezing meat reduces the concentration of Campylobacter.
- Some marination ingredients have shown reductions in *Campylobacter* on meat.

Available evidence for the reduction of *Campylobacter* was primarily focused on *C. jejuni* and *C. coli*. Interventions aimed at foodborne pathogens such as irradiation or thorough cooking are effective in eliminating *Campylobacter* on meat. Hazard-based interventions, good agriculture practices and hygienic practices for the general control of foodborne pathogens may be effective for the reduction of *Campylobacter*. There are no interventions that **specifically** control *Campylobacter* on meat.

The experts recognize further data gaps exist and that new technologies may offer promising approaches to reducing *Campylobacter* on chicken meat. Further global changes to the industry, the growth of global populations, climate change, and increased demand for animal protein in specific regions will guide the need for further control measure assessments.

## Introduction

### 1.1 REQUEST FROM CODEX

Campylobacteriosis is among the most frequently reported foodborne diseases worldwide (Havelaar *et al.*, 2015; Tack *et al.*, 2019). In response to the requests from Codex for scientific advice, FAO and WHO have undertaken risk assessments of foodborne pathogens in various foods since 1999 (FAO and WHO, 1999). In the past, the Joint FAO/WHO Expert Meetings on Microbiological Risk Assessment (JEMRA) have conducted risk assessments of *Campylobacter* spp. in broiler chickens (FAO and WHO, 2008, 2009b) and evaluated intervention measures being used in the production of chicken meat (FAO and WHO, 2009a).

In its report on the global burden of foodborne disease, the World Health Organization (WHO) estimated that in 2010 foodborne *Campylobacter* spp. caused more than 95 million illnesses, 21 374 deaths, and nearly 2 142 000 DALYs (Havelaar *et al.*, 2015). While numerous potential vehicles of transmission exist, commercial chicken meat has been identified as one of the most important food transmission vehicles for *Campylobacter* spp.

At its 52nd session in 2022, the Codex Committee on Food Hygiene (CCFH) requested that JEMRA collate relevant scientific information on *Salmonella* and *Campylobacter* in chicken meat in preparation for a potential update of the existing *Guidelines for the Control of Campylobacter and Salmonella in Chicken Meat* (CXG 78-2011).

To meet the request of the CCFH, FAO and WHO convened this expert meeting on the pre- and postharvest control of *Campylobacter* spp. in broiler chicken meat from 6 to 10 February 2023 at FAO headquarters, Rome. The goal of the meeting was to gather and evaluate recent data, evidence and scientific opinions on the topic. The output aim was to provide expert opinion on intervention strategies for the control of *Campylobacter* in pre- and post-production chicken meat.

### **1.2 CONSUMPTION AND PRODUCTION**

#### 1.2.1 Global consumption of chicken meat

According to FAO, the global poultry population was 27.9 billion head in 2019, with chickens accounting for 93 percent. Poultry refers to all domesticated birds raised for meat, eggs or feathers, while chicken meat (broiler chicken) production includes species of chickens raised for meat consumption. The number of chickens worldwide has more than doubled since 1990 with poultry accounting for more than 40 percent of all meat produced for human consumption (OECD and FAO, 2021).

*The Agricultural Outlook 2021–2030* suggests that the global meat supply will continue to expand over the next projection period reaching 374 million tons by 2030 (OECD and FAO, 2021). The overall global expansion of meat production is driven by production in China, the Americas and Africa. China alone is expected to account for the greatest increase in meat production followed by Brazil and the United States of America. The global consumption of meat protein from any animal species is also projected to increase by 14 percent by 2030, from the 2018–2020 projections (OECD and FAO, 2021). Factors influencing meat consumption include population growth and demographics, urbanization, income, meat prices, cultural norms, environmental aspects, animal welfare and people's personal health.

Chicken meat production and availability are expected to grow by 18 percent globally due to the relative affordability for middle-income consumers. International trade has also helped improve access to chicken products worldwide and increasingly so in Asia and the Near East where demand has outpaced production (OECD and FAO, 2021). Reasons for broiler chicken production growth are multifactorial and include: 1) the short birth to slaughter time compared to other meat-producing animals, 2) the ability for lower income families to gain a higher quality of life by raising birds in small-farm holdings, and 3) the overall increase in chicken meat demand as a result of population growth and growth in income levels.

As a result of the continued growth of the broiler chicken market, it is necessary to review pre- and post-production practices from hatchlings to final chicken meat products to help ensure the safety and quality of this important protein source for the world's population.

## 1.2.2 Intensification of broiler chicken production to meet growing needs

In many high-income countries, broiler chicken production has been intensified by increasing farm and flock sizes, developing rapidly growing chicken breeds, creating a global supply of parental stocks, and investing in optimal feeding (Li *et al.*, 2021; Jeni *et al.*, 2021). Typically, commercial farms receive day-old chicks, raise them to a designated age of slaughter and transport the whole flock to an abattoir. In many low- and middle-income countries, small to medium-size enterprises, backyard farming and live-bird markets dominate domestic broiler chicken meat production (Delabouglise *et al.*, 2019; Wong *et al.*, 2017). Biosecurity is a challenge for the entire chicken production industry but especially so for enterprises with birds raised in backyard systems and sold at live-bird markets. These less intensive chicken production systems rarely apply strict biosecurity and good husbandry practices. When farms are transitioning to increased intensification, the education and motivation of farmers in management and biosecurity is crucial.

In the commercial systems, market forces such as an increased demand for chicken meat products, demand for organically raised products, animal welfare considerations, and housing requirements all need to be taken into account when designing *Campylobacter* spp. control strategies. Such control strategies should be designed, implemented and monitored with consideration to specific production conditions.

### **1.3 SCOPE OF THIS REPORT**

The purpose of the experts meeting was to collect, review and discuss relevant measures for control of *Campylobacter* spp. in broiler chicken production, from the primary production stage to consumption.

The scope of the meeting included, but was not limited to, aspects of primary broiler chicken production, processing, distribution, product handling, preparation, retail and consumption of chicken meat. Emphasis was placed on the identification and evaluation of control measures to reduce campylobacteriosis associated with consumption of chicken meat, taking into consideration the effectiveness and practicalities of measures.

The objectives of this meeting were:

• To review publicly available literature and guidelines from competent authorities and industry associations (e.g. compliance guidelines, code of

practices, and so on) to assess the current state of knowledge in controlling *Campylobacter* spp. in chicken meat.

• To review mitigation/intervention measures being used at different points along the food chain and assess their effectiveness at reducing *Campylobacter* spp. in chicken meat.

### **1.4 LITERATURE SURVEY**

A review of the available scientific literature targeting changes in knowledge since 2008 was used to develop a bibliography. Scientific articles were selected from two databases (*Web of Science* and *PubMed*), and as there was a need to consider studies published in languages other than English, data from member countries and expert opinions were also relied upon.

The records from *Web of Science* (n = 2498) and *PubMed* (n = 1964) were added into *Distiller*. The function "Duplication Detection" was used by comparing the Title, Author and Abstract. A total of 915 duplicate articles were found. The "Smart Quarantine" feature in *Distiller* was used to remove these duplicates resulting in 3547 (2498 + 1964 – 915) publications which were used to establish the working database for the meeting. The search was carried out on 23 October 2022. The keywords used for searching the literature are detailed in Annex 1.

The database was further refined using a two-step process for the relevance screening and confirmation of the 3547 articles, which were prepared in a dataset for the experts to review. The details of this procedure were presented to the experts for their further review and are included in Annex 2.

The experts had a further review of these publications during the meeting, to finalize whether the literature was considered control measures or not. They provided recommendations and conclusions based on the included scientific literature as well as their expert opinions. Literature with applicability to a commercial setting was considered. The final reviewed papers, not necessarily cited, are listed in the bibliography at the end of the report.

## **Control measures**

### 2.1 CONTROLS DURING PRIMARY PRODUCTION

### 2.1.1 Biosecurity, management and awareness of *Campylobacter* prevalence addressed in approaches to control *Campylobacter* spp.

To complete the assessment of control measures during primary production, experts reviewed 45 papers from the literature survey (listed in Annex 4.1.1). The following sections highlight the findings.

#### • Biosecurity

Biosecurity measures appropriate for the control of *Campylobacter* cover multiple factors, including good production practices such as strict biosecurity and sanitation (Hansson *et al.*, 2010; Henry *et al.*, 2011; Jonsson *et al.*, 2012; Hasan *et al.*, 2020; Sandberg *et al.*, 2017; Hertogs *et al.*, 2021; Schweitzer *et al.*, 2021; Golden and Mishra, 2020; Georgiev, Beauvais and Guitian, 2017). These measures remain the single most effective tool to reduce *Campylobacter* contamination at all stages of primary production and should form the foundation of any intervention strategy.

#### Management

The partial depopulation (thinning) has been found to have an opposite effect with an increased risk of breaching biosecurity measures and is evidenced to be a contributing factor to *Campylobacter* contamination of broiler chicken flocks (Crotta, Georgiev and Guitian, 2016; EFSA, 2020; van Wagenberg *et al.*, 2016;

Smith *et al.*, 2016; Allen *et al.*, 2008; Higham *et al.*, 2018; Georgiev, Beauvais and Guitian, 2017). Estimates of relative risk reductions from discontinued thinning range from 2–33 percent (Crotta, Georgiev and Guitian, 2016; Georgiev, Beauvais and Guitian, 2017; van Wagenberg *et al.*, 2016). Alternately, adequate cleaning and disinfection practices, as well as a "resting period", to decontaminate housing between flocks have been associated with lower *Campylobacter* contamination of subsequent flocks (Zbrun *et al.*, 2021; Bailey *et al.*, 2022; Reichelt *et al.*, 2022; Agunos *et al.*, 2014).

#### Awareness of Campylobacter prevalence in primary production

*Campylobacter* prevalence amongst broiler flocks varies by season and is dependent on the region. Temperature and humidity in countries with both temperate and warm climates has been associated with changes in prevalence (Urdaneta *et al.*, 2023; Guerin *et al.*, 2008; Chowdhury *et al.*, 2013; Iannetti *et al.*, 2020; Sindiyo *et al.*, 2018; Seman, Gregova and Korin, 2020; Higham *et al.*, 2018; Wanja *et al.*, 2022). In addition, ventilation type was associated with *Campylobacter* prevalence in a Spanish study (Urdaneta *et al.*, 2023). Older birds are more likely to be infected by *Campylobacter*, though evidence regarding prevalence of *Campylobacter* amongst conventional *versus* organic flocks, as well as breed, flock size and stocking density was mixed (Iannetti *et al.*, 2020; Rawson, Dawkins and Bonsall, 2019; Lassen *et al.*, 2022; McKenna *et al.*, 2018; Borck Høg *et al.*, 2016; Williams *et al.*, 2013; Russell *et al.*, 2021; Babacan *et al.*, 2020; Griekspoor *et al.*, 2013; van Wagenberg *et al.*, 2016).

There is a limited number of studies investigating the effect of conventional versus extensive and antibiotic-free production systems on Campylobacter species presence, antibiotic resistance and animal welfare (Hansson et al., 2010; Iannetti et al., 2020; Babacan et al., 2020; Bailey et al., 2019; Bull et al., 2008; El-Shibiny, Connerton, P.L. and Connerton, I.F., 2005). Of those studies reviewed, five reported lower Campylobacter prevalence in flocks with higher welfare scores, amongst both conventional and extensive flocks (Bull et al., 2008; Williams et al., 2013; Colles et al., 2016; Ianetti et al., 2020; Di Marcantonio et al., 2022). Older birds raised in extensive production systems were more frequently colonized by C. coli (El-Shibiny, Connerton, P.L. and Connerton, I.F., 2005; Babacan et al., 2020). Organically raised flocks were associated with lower levels of Antimicrobial Resistance (AMR) in primary production (Luangtongkum et al., 2006; Bailey et al., 2019; Hansson et al., 2021). More research is needed to understand complex interactions between rearing system, chicken age/breed and welfare in determining Campylobacter prevalence and risk to human health (Ianetti et al., 2020; Di Marcantonio et al., 2022).

## • Sources of *Campylobacter* contamination into primary production settings and management interventions

The age of a broiler house, number of houses on the property, house layout including ante-room(s), water sources and drinker type, proximity to other farms, livestock and other animals were all identified as risk factors for contaminating the birds by various studies (Torralbo *et al.*, 2014; Borck Høg *et al.*, 2016; Torralbo *et al.*, 2014; van Wagenberg *et al.*, 2016; Hertogs *et al.*, 2021). Genomic typing data indicates some *Campylobacter* lineages are associated with specific livestock species (including broiler chickens) and wild birds, with little overlap between the *Campylobacter* lineage types (Colles *et al.*, 2008; De Haan *et al.*, 2010; Griekspoor *et al.*, 2013; Sheppard *et al.*, 2010). *Campylobacter* prevalence in wild birds appears to be associated with feeding guild, with *Campylobacter* genotype overriding geographic signals (Griekspoor *et al.*, 2013; Hald *et al.*, 2016). Transport was assessed by a few studies. The effects of transport on *Campylobacter* presence in broiler flocks were mixed with no clear indication of an increase or decrease (Mendes *et al.*, 2020; Sevilla-Navarro *et al.*, 2020).

The impact of litter type on *Campylobacter* colonization of broiler flocks is of interest. Some evidence suggests a high incidence of foot lesions derived from wet litter conditions, which may in turn be predictive of higher *Campylobacter* contamination load in the flock (Williams *et al.*, 2013; Alpigiani *et al.*, 2017). Litter quality is included in the internationally recognized WelfareQuality\* protocol and was considered a risk factor in some of the previously mentioned studies investigating links between flock welfare and *Campylobacter* prevalence (Ianetti *et al.*, 2016; Di Marcantonio *et al.*, 2022). The practice of re-use and treatment of litter between flocks varies globally, however no recommendations for best practice can be made at the current time (Rothrock *et al.*, 2008; Bailey *et al.*, 2022).

There was evidence from Denmark and Iceland that fly screens were associated with lower prevalence of *Campylobacter* in flocks if high level of biosecurity measures were already in place (Tustin *et al.*, 2010); however, no further evidence is available since the original publication.

There is sparse information regarding the role of hatcheries and breeder flocks as sources of transmission of *Campylobacter* to primary production flocks. Vertical transmission of *Campylobacter* through eggs is largely rejected by the scientific community, though there are occasional reports of *Campylobacter* detected in the reproductive tract of breeding birds (Battersby *et al.*, 2016). Broiler flocks are usually colonized with different strains than the parent flock (Al Hakeem *et al.*, 2022). *Campylobacter* DNA has been detected from commercial broiler

chicks less than 8 days old, suggesting that early routes of contamination might have an impact in some instances, however viable culture could not be recovered (Colles *et al.*, 2021). Vertical transmission remains a topic of investigation for some studies (Lu *et al.*, 2021). To date, quality evidence to support this as a transmission route is lacking.

Many risk factors contribute to *Campylobacter* spp. colonization of broiler flocks, indicating the difficulties in maintaining effective countermeasures against its entry into the broiler environment (Natsos *et al.*, 2020). Studies have indicated that horizontal transmission from environmental sources is the most significant cause of dissemination into primary production flocks. Newer precision based genotyping studies will continue to provide further evidence to understand and refine all forms of transmission routes.

It is challenging to determine the extent of impact that farm personnel or professionals visiting farms as a control measure. One study concluded disinfection of broiler houses by unskilled personnel was a risk factor for *Campylobacter* colonization (Natsos *et al.*, 2020) whereas another study found no effect on *Campylobacter* prevalence (Näther *et al.* 2009). In a UK specific study, financial incentive for farmers resulted in farmers producing a greater number of flocks with lower levels of *Campylobacter* contamination (Higham *et al.*, 2018). In a US-based questionnaire study of broiler industry stakeholders, it was concluded that many survey participants were unfamiliar with *Campylobacter* specific risk factors (Hwang and Singer, 2020). Without further assessment, the impact of personnel on the introduction and levels of *Campylobacter* contamination is inconclusive.

#### Conclusions

Strict biosecurity and sanitation measures are a key requirement by which to control *Campylobacter* and should form the foundation for any on-farm based interventions. There is some evidence that management practices can influence *Campylobacter* population; however, other impact factors such as seasonal *Campylobacter* population variation in specific geographical regions, breed of birds, and age of birds at slaughter cannot be easily controlled by management practices alone. Transmission route analyses will be helped by tools such as rapid molecular assays and genomic sequencing that will in turn contribute evidence used to guide future intervention recommendations and refinement of biosecurity measures. The genomic profiling of *Campylobacter* strain lineages in broiler chicken production will also enable assessment of impact when studying interactions between the chickens and environmental microbiomes.

Important considerations for on-farm assessments include:

- whether it is necessary to partially depopulate ("thin") flocks;
- down time between flocks;
- litter management and welfare of flocks;
- housing, including age/ability to disinfect, ventilation and layout (ante-room[s]); and
- training and compliance of staff and visitors on strict biosecurity measures.

# 2.1.2 Vaccine- and bacteriophage-based approaches for the control of *Campylobacter* spp.

Vaccines are agents or portions of a bacterial cell that are used to prevent *Campylobacter* colonization in broiler birds and to decrease the potential for *Campylobacter* transmission to meat products. Vaccines against *Campylobacter* have been investigated over many years. A number of different vaccine types that have been investigated include attenuated, subunit, inactivated, conjugate, DNA, vector and recombinant vaccines. While efforts in developing effective *Campylobacter* vaccines have been made, there is currently no single vaccine approved for use in broiler bird production.

Bacteriophages, the viruses that specifically infect target bacteria, thereby killing the host bacterial cell, have also been investigated for their potential to control *Campylobacter*. The field of *Campylobacter* specific bacteriophages (phages) is relatively limited with very few published data available.

#### 2.1.2.1 Vaccination-based approaches to control Campylobacter

The experts reviewed 23 papers related to vaccine studies for *Campylobacter* control with 14 directly applicable to the current review (Annex 4.1.2). The following sections highlight evidence pertaining to vaccination as an approach to *Campylobacter* control in broiler birds.

#### • Quality of evidence

Currently, there are no commercial vaccines for *Campylobacter* readily available. Potential candidates (e.g. reverse vaccinology approach) are on the horizon; however, most trials at present are in a "proof-of-concept phase" or have reported findings from laboratory or pilot-scale level studies. The quality of the papers was satisfactory in terms of methodology and study design that incorporate controls and challenge groups. Assessed trial studies also had a satisfactory level of statistical analysis to determine measurable effects. A range of vaccine methods and types (glycans, inserted proteins, flagellin, subunit, *in ovo*, and live attenuated) were included in the literature assessed (Layton *et al.*, 2011; Jeon *et al.*, 2022;

Gloanec *et al.*, 2022). Some approaches included an alternative vector expressing *Campylobacter* proteins to deliver antigens to the poultry host, e.g. *Salmonella* (Buckley *et al.*, 2010; Sahin *et al.*, 2015) and *Eimeria* (Clark *et al.*, 2012), yet conclusions could not be reached on the approaches as control measures.

The literature reviewed was in the proof-of-concept stage, with no papers indicating any commercial-level trials or follow-up studies, demonstrating the infancy of the field of vaccine development for *Campylobacter* control in broilers.

#### • The measure of effectiveness

The measure of effectiveness to control *Campylobacter* was reported as log level reductions in the ceca or gut (quantitative) or number of birds positive following vaccination and challenge (prevalence). Reported vaccine effects varied from no reductions of *Campylobacter* noted (Meunier *et al.*, 2018; Mauri *et al.*, 2021; Vandeputte *et al.*, 2019) to low level reductions of 1–2 logs or less (Łaniewski *et al.*, 2014; Buckley *et al.*, 2010; Chintoan-Uta, Cassady-Cain and Stevens, 2016; Gloanec *et al.*, 2022), while a few studies reported high level reductions (up to 4 logs) (Layton *et al.*, 2011; Cui *et al.* 2022; Nothaft *et al.*, 2021).

There was evidence that some of the candidate vaccines appear to be promising; however, further research is warranted to examine the candidates on a larger scale, e.g. commercial farms (Cui *et al.* 2022; Layton *et al.*, 2011). There is also limited evidence as to the practicality of vaccine production for these candidates on a larger scale or as to wider industry vaccination strategy.

Some candidate vaccines were found to induce cellular or humoral response in the chicken host, meaning that the host immune response recognized *Campylobacter*, but in terms of elimination of *Campylobacter* from the chicken host this did not translate to stable (rather than transient) caecal colonization or reductions levels (> 2 logs) (Radomska *et al.*, 2016; Cui *et al.* 2022; Gloanec *et al.*, 2022), suggesting that the effects were not consistent and unlikely to be permanent.

Most studies assessed used a homologous challenge approach (i.e. single strain, typically a *C. jejuni*) with no studies using a heterologous challenge (i.e. using diverse *Campylobacter* strains). No studies to date investigated the ability of a vaccine candidate to provide cross protection against diverse *Campylobacter* species. Additional studies are warranted to understand the levels and types of cross protection provided by potential vaccine candidates.

#### • Scalability or applicability (practicability)

Further studies are required for extensive scale application of candidate vaccines, i.e. farm-level production. All studies reviewed were currently at the lab or pilot-scale

(research farm) level only with no indication for larger pilot-scale studies such as commercial farm or production-level studies and no indication of commercial vaccine trials. Such large-scale trials will be necessary to assess the scalability and potential impact of candidate vaccines for full implementation in broiler production.

#### • Geographical representation of the studies

All vaccine candidate studies were conducted only in high-income countries or regions of the world; therefore, the translation of these studies to underdeveloped regions will warrant additional studies in specific regions to assess efficacy and practicality.

#### Conclusion

Vaccines for *Campylobacter* control continues to be an emerging field and warrants further investigation and investment to take proof-of-concept studies to commercial trial phases.

## 2.1.2.2 Bacteriophage-based approaches for the control of *Campylobacter* spp.

The experts reviewed ten papers (Annex 4.1.2) on the use of bacteriophages as an intervention measure for *Campylobacter* colonization in broilers.

#### • Quality of evidence

There are currently no commercial products available for use in primary production. The bacteriophages used in the studies reviewed varied among studies, with some well-characterized bacteriophages. Of the six experimental studies reviewed, all had treatment groups and control groups, allowing assessment of the effect of phage treatment in a statistically sound manner.

#### • The measure of effectiveness

Some studies used a single phage strain (Fischer *et al.*, 2013; Furuta *et al.*, 2017; El-Shibiny *et al.*, 2009), whereas others used multiple (a cocktail of) phage strains (Carvalho *et al.*, 2010; Kittler *et al.*, 2013; Richards, Connerton, P.L. and Connerton, I.F., 2019; Hammerl *et al.*, 2014; D'Angelantonio *et al.*, 2021; Chinivasagam *et al.*, 2020). Treatment was typically via drinking water or gavage. The effects of phage treatment on reductions of *Campylobacter* ranged from a 1 log to a > 3 log reduction of *Campylobacter* colonization in faecal droppings or caecal contents after necropsy. The effect was most evident a few days post treatment; however, *Campylobacter* colonization levels increased over time post treatment. In some studies, *Campylobacter* strains were screened for resistance to phages after treatment. Resistance was reported; however, these strains did not

overgrow susceptible strains. In some studies, there was considerable variation between groups/barns used in the study. A reduction of 3 log is considered relevant for public health, based on a risk assessment by Rosenquist *et al.* (2003).

Over the last decade or more, there has not appeared to be any technical progress in the experimental approaches used for the application of phages for *Campylobacter* control. Studies do not address the variability between *Campylobacter* strains and the applicability or practicality of phages for field or commercial purposes.

## • Scalability or applicability (practicality) and geographical representation studies

Except for two studies performed on-farm (Australia and Germany) (Chinivasagam *et al.*, 2020; Kittler *et al.*, 2013) with naturally colonized flocks, studies were performed in experimental, controlled settings. Applicability could therefore not be assessed for commercial settings. One field study was conducted in Australia (Chinivasagam *et al.*, 2020); all other studies were performed in Europe, and therefore there is no indication of how bacteriophage approaches would be applicable in other regions.

Phages are variable depending on the strain type and vary in the level of activity. Additionally, the potential for *Campylobacter* to develop resistance to the phage is considerable. Bacteriophage approaches to control *Campylobacter* are not clear. In conclusion, no recommendation on the use of bacteriophages can be made at this time.

# 2.1.3 Feed and drinking water additives approaches for the control of *Campylobacter* spp.

Feed and water additives are chemical or biological supplements that can be added to water or feed formulations. Additives as a control measure of *Campylobacter* have been assessed for effectiveness and scalability. In-feed and in-water additives applied at the primary stage of production (farms) were reviewed.

#### • Quality of evidence

The experts reviewed a total of 89 papers by first screening and then further subdividing them into relevant subheadings of organic acids, probiotics and plant-based additives. The quality of evidence varied with most of the available literature containing laboratory and pilot stage studies. Although some studies have provided promising results as to the effects of organic acid additives, extrapolation of the evidence to the commercial scale of production systems could not readily be made.

#### • Organic acids: measure of effectiveness

Seven papers reported on the impacts of medium chain fatty acids (caprylic acid) supplementation in feed. The supplementation of medium chain fatty acids in feed is considered a promising tool for the reduction of *Campylobacter* colonization in commercial broiler flocks (Van Gerwe *et al.*, 2010). Therapeutic supplementation of caprylic acid at 0.7 percent in the feed for 3 or 7 days before slaughter showed a consistent 3 to 4 log reduction in caecal *C. jejuni* counts in market-aged broiler chickens (De Los Santos *et al.*, 2010). The results of this study indicate that caprylic acid's ability to reduce *Campylobacter* does not appear to be due to changes in caecal microflora or a decrease in intestinal pH. It is hypothesized that caprylic acid may compromise the outer membrane determinants in *Campylobacter* which are needed for bacterial adaptation to host environment and colonization. It is also possible that caprylic acid has a direct inhibitory effect on the expression of virulence factors necessary for *C. jejuni* colonization in chicks.

The in-water supplementation of caprylic acid at various concentration levels did not reduce *Campylobacter* shedding due to the changes in adsorption and caprylic concentrations reaching the lower intestinal tract (Metcalf *et al.*, 2011). The addition of sorbic acid, benzoic acid, propionic acid, and acetic acid in drinking water reduced the *Campylobacter* shedding reaching a maximum 2 log reduction during the rearing period (Gharib Naseri, Rahimi and Khaki, 2012); however, the drinking water additive failed to diminish *Campylobacter* colonization in the intestinal colonic and caecal contents at the time of slaughter (Szott *et al.*, 2022). Therefore, the benefits of drinking water treatment to controlling *Campylobacter* shedding and transmission was not significantly evidenced and would need further investigation.

Other organic acid and medium chain fatty acids showed *in vitro* study reductions (Greene *et al.*, 2022); however, the efficacy in *in vivo* pilot and commercial scale studies were not performed, as there were negative bird performance effects in treatment groups. Ferric tyrosine added to the feed at 0.02, 0.05 and 0.20 g/kg can reduce *Campylobacter* in the ceca at 42 days up to 3 log (Currie *et al.*, 2018), but other studies showed lower reductions (i.e. 2 log reduction) (Khattak *et al.*, 2018).

#### Probiotics: measure of effectiveness

The use of in-feed probiotic additives as a measure to control *Campylobacter* in birds at the primary production level was addressed in 19 papers assessed by the experts. Probiotic bacteria are thought to reduce pathogenic bacteria (including *Campylobacter*) in the gut through competition for shared attachment sites in the mucosa (Lu and Walker, 2001) or through production of antimicrobial

metabolites (Oelschlaeger, 2010; Neal-McKinney *et al.*, 2012). The overall lasting reduction effect by probiotic treatment on *Campylobacter* presence was inconsistent across the literature. Administration of feed additives at specific stages of growth demonstrated some reduction of *Campylobacter* presence at midgrowth stage (day 18–21) (Ghareeb *et al.*, 2012); however, those reductions were not sustained to the end of production. Some pilot studies demonstrated that administration of *Lactobacillus casei* overexpresses myosin-cross-reactive antigen (LC) in the feed and reduced *C. jejuni* colonization in the cecum, ileum and jejunum, by more than 1 log CFU/g when compared to the no-probiotic control group (Tabashsum *et al.*, 2020). The strain was able to generate bioactive compounds including conjugated linoleic acid. Commercial or large-scale trials would be required to demonstrate applicability to large-scale farming. The literature predominantly reported on *C. jejuni* strains. Probiotic administration did not demonstrate an effect in reducing *C. coli* levels in caeca (Mortada *et al.*, 2020).

#### • Plant-based additives: measures of effectiveness

Plant-based molecules added to feed and water vary in type, concentration of administration and timing of administration. Twenty-five papers addressed plant-based additives as a strategy for reducing *Campylobacter* within the lower gut of birds. Molecules such as *trans*-cinnamaldehyde, eugenol, carvacrol, and thymol demonstrated *Campylobacter* reduction of up to 4 logs *in vitro* (Kollanoor Johny *et al.*, 2010) but to a lower extent (approximately a 2 log reduction) *in vivo* (Allaoua *et al.*, 2022). In addition, other plant-molecules, such as glycyrrhiza glabra (licorice) extract, beta-resorcylic acid, and thymol were tested in pilot *in vivo* studies, showing limited efficacy in the reduction of *Campylobacter* in caecal/faecal counts (~1–2 log reduction) (Wagle *et al.*, 2017; Ibrahim *et al.*, 2020; Epps *et al.* 2015).

#### • Other additives and combination of additives: measure of effectiveness

In papers applying multiple strategies, it is difficult to assess the specific impact of individual components, and the overall effect cannot be attributed to the administration of an additive molecule. For example, up to 1 log CFU/g fewer *Campylobacter* were recovered from broilers administered either direct-fed *Bacillus* culture, prebiotic, refined functional carbohydrates (including mannooligosaccharides (MOS), b-glucans, and mannose), or symbiotic when compared to the untreated control group (Froebel *et al.*, 2020). Refined functional carbohydrates, including MOS, b-glucans and mannose, are important to the pathogen inhibiting functionality of prebiotics (Oyofo *et al.*, 1989; Spring *et al.*, 2000). These carbohydrates are thought to bind to bacterial surface adhesins thereby inactivating the bacteria's ability to infect the host (Fernandez *et al.*, 2002; Walker *et al.*, 2018).

Furthermore, the reported reduction effect of *Campylobacter* using a combination of in-feed additives (microencapsulated propionic, sorbic acids and pure botanicals) at different concentrations administered at various stages of growth resulted in a synergistic action between organic acids and pure botanicals (Grilli *et al.*, 2013). Greene *et al.* (2022) reported the anti-*Campylobacter* properties of nine compounds belonging to organic acids, medium chain fatty acids and essential oils in *in vitro* experiments; however, it was concluded that while the tested compounds are effective against *Campylobacter* in laboratory model studies, the negative effects on broiler performance metrics deem the approach ineligible for use *in vivo*.

Further large-scale *in vivo* studies are required to adequately evaluate the efficacy of a multi-additive approach to reducing *Campylobacter* in the lower gut of birds at the end stage of primary production.

The field of other natural additives is expanding, and further work is required to evaluate the impacts on *Campylobacter* reduction in experimental, pilot and large-scale commercial trials. Other additive candidates include whey powder, which when applied in feed, may affect the physico-chemical integrity of intestinal digesta and improve the balance of intestinal microbiota (Tsiouris *et al.*, 2020; Wilson *et al.*, 2018) However, the existing evidence showed that the effect on the reduction of *Campylobacter* in caecal contents was limited based on studies conducted under both experimental and field settings.

#### • Scalability, applicability, and geographical representation

The scalability and applicability of in-feed and in-water additives is limited at this time. Smaller scale *in vivo* studies were conducted in some investigations; however, large-scale commercial-level studies are required to assess the efficacy of interventions at scale.

Primary-source pilot studies were predominantly conducted in one region. Multiregional and commercial-scale studies were not identified in the assessed literature, limiting extrapolation of findings beyond the pilot study site.

#### Conclusions

The evidence for clear reductions to *Campylobacter* colonization in the lower gut of birds at various stages of primary production was inconsistent. The use of caprylic acid (in-feed) demonstrated promise in pilot studies, yet further commercial-scale evaluation is needed to assess broad-scale application. With the evidence available, probiotic and plant-based in-feed additives showed limited and inconsistent reductions to *Campylobacter* in pilot studies.

To fully assess additives as an intervention strategy at the primary production

stage, important considerations for assessment are:

- fully addressing the start and end of the additive administration and timing of measured reduction levels;
- addressing whether measured reductions are calculated on naturally contaminated birds or challenged birds; and
- compulsorily measuring intervention impact at the end of the rearing cycle in commercial scale conditions to appropriately assess the true reduction of *Campylobacter*.

### 2.2 CONTROL MEASURES DURING PROCESSING

Experts reviewed 141 papers from the literature survey related to *Campylobacter* control during broiler processing. These papers include both primary studies and systematic reviews, listed in Annex 4.2.

In addition to control measures developed for the primary production stage, a range of processing steps and interventions during the primary and secondary processing stages have been developed to reduce *Campylobacter* contamination on/in broiler chicken carcasses and parts. Such intervention measures include general hygienic improvements, technological advancement in processing operations, and decontamination strategies applied throughout or at specific processing steps. In this section, the effectiveness of different processing steps and interventions to reduce the prevalence and/or concentration of *Campylobacter* were critically reviewed considering the quality of evidence, practicability of the evaluated measures, and the geographical representation.

#### • Effectiveness of good hygiene practices

Good hygienic practices (GHP) are important in minimizing *Campylobacter* contamination during processing. Once *Campylobacter* is introduced in the slaughter line, it can spread to the chicken meat, especially at the defeathering and evisceration steps (Cools *et al.*, 2005). Some *Campylobacter* strains may also survive after cleaning and disinfection in a slaughter environment, thus persisting longer and becoming a source of cross-contamination over time (García-Sánchez *et al.*, 2017). The implementation and compliance of GHPs as well as following Hazard Analysis and Critical Control Points (HACCP) principles at the processing establishment is a systematic way to maintain cleanliness, control the presence and levels of *Campylobacter*, and minimize contamination during chicken processing. The rigorous implementation of systematic approaches can significantly enhance the integrity and wholesomeness of a food safety management system in broiler

processing, thereby leading to the reduction of *Campylobacter* contamination in broiler chicken products and a subsequent decrease in associated public health risks linked to chicken consumption (EFSA Panel on Biological Hazards, 2011).

The effectiveness of good hygiene practices at the processing establishment is in part dependent upon incoming flock prevalence and concentration of *Campylobacter* in the gastrointestinal tract and on the bird. Elevated levels of *Campylobacter* on the exterior and within the intestinal contents of broilers can exert greater pressure on a good hygiene management system. A study investigating risk factors for *Campylobacter* contamination on broiler carcasses after slaughter found a significant correlation between the presence and quantity of *Campylobacter* in caecal contents and on birds arriving at the processing establishment (Hue *et al.*, 2010). Similar finding was evidenced in an EU-wide baseline survey of *Campylobacter* on broiler carcasses, revealing an approximately thirtyfold increase in the probability of obtaining a *Campylobacter*-positive carcass post slaughter from a batch of live birds colonized with the pathogen (EFSA, 2010). Good hygiene practices alone during slaughter and processing will have limited impact on reduction levels if incoming bird contamination is high.

#### • Effectiveness of a multihurdle approach

Employing a combination of processing effects that include physical and/ or chemical interventions can enhance the impact of Campylobacter control measures. It is common to employ a multihurdle approach for a synergistic reduction of Campylobacter contamination in chicken processing. For example, processing steps such as scalding and chilling of carcasses have been shown to decrease the number of Campylobacter (Rasschaert et al., 2020). In some countries, decontaminants, such as chlorine, added to the chilling water also effectively reduce the total bacterial load on carcasses (Bucher et al., 2015). Furthermore, following an implementation of crust freezing combined with logistic slaughter, a lower prevalence was observed in fresh, chilled broiler meat in Denmark. Recently published research has evaluated the cumulative effect of combining two or more chemical and/or physical interventions in a risk assessment, which revealed that the multihurdle approach can advance the control of Campylobacter contamination in chicken products (Dogan et al., 2019). Additionally, alternative approaches in processing were evaluated. The ultrasonication of chicken drumsticks was more effective in combination with chemical processing aids, compared to chemical treatments used on their own (Koolman et al., 2014a; 2014b), or in combination with a vacuum and a water resonance system (Vetchapitak et al., 2020). Greater effectiveness was not guaranteed with the use of crust freezing and UV in combination, giving no additional benefit over the use of crust freezing alone (Haughton et al., 2012). Some may not yet be practical in a commercial plant or when used alone (Gunther, Phillips and Sommers, 2016; Isohanni and Lyhs, 2009).

#### 2.2.1 Effectiveness of specific processing procedures

#### • Logistic slaughter

Logistic slaughter employs a method of scheduling known colonized flocks to be slaughtered only after non-colonized flocks, at the end of a slaughter day or week. Logistic slaughter can reduce *Campylobacter* cross-contamination between flocks through the exposure to contaminated equipment (e.g. processing lines), as evidenced by a study where fewer positive samples were recovered from birds slaughtered after *Campylobacter* negative flocks compared to *Campylobacter* positive flocks (Reich *et al.*, 2008). Norway, Iceland and Denmark have fully implemented logistic slaughter in national *Campylobacter* control programmes (Rosenquist *et al.*, 2009). However, the impact of logistic slaughter on reducing the risk of *Campylobacter* infections in humans has not been fully evaluated (Nauta *et al.*, 2009).

#### • Qualitative and quantitative target incentives

Setting qualitative and quantitative targets for *Campylobacter* at the processing plant may be used to optimize process control. Setting quantitative limits (e.g. *Campylobacter* colonies per gram of fresh meat) could provide incentive to reduce the occurrence of highly contaminated carcasses. In New Zealand, setting a quantitative regulatory limit for chilled chicken carcasses was introduced to reduce *Campylobacter* levels in processed chicken. This measure was also used to evaluate the effectiveness of implemented processing measures on reduction levels of *Campylobacter* (Wagenaar, French and Havelaar, 2013). These practices are further supported by a risk assessment of *Campylobacter* in chicken. Reducing the levels of *Campylobacter* on meat prior to reaching the consumer would be effective in reducing campylobacteriosis incidence (Nauta *et al.*, 2009).

#### • Scalding

Reduction in *Campylobacter* contamination on chicken carcasses can occur at the scalding step through physical removal and cell inactivation under elevated temperatures (hard scalding). Based on a recent systematic review and meta-analysis, hard scalding with higher temperature exerts a consistent log reduction (mean: 1.85 log CFU, 95 percent CI [1.60, 2.09]) (Dogan *et al.*, 2022), while soft scalding with lower temperature was associated with a lower log reduction. Decreased prevalence of *Campylobacter* on broiler carcasses associated with scalding has been consistently observed (OR: 0.18, 95 percent CI [0.11, 0.30]). Furthermore, a multihurdle approach of scalding, and using chemical additives, such as chloride,

had a greater reduction in both concentration and prevalence at the end of this processing step (Berrang and Dickens, 2000; Berrang *et al.*, 2003; Berrang, Windham and Meinersmann, 2011). These findings were primarily based on studies measuring the naturally occurring contamination of *Campylobacter* before and after the scalding step in the pilot or processing scale.

#### • Defeathering and evisceration

The defeathering and evisceration steps during processing are associated with an increased prevalence and concentration of carcass Campylobacter contamination, implying these are critical points in broiler processing for Campylobacter contamination of downstream products (Dogan et al., 2022; Gruntar et al., 2015; Zweifel, Althaus and Stephan, 2015; Huang et al., 2017; Reich et al., 2008; Gichure et al., 2022). The increase in carcass contamination after evisceration is due to the leakage of faecal contents from pressure applied onto carcasses and visceral rupture. Primary studies suggest an overall increase in Campylobacter concentration (mean: 0.88 log CFU, 95 percent CI [0.54, 1.23]) and prevalence (OR: 5.8, 95 percent CI [1.15, 29.42]) following the defeathering step, with similar increases observed after evisceration. However, variation was observed between individual studies. The variation may be related to differences in initial contamination levels and procedural differences between processing establishment (Luning et al., 2011; Zweifel, Althaus and Stephan, 2015; Pacholewicz et al., 2016). Interventions such as organic acid injection into the cloaca and cloacal plugging have been tested and seem to be promising approaches (Berrang et al., 2001; Berrang, Smith and Hinton 2006a, 2006b; Berrang, Windham and Meinersmann, 2011). However, the feasibility of their applications on a broader scale for the industry is uncertain.

#### Chilling during processing

During immersion chilling, broiler carcasses are dipped into cold water with or without aids to cool down processed carcasses as rapidly as possible, thereby minimizing microbial growth in the cold-water bath. The reports on the effect of immersion chilling on *Campylobacter* contamination of carcasses vary. In some studies, a decrease in *Campylobacter* concentration was consistently observed regardless of the use of processing aids to cool the carcasses (0.70–4.12 log CFU) (Dogan *et al.*, 2022). Immersion chilling without chemical additives decreased *Campylobacter* 1.25 log CFU (95 percent CI [0.96, 1.55]) – a mean of 1.95 log CFU, (95 percent CI [1.46, 2.45]) when the chemical additive chloride was added, indicating that the addition of chlorine may control the concentration of *Campylobacter* on carcasses post chilling without chemical aids with an increase

in prevalence observed (OR: 1.55, 95 percent CI [0.81, 2.96]) (Dogan *et al.*, 2022). The increased prevalence may be explained by a cross-contamination effect of dipping carcasses into contaminated chilling water. Still, immersion chilling with chemical additives may compensate for the water contamination effect and minimally reduced prevalence (OR: 0.71, 95 percent CI [0.28, 0.61]) (Dogan *et al.*, 2022). Adding chemical processing aids to chilling water can control the transmission of *Campylobacter* across carcasses during the immersion chilling process.

As an alternative approach, air chilling presents some advantages over immersion chilling in product quality, thus gaining popularity in broiler processing. In terms of *Campylobacter* control, most studies reported a slight decrease in *Campylobacter* concentration associated with air chilling, showing a mean log reduction of 1 log CFU based on studies conducted at pilot or processing scale. However, most studies reported no significant effect based on studies evaluating challenged or naturally occurring contamination at either pilot or processing scale (Dogan *et al.*, 2022).

As with control measures to reduce *Campylobacter* during chilling, immersion chilling is more effective than air chilling in terms of reducing the exterior surface concentration of *Campylobacter* on broiler chicken, regardless of the use of processing aids.

#### • Alternative Campylobacter reduction approaches

Freezing and ionizing (gamma) irradiation have been shown to be effective in reducing *Campylobacter* contamination on chicken carcasses but are not recommended as the primary method of pathogen control (Umaraw *et al.*, 2017; Gellynck *et al.*, 2008; Gunther *et al.*, 2019; Cox and Pavic, 2010). Experimental trials have shown that crust freezing of chicken meat portions may reduce *Campylobacter* by up to 1.5 log CFU with minimal impact on the colour of treated skin (Haughton *et al.*, 2012). In addition, flesh temperature may be maintained above – 2 °C (Burfoot *et al.*, 2016) consistent with the EU target for selling fresh, rather than frozen, meat.

Steam, ultrasonication, high-intensity light pulse, visible light, and UV-C among others have shown promise either at the laboratory or pilot scale, but their impact is unknown at commercial scale (Sorro *et al.*, 2020; Chun *et al.*, 2010; Koolman *et al.*, 2014a, 2014b; Vetchapitak *et al.*, 2020; Haughton *et al.*, 2012; Moazzami, Fernström and Hansson, 2021).

20
#### Chemical additives as a Campylobacter reduction approach

In the countries and regions where chemical processing aids are permitted for use in chicken processing, common examples include chlorine, chlorine derivatives (e.g. chlorine dioxide, acidified sodium chlorite), peroxyacetic acids (PAA), organic acids (e.g. lactic acid [LA]), and quaternary ammonium compound (e.g. cetylpyridinium chloride [CPC]) (Ebel, Williams and Tameru, 2019). These chemical aids are added to water used for spray and/or immersion, with the aim of reducing Campylobacter on broiler carcasses and parts. Studies have documented variation in the effectiveness of chemical processing aids, reporting factors such as initial contamination, amount of organic matter on the bird and carcass and chemical application conditions responsible for variation in *Campylobacter* reduction levels. Among the different classes of chemical processing aids, chlorine may be less effective as an intervention in decontaminating carcasses containing *Campylobacter* in comparison to PAA, during the pre-chill, chill or post-chill immersion or spray application step (Cano, Meneses and Chaves, 2021). Organic acid such as lactic acid is a chemical control option considered during post-chill treatments, as it has fewer occupational concerns. Peroxyacetic acids and lactic acid demonstrate comparable reduction levels in Campylobacter contamination in immersion studies, while the comparison for spray applications is inconclusive. Effectiveness of PAA was comparable to or higher than that of CPC, depending on specific product type and treatment conditions (Cano, Meneses and Chaves, 2021; Dogan et al., 2022). Unlike physical processing effects, the effectiveness of chemical processing aids has mostly been investigated in challenge studies conducted in laboratory settings or pilot plants, which may introduce uncertainty when extrapolating the impact of such aids on naturally occurring contamination at a broader processing scale.

Marinades are commonly used to enhance the sensory characteristics of meat, yet studies have also demonstrated their potential to reduce *Campylobacter* on chicken meat. Marinades typically contain water, salt, spices, sugar, wine, and acidic ingredients like vinegar, lemon, or lime juice (Lopes, Da Silva and Tondo, 2022). The use of marinades for chicken cuts such as breast, fillets and wings reduced *Campylobacter* contamination from 0.5 to 3 log CFU (Park, Hong and Yoon, 2014; Birk *et al.*, 2010; Zakarienė *et al.*, 2015; Isohanni *et al.*, 2010), with some studies reporting reductions of up to 6 log CFU (Thanissery and Smith, 2014). pH was identified as the most significant factor influencing *Campylobacter* inactivation, although other factors such as specific ingredients' compounds and storage temperature also played a role. Currently available evidence was collected from studies conducted in laboratory settings using inoculated chicken cuts and parts.

### 2.3 CONTROLS DURING POST-PROCESSING, RETAIL AND CONSUMER LEVELS

The experts reviewed 77 papers, of which 19 were strictly related to post-processing stages of chicken meat production. Control measures were evaluated and highlighted in the next section (Annex 4.3).

### 2.3.2 Effectiveness of specific retail/post-processing measures

#### • Freezing of meat product

Laboratory-based studies found that freezing chicken meat for 1 week at 20 °C significantly decreased *Campylobacter* counts on meat by 1.73 log CFU/g (Bolton *et al.*, 2014). Continued storage resulted in additional reductions, but the greatest reductions occurred after the first week. Other studies have also reported freezing at – 20 °C resulting in reductions of 2.2–2.6 logs within 9 days of storage (El-Shibiny, Connerton, P. and Connerton, I., 2009). Variability was noted in the survival of strains examined, suggesting that *Campylobacter* strains have varying levels of freezing resistance. Where studies of contaminated cooked product were undertaken, low level contamination (50 CFU/g) could not be detected on 70 percent of frozen samples after storage for 7 days and was detectable at reduced levels on 92.5 percent of samples inoculated with 500 CFU/g, suggesting that the organism can survive freezing (Eideh and AlQadiri, 2011), posing a potential health risk where cross-contamination of cooked product occurs. Freezing is an effective control measure for reducing *Campylobacter*; however, this measure does not reduce levels to zero.

#### Cross-contamination awareness

Prevention of cross-contamination by replacing cutlery and cutting boards used for food preparation after handling raw chicken and the prevention of hand contact with other foodstuffs after handling contaminated chicken meat considerably reduced *Campylobacter* contamination to other foods (Jong *et al.*, 2008; Verhoeff-Bakkenes *et al.*, 2008). Washing cutting boards and cutlery in hot water and detergent have greater effects on reducing *Campylobacter* contamination than using cold water (Jong *et al.*, 2008). Reducing cross-contamination between materials contaminated with *Campylobacter* and other foods that may not receive further treatment (e.g. salad preparation) is especially important to preventing risks of infection (Verhoeff-Bakkenes *et al.*, 2008).

#### Consumer education

A large study showed that international recommendations to consumers is difficult. Particularly in terms of the lack of easy "rule-of-thumb" guidelines or tools to check safe cooking conditions at home. The study concluded that regional differences in contamination levels, food culture and regional economies are challenges to developing international recommendations that ensure consumer food safety specific to *Campylobacter* infection risk that could be easily implemented by each country (Langsrud *et al.*, 2020).

Evaluation of web-based information interventions designed and tested on participant motivation and intentions to cook more safely was conducted (Nauta *et al.*, 2008). The intervention supported by the emotion "disgust" was selected as the most promising information intervention. Alone, this intervention has no effect on infection risk but together with instructions for meal preparation, it was found that the risk decreased (Nauta *et al.*, 2008).

#### Control measures in commercial kitchens

Examination of *Campylobacter* contamination in restaurant kitchens reported that the prevalence of Campylobacter increased after raw meat handling. The prevalence of Campylobacter on food contact surfaces also increased, and in some cases the organism was present on food surfaces prior to raw meat preparation (Lai et al., 2021). Washing of boards and utensils in hot water and detergent was reported to reduce contamination, but poor washing hygiene practices did not reduce contamination (Lai et al., 2021; Jong et al., 2008). For commercial kitchens, poor sanitation offers greater opportunities for cross-contamination and Campylobacter transmission during preparation of chicken dishes (Lai et al., 2021). Use of neutral electrolyzed water (NEW), quaternary ammonium (QUAT), and lactic acid-based solutions on the surfaces of inoculated scarred polypropylene and wooden food-cutting boards reduced C. jejuni levels (Al-Qadiri et al., 2016). The reduction was dependent on contact time and the type of surface. These agents were found to cause reductions of 3 log on boards within the first minute of application and  $> 5 \log$  reductions within 5 minutes of application. Reduction levels were lower on boards made of wood compared to polypropylene.

In conclusion, the use of good hygiene practices and appropriate training of food handlers in food preparation kitchens that cater to mass events (restaurants) are essential to reduce the risk of cross-contamination between raw meats and finished (cooked products), and ultimately reduce the exposure of *Campylobacter* to consumers. In addition, the use of appropriate sanitizing agents and washing procedures for chopping boards, surfaces and kitchen tools is essential to reduce the risk of cross-contamination and human exposure.

#### • Heating

Studies using naturally contaminated chicken meat provide a more realistic model of treatments or interventions (e.g. cooking or processing) (Vaz *et al.* 2021).

Cooking at 70 °C was found to reduce *Campylobacter* levels by > 1.5 logs within 30 minutes (Vaz *et al.*, 2021), but the organism survival persisted following initial reductions indicating that a core temperature is necessary to reduce *Campylobacter* to negligible levels. Heating to a core temperature > 74 °C has been shown to be required to reduce *Campylobacter* levels in cooked products (Sampers *et al.*, 2010; FAO and WHO, 2011).

#### • Marinades post-processing

As with marination at broiler meat processing stage, marination of chicken products post-processing (e.g. consumer-based level) can reduce consumer risk of *Campylobacter* exposure. Some studies suggest *Campylobacter* reductions of 1–2 logs were possible using marinades, but such reductions are dependent on the concentration and type of ingredients used in the marinade (e.g. tartaric acid, oregano, thymol; essential oils). In one study, the effects of thyme oil extract in the marinade appeared to enhance the killing effect of *Campylobacter* resulting in a > 1 log reduction in inoculated chicken wings held at 4 °C for 168 h (Zakarienė *et al.*, 2015).

The spread of tartaric acid solutions (2, 4, 6 and 10 percent) on chicken meat was found to reduce the level of *Campylobacter* by 0.5 to 2 logs after storage for 3 days at 4 °C, but the studies were carried out *in vitro* and significant variation was observed among the 14 strains exposed to this acid (Birk *et al.*, 2010).

Apple-based edible films containing carvacrol and cinnamaldehyde were evaluated for bactericidal activity against antibiotic resistant and susceptible *C. jejuni* strains on chicken meat. Retail chicken breast samples inoculated with antibiotic resistant strains and a susceptible strain were wrapped in apple films containing cinnamaldehyde or carvacrol at 0.5 percent, 1.5 percent and 3 percent concentrations, and incubated at 4 or 23 °C for 72 h. The antimicrobial films exhibited dose- and temperature-dependent bactericidal activity against all tested strains. The films with cinnamaldehyde were more effective than carvacrol films, and reductions at 23 °C were greater than those at 4 °C. Populations of all strains were reduced to below detection level at 23 °C after 72 h exposure. At 4 °C cinnamaldehyde exposure reduced *Campylobacter* levels between 0.2 to 2.5 logs and 1.8 to 6.0 logs at concentrations of 1.5 percent and 3.0 percent, respectively (Mild *et al.*, 2011).

Interactions of *Campylobacter* with the microbial community throughout pre- and post-production processing are not yet well identified. A better understanding of the interactions between *Campylobacter* and the surrounding microbiota may further guide control measures support information used in qualitative and quantitative microbial risk assessments (QMRA) (Chintoan-Uta *et al.*, 2022; Ijaz *et al.*, 2018).

### Review of the Code of Practice

The expert committee was tasked with a review of the current code of practice with the view of assessing whether the current code of practice requires updates in light of new information or research or changes that require addressing or revision.

The experts evaluated the impact or efficacy of control measures relevant to *Campylobacter* spp. in the broiler production chain, noting the variability of the impact reviewed and recommended revisions to the *Guidelines for the Control of Campylobacter and Salmonella in Chicken Meat (CXG 78-2011)*, paragraphs 1 to 115, based on the currently available evidence (Annex 3).

### Concluding remarks

Since the first publications in the early 1970s showed an association between "Vibrios" (now named *Campylobacters*) in poultry and stool samples of humans with enteritis, tremendous efforts have been made to understand the infection from both the pathogen and the animal host side. Within broiler chicken production systems, various interventions throughout the process have been developed and evaluated. Although some successes have been reported, effective *Campylobacter* interventions are still very limited. Continuous research, innovation and collaboration among stakeholders are essential to further improve intervention strategies. The currently available interventions in broiler chicken pre- and post-production systems have been reviewed in this report as control measures for *Campylobacter*. As most of the burden of campylobacteriosis in humans is caused by *C. jejuni* and *C. coli*, the majority of available evidence is primarily reported on *C. jejuni* and *C. coli* species. For this report, the experts concentrated on the review and assessment of intervention measures to these two species.

Several interventions have been proposed and tested, including the use of bacteriophages, vaccination, probiotics, and disinfectants, among others. Studies have also explored the effects of processing techniques, such as air chilling versus water chilling.

Overall, the results of these studies have been mixed, with some interventions showing promise in reducing the prevalence of *Campylobacter* in specific stages of production, while others have had little to no effect. It is important to note

that interventions that have been successful in one study may not necessarily be effective in other settings, as the prevalence and characteristics of *Campylobacter* can vary widely between different processing facilities.

The broader interventions at the primary production stage include precise biosecurity measures, such as strict hygiene practices and control of wildlife access to chicken farms. Targeted hazard-based interventions in the post-processing stages of production and good agriculture practices and hygienic practices for the general control of foodborne pathogens throughout the production system may be effective for the reduction of *Campylobacter*; however, there are no interventions that **specifically** control *Campylobacter* on meat.

The experts recognize further data gaps exist and that some new technologies may offer promising approaches to reducing *Campylobacter* in pre- and post-production of chicken meat. Further global changes to the industry, growing of global populations, climate change, and increased demand for animal protein in specific regions will guide the need for further control measure assessments.

In conclusion, while interventions to reduce *Campylobacter* in chicken processing have shown some promise, further research is needed to identify effective interventions that can be implemented on a large scale. It is also important to note that a multifaceted approach involving both pre- and post-harvest interventions is necessary to effectively control *Campylobacter* in chicken production systems, and an implementation of robust monitoring and surveillance systems to track *Campylobacter* throughout the system is crucial to evaluating the effectiveness of interventions and identifying emerging trends.

### Intervention summary synopsis

A number of interventions were identified through primary production to post-processing stages. The expert consultation concluded the following in each stage of production:

## Assessment of primary production interventions for the control of *Campylobacter*

#### Biosecurity and management approaches

- Using strict biosecurity measures (hygiene practices and sanitation) can enhance the control of *Campylobacter* in broiler flocks.
- Risk factors for *Campylobacter* contamination at primary production establishments, such as partial depopulation, litter management, down period length, proximity to other livestock and slaughter age can help guide intervention strategies.

#### Vaccination-based approaches

- Currently, there are no commercial vaccines for *Campylobacter* readily available for any stage of primary production, and vaccination studies were limited to *C. jejuni* only.
- Several potential vaccine candidates are in the proof-of-concept phase but cannot yet be considered as an intervention.
- Some vaccines induced a cellular or humoral response in the chicken host, but this did not always translate to reduced caecal colonization by *Campylobacter* in pilot studies.

#### Bacteriophage-based approaches

- There are currently no commercial products available for use in primary production.
- The effects of phage therapy may be transitory and prone to resistance.

#### Feed and water additives

#### **Organic acids**

- In feed, short- and medium-chain fatty acids, and in particular, caprylic acid show promise as feed additives in reducing *Campylobacter* in pilot studies.
- In water, organic acids reduced *Campylobacter* in caecal/faecal specimens at the end of the primary production period; however, the effects were not sustained to the end of production in pilot studies.

#### Probiotics

• In feed, there is inconsistent evidence on the efficacy of probiotics as an intervention for reducing *Campylobacter* in broilers at primary production level.

#### Plant-based additives

• In feed, the efficacy of some plant-based molecules in *in vivo* pilot studies showed limited reduction of *Campylobacter* in caecal/faecal specimens at the end of the primary production period.

#### Assessment of processing interventions for the control of Campylobacter

- Good hygiene practices during processing are important in minimizing *Campylobacter* contamination on meat.
- The effectiveness of interventions during processing is dependent upon the incoming flock prevalence and concentration of *Campylobacter* in the gastrointestinal tract and on the bird.
- The impact of processing practices can be enhanced by a combination of a multihurdle approach, processing effects, physical, and/or chemical interventions.

#### **Processing procedures**

- Logistic slaughter scheduling can reduce *Campylobacter* cross-contamination.
- Scalding reduces the carcass surface concentration and prevalence of *Campylobacter*; however, the effect depends on the temperature, and dilution effect.

- Defeathering and evisceration may increase both prevalence and concentration of *Campylobacter* on carcasses.
- Immersion chilling can reduce the carcass concentration of *Campylobacter*; however, this is dependent on the initial *Campylobacter* load of incoming birds.
- In combination with processing aids, immersion chilling may reduce the carcass prevalence of *Campylobacter*.
- Air chilling may reduce concentration of *Campylobacter*, but the efficacy of air chilling in reducing the prevalence of *Campylobacter* when used without other processing aids is inconclusive.

#### Physical

- Irradiation is effective at eliminating *Campylobacter* on meat.
- Freezing meat reduces the concentration of *Campylobacter*.
- Steam, ultrasonication, high-intensity light pulse, visible light, UV-C and other technologies have shown promise either at the laboratory or pilot scale, but their impact is unknown at commercial scale.

#### Chemical

- Processing aids such as chlorine derivatives, peroxyacetic acids, and organic acids added to water used for washing and or dipping may reduce *Campylobacter* on carcasses.
- Some marination ingredients have shown reductions in *Campylobacter* on meat.

# Assessment of post-processing interventions for the control of *Campylobacter*

- Thorough cooking is effective at eliminating *Campylobacter* on meat.
- The application of good hygiene practices is important in reducing *Campylobacter* on meat.
- Freezing meat reduces the concentration of *Campylobacter*.
- Some marination ingredients have shown reductions in *Campylobacter* on meat.



### References

- Agunos, A., Waddell, L., Léger, D. & Taboada, E. 2014. A systematic review characterizing on-farm sources of *Campylobacter* spp. for broiler chickens. *PLoS ONE*, 9(8): e104905. https://doi.org/10.1371/journal. pone.0104905
- Al Hakeem, W.G., Fathima, S., Shanmugasundaram, R. & Selvaraj, R.K. 2022. *Campylobacter jejuni* in poultry: pathogenesis and control strategies. *Microorganisms*, 10(11): 2134. https://doi.org/10.3390/ microorganisms10112134
- Al-Qadiri, H.M., Ovissipour, M., Al-Alami, N., Govindan, B.N., Shiroodi,
   S.G. & Rasco, B. 2016. Efficacy of neutral electrolyzed water, quaternary ammonium and lactic acid-based solutions in controlling microbial contamination of food cutting boards using a manual spraying technique. *Journal of Food Science*, 81(5). https://doi.org/10.1111/1750-3841.13275
- Allaoua, M., Bonnafé, E., Etienne, P., Noirot, V., Gabarrou, J.-F., Castinel, A., Pascal, G. et al. 2022. A carvacrol-based product reduces *Campylobacter jejuni* load and alters microbiota composition in the caeca of chickens. *Journal of Applied Microbiology*, 132(6): 4501–4516. https://doi. org/10.1111/jam.15521
- Allen, V.M., Weaver, H., Ridley, A.M., Harris, J.A., Sharma, M., Emery, J., Sparks, N., Lewis, M. & Edge, S. 2008. Sources and spread of thermophilic *Campylobacter* spp. during partial depopulation of broiler chicken flocks. *Journal of Food Protection*, 71(2): 264–270. https://doi. org/10.4315/0362-028X-71.2.264

- Alpigiani, I., Abrahantes, J.C., Michel, V., Huneau-Salaün, A., Chemaly, M., Keeling, L.J., Gervelmeyer, A. et al. 2017. Associations between animal welfare indicators and *Campylobacter* spp. in broiler chickens under commercial settings: A case study. *Preventive Veterinary Medicine*, 147: 186–193. https://doi.org/10.1016/j.prevetmed.2017.09.005
- Babacan, O., Harris, S.A., Pinho, R.M., Hedges, A., Jørgensen, F. & Corry, J.E.L. 2020. Factors affecting the species of *Campylobacter* colonizing chickens reared for meat. *Journal of Applied Microbiology*, 129(4): 1071–1078. https://doi.org/10.1111/jam.14651
- Bailey, M.A., Bourassa, D.V., Krehling, J.T., Munoz, L., Chasteen, K.S., Escobar, C. & Macklin, K.S. 2022. Effects of common litter management practices on the prevalence of *Campylobacter jejuni* in broilers. *Animals*, 12(7): 858. https://doi.org/10.3390/ani12070858
- Bailey, M.A., Taylor, R.M., Brar, J.S., Corkran, S.C., Velásquez, C., Novoa Rama, E., Oliver, H.F. & Singh, M. 2019. Prevalence and antimicrobial resistance of *Campylobacter* from antibiotic-free broilers during organic and conventional processing. *Poultry Science*, 98(3): 1447–1454. https:// doi.org/10.3382/ps/pey486
- Battersby, T., Whyte, P. & Bolton, D.J. 2016. The pattern of *Campylobacter* contamination on broiler farms; external and internal sources. *Journal of Applied Microbiology*, 120(4): 1108–1118. https://doi.org/10.1111/ jam.13066
- Berrang, M., Meinersmann, R., Buhr, R., Reimer, N., Philips, R. & Harrison, M. 2003. Presence of *Campylobacter* in the respiratory tract of broiler carcasses before and after commercial scalding. *Poultry Science*, 82(12): 1995–1999. https://doi.org/10.1093/ps/82.12.1995
- Berrang, M.E. & Dickens, J.A. 2000. Presence and level of *Campylobacter* spp. on broiler carcasses throughout the processing plant. *Journal of Applied Poultry Research*, 9(1): 43–47. https://doi.org/10.1093/japr/9.1.43
- Berrang, M.E., Buhr, R.J., Cason, J.A. & Dickens, J.A. 2001. Broiler carcass contamination with *Campylobacter* from feces during defeathering. *Journal of Food Protection*, 64(12): 2063–2066. https://doi. org/10.4315/0362-028X-64.12.2063
- Berrang, M.E., Meinersmann, R.J., Cox, N.A. & Fedorka-Cray, P.J. 2011. Application of chlorine dioxide to lessen bacterial contamination during broiler defeathering. *Journal of Applied Poultry Research*, 20(1): 33–39. https://doi.org/10.3382/japr.2010-00178

- Berrang, M.E., Smith, D.P. & Hinton, A. 2006a. Application of distilled white vinegar in the Cloaca to counter the increase in *Campylobacter* numbers on broiler skin during feather removal. *Journal of Food Protection*, 69(2): 425–427. https://doi.org/10.4315/0362-028X-69.2.425
- Berrang, M.E., Smith, D.P. & Hinton, A. 2006b. Organic acids placed into the Cloaca to reduce *Campylobacter* contamination of broiler skin during defeathering. *Journal of Applied Poultry Research*, 15(2): 287–291. https:// doi.org/10.1093/japr/15.2.287
- Berrang, M.E., Windham, W.R. & Meinersmann, R.J. 2011. Campylobacter, Salmonella, and Escherichia coli on broiler carcasses subjected to a high pH scald and low pH postpick chlorine dip. Poultry Science, 90(4): 896– 900. https://doi.org/10.3382/ps.2010-00900
- Birk, T., Grønlund, A.C., Christensen, B.B., Knøchel, S., Lohse, K. & Rosenquist, H. 2010. Effect of organic acids and marination ingredients on the survival of *Campylobacter jejuni* on Meat. *Journal of Food Protection*, 73(2): 258–265. https://doi.org/10.4315/0362-028X-73.2.258
- Bolton, D., Meredith, H., Walsh, D. & McDowell, D. 2014. Poultry food safety control interventions in the domestic kitchen. *Journal of Food Safety*, 34(1): 34–41. https://doi.org/10.1111/jfs.12092
- Borck Høg, B., Sommer, H.M., Larsen, L.S., Sørensen, A.I.V., David, B., Hofshagen, M. & Rosenquist, H. 2016. Farm specific risk factors for *Campylobacter* colonisation in Danish and Norwegian broilers. *Preventive Veterinary Medicine*, 130: 137–145. https://doi.org/10.1016/ j.prevetmed.2016.04.002
- Bucher, O., Waddell, L., Greig, J. & Smith, B.A. 2015. Systematic review-metaanalysis of the effect of chilling on *Campylobacter* spp. during primary processing of broilers. *Food Control*, 56: 211–217. https://doi.org/10.1016/ j.foodcont.2015.03.032
- Buckley, A.M., Wang, J., Hudson, D.L., Grant, A.J., Jones, M.A., Maskell, D.J.
   & Stevens, M.P. 2010. Evaluation of live-attenuated *Salmonella* vaccines expressing *Campylobacter* antigens for control of *C. jejuni* in poultry. *Vaccine*, 28(4): 1094–1105. https://doi.org/10.1016/j.vaccine.2009.10.018
- Bull, S.A., Thomas, A., Humphrey, T., Ellis-Iversen, J., Cook, A.J., Lovell, R.
  & Jorgensen, F. 2008. Flock health indicators and *Campylobacter* spp. in commercial housed broilers reared in Great Britain. *Applied and Environmental Microbiology*, 74(17): 5408–5413. https://doi.org/10.1128/ AEM.00462-08

- Burfoot, D., Hall, J., Nicholson, K., Holmes, K., Hanson, C., Handley, S.
  & Mulvey, E. 2016. Effect of rapid surface cooling on *Campylobacter* numbers on poultry carcasses. *Food Control*, 70: 293–301. https://doi.org/10.1016/j.foodcont.2016.05.041
- Cano, C., Meneses, Y. & Chaves, B.D. 2021. Application of peroxyacetic acid for decontamination of raw poultry products and comparison to other commonly used chemical antimicrobial interventions: a review. *Journal* of Food Protection, 84(10): 1772–1783. https://doi.org/10.4315/JFP-21-107
- Chinivasagam, H.N., Estella, W., Maddock, L., Mayer, D.G., Weyand, C., Connerton, P.L. & Connerton, I.F. 2020. Bacteriophages to control *Campylobacter* in commercially farmed broiler chickens, in Australia. *Frontiers in Microbiology*, 11: 632. https://doi.org/10.3389/ fmicb.2020.00632
- Chintoan-Uta, C., Wisedchanwet, T., Glendinning, L., Bremner, A., Psifidi, A., Vervelde, L., Watson, K., Watson, M. & Stevens, M.P. 2020. Role of cecal microbiota in the differential resistance of inbred chicken lines to colonization by *Campylobacter jejuni*. *Applied and Environmental Microbiology*, 86(7): e02607-19. https://doi.org/10.1128/AEM.02607-19
- Chowdhury, S., Themudo, G.E., Sandberg, M. & Ersbøll, A.K. 2013. Spatiotemporal patterns of *Campylobacter* colonization in Danish broilers. *Epidemiology and Infection*, 141(5): 997–1008. https://doi.org/10.1017/ S0950268812001446
- Chun, H.H., Kim, J.Y., Lee, B.D., Yu, D.J. & Song, K.B. 2010. Effect of UV-C irradiation on the inactivation of inoculated pathogens and quality of chicken breasts during storage. *Food Control*, 21(3): 276–280. https://doi.org/10.1016/j.foodcont.2009.06.006
- Clark, J.D., Oakes, R.D., Redhead, K., Crouch, C.F., Francis, M.J., Tomley, F.M. & Blake, D.P. 2012. Eimeria species parasites as novel vaccine delivery vectors: Anti-*Campylobacter jejuni* protective immunity induced by Eimeria tenella-delivered CjaA. *Vaccine*, 30(16): 2683–2688. https://doi. org/10.1016/j.vaccine.2012.02.002
- Colles, F.M., Cain, R.J., Nickson, T., Smith, A.L., Roberts, S.J., Maiden, M.C.J., Lunn, D. & Dawkins, M.S. 2016. Monitoring chicken flock behaviour provides early warning of infection by human pathogen *Campylobacter*. *Proceedings of the Royal Society B: Biological Sciences*, 283(1822): 20152323. https://doi.org/10.1098/rspb.2015.2323

- Colles, F.M., Dingle, K.E., Cody, A.J. & Maiden, M.C.J. 2008. Comparison of *Campylobacter* populations in wild geese with those in starlings and free-range poultry on the same farm. *Applied and Environmental Microbiology*, 74(11): 3583–3590. https://doi.org/10.1128/AEM.02491-07
- Colles, F.M., Hedges, S.J., Dixon, R., Preston, S.G., Thornhill, P., Barfod, K.K., Gebhardt-Henrich, S.G. et al. 2021. Parallel sequencing reveals *Campylobacter* spp. in commercial meat chickens less than 8 days old. *Applied and Environmental Microbiology*, 87(23): e01060-21. https://doi. org/10.1128/AEM.01060-21
- Cools, I., Uyttendaele, M., Cerpentier, J., D'Haese, E., Nelis, H.J. & Debevere, J. 2005. Persistence of *Campylobacter jejuni* on surfaces in a processing environment and on cutting boards. *Letters in Applied Microbiology*, 40(6): 418–423. https://doi.org/10.1111/j.1472-765X.2005.01694.x
- **Cox, J.M. and Pavic, A.,** 2010. Advances in enteropathogen control in poultry production. *Journal of Applied Microbiology*, 108(3): 745–755. 10.1111/ j.1365-2672.2009.04456.x
- Crotta, M., Georgiev, M. & Guitian, J. 2017. Quantitative risk assessment of *Campylobacter* in broiler chickens – Assessing interventions to reduce the level of contamination at the end of the rearing period. *Food Control*, 75: 29–39. https://doi.org/10.1016/j.foodcont.2016.12.024
- Currie, D., Green, M., Dufailu, O.A., Matthaios, P., Soultanas, P., McCartney, E., Lester, H. et al. 2018. Dietary supplementation with ferric tyrosine improves zootechnical performance and reduces caecal *Campylobacter* spp. load in broilers. *British Poultry Science*, 59(6): 646–653. https://doi. org/10.1080/00071668.2018.1507015
- D'Angelantonio, D., Scattolini, S., Boni, A., Neri, D., Di Serafino, G., Connerton, P., Connerton, I. *et al.* 2021. Bacteriophage therapy to reduce colonization of *Campylobacter jejuni* in broiler chickens before slaughter. *Viruses*, 13(8): 1428. https://doi.org/10.3390/v13081428
- De Haan, C.P.A., Kivistö, R., Hakkinen, M., Rautelin, H. & Hänninen, M.
   L. 2010. Decreasing trend of overlapping multilocus sequence types between human and chicken *Campylobacter jejuni* isolates over a decade in Finland. *Applied and Environmental Microbiology*, 76(15): 5228–5236. https://doi.org/10.1128/AEM.00581-10

- De Los Santos, F.S., Hume, M., Venkitanarayanan, K., Donoghue, A.M., Hanning, I., Slavik, M.F., Aguiar, V.F. et al. 2010. Caprylic acid reduces enteric Campylobacter colonization in market-aged broiler chickens but does not appear to alter cecal microbial populations. Journal of Food Protection, 73(2): 251–257. https://doi.org/10.4315/0362-028X-73.2.251.
- Delabouglise, A., Nguyen-Van-Yen, B., Thanh, N.T.L., Xuyen, H.T.A., Tuyet, P.N., Lam, H.M. & Boni, M.F. 2019. Poultry population dynamics and mortality risks in smallholder farms of the Mekong river delta region. *BMC Veterinary Research*, 15(1): 205. https://doi.org/10.1186/s12917-019-1949-y
- Di Marcantonio, L., Marotta, F., Vulpiani, M.P., Sonntag, Q., Iannetti, L., Janowicz, A., Serafino, G.D., Di Giannatale, E. & Garofolo, G. 2022. Investigating the cecal microbiota in broiler poultry farms and its potential relationships with animal welfare. *Research in Veterinary Science*, 144: 115–125. https://doi.org/10.1016/j.rvsc.2022.01.020
- Dogan, O.B., Aditya, A., Ortuzar, J., Clarke, J. & Wang, B. 2022. A systematic review and meta-analysis of the efficacy of processing stages and interventions for controlling *Campylobacter* contamination during broiler chicken processing. *Comprehensive Reviews in Food Science and Food Safety*, 21(1): 227–271. 10.1111/1541-4337.12860
- Dogan, O.B., Clarke, J., Mattos, F. & Wang, B. 2019. A quantitative microbial risk assessment model of *Campylobacter* in broiler chickens: Evaluating processing interventions. *Food Control*, 100: 97–110. https://doi. org/10.1016/j.foodcont.2019.01.003
- **Ebel, E.D., Williams, M.S. & Tameru, B.** 2019. Relatedness of *Salmonella* contamination frequency on chicken carcasses and parts when processed in the same establishment. *Food Control*, 100: 198–203. https://doi.org/10.1016/j.foodcont.2018.12.031
- EFSA (European Food Safety Authority). 2010. Analysis of the baseline survey on the prevalence of *Campylobacter* in broiler batches and of *Campylobacter* and *Salmonella* on broiler carcasses, in the EU, 2008. *EFSA Journal*, 8(8): 1522. https://doi.org/10.2903/j.efsa.2010.1522
- **EFSA Panel on Biological Hazards (BIOHAZ).** 2011. Scientific opinion on *Campylobacter* in broiler meat production: control options and performance objectives and/or targets at different stages of the food chain. *EFSA Journal* 2011, 9(4): 2105. https://doi.org/10.2903/ j.efsa.2011.2105

- **EFSA.** 2020. Update and review of control options for *Campylobacter* in broilers at primary production. *EFSA Journal 2020*, 18(4): 6090.
- Eideh, A.M.F. & Al-Qadiri, H.M. 2011. Effect of refrigerated and frozen storage on the survival of *Campylobacter jejuni* in cooked chicken meat breast. *Journal of Food Science*, 76(1). https://doi.org/10.1111/j.1750-3841.2010.01924.x
- El-Shibiny, A., Connerton, P. & Connerton, I. 2009. Survival at refrigeration and freezing temperatures of *Campylobacter coli* and *Campylobacter jejuni* on chicken skin applied as axenic and mixed inoculums. *International Journal of Food Microbiology*, 131(2–3): 197–202. https://doi.org/10.1016/j.ijfoodmicro.2009.02.024
- El-Shibiny, A., Connerton, P.L. & Connerton, I.F. 2005. Enumeration and diversity of *Campylobacters* and Bacteriophages isolated during the rearing cycles of free-range and organic chickens. *Applied and Environmental Microbiology*, 71(3): 1259–1266. https://doi.org/10.1128/ AEM.71.3.1259-1266.2005
- El-Shibiny, A., Scott, A., Timms, A., Metawea, Y., Connerton, P. & Connerton, I. 2009. Application of a group II *Campylobacter* bacteriophage to reduce strains of *Campylobacter jejuni* and *Campylobacter coli* colonizing broiler chickens. *Journal of Food Protection*, 72(4): 733–740. https://doi. org/10.4315/0362-028X-72.4.733
- Epps, S.V.R., Harvey, R.B., Byrd, J.A., Petrujkić, B.T., Sedej, I., Beier, R.C., Phillips, T.D. *et al.* 2015. Comparative effect of thymol or its glucose conjugate, thymol-β- d -glucopyranoside, on *Campylobacter* in avian gut contents. *Journal of Environmental Science and Health, Part B*, 50(1): 55–61. https://doi.org/10.1080/03601234.2015.965634E
- FAO & WHO (World Health Organization). 1999. Codex Alimentarius Commission. Report of the thirty second session of the Codex Committee on Food Hygiene. Washington DC, 29 November-4 December 1999. Rome. https://www.fao.org/fao-who-codexalimentarius/sh-proxy/ en/?lnk=1&url=https%253A%252F%252Fworkspace.fao.org%252Fsites %252Fcodex%252FMeetings%252FCX-712-32%252FAl01\_13e.pdf
- FAO & WHO. 2008. Risk assessment of *Campylobacter* spp. in broiler chickens. Interpretative summary. Microbiological Risk Assessment Series, No. 11. Rome. https://www.fao.org/3/a1468e/a1468e.pdf

- FAO & WHO. 2009a. Salmonella and Campylobacter in chicken meat. Meeting report. Microbiological Risk Assessment Series, No. 19. Rome. https:// www.fao.org/3/i1133e/i1133e.pdf
- FAO & WHO. 2009b. Risk assessment of *Campylobacter* spp. in broiler chickens. Technical report. Microbiological Risk Assessment Series, No. 12. Rome. https://www.fao.org/3/a1469e/a1469e.pdf
- FAO & WHO. 2011. Codex *Guidelines for the Control of Campylobacter and* Salmonella in Chicken Meat (CXG 78-2011). Rome. https://www.fao.org/ fao-who-codexalimentarius/sh-proxy/de/?lnk=1&url=https%253A% 252F%252Fworkspace.fao.org%252Fsites%252Fcodex%252FStandards% 252FCXG%2B78-2011%252FCXG\_078e.pdf
- Fernandez, F., Hinton, M. & Gils, B.V. 2002. Dietary mannan-oligosaccharides and their effect on chicken caecal microflora in relation to *Salmonella* Enteritidis colonization. *Avian Pathology*, 31(1): 49–58. https://doi. org/10.1080/03079450120106000
- Fischer, S., Kittler, S., Klein, G. & Glünder, G. 2013. Impact of a single phage and a phage cocktail application in broilers on reduction of *Campylobacter jejuni* and development of resistance. *PLoS ONE*, 8(10): e78543. https://doi.org/10.1371/journal.pone.0078543
- Froebel, L.K., Jalukar, S., Lavergne, T.A., Coufal, C.D. & Duong, T. 2020. Administration of direct-fed Bacillus cultures and refined functional carbohydrates to broiler chickens improves growth performance and promotes positive shifts in gastrointestinal microbiota. *Journal* of Applied Poultry Research, 29(4): 765–774. https://doi.org/10.1016/ j.japr.2020.06.004
- Furuta, M., Nasu, T., Umeki, K., Hoang Minh, D., Honjoh, K.-I. & Miyamoto, T. 2017. Characterization and application of Lytic bacteriophages against *Campylobacter jejuni* isolated from poultry in Japan. *Biocontrol Science*, 22(4): 213–221. https://doi.org/10.4265/bio.22.213
- García-Sánchez, L., Melero, B., Jaime, I., Hänninen, M.-L., Rossi, M. & Rovira, J. 2017. *Campylobacter jejuni* survival in a poultry processing plant environment. *Food Microbiology*, 65: 185–192. https://doi.org/10.1016/ j.fm.2017.02.009
- Gellynck, X., Messens, W., Halet, D., Grijspeerdt, K., Hartnett, E. & Viaene, J. 2008. Economics of reducing *Campylobacter* at different levels within the Belgian Poultry Meat Chain. *Journal of Food Protection*, 71(3): 479–485. https://doi.org/10.4315/0362-028X-71.3.479

- Georgiev, M., Beauvais, W. & Guitian, J. 2017. Effect of enhanced biosecurity and selected on-farm factors on *Campylobacter* colonization of chicken broilers. *Epidemiology and Infection*, 145(3): 553–567. https://doi. org/10.1017/S095026881600251X
- Ghareeb, K., Awad, W.A., Mohnl, M., Porta, R., Biarnés, M., Böhm, J. & Schatzmayr, G. 2012. Evaluating the efficacy of an avian-specific probiotic to reduce the colonization of *Campylobacter jejuni* in broiler chickens. *Poultry Science*, 91(8): 1825–1832. https://doi.org/10.3382/ ps.2012-02168
- Gharib Naseri, K., Rahimi, S. & Khaki, P. 2012. Comparison of the effects of probiotic, organic acid and medicinal plant on *Campylobacter jejuni* challenged broiler chickens. *Journal of Agricultural Science and Technology*, 14(7): 1485–1496.
- Gichure, J.N., Kamau Njage, P.M., Wambui, J.M., Dykes, G.A., Buys, E.M.
   & Coorey, R. 2022. Systematic-review and meta-analysis on effect of decontamination interventions on prevalence and concentration of *Campylobacter* spp. during primary processing of broiler chickens. *Food Microbiology*, 102: 103923. https://doi.org/10.1016/j.fm.2021.103923
- Gloanec, N., Dory, D., Quesne, S., Béven, V., Poezevara, T., Keita, A., Chemaly, M. & Guyard-Nicodème, M. 2022. Impact of DNA prime/protein boost vaccination against *Campylobacter jejuni* on immune responses and gut microbiota in chickens. *Vaccines*, 10(6): 981. https://doi.org/10.3390/ vaccines10060981
- Golden, C.E. & Mishra, A. 2020. Prevalence of Salmonella and Campylobacter spp. in alternative and conventionally produced chicken in the United States: A systematic review and meta-analysis. Journal of Food Protection, 83(7): 1181–1197. https://doi.org/10.4315/JFP-19-538
- Greene, G., Koolman, L., Whyte, P., Lynch, H., Coffey, A., Lucey, B., Egan, J., O'Connor, L. & Bolton, D. 2022. The efficacy of organic acid, medium chain fatty acid and essential oil based broiler treatments; *in vitro* anti-*Campylobacter jejuni* activity and the effect of these chemical-based treatments on broiler performance. *Journal of Applied Microbiology*, 132(1): 687–695. https://doi.org/10.1111/jam.15204
- Griekspoor, P., Colles, F.M., McCarthy, N.D., Hansbro, P.M., Ashhurst-Smith, C., Olsen, B., Hasselquist, D., Maiden, M.C.J. & Waldenström, J. 2013. Marked host specificity and lack of phylogeographic population structure of *Campylobacter jejuni* in wild birds. *Molecular Ecology*, 22(5): 1463– 1472. https://doi.org/10.1111/mec.12144

- Grilli, E., Vitari, F., Domeneghini, C., Palmonari, A., Tosi, G., Fantinati, P., Massi, P. & Piva, A. 2013. Development of a feed additive to reduce caecal *Campylobacter jejuni* in broilers at slaughter age: from *in vitro* to *in vivo*, a proof of concept. *Journal of Applied Microbiology*, 114(2): 308–317. https://doi.org/10.1111/jam.12053
- Gruntar, I., Biasizzo, M., Kušar, D., Pate, M. & Ocepek, M. 2015. Campylobacter jejuni contamination of broiler carcasses: Population dynamics and genetic profiles at slaughterhouse level. Food Microbiology, 50: 97–101. https://doi.org/10.1016/j.fm.2015.03.007
- Guerin, M.T., Martin, S.W., Reiersen, J., Berke, O., McEwen, S.A., Friðriksdóttir, V., Bisaillon, J.-R. & Lowman, R. 2008. Temperaturerelated risk factors associated with the colonization of broiler-chicken flocks with *Campylobacter* spp. in Iceland, 2001–2004. *Preventive Veterinary Medicine*, 86(1–2): 14–29. https://doi.org/10.1016/ j.prevetmed.2008.02.015
- Gunther, N.W., Abdul-Wakeel, A., Scullen, O.J. & Sommers, C. 2019. The evaluation of gamma irradiation and cold storage for the reduction of *Campylobacter jejuni* in chicken livers. *Food Microbiology*, 82: 249–253. https://doi.org/10.1016/j.fm.2019.02.014
- Gunther, N.W., Phillips, J.G. & Sommers, C. 2016. The effects of 405-nm visible light on the survival of *Campylobacter* on chicken skin and stainless steel. *Foodborne Pathogens and Disease*, 13(5): 245–250. https://doi. org/10.1089/fpd.2015.2084
- Hald, T., Aspinall, W., Devleesschauwer, B., Cooke, R., Corrigan, T., Havelaar, A.H., Gibb, H.J., Torgerson, P.R., Kirk, M.D., Angulo, F.J. & Lake, R.J., 2016. World Health Organization estimates of the relative contributions of food to the burden of disease due to selected foodborne hazards: a structured expert elicitation. *PLoS ONE*, 11(1): e0145839. https://doi.org/10.1371/journal.pone.0145839
- Hammerl, J.A., Jäckel, C., Alter, T., Janzcyk, P., Stingl, K., Knüver, M.T. & Hertwig, S. 2014. Reduction of *Campylobacter jejuni* in broiler chicken by successive application of group II and group III phages. *PLoS ONE*, 9(12): e114785. https://doi.org/10.1371/journal.pone.0114785
- Hansson, I., Engvall, E.O., Vågsholm, I. & Nyman, A. 2010. Risk factors associated with the presence of *Campylobacter*-positive broiler flocks in Sweden. *Preventive Veterinary Medicine*, 96(1–2): 114–121. https://doi. org/10.1016/j.prevetmed.2010.05.007

- Hansson, I., Tamminen, L.-M., Frosth, S. Fernström, L.-L., Emanuelson, U. & Boqvist, S. 2021. Occurrence of *Campylobacter* spp. in Swedish calves, common sequence types and antibiotic resistance patterns. *Journal of Applied Microbiology*, 130: 2111–2122. https://doi.org/10.1111/jam.14914
- Hasan, Md.M., Talukder, S., Mandal, A.K., Tasmim, S.T., Parvin, Mst.S., Ali, Md.Y., Sikder, M.H. & Islam, Md.T. 2020. Prevalence and risk factors of *Campylobacter* infection in broiler and cockerel flocks in Mymensingh and Gazipur districts of Bangladesh. *Preventive Veterinary Medicine*, 180: 105034. https://doi.org/10.1016/j.prevetmed.2020.105034
- Haughton, P.N., Lyng, J., Cronin, D., Fanning, S. & Whyte, P. 2012. Effect of crust freezing applied alone and in combination with ultraviolet light on the survival of *Campylobacter* on raw chicken. *Food Microbiology*, 32(1): 147–151. https://doi.org/10.1016/j.fm.2012.05.004
- Havelaar, A.H., Kirk, M.D., Torgerson, P.R., Gibb, H.J., Hald, T., Lake, R.J., Praet, N. et al. 2015. World Health Organization global estimates and regional comparisons of the burden of foodborne disease in 2010. PLOS Medicine, 12(12): e1001923. https://doi.org/10.1371/journal.pmed.1001923
- Henry, I., Reichardt, J., Denis, M. & Cardinale, E. 2011. Prevalence and risk factors for *Campylobacter* spp. in chicken broiler flocks in Reunion Island (Indian Ocean). *Preventive Veterinary Medicine*, 100(1): 64–70. https:// doi.org/10.1016/j.prevetmed.2011.03.007
- Hertogs, K., Haegeman, A., Schaumont, D., Gelaude, P., De Zutter, L., Dewulf, J., Heyndrickx, M. & Rasschaert, G. 2021. Contamination sources and transmission routes for *Campylobacter* on (mixed) broiler farms in Belgium, and comparison of the gut microbiota of flocks colonized and uncolonized with *Campylobacter*. *Pathogens*, 10(1): 66. https://doi. org/10.3390/pathogens10010066
- Higham, L.E., Scott, C., Akehurst, K., Dring, D., Parnham, A., Waterman,
  M. & Bright, A. 2018. Effects of financial incentives and cessation of thinning on prevalence of *Campylobacter*: a longitudinal monitoring study on commercial broiler farms in the UK. *Veterinary Record*, 183(19): 595–595. https://doi.org/10.1136/vr.104823
- Higham, L.E., Scott, C., Akehurst, K., Dring, D., Parnham, A., Waterman,
  M. & Bright, A. 2018. Effects of financial incentives and cessation of thinning on prevalence of *Campylobacter*: a longitudinal monitoring study on commercial broiler farms in the UK. *Veterinary Record*, 183(19): 595–595. https://doi.org/10.1136/vr.104823

- Huang, J., Zang, X., Zhai, W., Guan, C., Lei, T. & Jiao, X. 2017. Quantitative analysis of *Campylobacter* spp. contamination in chicken slaughtering lines by "label tracking method" in eastern China. *Food Control*, 80: 67–73. https://doi.org/10.1016/j.foodcont.2017.03.052
- Hue, O., Le Bouquin, S., Laisney, M.-J., Allain, V., Lalande, F., Petetin, I., Rouxel, S. *et al.* 2010. Prevalence of and risk factors for *Campylobacter* spp. contamination of broiler chicken carcasses at the slaughterhouse. *Food Microbiology*, 27(8): 992–999. https://doi.org/10.1016/ j.fm.2010.06.004
- Hwang, H. & Singer, R.S. 2020. Survey of the U.S. broiler industry regarding pre- and postharvest interventions targeted to mitigate *Campylobacter* contamination on broiler chicken products. *Journal of Food Protection*, 83(7): 1137–1148. https://doi.org/10.4315/JFP-19-527
- Iannetti, L., Neri, D., Santarelli, G.A., Cotturone, G., Podaliri Vulpiani, M., Salini, R., Antoci, S. et al. 2020. Animal welfare and microbiological safety of poultry meat: Impact of different at-farm animal welfare levels on at-slaughterhouse Campylobacter and Salmonella contamination. Food Control, 109: 106921. https://doi.org/10.1016/ j.foodcont.2019.106921
- Ibrahim, D., Sewid, A.H., Arisha, A.H., Abd El-fattah, A.H., Abdelaziz, A.M., Al-Jabr, O.A. & Kishawy, A.T.Y. 2020. Influence of Glycyrrhiza glabra extract on growth, gene expression of gut integrity, and *Campylobacter jejuni* colonization in broiler chickens. *Frontiers in Veterinary Science*, 7: 612063. https://doi.org/10.3389/fvets.2020.612063
- Ijaz, U.Z., Sivaloganathan, L., McKenna, A., Richmond, A., Kelly, C., Linton, M., Stratakos, A.Ch. *et al.* 2018. Comprehensive longitudinal microbiome analysis of the chicken cecum reveals a shift from competitive to environmental drivers and a window of opportunity for *Campylobacter. Frontiers in Microbiology*, 9: 2452. https://doi. org/10.3389/fmicb.2018.02452
- Isohanni, P., Alter, T., Saris, P. & Lyhs, U. 2010. Wines as possible meat marinade ingredients possess antimicrobial potential against *Campylobacter. Poultry Science*, 89(12): 2704–2710. https://doi. org/10.3382/ps.2009-00521
- Isohanni, P.M.I. & Lyhs, U. 2009. Use of ultraviolet irradiation to reduce *Campylobacter jejuni* on broiler meat. *Poultry Science*, 88(3): 661–668. https://doi.org/10.3382/ps.2008-00259

- Jeni, R.E., Dittoe, D.K., Olson, E.G., Lourenco, J., Corcionivoschi, N., Ricke, S.C. & Callaway, T.R. 2021. Probiotics and potential applications for alternative poultry production systems. *Poultry Science*, 100(7): 101156. https://doi.org/10.1016/j.psj.2021.101156
- Jeon, B., Saisom, T., Sasipreeyajan, J. & Luangtongkum, T. 2022. Liveattenuated oral vaccines to reduce *Campylobacter* colonization in poultry. *Vaccines*, 10(5): 685. https://doi.org/10.3390/vaccines10050685
- Jong, A.E.I., Verhoeff-Bakkenes, L., Nauta, M.J. & De Jonge, R. 2008. Crosscontamination in the kitchen: effect of hygiene measures. *Journal of Applied Microbiology*, 105(2): 615–624. https://doi.org/10.1111/j.1365-2672.2008.03778.x
- Jonsson, M.E., Chriél, M., Norström, M. & Hofshagen, M. 2012. Effect of climate and farm environment on *Campylobacter* spp. colonisation in Norwegian broiler flocks. *Preventive Veterinary Medicine*, 107(1–2): 95–104. https://doi.org/10.1016/j.prevetmed.2012.05.002
- Khattak, F., Paschalis, V., Green, M., Houdijk, J.G.M., Soultanas, P. & Mahdavi, J. 2018. TYPLEX<sup>®</sup> Chelate, a novel feed additive, inhibits *Campylobacter jejuni* biofilm formation and cecal colonization in broiler chickens. *Poultry Science*, 97(4): 1391–1399. https://doi.org/10.3382/ps/pex413
- Kittler, S., Fischer, S., Abdulmawjood, A., Glünder, G. & Klein, G. 2013. Effect of bacteriophage application on *Campylobacter jejuni* loads in commercial broiler flocks. *Applied and Environmental Microbiology*, 79(23): 7525–7533. https://doi.org/10.1128/AEM.02703-13
- Kollanoor Johny, A., Darre, M.J., Donoghue, A.M., Donoghue, D.J.
  & Venkitanarayanan, K. 2010. Antibacterial effect of transcinnamaldehyde, eugenol, carvacrol, and thymol on *Salmonella* Enteritidis and *Campylobacter jejuni* in chicken cecal contents in vitro. *Journal of Applied Poultry Research*, 19(3): 237–244. https://doi.org/10.3382/japr.2010-00181
- Koolman, L., Whyte, P., Meade, J., Lyng, J. & Bolton, D. 2014a. A combination of chemical and ultrasonication treatments to reduce *Campylobacter jejuni* on raw poultry. *Food and Bioprocess Technology*, 7(12): 3602–3607. https://doi.org/10.1007/s11947-014-1370-2
- Lai, H., Tang, Y., Ren, F., Li, Z., Li, F., Cui, C., Jiao, X. & Huang, J. 2021. An investigation into the critical factors influencing the spread of *Campylobacter* during chicken handling in commercial kitchens in China. *Microorganisms*, 9(6): 1164. https://doi.org/10.3390/ microorganisms9061164

- Langsrud, S., Sørheim, O., Skuland, S.E., Almli, V.L., Jensen, M.R., Grøvlen, M.S., Ueland, Ø. & Møretrø, T. 2020. Cooking chicken at home: Common or recommended approaches to judge doneness may not assure sufficient inactivation of pathogens. *PLoS ONE*, 15(4): e0230928. https:// doi.org/10.1371/journal.pone.0230928
- Łaniewski, P., Kuczkowski, M., Chrząstek, K., Woźniak, A., Wyszyńska, A., Wieliczko, A. & Jagusztyn-Krynicka, E.K. 2014. Evaluation of the immunogenicity of *Campylobacter jejuni* CjaA protein delivered by *Salmonella* enterica sv. Typhimurium strain with regulated delayed attenuation in chickens. *World Journal of Microbiology and Biotechnology*, 30(1): 281–292. https://doi.org/10.1007/s11274-013-1447-5
- Lassen, B., Helwigh, B., Kahl Petersen, C. & Ellis-Iversen, J. 2022. Systematic review of products with potential application for use in the control of *Campylobacter* spp. in organic and free-range broilers. *Acta Veterinaria Scandinavica*, 64(1): 24. https://doi.org/10.1186/s13028-022-00644-z
- Layton, S.L., Morgan, M.J., Cole, K., Kwon, Y.M., Donoghue, D.J., Hargis, B.M. & Pumford, N.R. 2011. Evaluation of Salmonella-vectored Campylobacter peptide epitopes for reduction of Campylobacter jejuni in broiler chickens. Clinical and Vaccine Immunology, 18(3): 449–454. https://doi.org/10.1128/CVI.00379-10
- Li, S., He, Y., Mann, D.A. & Deng, X. 2021. Global spread of *Salmonella* Enteritidis via centralized sourcing and international trade of poultry breeding stocks. *Nature Communications*, 12(1): 5109. https://doi. org/10.1038/s41467-021-25319-7
- Lopes, S.M., Da Silva, D.C. & Tondo, E.C. 2022. Bactericidal effect of marinades on meats against different pathogens: a review. *Critical Reviews in Food Science and Nutrition*, 62(27): 7650–7658. https://doi.org/10.1080/104083 98.2021.1916734
- Lu, L. & Walker, W.A. 2001. Pathologic and physiologic interactions of bacteria with the gastrointestinal epithelium. *The American Journal of Clinical Nutrition*, 73(6): 1124S-1130S. https://doi.org/10.1093/ajcn/73.6.1124S
- Lu, T., Marmion, M., Ferone, M., Wall, P. & Scannell, A.G.M. 2021. On farm interventions to minimise *Campylobacter* spp. contamination in chicken. *British Poultry Science*, 62(1): 53–67. https://doi.org/10.1080/00071668.2 020.1813253

- Luangtongkum, T., Morishita, T.Y., Ison, A.J., Huang, S., McDermott, P.F. & Zhang, Q. 2006. Effect of conventional and organic production practices on the prevalence and antimicrobial resistance of *Campylobacter* spp. in poultry. *Applied and Environmental Microbiology*, 72(5): 3600–3607. doi: 10.1128/AEM.72.5.3600-3607.2006
- Luning, P.A., Jacxsens, L., Rovira, J., Osés, S.M., Uyttendaele, M. & Marcelis, W.J. 2011. A concurrent diagnosis of microbiological food safety output and food safety management system performance: Cases from meat processing industries. *Food Control*, 22(3–4): 555–565. https://doi. org/10.1016/j.foodcont.2010.10.003
- Mauri, M., Sannasiddappa, T.H., Vohra, P., Corona-Torres, R., Smith, A.A., Chintoan-Uta, C., Bremner, A. *et al.* 2021. Multivalent poultry vaccine development using protein glycan coupling technology. *Microbial Cell Factories*, 20(1): 193. https://doi.org/10.1186/s12934-021-01682-4
- McKenna, A., Ijaz, U.Z., Kelly, C., Linton, M., Sloan, W.T., Green, B.D., Lavery, U. et al. 2020. Impact of industrial production system parameters on chicken microbiomes: mechanisms to improve performance and reduce *Campylobacter. Microbiome*, 8(1): 128. https://doi.org/10.1186/s40168-020-00908-8
- Mendes, Â.J., Santos-Ferreira, N.L., Costa, F.M., Lopes, E.P., Freitas-Silva,
   J., Inácio, Â.S., Moreira, F. & Martins Da Costa, P. 2020. External
   contamination of broilers by *Campylobacter* spp. increases from the farm
   to the slaughterhouse. *British Poultry Science*, 61(4): 400–407. https://doi.
   org/10.1080/00071668.2020.1736264
- Metcalf, J.H., Donoghue, A.M., Venkitanarayanan, K., Reyes-Herrera, I., Aguiar, V.F., Blore, P.J. & Donoghue, D.J. 2011. Water administration of the medium-chain fatty acid caprylic acid produced variable efficacy against enteric *Campylobacter* colonization in broilers 1, 2. *Poultry Science*, 90(2): 494–497. https://doi.org/10.3382/ps.2010-00891
- Meunier, M., Guyard-Nicodème, M., Vigouroux, E., Poezevara, T., Béven, V., Quesne, S., Amelot, M. et al. 2018. A DNA prime/protein boost vaccine protocol developed against *Campylobacter jejuni* for poultry. *Vaccine*, 36(16): 2119–2125. https://doi.org/10.1016/j.vaccine.2018.03.004
- Mild, R.M., Joens, L.A., Friedman, M., Olsen, C.W., McHugh, T.H., Law, B. & Ravishankar, S. 2011. Antimicrobial edible apple films inactivate antibiotic resistant and susceptible *Campylobacter jejuni* strains on chicken breast. *Journal of Food Science*, 76(3). https://doi.org/10.1111/ j.1750-3841.2011.02065.x

- Moazzami, M., Fernström, L.-L. & Hansson, I. 2021. Reducing *Campylobacter jejuni, Enterobacteriaceae* and total aerobic bacteria on transport crates for chickens by irradiation with 265-nm ultraviolet light (UV–C LED). *Food Control*, 119: 107424. https://doi.org/10.1016/ j.foodcont.2020.107424
- Mortada, M., Cosby, D.E., Shanmugasundaram, R. & Selvaraj, R.K. 2020. In vivo and in vitro assessment of commercial probiotic and organic acid feed additives in broilers challenged with *Campylobacter coli. Journal of Applied Poultry Research*, 29(2): 435–446. https://doi.org/10.1016/ j.japr.2020.02.001
- Näther, G., Alter, T., Martin, A. & Ellerbroek, L. 2009. Analysis of risk factors for *Campylobacter* species infection in broiler flocks. *Poultry Science*, 88(6): 1299–1305. https://doi.org/10.3382/ps.2008-00389
- Natsos, G., Mouttotou, N.K., Magiorkinis, E., Ioannidis, A., Rodi-Burriel, A., Chatzipanagiotou, S. & Koutoulis, K.C. 2020. Prevalence of and risk factors for *Campylobacter* spp. colonization of broiler chicken flocks in Greece. *Foodborne Pathogens and Disease*, 17(11): 679–686. https://doi. org/10.1089/fpd.2020.2795
- Nauta, M. J., Fischer, A. R., Van Asselt, E. D., De Jong, A. E., Frewer, L. J., & De Jonge, R. 2008. Food safety in the domestic environment: the effect of consumer risk information on human disease risks. *Risk Analysis: An International Journal*, 28(1): 179–192. 10.1111/j.1539-6924.2008.01012.x
- Nauta, M., Hill, A., Rosenquist, H., Brynestad, S., Fetsch, A., Van Der Logt, P., Fazil, A. et al. 2009. A comparison of risk assessments on *Campylobacter* in broiler meat. *International Journal of Food Microbiology*, 129(2): 107–123. https://doi.org/10.1016/j.ijfoodmicro.2008.12.001
- Neal-McKinney, J.M., Lu, X., Duong, T., Larson, C.L., Call, D.R., Shah, D.H. & Konkel, M.E. 2012. Production of organic acids by probiotic lactobacilli can be used to reduce pathogen load in poultry. *PLoS ONE*, 7(9): e43928. https://doi.org/10.1371/journal.pone.0043928
- **Oelschlaeger, T.A.** 2010. Mechanisms of probiotic actions A review. *International Journal of Medical Microbiology*, 300(1): 57–62. https://doi. org/10.1016/j.ijmm.2009.08.005
- Oyofo, B.A., DeLoach, J.R., Corrier, D.E., Norman, J.O., Ziprin, R.L. & Mollenhauer, H.H. 1989. Prevention of *Salmonella* typhimurium colonization of broilers with D-mannose. *Poultry Science*, 68(10): 1357– 1360. https://doi.org/10.3382/ps.0681357

- Organisation for Economic Co-operation and Development (OECD) & FAO. 2021. OECD-FAO Agricultural Outlook 2021-2030. Paris, OECD Publishing. https://www.fao.org/publications/card/en/c/CB5332EN/
- Pacholewicz, E., Sura Barus, S.A., Swart, A., Havelaar, A.H., Lipman, L.J.A. & Luning, P.A. 2016. Influence of food handlers' compliance with procedures of poultry carcasses contamination: A case study concerning evisceration in broiler slaughterhouses. *Food Control*, 68: 367–378. https://doi.org/10.1016/j.foodcont.2016.04.009
- Park, N.Y., Hong, S.H. & Yoon, K.S. 2014. Effects of commercial marinade seasoning and a natural blend of cultured sugar and vinegar on *Campylobacter jejuni* and *Salmonella* Typhimurium and the texture of chicken breasts. *Poultry Science*, 93(3): 719–727. https://doi.org/10.3382/ ps.2013-03595
- Rasschaert, G., De Zutter, L., Herman, L. & Heyndrickx, M. 2020. *Campylobacter* contamination of broilers: the role of transport and slaughterhouse. *International Journal of Food Microbiology*, 322: 108564. https://doi.org/10.1016/j.ijfoodmicro.2020.108564
- Rawson, T., Dawkins, M.S. & Bonsall, M.B. 2019. A mathematical model of *Campylobacter* dynamics within a broiler flock. *Frontiers in Microbiology*, 10: 1940. https://doi.org/10.3389/fmicb.2019.01940
- Reich, F., Atanassova, V., Haunhorst, E. & Klein, G. 2008. The effects of *Campylobacter* numbers in caeca on the contamination of broiler carcasses with *Campylobacter*. *International Journal of Food Microbiology*, 127(1–2): 116–120. https://doi.org/10.1016/j.ijfoodmicro.2008.06.018
- Reichelt, B., Szott, V., Epping, L., Semmler, T., Merle, R., Roesler, U. & Friese, A. 2022. Transmission pathways of *Campylobacter* spp. at broiler farms and their environment in Brandenburg, Germany. *Frontiers in Microbiology*, 13: 982693. https://doi.org/10.3389/fmicb.2022.982693
- Richards, P.J., Connerton, P.L. & Connerton, I.F. 2019. Phage biocontrol of *Campylobacter jejuni* in chickens does not produce collateral effects on the gut microbiota. *Frontiers in Microbiology*, 10: 476. https://doi. org/10.3389/fmicb.2019.00476
- Rosenquist, H., Boysen, L., Galliano, C., Nordentoft, S., Ethelberg, S. & Borck,
  B. 2009. Danish strategies to control *Campylobacter* in broilers and
  broiler meat: facts and effects. *Epidemiology and Infection*, 137(12): 1742–1750. https://doi.org/10.1017/S0950268809002684

- Rosenquist, H., Nielsen, N.L., Sommer, H.M., Nørrung, B. & Christensen, B.B. 2003. Quantitative risk assessment of human campylobacteriosis associated with thermophilic *Campylobacter* species in chickens. *International Journal of Food Microbiology*, 83(1): 87–103. https://doi. org/10.1016/S0168-1605(02)00317-3
- Rothrock, M.J., Cook, K.L., Warren, J.G. & Sistani, K. 2008. The effect of alum addition on microbial communities in poultry litter. *Poultry Science*, 87(8): 1493–1503. https://doi.org/10.3382/ps.2007-00491
- Russell, K.M., Smith, J., Bremner, A., Chintoan-Uta, C., Vervelde, L., Psifidi, A. & Stevens, M.P. 2021. Transcriptomic analysis of caecal tissue in inbred chicken lines that exhibit heritable differences in resistance to *Campylobacter jejuni*. *BMC Genomics*, 22(1): 411. https://doi. org/10.1186/s12864-021-07748-2
- Sahin, O., Kassem, I.I., Shen, Z., Lin, J., Rajashekara, G. & Zhang, Q. 2015. *Campylobacter* in poultry: ecology and potential interventions. *Avian Diseases*, 59(2): 185–200. https://doi.org/10.1637/11072-032315-Review
- Sampers, I., Habib, I., De Zutter, L., Dumoulin, A. & Uyttendaele, M. 2010. Survival of *Campylobacter* spp. in poultry meat preparations subjected to freezing, refrigeration, minor salt concentration, and heat treatment. *International Journal of Food Microbiology*, 137(2–3): 147–153. https:// doi.org/10.1016/j.ijfoodmicro.2009.11.013
- Sandberg, M., Dahl, J., Lindegaard, L.L. & Pedersen, J.R. 2017. Compliance/ non-compliance with biosecurity rules specified in the Danish Quality Assurance system (KIK) and *Campylobacter*- positive broiler flocks 2012 and 2013. *Poultry Science*, 96(1): 184–191. https://doi.org/10.3382/ps/ pew277
- Schweitzer, P.M., Susta, L., Varga, C., Brash, M.L. & Guerin, M.T. 2021. Demographic, husbandry, and biosecurity factors associated with the presence of *Campylobacter* spp. in small poultry flocks in Ontario, Canada. *Pathogens*, 10(11): 1471. https://doi.org/10.3390/ pathogens10111471
- Seman, M., Gregova, G. & Korim, P. 2020. Comparison of Campylobacter spp. and flock health indicators of broilers in Iceland. Annals of Agricultural and Environmental Medicine, 27(4): 579–584. https://doi.org/10.26444/ aaem/127181

- Sevilla-Navarro, S., Marin, C., Cortés, V., García, C. & Catalá-Gregori, P. 2020. *Campylobacter* prevalence and risk factors associated with exceeding allowable limits in poultry slaughterhouses in Spain. *Veterinary Record*, 186(13): 415–415. https://doi.org/10.1136/vr.105558
- Sheppard, S.K., Colles, F., Richardson, J., Cody, A.J., Elson, R., Lawson, A., Brick, G. et al. 2010. Host association of Campylobacter genotypes transcends geographic variation. Applied and Environmental Microbiology, 76(15): 5269–5277. https://doi.org/10.1128/AEM.00124-10
- Sindiyo, E., Maganga, R., Thomas, K.M., Benschop, J., Swai, E., Shirima, G. & Zadoks, R.N. 2018. Food safety, health management, and biosecurity characteristics of poultry farms in Arusha City, Northern Tanzania, along a gradient of intensification. *The East African Health Research Journal*, 2(2): 168–180. https://doi.org/10.24248/EAHRJ-D-18-00034
- Smith, S., Messam, L.L.McV., Meade, J., Gibbons, J., McGill, K., Bolton, D. & Whyte, P. 2016. The impact of biosecurity and partial depopulation on *Campylobacter* prevalence in Irish broiler flocks with differing levels of hygiene and economic performance. *Infection Ecology & Epidemiology*, 6(1): 31454. https://doi.org/10.3402/iee.v6.31454
- Spring P., Wenk C., Dawson K.A., Newman K.E. 200. The effects of dietary mannanoligosaccharides on cecal parameters and the concentrations of enteric bacteria in the ceca of *Salmonella*-challenged broiler chicks. *Poultry Science*, 79(2): 205–211. https://doi.org/10.1093/ps/79.2.205
- Szott, V., Peh, E., Friese, A., Roesler, U., Kehrenberg, C., Ploetz, M. & Kittler, S. 2022. Antimicrobial effect of a drinking water additive comprising four organic acids on *Campylobacter* load in broilers and monitoring of bacterial susceptibility. *Poultry Science*, 101(12): 102209. https://doi. org/10.1016/j.psj.2022.102209
- Tabashsum, Z., Peng, M., Alvarado-Martinez, Z., Aditya, A., Bhatti, J., Romo, P.B., Young, A. & Biswas, D. 2020. Competitive reduction of poultry-borne enteric bacterial pathogens in chicken gut with bioactive *Lactobacillus casei*. *Scientific Reports*, 10(1): 16259. https://doi. org/10.1038/s41598-020-73316-5
- Tack, D.M., Marder, E.P., Griffin, P.M., Cieslak, P.R., Dunn, J., Hurd, S.,
   Scallan, E. *et al.* 2019. Preliminary incidence and trends of infections with pathogens transmitted commonly through food foodborne diseases active surveillance network, 10 U.S. sites, 2015–2018. *MMWR*. *Morbidity and Mortality Weekly Report*, 68(16): 369–373. https://doi.org/10.15585/mmwr.mm6816a2

- Thanissery, R. & Smith, D.P. 2014. Marinade with thyme and orange oils reduces Salmonella Enteritidis and Campylobacter coli on inoculated broiler breast fillets and whole wings. Poultry Science, 93(5): 1258–1262. https:// doi.org/10.3382/ps.2013-03697
- Torralbo, A., Borge, C., Allepuz, A., García-Bocanegra, I., Sheppard, S.K., Perea, A. & Carbonero, A. 2014. Prevalence and risk factors of *Campylobacter* infection in broiler flocks from southern Spain. *Preventive Veterinary Medicine*, 114(2): 106–113. https://doi.org/10.1016/ j.prevetmed.2014.01.019
- Tsiouris, V., Kontominas, M.G., Filioussis, G., Chalvatzi, S., Giannenas, I.,
  Papadopoulos, G., Koutoulis, K., Fortomaris, P. & Georgopoulou,
  I. 2020. The effect of whey on performance, gut health and bone morphology parameters in broiler chicks. *Foods*, 9(5): 588. doi: 10.3390/ foods9050588
- Tustin, J., Laberge, K., Michel, P., Reiersen, J., Dađadóttir, S., Briem, H., Harđardóttir, H. et al. 2011. A national epidemic of Campylobacteriosis in Iceland, lessons learned. Zoonoses and Public Health, 58(6): 440–447. https://doi.org/10.1111/j.1863-2378.2010.01387.x
- Umaraw, P., Prajapati, A., Verma, A.K., Pathak, V. & Singh, V.P. 2017. Control of *campylobacter* in poultry industry from farm to poultry processing unit: A review. *Critical Reviews in Food Science and Nutrition*, 57(4): 659–665. https://doi.org/10.1080/10408398.2014.935847
- Urdaneta, S., Lorca-Oró, C., Dolz, R., López-Soria, S. & Cerdà-Cuéllar, M. 2023. In a warm climate, ventilation, indoor temperature and outdoor relative humidity have significant effects on *Campylobacter* spp. colonization in chicken broiler farms which can occur in only 2 days. *Food Microbiology*, 109: 104118. https://doi.org/10.1016/ j.fm.2022.104118
- Van Gerwe, T., Bouma, A., Klinkenberg, D., Wagenaar, J.A., Jacobs-Reitsma, W.F. & Stegeman, A. 2010. Medium chain fatty acid feed supplementation reduces the probability of *Campylobacter jejuni* colonization in broilers. *Veterinary Microbiology*, 143(2–4): 314–318. https://doi.org/10.1016/j.vetmic.2009.11.029
- van Wagenberg, C.P.A., Van Horne, P.L.M., Sommer, H.M. & Nauta, M.J. 2016. Cost-effectiveness of *Campylobacter* interventions on broiler farms in six European countries. *Microbial Risk Analysis*, 2–3: 53–62. https://doi. org/10.1016/j.mran.2016.05.003

- Vandeputte, J., Martel, A., Van Rysselberghe, N., Antonissen, G., Verlinden, M., De Zutter, L., Heyndrickx, M. et al. 2019. In ovo vaccination of broilers against *Campylobacter jejuni* using a bacterin and subunit vaccine. *Poultry Science*, 98(11): 5999–6004. https://doi.org/10.3382/ps/pez402
- Vaz, C.S.L., Voss-Rech, D., Rebelatto, R., Duarte, S.C., Coldebella, A. & Bessa, M.C. 2021. Survival of *Campylobacter jejuni* in chicken at refrigeration and cooking temperatures. *Pesquisa Agropecuária Brasileira*, 56: e02405. https://doi.org/10.1590/s1678-3921.pab2021.v56.02405
- Verhoeff-Bakkenes, L., Beumer, R.R., De Jonge, R., Van Leusden, F.M. & De Jong, A.E.I. 2008. Quantification of *Campylobacter jejuni* crosscontamination via hands, cutlery, and cutting board during preparation of a chicken fruit salad. *Journal of Food Protection*, 71(5): 1018–1022. https://doi.org/10.4315/0362-028X-71.5.1018
- Vetchapitak, T., Shinki, T., Sasaki, S., Taniguchi, T., Luangtongkum, T. & Misawa, N. 2020. Evaluation of chemical treatment combined with vacuum and ultrasonication with a water resonance system for reducing *Campylobacter* on naturally contaminated chicken carcasses. *Food Control*, 112: 107087. https://doi.org/10.1016/j.foodcont.2020.107087
- Wagenaar, J.A., French, N.P. & Havelaar, A.H. 2013. Preventing *Campylobacter* at the source: Why is it so difficult? *Clinical Infectious Diseases*, 57(11): 1600–1606. https://doi.org/10.1093/cid/cit555
- Wagle, B.R., Arsi, K., Upadhyay, A., Shrestha, S., Venkitanarayanan, K., Donoghue, A.M. & Donoghue, D.J. 2017. β-Resorcylic Acid, a Phytophenolic compound, reduces *Campylobacter jejuni* in postharvest poultry. *Journal of Food Protection*, 80(8): 1243–1251. https://doi. org/10.4315/0362-028X.JFP-16-475
- Walker, G.K., Jalukar, S. & Brake, J. 2018. The effect of refined functional carbohydrates from enzymatically hydrolyzed yeast on the transmission of environmental *Salmonella* Senftenberg among broilers and proliferation in broiler housing. *Poultry Science*, 97(4): 1412-1419. https://doi.org/10.3382/ps/pex430
- Wanja, D.W., Mbuthia, P.G., Aboge, G.O. & Bebora, L.C. 2022. Seasonal prevalence and molecular identification of thermophilic *Campylobacter* from chicken, cattle, and respective drinking water in Kajiado County, Kenya. *International Journal of Microbiology*, 2022: 1–11. https://doi. org/10.1155/2022/1526641

- Williams, L.K., Sait, L.C., Trantham, E.K., Cogan, T.A. & Humphrey, T.J. 2013. Campylobacter infection has different outcomes in fast- and slowgrowing broiler chickens. Avian Diseases, 57(2): 238–241. https://doi. org/10.1637/10442-110212-Reg.1
- Wong, J.T., de Bruyn, J., Bagnol, B., Grieve, H., Li, M., Pym, R. & Alders, R.G. 2017. Small-scale poultry and food security in resource-poor settings: A review. *Global Food Security*, 15: 43–52. https://doi.org/10.1016/ j.gfs.2017.04.003
- Zakarienė, G., Rokaitytė, A., Ramonaitė, S., Novoslavskij, A., Mulkytė, K., Zaborskienė, G. & Malakauskas, M. 2015. The antimicrobial effect of spice-based marinades against *Campylobacter jejuni* on contaminated fresh broiler wings. *Journal of Food Science*, 80(3). https://doi. org/10.1111/1750-3841.12788
- Zbrun, M.V., Rossler, E., Olivero, C.R., Soto, L.P., Zimmermann, J.A., Frizzo, L.S. & Signorini, M.L. 2021. Possible reservoirs of thermotolerant *Campylobacter* at the farm between rearing periods and after the use of enrofloxacin as a therapeutic treatment. *International Journal of Food Microbiology*, 340: 109046. https://doi.org/10.1016/ j.ijfoodmicro.2021.109046
- Zweifel, C., Althaus, D. & Stephan, R. 2015. Effects of slaughter operations on the microbiological contamination of broiler carcasses in three abattoirs. *Food Control*, 51: 37–42. https://doi.org/10.1016/j.foodcont.2014.11.002

# Annex

MEASURES FOR THE CONTROL OF CAMPYLOBACTER SPP. IN CHICKEN MEAT
## THE KEYWORDS USED FOR SEARCHING THE CONTROL MEASURES OF *CAMPYLOBACTER* SPP. IN POULTRY

Parameter	Search
1 (poultry)	poultry
	OR chicken*
	OR hen*
	OR broiler*
	OR "Gallus gallus"
	OR "Gallus domesticus"
	OR "G gallus"
	OR "G domesticus"
	OR "Gallus gallus domesticus"
	OR "G gallus domesticus"
	OR duck*
	OR turkey*
	OR goose
	OR geese
	OR guineafowl*
	OR pigeon*
	OR "quail"
	NOT layer*
2 (Salmonella)	Campy or campylobacter
3 (intervention)	intervention*
	OR antibiotic*
	OR antimicrobial*
	OR antibacterial*
	OR bacteriophage*
	OR bifidobac*
	OR biosecur*
	OR boning
	OR chlorine
	OR chill*
	OR "competitive exclusion"
	OR contamination

3 (intervention)

OR control OR cool\* OR cut\* OR debon\* OR decontaminat\* OR decreas\* OR dehid\* OR dehair\* OR disinfect\* OR dress\* OR efficacy OR eviserat\* OR fabricat\* OR grind\* OR "hot water" OR hygiene OR immunis\* OR immuniz\* OR inactiv\* OR irradiat\* OR lactob\* OR "lactic acid bacteria" OR mitigat\* OR pasteuriz\* OR phage\* OR probiotic\* OR reduce\* **OR** reducing OR reduction OR rins\* OR skin\* OR "sodium chlorate" OR spray\* OR steam OR treatment\* OR storage OR trial

4

Title/Abstract (1 AND 2 AND 3)

OR trim\* OR vaccin\* OR vaccum\* OR wash\*

# THE QUESTIONS FOR THE TWO-STEP RELEVANCE SCREENING AND CONFIRMATION

## A2.1 Relevance screening

Question	Options	Key definitions
1. Does this citation describe research	<ul> <li>Yes, primary research</li> <li>Yes, systematic review/</li> </ul>	<b>Primary research</b> is a collection of new data in a single study.
evaluating the efficacy and/or effectiveness (including costs or practically of implementation) of <u>interventions</u> to control	<ul> <li>meta-analysis</li> <li>Yes, risk assessment, risk profile, or other risk-based tool (e.g. cost-benefit analysis)</li> <li>No, or it is a narrative literature review on the</li> </ul>	<b>Risk assessment</b> is a scientifically based process consisting of the following steps: (i) hazard identification, (ii) hazard characterization, (iii) exposure assessment, and (iv) risk characterization.
Campylobacter in poultry at any stage from the primary production to consumption? Selections 1–3 will pass the citation to the next review stage and the	literature review on the subject (exclude)	<b>Risk profile</b> presents the current state of knowledge related to a food safety issue, describes potential options that have been identified to date (if any), and the food safety policy context that will influence further possible actions. Other risk-based tools could include cost-benefit analyses, risk ranking, or risk
		prioritizations. Systematic review is a structured review of a clearly defined question with a transparent search strategy, relevance screening process, data extraction, risk-of-bias assessment and synthesis of results. Meta-analysis is a statistical technique that can be used on data collected in a systematic review. Exclude research on feral animals (e.g. feral pigs not produced for human consumption), and <i>in vitro</i> lab experiments.

2. What commodity is investigated?	<ul> <li>Chicken (Gallus gallus domesticus)</li> <li>Duck (Cairina sp)</li> <li>Turkey (Meleagris)</li> <li>Goose (Anser anser)</li> <li>Guineafowl (Numida meleagris)</li> <li>Pigeon (Columba livia</li> </ul>
	domestica)
	• Quail (Coturnix japonica)
	• Other

## A2.2 Relevance confirmation

Question	Options	Key definitions
Did the study investigate outcomes other than <i>Campylobacter</i> ?	<ul> <li>Yes, <i>E. coli</i> (generic and/or pathogenic strains)</li> <li>Yes, <i>Salmonella</i></li> <li>Yes, other bacteria</li> <li>No</li> </ul>	
In what setting was the study carried out?	<ul> <li>Commercial/field conditions</li> <li>Research farm/pilot plant</li> <li>Smallholder farm/ abattoir conditions</li> <li>Laboratory conditions</li> <li>Not reported</li> </ul>	
In what country was the study conducted?	<ul> <li>The information is in the abstract, which is: (COMMENT)</li> <li>Cannot tell from the abstract</li> </ul>	Specify country name only (not subregions, states, provinces, etc.)
How much logarithm reduction?	<ul> <li>The information is in the abstract, which is (COMMENT)</li> <li>Cannot tell from the abstract</li> <li>Other way to reflect the efficiency (COMMENT)</li> </ul>	

Is it at farm?	What is the intervention?
<ul><li>Yes</li><li>No</li></ul>	<ul> <li>Biosecurity/ management practices</li> <li>Vaccination</li> <li>Antimicrobials</li> <li>Competitive exclusion/ probiotics</li> <li>Feed/water acidification</li> <li>Feed characteristics/ management</li> <li>Bacteriophages</li> <li>Other (COMMENT)</li> </ul>
Is it from transport to slaughter? • Yes • No	What is the intervention? (COMMENT)
Is it processing? • Yes • No	<ul> <li>What is the intervention?</li> <li>Segregated/logistic slaughter</li> <li>Cleaning/disinfection of equipment/ environments</li> <li>Carcass/product washes, rinses, sprays</li> <li>Standard processing procedures/good hygienic practices (GHP)</li> <li>Irradiation</li> <li>Modified packaging</li> <li>Bacteriophages</li> <li>Other (COMMENT)</li> </ul>
Is it from post- processing to consumer? • Yes • No	<ul> <li>What is the intervention?</li> <li>Biosecurity/ management practices</li> <li>Vaccination</li> <li>Antimicrobials</li> <li>Competitive exclusion/ probiotics</li> <li>Feed/water acidification</li> <li>Feed characteristics/ management</li> <li>Bacteriophages</li> <li>Other (COMMENT)</li> </ul>

Antimicrobials: Examples include: Fluroquinolones, cephalosporins, gentamicin, ampicillin, tetracyclines, spectinomycin, ciprofloxacin, ceftriaxone. These may be administered via feed.

**Biosecurity:** This includes, but is not limited to, sanitation, biosafety, disinfection, hygiene and hygiene barriers, allin-all-out production, depopulation, staff and the environment, litter testing and treatment, pest control, etc.

#### **Competitive exclusion:** This may also be referred to as probiotics, prebiotics, and synbiotics. It may include Lactobacillus spp., bacteroides, Bifidobacterium spp., Enterococcus faecium, Aspergillus oryzae, and Saccharomyces spp. (S. cerevisiae, S. boulardii). It may be caecal contents or other materials from animals or the environment that contain many different or unknown bacterial species.

Feed/water acidification: This is the addition of organic acids, such as lactic acid, to feed or water. It would include "nutraceuticals" such as copper, chromium, zinc, betaine or carnitine.

Feed management: These are, for example, comparisons of coarse/ finely ground feed, fermented feed, or liquid feed.

#### Segregated/logistic slaughter: For example, slaughtering/processing of more highly contaminated lots after less contaminated lots

Standard processing procedures/good hygiene practices (GHP) refer to steps such as singeing, dehiding, cooling, chilling, etc.

## **REVIEW OF THE CODEX GUIDELINES**

Recommended revisions to the *Guidelines for the Control of* Campylobacter *and* Salmonella *in Chicken Meat (GXG 78-2011)*, as they relate specifically to the control of *Campylobacter* spp.

Para.	CAC/GL 78-2011	JEMRA Recommendations
5.	<ul> <li>The Guidelines are presented in a flow diagram format so as to enhance practical application of a primary production-to-consumption approach to food safety. This format:</li> <li>demonstrates differences and commonalities in approach for control measures for <i>Campylobacter</i> and <i>Salmonella</i>;</li> <li>; and</li> <li>facilitates development of HACCP plans at individual premises and national levels.</li> </ul>	• To consider updating the last bullet to read: "Facilitate development of HACCP and <b>risk assessment</b> plans at individual premises"
9.	Scope	<ul> <li>Consider clarifying the scope to include: thermotolerant Campylobacter and non-Typhoidal Salmonella.</li> </ul>
12.	The Guidelines systematically present GHP-based control measures and examples of hazard- based control measures. GHP is a prerequisite to making choices on hazard-based control measures Government and industry can use choices on hazard-based control measures to inform decisions on critical control points (CCPs) when applying HACCP principles to a particular food process.	<ul> <li>Consider the following addition:</li> <li>"(CCPs) and relative risk reduction when applying HACCP and risk assessment principles to a particular food process."</li> </ul>

Para.	CAC/GL 78-2011	JEMRA Recommendations
4.	Definitions	Support the following recommendation from the JEMRA meeting on <i>Salmonella</i> controls:
		<ul> <li>To consider including a definition for a production lot as per the Guidelines on the management of biological foodborne outbreaks.</li> </ul>
		• Lot: A definite quantity of ingredients or of a food that is intended to have uniform character and quality, within specified limits, is produced, packaged and labelled under the same conditions, and is assigned a unique reference identification by the food business operator. It may also be referred to as a "batch".
16.	<ul> <li>5. PRINCIPLES APPLYING TO CONTROL OF CAMPYLOBACTER AND SALMONELLA IN CHICKEN MEAT</li> <li>Overarching</li> <li>i. The principles of food safety risk analysis should be incorporated wherever possible and appropriate in the control of Campylobacter and Salmonella in chicken meat from primary production to consumption.</li> <li>ii. Wherever possible and practical, Competent Authorities</li> </ul>	<ul> <li>Consider the following editorial change: "5i) The principles of food safety risk analysis should be incorporated to the extent possible and as appropriate" and "5ii) To the extent possible and as appropriate, Competent Authorities should"</li> </ul>
18.	Food Safety Risk Profile for <i>Salmonella</i> species in broiler (young) chicken, June 2007.	Support the following recommendation from the JEMRA meeting on <i>Salmonella</i> controls:
	Food Safety Risk Profile for Campylobacter species in broiler	• To verify that the links referenced in the footnote are current and active.
	(young) chicken, suite 2007.	<ul> <li>IO evaluate paragraph I8 and to consider updating it, if needed.</li> </ul>

Para.	CAC/GL 78-2011	JEMRA Recommendations
Section 7	PRIMARY PRODUCTION-TO- CONSUMPTION APPROACH TO CONTROL MEASURES	Consider the following updates to account for religious practices that do not include stunning:
		<ul> <li>Process Flow Diagram 2: Step 14         <ul> <li>slaughter: B1) with stunning B2)</li> <li>without stunning, then A) Hang,</li> <li>then B1) split into Gas and Electrical</li> </ul> </li> </ul>
		Consider tick marks in <i>Campylobacter</i> column in the summary table for:
		<ul> <li>Receive at Slaughterhouse (use Establishment as a term)</li> </ul>
		<ul> <li>Dress (interventions validated for Campylobacter during dressing (decontamination)</li> </ul>
		<ul> <li>Portion (interventions directed at portions/parts)</li> </ul>
24. Control of Salmone is streng of a com personne particula measure should be with rele	Control of Campylobacter and Salmonella in grandparent flocks is strengthened by the application of a combination of biosecurity and personnel hygiene measures. The particular combination of control	Support the following recommendation from the JEMRA meeting on <i>Salmonella</i> controls:
		<ul> <li>Consider including a definition for biosecurity that includes personal hygiene.</li> </ul>
	should be determined in consultation with relevant stakeholders.	<ul> <li>Consider aligning with the WOAH definition: https://www.woah.org/ fileadmin/Home/eng/Health_ standards/tahc/current/glossaire. pdf.</li> </ul>
		<ul> <li>Consider changing the text to read</li> <li>"by the application of effective biosecurity measures."</li> </ul>
32.	Personnel involved in the transportation of day-old chicks to parent flocks should not enter any livestock buildings and should prevent cross-contamination of day-old chicks during loading and unloading.	Consider updating the text to read:     "transportation of day-old chicks     to parent flock establishments"
36.	Personnel involved in the transportation of day-old chicks should not enter any livestock buildings.	<ul> <li>Consider changing "livestock buildings" to "livestock establishments."</li> </ul>

Para.	CAC/GL 78-2011	JEMRA Recommendations
40.	The use of fly screens to reduce or eliminate fly infestation in broiler houses has been shown to decrease the percentage of <i>Campylobacter</i> spppositive flocks from 51.4% to 15.4%.	<ul> <li>Consider revising this guidance as there has not been any clear evidence since the initial studies on effectiveness.</li> <li>Consider eliminating the percentages and leave the remainder of the statement.</li> </ul>
		<ul> <li>Consider adding a statement to include fly screens, in combination with high biosecurity measures.</li> </ul>
41.	Full depopulation of the flock should be carried out where possible. Where this is not practicable and partial depopulation is practised, particular attention should be paid to strict biosecurity and hygiene of catchers and the equipment they use.	<ul> <li>Consider adding the following to statement paragraph 41: "Partial depopulation has been shown to be a risk factor for the increase of <i>Campylobacter</i> contamination" as there are several reports on thinning/depopulation and <i>Campylobacter</i> contamination due to biosecurity deficiencies.</li> </ul>
45.	Where appropriate to the national situation, information about the flock, in particular about <i>Salmonella</i> and/or <i>Campylobacter</i> status should be provided in a timely manner to enable logistic slaughter and/ or channelling of poultry meat to treatment.	<ul> <li>Consider replacing the word "treatment" with "reduced risk processing", "intervention" or "custom processing".</li> </ul>
54.	Washing with abundant potable running water	<ul> <li>Support the recommendations from the JEMRA meeting on Salmonella controls:</li> <li>Consider replacing potable water with fit for purpose water to align with CXG1-1969, paragraph 70. Text should be adjusted to fit for purpose water.</li> </ul>
59	Other factors that should be taken into account when designing process control systems that minimize contamination during scalding include: • degree of agitation; • use of multistaged tanks; •; • tanks being cleaned and disinfected at least daily; and • hygiene measures applied to reused/recycled water.	<ul> <li>Consider replacing "daily" with "tanks being cleaned and disinfected at an adequate frequency (e.g. end of shift)"</li> <li>Consider adding the following bullet: "Directed water scalders"</li> </ul>

Para.	CAC/GL 78-2011	JEMRA Recommendations
63.	9.4.1.5 Crop removal 63. Where possible, crops should be extracted in a manner that is likely to limit carcass contamination.	• Consider adding the following to section 9.4.1.5: "The use of cropper systems allows the release of accumulated dirty water on the carcass cavity, so efforts to remove collected water prior to chilling should be considered."
67.	Carcass washing systems with 13 washers using water with 25-35 ppm total chlorine have been shown to reduce levels of <i>Campylobacter</i> by about 0.5 log <sub>10</sub> CFU/ml of whole carcass rinse sample. Post-wash sprays using Acidified Sodium Chlorite (ASC) or TSP may further reduce <i>Campylobacter</i> levels by an average of 1.3 log <sub>10</sub> CFU/ml or 1.0 log <sub>10</sub> CFU/ml of whole carcass rinse samples respectively.	Consider adding the following statement: "Carcass washes with 400 ppm paracetic acid (PAA), showed 1.2 log reductions of Campylobacter prior to chilling." (Cano, Menses and Chaves, 2021; Chousalkar et al., 2019; De Villena et al., 2022; Dogan et al., 2019, 2022)
69.	An on-line reprocessing spray system incorporating ASC has been shown to reduce Campylobacter in the whole carcass rinse sample by about 2.1 log <sub>10</sub> CFU/ml and to reduce the prevalence of <i>Salmonella</i> positive carcasses from 37% to 10%.	Consider adding the following statement: "Inside and outside bird washers used for online-reprocessing at 100 ppm of PAA showed 0.5 log reductions of Campylobacter." (Cano, Menses and Chaves, 2021; Chousalkar et al., 2019; De Villena et al., 2022; Dogan et al., 2019)
70.	Dipping carcasses in 10% TSP reduced <i>Campylobacter</i> by 1.7 log <sub>10</sub> CFU/g neck skin and the MPN of <i>Salmonella</i> was reduced from 1.92 log <sub>10</sub> CFU/g neck skin to undetectable levels.	<ul> <li>Consider removing the recommendation for TSP since it may not be as commonly used at this time.</li> <li>Consider adding the following statement: "Dip treatments in 200 ppm of PAA showed 1.4 log reductions of Campylobacter in carcass rinses." (Cano, Menses and Chaves, 2021; Chousalkar et al., 2019; De Villena et al., 2022; Dogan et al., 2019)</li> </ul>

Para.	CAC/GL 78-2011	JEMRA Recommendations
78.	The use of chlorine in the chill tank may not act as a decontaminating agent by acting directly on the contaminated carcass. However, there would be a washing off effect by the water itself, and the addition of chlorine at a level sufficient to maintain a free residual in the water would then inactivate <i>Campylobacter</i> and <i>Salmonella</i> washed off, preventing re-attachment and cross- contamination.	<ul> <li>Consider adding the following statement: "Immersion chillers using 225 ppm of PAA showed reductions of 1.18 log in concentration and a 76.5% reduction in prevalence of Campylobacter." (Cano, Menses and Chaves, 2021; Chousalkar et al., 2019; De Villena et al., 2022; Dogan et al., 2022)</li> </ul>
81.	Forced air chilling (blast chilling) may reduce the concentration of <i>Campylobacter</i> on chicken carcasses by 0.4 log <sub>10</sub> CFU/carcass.	<ul> <li>Consider updating the value and excluding the unit of measure: "up to 1 log." (ANSES, 2018)</li> </ul>
82.	Immersion chilling has been shown to reduce concentrations of <i>Campylobacter</i> by 1.1–1.3 log <sub>10</sub> CFU/ ml of carcass rinse.	<ul> <li>Consider updating this text to: <i>"Immersion chilling reduces</i> <i>concentrations of Campylobacter,</i> <i>and with a combination of</i> <i>processing aids can result in a</i> <i>higher log reduction."</i> (Dogan et al., 2022)</li> </ul>
84.	Immersing whole carcasses in 600-800 ppm ASC at pH 2.5 to 2.7 for 15 seconds immediately post- chill, has been shown to reduce <i>Campylobacter</i> by 0.9-1.2 log <sub>10</sub> CFU/ ml of whole carcass rinse sample.	<ul> <li>Consider adding the following text: "Post chill tank interventions using sprays of PAA at 1 000 ppm showed up to 2.1 log reductions of Campylobacter." (Cano, Menses and Chaves, 2021; Chousalkar et al., 2019; De Villena et al., 2022; Dogan et al., 2022)</li> </ul>
9.10.1	For Campylobacter	<ul> <li>Consider adding a new paragraph for Campylobacter and the following text: "Immersion of chicken wings in 1 000 ppm of PAA for 30 s has been shown to reduce 2.3 logs of Campylobacter" (Cano, Menses and Chaves, 2021; Chousalkar et al., 2019; De Villena et al., 2022; Dogan et al., 2022)</li> </ul>

Para.	CAC/GL 78-2011	JEMRA Recommendations
94.	Freezing of naturally contaminated carcasses followed by 31 days of storage at -20 degrees C has been shown to reduce <i>Campylobacter</i> by 0.7 to 2.9 log <sub>10</sub> CFU/g.	<ul> <li>Consider updating this statement to read: "Freezing of carcasses and portions contaminated with Campylobacter followed by storage at -20 degrees C has been shown to reduce Campylobacter by up to 2 logs" (Dogan et al., 2022).</li> </ul>
95.	Crust freezing using continuous carbon dioxide belt freezing of skinless breast fillets has been shown to give a reduction of <i>Campylobacter</i> of $0.4 \log_{10}$ CFU/fillet.	<ul> <li>Consider updating the statement to include all chicken meat products and not just skinless breast fillets.</li> </ul>
108.	Chicken meat should be cooked according to a process that is capable of achieving at least a 7 log reduction in both <i>Campylobacter</i> and <i>Salmonella</i> .	<ul> <li>Consider updating the text to read: "Chicken meat should be cooked according to a process that is capable of reaching an internal temperature that can inactivate Salmonella and Campylobacter, for example 74 °C."</li> </ul>
115.	Chicken meat should be cooked according to a process that is capable of achieving at least a 7 log reduction in both <i>Campylobacter</i> and <i>Salmonella</i> .	<ul> <li>Consider updating the text to read: "Chicken meat should be cooked according to a process that is capable of reaching an internal temperature that can inactivate Salmonella and Campylobacter, for example 74 °C."</li> </ul>

### **References:**

- ANSES. 2018. State of knowledge relating to the contamination of broilers with *Campylobacter* and assessment of the impact of interventions at different stages of the food chain in France. *Collective Expert Appraisal Report*. 2016-SA-0183. Maisons-Alfort Cedex, France. https://www.anses.fr/en/ system/files/BIORISK2016SA0183RaEN.pdf
- Cano, C., Meneses, Y. & Chaves, B.D. 2021. Application of Peroxyacetic Acid for decontamination of raw poultry products and comparison to other commonly used chemical antimicrobial interventions: A review. *Journal* of Food Protection, 84(10): 1772–1783. https://doi.org/10.4315/JFP-21-107

- Chousalkar, K., Sims, S., McWhorter, A., Khan, S. & Sexton, M. 2019. The effect of sanitizers on microbial levels of chicken meat collected from commercial processing plants. *International Journal of Environmental Research and Public Health*, 16(23): 4807. https://doi.org/10.3390/ijerph16234807
- De Villena, J.F., Vargas, D.A., Bueno López, R., Chávez-Velado, D.R., Casas, D.E., Jiménez, R.L. & Sanchez-Plata, M.X. 2022. Bio-mapping indicators and pathogen loads in a commercial broiler processing facility operating with high and low antimicrobial intervention levels. *Foods*, 11(6): 775. https://doi.org/10.3390/foods11060775
- Dogan, O.B., Aditya, A., Ortuzar, J., Clark, J. & Wang, B. 2022. A systematic review and meta-analysis of the efficacy of processing stages and interventions for controlling Campylobacter contamination during broiler chicken processing. *Comprehensive Reviews in Food Science and Food Safety*, 21(1): 227–271. https://doi.org/10.1111/1541-4337.12860
- Dogan, O.B., Clarke, J., Mattos, F. & Wang, B. 2019. A quantitative microbial risk assessment model of Campylobacter in broiler chickens: Evaluating processing interventions. *Food Control*, 100: 97–110. https://doi. org/10.1016/j.foodcont.2019.01.003

## **BIBLIOGRAPHY USED FOR SCOPING REVIEW**

## A 4.1 Primary

## A4.1.1 Biosecurity and management approaches

- Agunos, A., Waddell, L., Léger, D. & Taboada, E. 2014. A systematic review characterizing on-farm sources of *Campylobacter* spp. for broiler chickens. *PLoS ONE*, 9(8): e104905. https://doi.org/10.1371/journal. pone.0104905
- Al Hakeem, W.G., Fathima, S., Shanmugasundaram, R. & Selvaraj, R.K. 2022. *Campylobacter jejuni* in poultry: pathogenesis and control strategies. *Microorganisms*, 10(11): 2134. https://doi.org/10.3390/microorganisms 10112134
- Allen, V.M., Weaver, H., Ridley, A.M., Harris, J.A., Sharma, M., Emery, J., Sparks, N., Lewis, M. & Edge, S. 2008. Sources and spread of thermophilic *Campylobacter* spp. during partial depopulation of broiler chicken flocks. *Journal of Food Protection*, 71(2): 264–270. https://doi. org/10.4315/0362-028X-71.2.264
- Alpigiani, I., Abrahantes, J.C., Michel, V., Huneau-Salaün, A., Chemaly, M., Keeling, L.J., Gervelmeyer, A. et al. 2017. Associations between animal welfare indicators and *Campylobacter* spp. in broiler chickens under commercial settings: A case study. *Preventive Veterinary Medicine*, 147: 186–193. https://doi.org/10.1016/j.prevetmed.2017.09.005
- Babacan, O., Harris, S.A., Pinho, R.M., Hedges, A., Jørgensen, F. & Corry, J.E.L. 2020. Factors affecting the species of *Campylobacter* colonizing chickens reared for meat. *Journal of Applied Microbiology*, 129(4): 1071–1078. https://doi.org/10.1111/jam.14651
- Bailey, M.A., Bourassa, D.V., Krehling, J.T., Munoz, L., Chasteen, K.S., Escobar, C. & Macklin, K.S. 2022. Effects of common litter management practices on the prevalence of *Campylobacter jejuni* in broilers. *Animals*, 12(7): 858. https://doi.org/10.3390/ani12070858

- Battersby, T., Whyte, P. & Bolton, D.J. 2016. The pattern of *Campylobacter* contamination on broiler farms; external and internal sources. *Journal of Applied Microbiology*, 120(4): 1108–1118. https://doi.org/10.1111/ jam.13066
- Borck Høg, B., Sommer, H.M., Larsen, L.S., Sørensen, A.I.V., David, B., Hofshagen, M. & Rosenquist, H. 2016. Farm specific risk factors for *Campylobacter* colonisation in Danish and Norwegian broilers. *Preventive Veterinary Medicine*, 130: 137–145. https://doi.org/10.1016/ j.prevetmed.2016.04.002
- Chowdhury, S., Themudo, G.E., Sandberg, M. & Ersbøll, A.K. 2013. Spatiotemporal patterns of *Campylobacter* colonization in Danish broilers. *Epidemiology and Infection*, 141(5): 997–1008. https://doi.org/10.1017/ S0950268812001446
- Colles, F.M., Cain, R.J., Nickson, T., Smith, A.L., Roberts, S.J., Maiden, M.C.J., Lunn, D. & Dawkins, M.S. 2016. Monitoring chicken flock behaviour provides early warning of infection by human pathogen *Campylobacter*. *Proceedings of the Royal Society B: Biological Sciences*, 283(1822): 20152323. https://doi.org/10.1098/rspb.2015.2323
- **Colles, F.M., Dingle, K.E., Cody, A.J. & Maiden, M.C.J.** 2008. Comparison of *Campylobacter* populations in wild geese with those in starlings and free-range poultry on the same farm. *Applied and Environmental Microbiology*, 74(11): 3583–3590. https://doi.org/10.1128/AEM.02491-07
- Crotta, M., Georgiev, M. & Guitian, J. 2017. Quantitative risk assessment of *Campylobacter* in broiler chickens – Assessing interventions to reduce the level of contamination at the end of the rearing period. *Food Control*, 75: 29–39. https://doi.org/10.1016/j.foodcont.2016.12.024
- Di Marcantonio, L., Marotta, F., Vulpiani, M.P., Sonntag, Q., Iannetti, L., Janowicz, A., Serafino, G.D., Di Giannatale, E. & Garofolo, G. 2022. Investigating the cecal microbiota in broiler poultry farms and its potential relationships with animal welfare. *Research in Veterinary Science*, 144: 115–125. https://doi.org/10.1016/j.rvsc.2022.01.020
- Georgiev, M., Beauvais, W. & Guitian, J. 2017. Effect of enhanced biosecurity and selected on-farm factors on *Campylobacter* colonization of chicken broilers. *Epidemiology and Infection*, 145(3): 553–567. https://doi. org/10.1017/S095026881600251X

- Golden, C.E. & Mishra, A. 2020. Prevalence of Salmonella and Campylobacter spp. in alternative and conventionally produced chicken in the United States: a systematic review and meta-analysis. Journal of Food Protection, 83(7): 1181–1197. https://doi.org/10.4315/JFP-19-538
- Griekspoor, P., Colles, F.M., McCarthy, N.D., Hansbro, P.M., Ashhurst-Smith,
  C., Olsen, B., Hasselquist, D., Maiden, M.C.J. & Waldenström, J. 2013.
  Marked host specificity and lack of phylogeographic population structure of *Campylobacter jejuni* in wild birds. *Molecular Ecology*, 22(5): 1463–1472. https://doi.org/10.1111/mec.12144
- Guerin, M.T., Martin, S.W., Reiersen, J., Berke, O., McEwen, S.A.,
  Friðriksdóttir, V., Bisaillon, J.-R. & Lowman, R. 2008. Temperaturerelated risk factors associated with the colonization of broiler-chicken flocks with *Campylobacter* spp. in Iceland, 2001–2004. *Preventive Veterinary Medicine*, 86(1–2): 14–29. https://doi.org/10.1016/ j.prevetmed.2008.02.015
- Hansson, I., Olsson Engvall, E., Vågsholm, I. & Nyman, A. 2010. Risk factors associated with the presence of *Campylobacter*-positive broiler flocks in Sweden. *Preventive Veterinary Medicine*, 96:114-121. doi: 10.1016/ j.prevetmed.2010.05.007
- Hasan, Md.M., Talukder, S., Mandal, A.K., Tasmim, S.T., Parvin, Mst.S., Ali, Md.Y., Sikder, M.H. & Islam, Md.T. 2020. Prevalence and risk factors of *Campylobacter* infection in broiler and cockerel flocks in Mymensingh and Gazipur districts of Bangladesh. *Preventive Veterinary Medicine*, 180: 105034. https://doi.org/10.1016/j.prevetmed.2020.105034
- Henry, I., Reichardt, J., Denis, M. & Cardinale, E. 2011. Prevalence and risk factors for *Campylobacter* spp. in chicken broiler flocks in Reunion Island (Indian Ocean). *Preventive Veterinary Medicine*, 100(1): 64–70. https:// doi.org/10.1016/j.prevetmed.2011.03.007
- Hertogs, K., Haegeman, A., Schaumont, D., Gelaude, P., De Zutter, L., Dewulf,
   J., Heyndrickx, M. & Rasschaert, G. 2021. Contamination sources and transmission routes for *Campylobacter* on (mixed) broiler farms in Belgium, and comparison of the gut microbiota of flocks colonized and uncolonized with *Campylobacter*. *Pathogens*, 10(1): 66. https://doi.org/10.3390/pathogens10010066

- Higham, L.E., Scott, C., Akehurst, K., Dring, D., Parnham, A., Waterman,
  M. & Bright, A. 2018. Effects of financial incentives and cessation of thinning on prevalence of *Campylobacter*: a longitudinal monitoring study on commercial broiler farms in the UK. *Veterinary Record*, 183(19): 595–595. https://doi.org/10.1136/vr.104823
- Higham, L.E., Scott, C., Akehurst, K., Dring, D., Parnham, A., Waterman,
  M. & Bright, A. 2018. Effects of financial incentives and cessation of thinning on prevalence of *Campylobacter* : a longitudinal monitoring study on commercial broiler farms in the UK. *Veterinary Record*, 183(19): 595–595. https://doi.org/10.1136/vr.104823
- Hwang, H. & Singer, R.S. 2020. Survey of the U.S. broiler industry regarding pre- and postharvest interventions targeted to mitigate *Campylobacter* contamination on broiler chicken products. *Journal of Food Protection*, 83(7): 1137–1148. https://doi.org/10.4315/JFP-19-527
- Iannetti, L., Neri, D., Santarelli, G.A., Cotturone, G., Podaliri Vulpiani, M., Salini, R., Antoci, S. et al. 2020. Animal welfare and microbiological safety of poultry meat: Impact of different at-farm animal welfare levels on at-slaughterhouse Campylobacter and Salmonella contamination. Food Control, 109: 106921. https://doi.org/10.1016/j.foodcont.2019.106921
- Jonsson, M.E., Chriél, M., Norström, M. & Hofshagen, M. 2012. Effect of climate and farm environment on *Campylobacter* spp. colonisation in Norwegian broiler flocks. *Preventive Veterinary Medicine*, 107(1–2): 95–104. https://doi.org/10.1016/j.prevetmed.2012.05.002
- Lassen, B., Helwigh, B., Kahl Petersen, C. & Ellis-Iversen, J. 2022. Systematic review of products with potential application for use in the control of *Campylobacter* spp. in organic and free-range broilers. *Acta Veterinaria Scandinavica*, 64(1): 24. https://doi.org/10.1186/s13028-022-00644-z
- McKenna, A., Ijaz, U.Z., Kelly, C., Linton, M., Sloan, W.T., Green, B.D., Lavery, U. et al. 2020. Impact of industrial production system parameters on chicken microbiomes: mechanisms to improve performance and reduce *Campylobacter. Microbiome*, 8(1): 128. https://doi.org/10.1186/s40168-020-00908-8
- Mendes, Â.J., Santos-Ferreira, N.L., Costa, F.M., Lopes, E.P., Freitas-Silva, J., Inácio, Â.S., Moreira, F. & Martins Da Costa, P. 2020. External contamination of broilers by *Campylobacter* spp. increases from the farm to the slaughterhouse. *British Poultry Science*, 61(4): 400–407. https://doi. org/10.1080/00071668.2020.1736264

74

- Näther, G., Alter, T., Martin, A. & Ellerbroek, L. 2009. Analysis of risk factors for *Campylobacter* species infection in broiler flocks. *Poultry Science*, 88(6): 1299–1305. https://doi.org/10.3382/ps.2008-00389
- Natsos, G., Mouttotou, N.K., Magiorkinis, E., Ioannidis, A., Rodi-Burriel, A., Chatzipanagiotou, S. & Koutoulis, K.C. 2020. Prevalence of and risk factors for *Campylobacter* spp. colonization of broiler chicken flocks in Greece. *Foodborne Pathogens and Disease*, 17(11): 679–686. https://doi. org/10.1089/fpd.2020.2795
- Rawson, T., Dawkins, M.S. & Bonsall, M.B. 2019. A mathematical model of *Campylobacter* dynamics within a broiler flock. *Frontiers in Microbiology*, 10: 1940. https://doi.org/10.3389/fmicb.2019.01940
- Reichelt, B., Szott, V., Epping, L., Semmler, T., Merle, R., Roesler, U. & Friese, A. 2022. Transmission pathways of *Campylobacter* spp. at broiler farms and their environment in Brandenburg, Germany. *Frontiers in Microbiology*, 13: 982693. https://doi.org/10.3389/fmicb.2022.982693
- Rothrock, M.J., Cook, K.L., Warren, J.G. & Sistani, K. 2008. The effect of alum addition on microbial communities in poultry litter. *Poultry Science*, 87(8): 1493–1503. https://doi.org/10.3382/ps.2007-00491
- Sandberg, M., Dahl, J., Lindegaard, L.L. & Pedersen, J.R. 2017. Compliance/ non-compliance with biosecurity rules specified in the Danish Quality Assurance system (KIK) and *Campylobacter*- positive broiler flocks 2012 and 2013. *Poultry Science*, 96(1): 184–191. https://doi.org/10.3382/ps/ pew277
- Schweitzer, P.M., Susta, L., Varga, C., Brash, M.L. & Guerin, M.T. 2021. Demographic, husbandry, and biosecurity factors associated with the presence of *Campylobacter* spp. in small poultry flocks in Ontario, Canada. *Pathogens*, 10(11): 1471. https://doi.org/10.3390/ pathogens10111471
- Sevilla-Navarro, S., Marin, C., Cortés, V., García, C. & Catalá-Gregori, P. 2020. *Campylobacter* prevalence and risk factors associated with exceeding allowable limits in poultry slaughterhouses in Spain. *Veterinary Record*, 186(13): 415–415. https://doi.org/10.1136/vr.105558
- Sheppard, S.K., Colles, F., Richardson, J., Cody, A.J., Elson, R., Lawson, A., Brick, G. et al. 2010. Host association of Campylobacter genotypes transcends geographic variation. Applied and Environmental Microbiology, 76(15): 5269–5277. https://doi.org/10.1128/AEM.00124-10

Smith, S., Messam, L.L.McV., Meade, J., Gibbons, J., McGill, K., Bolton, D. & Whyte, P. 2016. The impact of biosecurity and partial depopulation on *Campylobacter* prevalence in Irish broiler flocks with differing levels of hygiene and economic performance. *Infection Ecology & Epidemiology*, 6(1): 31454. https://doi.org/10.3402/iee.v6.31454

Torralbo, A., Borge, C., Allepuz, A., García-Bocanegra, I., Sheppard, S.K., Perea, A. & Carbonero, A. 2014. Prevalence and risk factors of *Campylobacter* infection in broiler flocks from southern Spain. *Preventive Veterinary Medicine*, 114(2): 106–113. https://doi.org/10.1016/ j.prevetmed.2014.01.019

- Urdaneta, S., Lorca-Oró, C., Dolz, R., López-Soria, S. & Cerdà-Cuéllar,
  M. 2023. In a warm climate, ventilation, indoor temperature and outdoor relative humidity have significant effects on *Campylobacter* spp. colonization in chicken broiler farms which can occur in only 2 days. *Food Microbiology*, 109: 104118. https://doi.org/10.1016/j.fm.2022.104118
- van Wagenberg, C.P.A., Van Horne, P.L.M., Sommer, H.M. & Nauta, M.J. 2016. Cost-effectiveness of *Campylobacter* interventions on broiler farms in six European countries. *Microbial Risk Analysis*, 2–3: 53–62. https://doi. org/10.1016/j.mran.2016.05.003
- Wanja, D.W., Mbuthia, P.G., Aboge, G.O. & Bebora, L.C. 2022. Seasonal prevalence and molecular identification of thermophilic *Campylobacter* from chicken, cattle, and respective drinking water in Kajiado County, Kenya. *International Journal of Microbiology*, 2022: 1–11. https://doi. org/10.1155/2022/1526641
- Williams, L.K., Sait, L.C., Trantham, E.K., Cogan, T.A. & Humphrey, T.J. 2013. Campylobacter infection has different outcomes in fast- and slowgrowing broiler chickens. Avian Diseases, 57(2): 238–241. https://doi. org/10.1637/10442-110212-Reg.1
- Zbrun, M.V., Rossler, E., Olivero, C.R., Soto, L.P., Zimmermann, J.A., Frizzo, L.S. & Signorini, M.L. 2021. Possible reservoirs of thermotolerant *Campylobacter* at the farm between rearing periods and after the use of enrofloxacin as a therapeutic treatment. *International Journal of Food Microbiology*, 340: 109046. https://doi.org/10.1016/ j.ijfoodmicro.2021.109046

76

## A4.1.2 Vaccine and phage-based approaches

### Vaccine

- Buckley, A.M., Wang, J., Hudson, D.L., Grant, A.J., Jones, M.A., Maskell, D.J.
   & Stevens, M.P. 2010. Evaluation of live-attenuated *Salmonella* vaccines expressing *Campylobacter* antigens for control of *C. jejuni* in poultry. *Vaccine*, 28(4): 1094–1105. https://doi.org/10.1016/j.vaccine.2009.10.018
- Chintoan-Uta, C., Cassady-Cain, R.L. & Stevens, M.P. 2016. Evaluation of flagellum-related proteins FliD and FspA as subunit vaccines against *Campylobacter jejuni* colonisation in chickens. *Vaccine*, 34(15): 1739– 1743. https://doi.org/10.1016/j.vaccine.2016.02.052
- Clark, J.D., Oakes, R.D., Redhead, K., Crouch, C.F., Francis, M.J., Tomley, F.M. & Blake, D.P. 2012. Eimeria species parasites as novel vaccine delivery vectors: Anti-*Campylobacter jejuni* protective immunity induced by Eimeria tenella-delivered CjaA. *Vaccine*, 30(16): 2683–2688. https://doi. org/10.1016/j.vaccine.2012.02.002
- Cui, Y., Guo, F., Guo, J., Cao, X., Wang, H., Yang, B., Zhou, H. et al. 2020. Immunization of chickens with the Enterobactin conjugate vaccine reduced *Campylobacter jejuni* colonization in the intestine. *Vaccines*, 8(4): 747. https://doi.org/10.3390/vaccines8040747
- Gloanec, N., Dory, D., Quesne, S., Béven, V., Poezevara, T., Keita, A., Chemaly, M. & Guyard-Nicodème, M. 2022. Impact of DNA prime/protein boost vaccination against *Campylobacter jejuni* on immune responses and gut microbiota in chickens. *Vaccines*, 10(6): 981. https://doi.org/10.3390/ vaccines10060981
- Jeon, B., Saisom, T., Sasipreeyajan, J. & Luangtongkum, T. 2022. Liveattenuated oral vaccines to reduce *Campylobacter* colonization in poultry. *Vaccines*, 10(5): 685. https://doi.org/10.3390/vaccines10050685
- Łaniewski, P., Kuczkowski, M., Chrząstek, K., Woźniak, A., Wyszyńska, A.,
   Wieliczko, A. & Jagusztyn-Krynicka, E.K. 2014. Evaluation of the immunogenicity of *Campylobacter jejuni* CjaA protein delivered by *Salmonella* enterica sv. Typhimurium strain with regulated delayed attenuation in chickens. *World Journal of Microbiology and Biotechnology*, 30(1): 281–292. https://doi.org/10.1007/s11274-013-1447-5

- Layton, S.L., Morgan, M.J., Cole, K., Kwon, Y.M., Donoghue, D.J., Hargis, B.M. & Pumford, N.R. 2011. Evaluation of *Salmonella*-vectored *Campylobacter* peptide epitopes for reduction of *Campylobacter jejuni* in broiler chickens. *Clinical and Vaccine Immunology*, 18(3): 449–454. https://doi.org/10.1128/CVI.00379-10
- Mauri, M., Sannasiddappa, T.H., Vohra, P., Corona-Torres, R., Smith, A.A., Chintoan-Uta, C., Bremner, A. et al. 2021. Multivalent poultry vaccine development using Protein Glycan Coupling Technology. *Microbial Cell Factories*, 20(1): 193. https://doi.org/10.1186/s12934-021-01682-4
- Meunier, M., Guyard-Nicodème, M., Vigouroux, E., Poezevara, T., Béven, V., Quesne, S., Amelot, M. et al. 2018. A DNA prime/protein boost vaccine protocol developed against *Campylobacter jejuni* for poultry. *Vaccine*, 36(16): 2119–2125. https://doi.org/10.1016/j.vaccine.2018.03.004
- Nothaft, H., Perez-Muñoz, M.E., Yang, T., Murugan, A.V.M., Miller, M., Kolarich, D., Plastow, G.S., Walter, J. & Szymanski, C.M. 2021. Improving chicken responses to glycoconjugate vaccination against *Campylobacter jejuni. Frontiers in Microbiology*, 12: 734526. https://doi. org/10.3389/fmicb.2021.734526
- Radomska, K.A., Vaezirad, M.M., Verstappen, K.M., Wösten, M.M.S.M., Wagenaar, J.A. & Van Putten, J.P.M. 2016. Chicken immune response after In Ovo immunization with Chimeric TLR5 activating flagellin of *Campylobacter jejuni*. *PLoS ONE*, 11(10): e0164837. https://doi. org/10.1371/journal.pone.0164837
- Thibodeau, A., Fravalo, P., Perron, A., Lewandowski, S.L.- & Letellier, A. 2017. Production and characterization of anti-*Campylobacter jejuni* IgY derived from egg yolks. *Acta Veterinaria Scandinavica*, 59(1): 80. https:// doi.org/10.1186/s13028-017-0346-4
- Vandeputte, J., Martel, A., Van Rysselberghe, N., Antonissen, G., Verlinden, M., De Zutter, L., Heyndrickx, M. et al. 2019. In ovo vaccination of broilers against *Campylobacter jejuni* using a bacterin and subunit vaccine. *Poultry Science*, 98(11): 5999–6004. https://doi.org/10.3382/ps/pez402

### Phage

Carvalho, C.M., Gannon, B.W., Halfhide, D.E., Santos, S.B., Hayes, C.M., Roe, J.M. & Azeredo, J. 2010. The in vivo efficacy of two administration routes of a phage cocktail to reduce numbers of *Campylobacter coli* and *Campylobacter jejuni* in chickens. *BMC Microbiology*, 10(1): 232. https:// doi.org/10.1186/1471-2180-10-232

- Chinivasagam, H.N., Estella, W., Maddock, L., Mayer, D.G., Weyand, C., Connerton, P.L. & Connerton, I.F. 2020. Bacteriophages to control *Campylobacter* in commercially farmed broiler chickens, in Australia. *Frontiers in Microbiology*, 11: 632. https://doi.org/10.3389/ fmicb.2020.00632
- D'Angelantonio, D., Scattolini, S., Boni, A., Neri, D., Di Serafino, G., Connerton, P., Connerton, I. *et al.* 2021. Bacteriophage therapy to reduce colonization of *Campylobacter jejuni* in broiler chickens before slaughter. *Viruses*, 13(8): 1428. https://doi.org/10.3390/v13081428
- El-Shibiny, A., Scott, A., Timms, A., Metawea, Y., Connerton, P. & Connerton, I. 2009. Application of a group II *Campylobacter* bacteriophage to reduce strains of *Campylobacter jejuni* and *Campylobacter coli* colonizing broiler chickens. *Journal of Food Protection*, 72(4): 733–740. https://doi. org/10.4315/0362-028X-72.4.733
- Fischer, S., Kittler, S., Klein, G. & Glünder, G. 2013. Impact of a single phage and a phage cocktail application in broilers on reduction of *Campylobacter jejuni* and development of resistance. *PLoS ONE*, 8(10): e78543. https://doi.org/10.1371/journal.pone.0078543
- Furuta, M., Nasu, T., Umeki, K., Hoang Minh, D., Honjoh, K.-I. & Miyamoto, T. 2017. Characterization and application of Lytic bacteriophages against *Campylobacter jejuni* isolated from poultry in Japan. *Biocontrol Science*, 22(4): 213–221. https://doi.org/10.4265/bio.22.213
- Hammerl, J.A., Jäckel, C., Alter, T., Janzcyk, P., Stingl, K., Knüver, M.T. & Hertwig, S. 2014. Reduction of *Campylobacter jejuni* in broiler chicken by successive application of group II and group III phages. *PLoS ONE*, 9(12): e114785. https://doi.org/10.1371/journal.pone.0114785
- Kittler, S., Fischer, S., Abdulmawjood, A., Glünder, G. & Klein, G. 2013. Effect of bacteriophage application on *Campylobacter jejuni* loads in commercial broiler flocks. *Applied and Environmental Microbiology*, 79(23): 7525–7533. https://doi.org/10.1128/AEM.02703-13
- Richards, P.J., Connerton, P.L. & Connerton, I.F. 2019. Phage biocontrol of *Campylobacter jejuni* in chickens does not produce collateral effects on the gut microbiota. *Frontiers in Microbiology*, 10: 476. https://doi. org/10.3389/fmicb.2019.00476

### A4.1.3 Feed and water additives approaches

#### Caprylic acid and other organic acids

- Currie, D., Green, M., Dufailu, O.A., Matthaios, P., Soultanas, P., McCartney, E., Lester, H. et al. 2018. Dietary supplementation with ferric tyrosine improves zootechnical performance and reduces caecal *Campylobacter* spp. load in broilers. *British Poultry Science*, 59(6): 646–653. https://doi. org/10.1080/00071668.2018.1507015
- De Los Santos, F.S., Donoghue, A.M., Venkitanarayanan, K., Dirain, M.L., Reyes-Herrera, I., Blore, P.J. & Donoghue, D.J. 2008. Caprylic acid supplemented in feed reduces Enteric *Campylobacter jejuni* colonization in ten-day-old broiler chickens. *Poultry Science*, 87(4): 800–804. https:// doi.org/10.3382/ps.2007-00280
- De Los Santos, F.S., Donoghue, A.M., Venkitanarayanan, K., Metcalf,
   J.H., Reyes-Herrera, I., Dirain, M.L., Aguiar, V.F., Blore, P.J. &
   Donoghue, D.J. 2009. The natural feed additive caprylic acid decreases
   *Campylobacter jejuni* colonization in market-aged broiler chickens.
   *Poultry Science*, 88(1): 61–64. https://doi.org/10.3382/ps.2008-00228
- De Los Santos, F.S., Donoghue, A.M., Venkitanarayanan, K., Reyes-Herrera, I., Metcalf, J.H., Dirain, M.L., Aguiar, V.F., Blore, P.J. & Donoghue, D.J. 2008. Therapeutic supplementation of caprylic acid in feed reduces *Campylobacter jejuni* colonization in broiler chicks. *Applied and Environmental Microbiology*, 74(14): 4564–4566. https://doi.org/10.1128/ AEM.02528-07
- De Los Santos, F.S., Hume, M., Venkitanarayanan, K., Donoghue, A.M., Hanning, I., Slavik, M.F., Aguiar, V.F. *et al.* 2010. Caprylic acid reduces enteric *Campylobacter* colonization in market-aged broiler chickens but does not appear to alter cecal microbial populations. *Journal of Food Protection*, 73(2): 251–257. https://doi.org/10.4315/0362-028X-73.2.251
- Greene, G., Koolman, L., Whyte, P., Lynch, H., Coffey, A., Lucey, B., Egan, J., O'Connor, L. & Bolton, D. 2022. The efficacy of organic acid, medium chain fatty acid and essential oil based broiler treatments; *in vitro* anti-*Campylobacter jejuni* activity and the effect of these chemical-based treatments on broiler performance. *Journal of Applied Microbiology*, 132(1): 687–695. https://doi.org/10.1111/jam.15204

- Hovorková, P. & Skřivanová, E. 2015. Use of Caprylic acid in broiler chickens: effect on *Campylobacter jejuni*. *Foodborne Pathogens and Disease*, 12(8): 712–718. https://doi.org/10.1089/fpd.2015.1978
- Khattak, F., Paschalis, V., Green, M., Houdijk, J.G.M., Soultanas, P. & Mahdavi, J. 2018. TYPLEX<sup>®</sup> Chelate, a novel feed additive, inhibits *Campylobacter jejuni* biofilm formation and cecal colonization in broiler chickens. *Poultry Science*, 97(4): 1391–1399. https://doi.org/10.3382/ps/pex413
- Metcalf, J.H., Donoghue, A.M., Venkitanarayanan, K., Reyes-Herrera, I., Aguiar, V.F., Blore, P.J. & Donoghue, D.J. 2011. Water administration of the medium-chain fatty acid caprylic acid produced variable efficacy against enteric *Campylobacter* colonization in broilers 1, 2. *Poultry Science*, 90(2): 494–497. https://doi.org/10.3382/ps.2010-00891

## Probiotics

- Arsi, K., Donoghue, A.M., Woo-Ming, A., Blore, P.J. & Donoghue, D.J. 2015. The efficacy of selected probiotic and prebiotic combinations in reducing *Campylobacter* colonization in broiler chickens. *Journal of Applied Poultry Research*, 24(3): 327–334. https://doi.org/10.3382/japr/pfv032
- Cean, A., Stef, L., Simiz, E., Julean, C., Dumitrescu, G., Vasile, A., Pet, E., Drinceanu, D. & Corcionivoschi, N. 2015. Effect of human isolated probiotic bacteria on preventing *Campylobacter jejuni* colonization of poultry. *Foodborne Pathogens and Disease*, 12(2): 122–130. https://doi. org/10.1089/fpd.2014.1849
- Dec, M., Nowaczek, A., Urban-Chmiel, R., Stępień-Pyśniak, D. & Wernicki, A. 2018. Probiotic potential of *Lactobacillus* isolates of chicken origin with anti-*Campylobacter* activity. *Journal of Veterinary Medical Science*, 80(8): 1195–1203. https://doi.org/10.1292/jyms.18-0092
- Froebel, L.K., Jalukar, S., Lavergne, T.A., Coufal, C.D. & Duong, T. 2020. Administration of direct-fed *Bacillus cultures* and refined functional carbohydrates to broiler chickens improves growth performance and promotes positive shifts in gastrointestinal microbiota. *Journal* of Applied Poultry Research, 29(4): 765–774. https://doi.org/10.1016/ j.japr.2020.06.004

- Ghareeb, K., Awad, W.A., Mohnl, M., Porta, R., Biarnés, M., Böhm, J. & Schatzmayr, G. 2012. Evaluating the efficacy of an avian-specific probiotic to reduce the colonization of *Campylobacter jejuni* in broiler chickens. *Poultry Science*, 91(8): 1825–1832. https://doi.org/10.3382/ ps.2012-02168
- Gharib Naseri, K., Rahimi, S. & Khaki, P. 2012. Comparison of the effects of probiotic, organic acid and medicinal plant on *Campylobacter jejuni* challenged broiler chickens. *Journal of Agricultural Science and Technology*, 14(7): 1485–1496.
- Gorain, C., Singh, A., Bhattacharyya, S., Kundu, A., Lahiri, A., Gupta, S. & Mallick, A.I. 2020. Mucosal delivery of live *Lactococcus lactis* expressing functionally active JlpA antigen induces potent local immune response and prevent enteric colonization of *Campylobacter jejuni* in chickens. *Vaccine*, 38(7): 1630–1642. https://doi.org/10.1016/j.vaccine.2019.12.064
- Guo, J.R., Dong, X.F., Liu, S. & Tong, J.M. 2018. High-throughput sequencing reveals the effect of Bacillus subtilis CGMCC 1.921 on the cecal microbiota and gene expression in ileum mucosa of laying hens. *Poultry Science*, 97(7): 2543–2556. https://doi.org/10.3382/ps/pey112
- Khan, M. 2019. Effect of newly characterized probiotic lactobacilli on weight gain, immunomodulation and gut microbiota of *Campylobacter jejuni* challenged broiler chicken. *Pakistan Veterinary Journal*, 39(04): 473–478. https://doi.org/10.29261/pakvetj/2019.051
- Massacci, F.R., Lovito, C., Tofani, S., Tentellini, M., Genovese, D.A., De Leo,
   A.A.P., Papa, P. et al. 2019. Dietary saccharomyces cerevisiae boulardii
   CNCM I-1079 positively affects performance and intestinal ecosystem in
   broilers during a *Campylobacter jejuni* infection. *Microorganisms*, 7(12):
   596. https://doi.org/10.3390/microorganisms7120596
- Mortada, M., Cosby, D.E., Shanmugasundaram, R. & Selvaraj, R.K. 2020. In vivo and in vitro assessment of commercial probiotic and organic acid feed additives in broilers challenged with *Campylobacter coli. Journal of Applied Poultry Research*, 29(2): 435–446. https://doi.org/10.1016/ j.japr.2020.02.001
- Nazef, L., Belguesmia, Y., Tani, A., Prévost, H. & Drider, D. 2008. Identification of lactic acid bacteria from poultry feces: evidence on anti-*Campylobacter* and anti-*Listeria* activities. *Poultry Science*, 87(2): 329–334. https://doi. org/10.3382/ps.2007-00282

- Saint-Cyr, M.J., Haddad, N., Taminiau, B., Poezevara, T., Quesne, S., Amelot, M., Daube, G. et al. 2017. Use of the potential probiotic strain Lactobacillus salivarius SMXD51 to control Campylobacter jejuni in broilers. International Journal of Food Microbiology, 247: 9–17. https:// doi.org/10.1016/j.ijfoodmicro.2016.07.003
- Śmiałek, M., Kowalczyk, J. & Koncicki, A. 2021. The use of probiotics in the reduction of *Campylobacter* spp. prevalence in poultry. *Animals*, 11(5): 1355. https://doi.org/10.3390/ani11051355
- Szott, V., Reichelt, B., Friese, A. & Roesler, U. 2022. A complex competitive exclusion culture reduces *Campylobacter jejuni* colonization in broiler chickens at slaughter age in vivo. *Veterinary Sciences*, 9(4): 181. https:// doi.org/10.3390/vetsci9040181
- Tabashsum, Z., Peng, M., Alvarado-Martinez, Z., Aditya, A., Bhatti, J., Romo, P.B., Young, A. & Biswas, D. 2020. Competitive reduction of poultry-borne enteric bacterial pathogens in chicken gut with bioactive *Lactobacillus casei*. *Scientific Reports*, 10(1): 16259. https://doi. org/10.1038/s41598-020-73316-5
- Ty, M., Taha-Abdelaziz, K., Demey, V., Castex, M., Sharif, S. & Parkinson, J. 2022. Performance of distinct microbial based solutions in a *Campylobacter* infection challenge model in poultry. *Animal Microbiome*, 4(1): 2. https://doi.org/10.1186/s42523-021-00157-6
- Willis, W.L. & Reid, L. 2008. Investigating the effects of dietary probiotic feeding regimens on broiler chicken production and *Campylobacter jejuni* presence. *Poultry Science*, 87(4): 606–611. https://doi.org/10.3382/ ps.2006-00458
- Yuri Kuznetsov, I.N. 2020. Evaluation lactic acid bacteria autostrains with anti-*Campylobacter jejuni* activity on broiler chickens productivity. *International Transaction Journal of Engineering*, Management: 11A15S: 15. https://doi.org/10.14456/ITJEMAST.2020.307

### The addition of plant-based molecules to feed

Allaoua, M., Bonnafé, E., Etienne, P., Noirot, V., Gabarrou, J.-F., Castinel, A., Pascal, G. et al. 2022. A carvacrol-based product reduces *Campylobacter jejuni* load and alters microbiota composition in the caeca of chickens. *Journal of Applied Microbiology*, 132(6): 4501–4516. https://doi. org/10.1111/jam.15521

- Allaoua, M., Etienne, P., Noirot, V., Carayon, J.-L., Téné, N., Bonnafé, E. & Treilhou, M. 2018. Pharmacokinetic and antimicrobial activity of a new carvacrol-based product against a human pathogen, *Campylobacter jejuni. Journal of Applied Microbiology*, 125(4): 1162–1174. https://doi. org/10.1111/jam.13915
- Arsi, K., Donoghue, A.M., Venkitanarayanan, K., Kollanoor-Johny, A., Fanatico, A.C., Blore, P.J. & Donoghue, D.J. 2014. The efficacy of the natural plant extracts, thymol and carvacrol against *Campylobacter* colonization in broiler chickens. *Journal of Food Safety*, 34(4): 321–325. https://doi.org/10.1111/jfs.12129
- Arsi, K., Donoghue, A.M., Woo-Ming, A., Blore, P.J. & Donoghue, D.J. 2015. The efficacy of selected probiotic and prebiotic combinations in reducing *Campylobacter* colonization in broiler chickens. *Journal of Applied Poultry Research*, 24(3): 327–334. https://doi.org/10.3382/japr/pfv032
- Epps, S.V.R., Harvey, R.B., Byrd, J.A., Petrujkić, B.T., Sedej, I., Beier, R.C., Phillips, T.D. *et al.* 2015. Comparative effect of thymol or its glucose conjugate, thymol-β- d -glucopyranoside, on *Campylobacter* in avian gut contents. *Journal of Environmental Science and Health, Part B*, 50(1): 55–61. https://doi.org/10.1080/03601234.2015.965634
- Friedman, M. 2017. Chemistry, antimicrobial mechanisms, and antibiotic activities of cinnamaldehyde against pathogenic bacteria in animal feeds and human foods. *Journal of Agricultural and Food Chemistry*, 65(48): 10406–10423. https://doi.org/10.1021/acs.jafc.7b04344
- Gharib Naseri, K., Rahimi, S. & Khaki, P. 2012. Comparison of the effects of probiotic, organic acid and medicinal plant on *Campylobacter jejuni* challenged broiler chickens. *Journal of Agricultural Science and Technology*, 14(7): 1485–1496.
- Hankel, J., Popp, J., Meemken, D., Zeiger, K., Beyerbach, M., Taube, V., Klein, G. & Visscher, C. 2018. Influence of lauric acid on the susceptibility of chickens to an experimental *Campylobacter jejuni* colonisation. *PLoS ONE*, 13(9): e0204483. https://doi.org/10.1371/journal.pone.0204483
- Ibrahim, D., Sewid, A.H., Arisha, A.H., Abd El-fattah, A.H., Abdelaziz, A.M., Al-Jabr, O.A. & Kishawy, A.T.Y. 2020. Influence of Glycyrrhiza glabra extract on growth, gene expression of gut integrity, and *Campylobacter jejuni* colonization in broiler chickens. *Frontiers in Veterinary Science*, 7: 612063. https://doi.org/10.3389/fvets.2020.612063

- Kollanoor Johny, A., Darre, M.J., Donoghue, A.M., Donoghue, D.J.
  & Venkitanarayanan, K. 2010. Antibacterial effect of transcinnamaldehyde, eugenol, carvacrol, and thymol on *Salmonella* Enteritidis and *Campylobacter jejuni* in chicken cecal contents in vitro. *Journal of Applied Poultry Research*, 19(3): 237–244. https://doi.org/10.3382/japr.2010-00181
- Kurekci, C., Al Jassim, R., Hassan, E., Bishop-Hurley, S.L., Padmanabha, J. & McSweeney, C.S. 2014. Effects of feeding plant-derived agents on the colonization of *Campylobacter jejuni* in broiler chickens. *Poultry Science*, 93(9): 2337–2346. https://doi.org/10.3382/ps.2014-03950
- Lee, N.-K., Jung, B.S., Na, D.S., Yu, H.H., Kim, J.-S. & Paik, H.-D. 2016. The impact of antimicrobial effect of chestnut inner shell extracts against *Campylobacter jejuni* in chicken meat. *LWT - Food Science and Technology*, 65: 746–750. https://doi.org/10.1016/j.lwt.2015.09.004
- Salaheen, S., Tabashsum, Z., Gaspard, S., Dattilio, A., Tran, T.H. & Biswas, D. 2018. Reduced *Campylobacter jejuni* colonization in poultry gut with bioactive phenolics. *Food Control*, 84: 1–7. https://doi.org/10.1016/ j.foodcont.2017.07.021
- Szott, V., Reichelt, B., Alter, T., Friese, A. & Roesler, U. 2020. In vivo efficacy of carvacrol on *Campylobacter jejuni* prevalence in broiler chickens during an entire fattening period. *European Journal of Microbiology and Immunology*, 10(3): 131–138. https://doi.org/10.1556/1886.2020.00011
- Tolnai, E., Fauszt, P., Fidler, G., Pesti-Asboth, G., Szilagyi, E., Stagel, A., Konya, J. et al. 2021. Nutraceuticals induced changes in the broiler gastrointestinal tract microbiota. *mSystems*, 6(2): e01124-20. https://doi. org/10.1128/mSystems.01124-20
- Wagle, B.R., Donoghue, A.M., Shrestha, S., Upadhyaya, I., Arsi, K., Gupta, A., Liyanage, R. et al. 2020. Carvacrol attenuates Campylobacter jejuni colonization factors and proteome critical for persistence in the chicken gut. Poultry Science, 99(9): 4566–4577. https://doi.org/10.1016/ j.psj.2020.06.020
- Zeiger, K., Popp, J., Becker, A., Hankel, J., Visscher, C., Klein, G. & Meemken, D. 2017. Lauric acid as feed additive – An approach to reducing *Campylobacter* spp. in broiler meat. *PLoS ONE*, 12(4): e0175693. https:// doi.org/10.1371/journal.pone.0175693

#### In water plant-based additives and organic acids

- Ghareeb, K., Awad, W.A., Mohnl, M., Porta, R., Biarnés, M., Böhm, J. & Schatzmayr, G. 2012. Evaluating the efficacy of an avian-specific probiotic to reduce the colonization of *Campylobacter jejuni* in broiler chickens. *Poultry Science*, 91(8): 1825–1832. https://doi.org/10.3382/ ps.2012-02168
- Gharib Naseri, K., Rahimi, S. & Khaki, P. 2012. Comparison of the effects of probiotic, organic acid and medicinal plant on *Campylobacter jejuni* challenged broiler chickens. *Journal of Agricultural Science and Technology*, 14(7): 1485–1496.
- Johny, A.K., Darre, M.J., Hoagland, T.A., Schreiber, D.T., Donoghue, A.M., Donoghue, D.J. & Venkitanarayanan, K. 2008. Antibacterial effect of trans-cinnamaldehyde on Salmonella Enteritidis and Campylobacter jejuni in chicken drinking water. Journal of Applied Poultry Research, 17(4): 490–497. https://doi.org/10.3382/japr.2008-00051
- Metcalf, J.H., Donoghue, A.M., Venkitanarayanan, K., Reyes-Herrera, I., Aguiar, V.F., Blore, P.J. & Donoghue, D.J. 2011. Water administration of the medium-chain fatty acid caprylic acid produced variable efficacy against enteric *Campylobacter* colonization in broilers 1, 2. *Poultry Science*, 90(2): 494–497. https://doi.org/10.3382/ps.2010-00891
- Robyn, J., Rasschaert, G., Hermans, D., Pasmans, F. & Heyndrickx, M. 2013. Is allicin able to reduce *Campylobacter jejuni* colonization in broilers when added to drinking water? *Poultry Science*, 92(5): 1408–1418. https://doi. org/10.3382/ps.2012-02863
- Sparks, N.H.C. 2009. The role of the water supply system in the infection and control of *Campylobacter* in chicken. *World's Poultry Science Journal*, 65(3): 459–474. https://doi.org/10.1017/S0043933909000324
- Szott, V., Peh, E., Friese, A., Roesler, U., Kehrenberg, C., Ploetz, M. & Kittler,
   S. 2022. Antimicrobial effect of a drinking water additive comprising four organic acids on *Campylobacter* load in broilers and monitoring of bacterial susceptibility. *Poultry Science*, 101(12): 102209. https://doi.org/10.1016/j.psj.2022.102209

## Other additives and combination of additives

- Froebel, L.K., Jalukar, S., Lavergne, T.A., Coufal, C.D. & Duong, T. 2020. Administration of direct-fed *Bacillus* cultures and refined functional carbohydrates to broiler chickens improves growth performance and promotes positive shifts in gastrointestinal microbiota. *Journal* of Applied Poultry Research, 29(4): 765–774. https://doi.org/10.1016/ j.japr.2020.06.004
- Greene, G., Koolman, L., Whyte, P., Lynch, H., Coffey, A., Lucey, B., Egan, J., O'Connor, L. & Bolton, D. 2022. The efficacy of organic acid, medium chain fatty acid and essential oil based broiler treatments; *in vitro* anti-*Campylobacter jejuni* activity and the effect of these chemical-based treatments on broiler performance. *Journal of Applied Microbiology*, 132(1): 687–695. https://doi.org/10.1111/jam.15204
- Grilli, E., Vitari, F., Domeneghini, C., Palmonari, A., Tosi, G., Fantinati, P., Massi, P. & Piva, A. 2013. Development of a feed additive to reduce caecal *Campylobacter jejuni* in broilers at slaughter age: from *in vitro* to *in vivo*, a proof of concept. *Journal of Applied Microbiology*, 114(2): 308–317. https://doi.org/10.1111/jam.12053
- Wilson, K.M., Bourassa, D.V., McLendon, B.L., Wilson, J.L. & Buhr, R.J. 2018. Impact of skip-a-day and every-day feeding programs for broiler breeder pullets on the recovery of *Salmonella* and *Campylobacter* following challenge. *Poultry Science*, 97(8): 2775–2784. https://doi.org/10.3382/ps/ pey150

## A4.2 Processing

A total of 141 studies were reviewed by the processing group, including primary studies, reviews, and risk assessments that evaluated the efficiency of interventions implemented during poultry processing for *Campylobacter* control.

Al-Qadiri, H.M., Ovissipour, M., Al-Alami, N., Govindan, B.N., Shiroodi,
 S.G. & Rasco, B. 2016. Efficacy of neutral electrolyzed water, quaternary ammonium and lactic acid-based solutions in controlling microbial contamination of food cutting boards using a manual spraying technique. *Journal of Food Science*, 81(5).

- Bai, Y., Ding, X., Zhao, Q., Sun, H., Li, T., Li, Z., Wang, H. *et al.* 2022. Development of an organic acid compound disinfectant to control foodborne pathogens and its application in chicken slaughterhouses. *Poultry Science*, 101(6): 101842. https://doi.org/10.1016/j.psj.2022.101842
- Bailey, M.A., Taylor, R.M., Brar, J.S., Corkran, S.C., Velásquez, C., Novoa Rama, E., Oliver, H.F. & Singh, M. 2019. Prevalence and antimicrobial resistance of *Campylobacter* from antibiotic-free broilers during organic and conventional processing. *Poultry Science*, 98(3): 1447–1454. https:// doi.org/10.3382/ps/pey486
- Balta, I., Linton, M., Pinkerton, L., Kelly, C., Ward, P., Stef, L., Pet, I. et al. 2021. The effect of natural antimicrobials on the *Campylobacter coli* T6SS+/– during in vitro infection assays and on their ability to adhere to chicken skin and carcasses. *International Journal of Food Microbiology*, 338: 108998. https://doi.org/10.1016/j.ijfoodmicro.2020.108998
- Bauermeister, L.J., Bowers, J.W.J., Townsend, J.C. & McKee, S.R. 2008. The microbial and quality properties of poultry carcasses treated with peracetic acid as an antimicrobial treatment. *Poultry Science*, 87(11): 2390–2398. https://doi.org/10.3382/ps.2008-00087
- Bauermeister, L.J., Bowers, J.W.J., Townsend, J.C. & Mckee, S.R. 2008. Validating the efficacy of peracetic acid mixture as an antimicrobial in poultry chillers. *Journal of Food Protection*, 71(6): 1119–1122. https://doi. org/10.4315/0362-028X-71.6.1119
- Berrang, M.E. & Bailey, J.S. 2009. On-line brush and spray washers to lower numbers of *Campylobacter* and *Escherichia coli* and presence of *Salmonella* on broiler carcasses during processing. *Journal of Applied Poultry Research*, 18(1): 74–78. https://doi.org/10.3382/japr.2008-00067
- Berrang, M.E., Meinersmann, R.J., Smith, D.P. & Zhuang, H. 2008. The effect of chilling in cold air or ice water on the microbiological quality of broiler carcasses and the population of *Campylobacter. Poultry Science*, 87(5): 992–998. https://doi.org/10.3382/ps.2007-00406
- Berrang, M.E., Windham, W.R. & Meinersmann, R.J. 2011. Campylobacter, Salmonella, and Escherichia coli on broiler carcasses subjected to a high pH scald and low pH postpick chlorine dip. Poultry Science, 90(4): 896– 900. https://doi.org/10.3382/ps.2010-00900
- Bertram, R., Kehrenberg, C., Seinige, D. & Krischek, C. 2019. Peracetic acid reduces *Campylobacter* spp. numbers and total viable counts on broiler breast muscle and drumstick skins during modified atmosphere package storage. *Poultry Science*, 98(10): 5064–5073. https://doi.org/10.3382/ps/pez266

- Birk, T., Grønlund, A.C., Christensen, B.B., Knøchel, S., Lohse, K. & Rosenquist, H. 2010. Effect of organic acids and marination ingredients on the survival of *Campylobacter jejuni* on Meat. *Journal of Food Protection*, 73(2): 258–265. https://doi.org/10.4315/0362-028X-73.2.258
- Bourassa, D.V., Harris, C.E., Bartenfeld Josselson, L.N. & Buhr, R.J. 2021. Assessment of stabilized hydrogen peroxide for use in reducing *Campylobacter* levels and prevalence on broiler chicken wings. *Journal of Food Protection*, 84(3): 449–455. https://doi.org/10.4315/JFP-20-356
- Boysen, L. & Rosenquist, H. 2009. Reduction of thermotolerant *Campylobacter* species on broiler carcasses following physical decontamination at slaughter. *Journal of Food Protection*, 72(3): 497–502. https://doi.org/10. 4315/0362-028X-72.3.497
- Bucher, O., Waddell, L., Greig, J. & Smith, B.A. 2015. Systematic review-metaanalysis of the effect of chilling on *Campylobacter* spp. during primary processing of broilers. *Food Control*, 56: 211–217. https://doi.org/10.1016/ j.foodcont.2015.03.032
- Burfoot, D. & Mulvey, E. 2011. Reducing microbial counts on chicken and turkey carcasses using lactic acid. *Food Control*, 22(11): 1729–1735. https://doi.org/10.1016/j.foodcont.2011.04.005
- Burfoot, D., Allen, V., Mulvey, E., Jewell, K., Harrison, D. & Morris, V. 2015. Reducing *Campylobacter* numbers on chicken carcasses using lactic acid in processing plants. *International Journal of Food Science & Technology*, 50(11): 2451–2457. https://doi.org/10.1111/ijfs.12912
- Burfoot, D., Hall, J., Nicholson, K., Holmes, K., Hanson, C., Handley, S. & Mulvey, E. 2016. Effect of rapid surface cooling on *Campylobacter* numbers on poultry carcasses. *Food Control*, 70: 293–301. https://doi. org/10.1016/j.foodcont.2016.05.041
- Burfoot, D., Mulvey, E., Jewell, K., Foy, E. & Howell, M. 2015. Effect of electrolysed water on *Campylobacter* numbers on poultry carcasses under practical operating conditions at processing plants. *Food Control*, 50: 472–476. https://doi.org/10.1016/j.foodcont.2014.09.019
- Cano, C., Meneses, Y. & Chaves, B.D. 2021. Application of peroxyacetic acid for decontamination of raw poultry products and comparison to other commonly used chemical antimicrobial interventions: A review. *Journal* of Food Protection, 84(10): 1772–1783. https://doi.org/10.4315/JFP-21-107

- Casagrande Proietti, P., Pergola, S., Bellucci, S., Menchetti, L., Miraglia, D. & Franciosini, M.P. 2018. Occurrence and antimicrobial susceptibility of *Campylobacter* spp. on fresh and refrigerated chicken meat products in Central Italy. *Poultry Science*, 97(8): 2895–2901. https://doi.org/10.3382/ ps/pey147
- Chaine, A., Arnaud, E., Kondjoyan, A., Collignan, A. & Sarter, S. 2013. Effect of steam and lactic acid treatments on the survival of *Salmonella* Enteritidis and *Campylobacter jejuni* inoculated on chicken skin. *International Journal of Food Microbiology*, 162(3): 276–282. https://doi.org/10.1016/ j.ijfoodmicro.2013.01.012
- Chen, X., Bauermeister, L.J., Hill, G.N., Singh, M., Bilgili, S.F. & Mckee, S.R. 2014. Efficacy of various antimicrobials on reduction of *Salmonella* and *Campylobacter* and quality attributes of ground chicken obtained from poultry parts treated in a postchill decontamination tank. *Journal of Food Protection*, 77(11): 1882–1888. https://doi.org/10.4315/0362-028X.JFP-14-114
- Coşansu, S. & Ayhan, K. 2010. Effects of lactic and acetic acid treatments on *Campylobacter jejuni* inoculated onto chicken leg and breast meat during storage at 4c and -18c. *Journal of Food Processing and Preservation*, 34: 98–113. https://doi.org/10.1111/j.1745-4549.2008.00320.x
- De Villena, J.F., Vargas, D.A., Bueno López, R., Chávez-Velado, D.R., Casas, D.E., Jiménez, R.L. & Sanchez-Plata, M.X. 2022. Bio-mapping indicators and pathogen loads in a commercial broiler processing facility operating with high and low antimicrobial intervention levels. *Foods*, 11(6): 775. https://doi.org/10.3390/foods11060775
- Deblais, L., Helmy, Y.A., Kumar, A., Antwi, J., Kathayat, D., Acuna, U.M., Huang, H. et al. 2019. Novel narrow spectrum benzyl thiophene sulfonamide derivatives to control Campylobacter. The Journal of Antibiotics, 72(7): 555–565. https://doi.org/10.1038/s41429-019-0168-x
- Dirks, B.P., Dobrynin, D., Fridman, G., Mukhin, Y., Fridman, A. & Quinlan, J.J. 2012. Treatment of raw poultry with nonthermal dielectric barrier discharge plasma to reduce *Campylobacter jejuni* and *Salmonella enterica*. *Journal of Food Protection*, 75(1): 22–28. https://doi.org/10.4315/0362-028X.JFP-11-153

90

- Dittoe, D.K., Feye, K.M., Peyton, B., Worlie, D., Draper, M.J. & Ricke, S.C. 2019. The addition of ViriditecTM Aqueous ozone to peracetic acid as an antimicrobial spray increases air quality while maintaining *Salmonella* Typhimurium, non-pathogenic *Escherichia coli*, and *Campylobacter jejuni* reduction on whole carcasses. *Frontiers in Microbiology*, 9: 3180. https:// doi.org/10.3389/fmicb.2018.03180
- Dogan, O.B., Aditya, A., Ortuzar, J., Clarke, J. & Wang, B. 2022. A systematic review and meta-analysis of the efficacy of processing stages and interventions for controlling *Campylobacter* contamination during broiler chicken processing. *Comprehensive Reviews in Food Science and Food Safety*, 21(1): 227–271. https://doi.org/10.1111/1541-4337.12860
- Dogan, O.B., Clarke, J., Mattos, F. & Wang, B. 2019. A quantitative microbial risk assessment model of *Campylobacter* in broiler chickens: Evaluating processing interventions. *Food Control*, 100: 97–110. https://doi. org/10.1016/j.foodcont.2019.01.003
- Duc, V.M., Kakiuchi, R., Obi, T., Asakura, H. & Chuma, T. 2022. The incidence of *Campylobacter* contamination levels through chicken-sashimi processing steps in a small-scale poultry processing plant applying the external stripping method. *Journal of Veterinary Medical Science*, 84(3): 414–419. https://doi.org/10.1292/jvms.21-0486
- Duffy, L.L., Osmond-McLeod, M.J., Judy, J. & King, T. 2018. Investigation into the antibacterial activity of silver, zinc oxide and copper oxide nanoparticles against poultry-relevant isolates of *Salmonella* and *Campylobacter*. *Food Control*, 92: 293–300. https://doi.org/10.1016/ j.foodcont.2018.05.008
- Ellerbroek, L.I., Lienau, J. -A. & Klein, G. 2010. *Campylobacter* spp. in broiler flocks at farm level and the potential for cross-contamination during slaughter. *Zoonoses and Public Health*, 57(7–8). https://doi.org/10.1111/ j.1863-2378.2009.01267.x
- Erega, A., Stefanic, P., Dogsa, I., Danevčič, T., Simunovic, K., Klančnik, A., Smole Možina, S. & Mandic Mulec, I. 2021. Bacillaene mediates the inhibitory effect of bacillus subtilis on *Campylobacter jejuni* biofilms. *Applied and Environmental Microbiology*, 87(12): e02955-20. https://doi. org/10.1128/AEM.02955-20

- Figueroa, G., Troncoso, M., López, C., Rivas, P. & Toro, M. 2009. Occurrence and enumeration of *Campylobacter* spp. during the processing of Chilean broilers. *BMC Microbiology*, 9(1): 94. https://doi.org/10.1186/1471-2180-9-94
- Foddai, A., Takeuchi-Storm, N., Høg, B.B., Kjeldgaard, J.S., Andersen, J.K. & Ellis-Iversen, J. 2022. Assessing *Campylobacter* cross-contamination of Danish broiler flocks at slaughterhouses considering true flock prevalence estimates and ad-hoc sampling. *Microbial Risk Analysis*, 21: 100214. https://doi.org/10.1016/j.mran.2022.100214
- Franz, E., Van Der Fels-Klerx, H.J., Thissen, J. & Van Asselt, E.D. 2012. Farm and slaughterhouse characteristics affecting the occurrence of *Salmonella* and *Campylobacter* in the broiler supply chain. *Poultry Science*, 91(9): 2376–2381. https://doi.org/10.3382/ps.2009-00367
- García-Sánchez, L., Melero, B., Jaime, I., Hänninen, M.-L., Rossi, M. & Rovira, J. 2017. *Campylobacter jejuni* survival in a poultry processing plant environment. *Food Microbiology*, 65: 185–192. https://doi.org/10.1016/ j.fm.2017.02.009
- Gellynck, X., Messens, W., Halet, D., Grijspeerdt, K., Hartnett, E. & Viaene, J. 2008. Economics of reducing campylobacter at different levels within the Belgian Poultry Meat Chain. *Journal of Food Protection*, 71(3): 479–485. https://doi.org/10.4315/0362-028X-71.3.479
- Gichure, J.N., Kamau Njage, P.M., Wambui, J.M., Dykes, G.A., Buys, E.M.
  & Coorey, R. 2022. Systematic-review and meta-analysis on effect of decontamination interventions on prevalence and concentration of *Campylobacter* spp. during primary processing of broiler chickens. *Food Microbiology*, 102: 103923. https://doi.org/10.1016/j.fm.2021.103923
- **Golden, C.E. & Mishra, A.** 2020. Prevalence of *Salmonella* and *Campylobacter* spp. in alternative and conventionally produced chicken in the United States: A systematic review and meta-analysis. *Journal of Food Protection*, 83(7): 1181–1197. https://doi.org/10.4315/JFP-19-538
- **González-Fandos, E., Maya, N. & Pérez-Arnedo, I.** 2015. Effect of propionic acid on *Campylobacter jejuni* attached to chicken skin during refrigerated storage. *International Microbiology*(18): 171–175. https://doi. org/10.2436/20.1501.01.247
- Gonzalez, S.V., Geornaras, I., Nair, M.N. & Belk, K.E. 2021. Evaluation of immersion and spray applications of antimicrobial treatments for reduction of *Campylobacter jejuni* on chicken wings. *Foods*, 10(4): 903. https://doi.org/10.3390/foods10040903
- Gruntar, I., Biasizzo, M., Kušar, D., Pate, M. & Ocepek, M. 2015. Campylobacter jejuni contamination of broiler carcasses: Population dynamics and genetic profiles at slaughterhouse level. Food Microbiology, 50: 97–101. https://doi.org/10.1016/j.fm.2015.03.007
- Guirin, G.F., Brusa, V., Adriani, C.D. & Leotta, G.A. 2020. Prevalence of *Campylobacter jejuni* and *Campylobacter coli* from broilers at conventional and kosher abattoirs and retail stores. *Revista Argentina de Microbiología*, 52(3): 217–220. https://doi.org/10.1016/j.ram.2019.07.002
- Gunther, N.W. 2010. Effects of polyphosphate additives on *Campylobacter* survival in processed chicken exudates. *Applied and Environmental Microbiology*, 76(8): 2419–2424. https://doi.org/10.1128/AEM.02865-09
- Gunther, N.W., Phillips, J.G. & Sommers, C. 2016. The effects of 405-nm visible light on the survival of *Campylobacter* on chicken skin and stainless steel. *Foodborne Pathogens and Disease*, 13(5): 245–250. https://doi. org/10.1089/fpd.2015.2084
- Hakeem, M.J. & Lu, X. 2021. Survival and control of *Campylobacter* in poultry production environment. *Frontiers in Cellular and Infection Microbiology*, 10: 615049. https://doi.org/10.3389/fcimb.2020.615049
- Haughton, P.N., Lyng, J., Cronin, D., Fanning, S. & Whyte, P. 2012. Effect of crust freezing applied alone and in combination with ultraviolet light on the survival of *Campylobacter* on raw chicken. *Food Microbiology*, 32(1): 147–151. https://doi.org/10.1016/j.fm.2012.05.004
- Haughton, P.N., Lyng, J.G., Cronin, D.A., Morgan, D.J., Fanning, S. &
  Whyte, P. 2011. Efficacy of UV light treatment for the microbiological decontamination of chicken, associated packaging, and contact surfaces. *Journal of Food Protection*, 74(4): 565–572. https://doi.org/10.4315/0362-028X.JFP-10-356
- Haughton, P.N., Lyng, J.G., Morgan, D.J., Cronin, D.A., Fanning, S. & Whyte,
   P. 2011. Efficacy of high-intensity pulsed light for the microbiological decontamination of chicken, associated packaging, and contact surfaces. *Foodborne Pathogens and Disease*, 8(1): 109–117.
- Haydar Ozdemir, G.I.C., Gorkem Cengiz, B.O. & Sen, E. 2019. Effect of lactic acid and steam treatments on *Campylobacter jejuni* on chicken skin. *Emirates Journal of Food and Agriculture*: 143. https://doi.org/10.9755/ ejfa.2019.v31.i2.1915

- Horiuchi, I., Kawata, H., Nagao, T., Imaohji, H., Murakami, K., Kino, Y., Yamasaki, H. et al. 2015. Antimicrobial activity and stability of weakly acidified chlorous acid water. *Biocontrol Science*, 20(1): 43–51. https://doi. org/10.4265/bio.20.43
- Hovorková, P. & Skřivanová, E. 2015. Use of caprylic acid in broiler chickens: effect on *Campylobacter jejuni*. *Foodborne Pathogens and Disease*, 12(8): 712–718. https://doi.org/10.1089/fpd.2015.1978
- Huang, J., Zang, X., Zhai, W., Guan, C., Lei, T. & Jiao, X. 2017. Quantitative analysis of *Campylobacter* spp. contamination in chicken slaughtering lines by "label tracking method" in eastern China. *Food Control*, 80: 67–73. https://doi.org/10.1016/j.foodcont.2017.03.052
- Iannetti, S., Calistri, P., Di Serafino, G., Marotta, F., Alessiani, A., Antoci, S., Neri, D. et al. 2020. Campylobacter jejuni and Campylobacter coli: prevalence, contamination levels, genetic diversity and antibiotic resistance in Italy. Veterinaria Italiana, 56(1). https://doi.org/10.12834/ VetIt.1819.9596
- Isohanni, P., Alter, T., Saris, P. & Lyhs, U. 2010. Wines as possible meat marinade ingredients possess antimicrobial potential against *Campylobacter*. *Poultry Science*, 89(12): 2704–2710. https://doi.org/10.3382/ps.2009-00521
- Isohanni, P.M.I. & Lyhs, U. 2009. Use of ultraviolet irradiation to reduce *Campylobacter jejuni* on broiler meat. *Poultry Science*, 88(3): 661–668. https://doi.org/10.3382/ps.2008-00259
- Ivanova, M., Singh, R., Dharmasena, M., Gong, C., Krastanov, A. & Jiang, X. 2014. Rapid identification of *Campylobacter jejuni* from poultry carcasses and slaughtering environment samples by real-time PCR. *Poultry Science*, 93(6): 1587–1597. https://doi.org/10.3382/ps.2013-03736
- Jackowska-Tracz, A. & Tracz, M. 2015. Effects of high hydrostatic pressure on *Campylobacter jejuni* in poultry meat. *Polish Journal of Veterinary Sciences*, 18(2): 261–266. https://doi.org/10.1515/pjvs-2015-0034
- Jiang, W., Li, K., Chiu, Y.-C., Waldman, C. & Shen, C. 2020. Inactivation of *Campylobacter jejuni* in moisture enhanced non-intact chicken patties by double pan-broiling as affected by cooking set-up temperature and pump rate. *LWT*, 133: 109938. https://doi.org/10.1016/j.lwt.2020.109938

- Kameyama, M., Chuma, T., Nishimoto, T., Oniki, H., Yanagitani, Y., Kanetou, R., Gotou, K. et al. 2012. Effect of cooled and chlorinated chiller water on *Campylobacter* and coliform counts on broiler Carcasses during chilling at a middle-size poultry processing plant. *Journal of Veterinary Medical Science*, 74(1): 129–133. https://doi.org/10.1292/jvms.11-0167
- Kataria, J., Vaddu, S., Rama, E.N., Sidhu, G., Thippareddi, H. & Singh,
  M. 2020. Evaluating the efficacy of peracetic acid on *Salmonella* and *Campylobacter* on chicken wings at various pH levels. *Poultry Science*, 99(10): 5137–5142. https://doi.org/10.1016/j.psj.2020.06.070
- Koolman, L., Whyte, P., Meade, J., Lyng, J. & Bolton, D. 2014a. A Combination of chemical and ultrasonication treatments to reduce *Campylobacter jejuni* on raw poultry. *Food and Bioprocess Technology*, 7(12): 3602–3607. https://doi.org/10.1007/s11947-014-1370-2
- Koolman, L., Whyte, P., Meade, J., Lyng, J. & Bolton, D. 2014b. Use of chemical treatments applied alone and in combination to reduce *Campylobacter* on raw poultry. *Food Control*, 46: 299–303. https://doi.org/10.1016/ j.foodcont.2014.05.041
- Kottawatta, K., Van Bergen, M., Abeynayake, P., Wagenaar, J., Veldman, K. & Kalupahana, R. 2017. *Campylobacter* in broiler chicken and broiler meat in Sri Lanka: influence of semi-automated vs. wet market processing on *Campylobacter* contamination of broiler neck skin samples. *Foods*, 6(12): 105. https://doi.org/10.3390/foods6120105
- Kumar, S., Singh, M., Cosby, D.E., Cox, N.A. & Thippareddi, H. 2020. Efficacy of peroxy acetic acid in reducing *Salmonella* and *Campylobacter* spp. populations on chicken breast fillets. *Poultry Science*, 99(5): 2655–2661. https://doi.org/10.1016/j.psj.2019.12.045
- Kure, C.F., Axelsson, L., Carlehög, M., Måge, I., Jensen, M.R. & Holck, A. 2020. The effects of a pilot-scale steam decontamination system on the hygiene and sensory quality of chicken carcasses. *Food Control*, 109: 106948. https://doi.org/10.1016/j.foodcont.2019.106948
- Lake, R.J., Horn, B.J., Dunn, A.H., Parris, R., Green, F.T. & McNickle, D.C. 2013. Cost-effectiveness of interventions to control *Campylobacter* in the New Zealand poultry meat food supply. *Journal of Food Protection*, 76(7): 1161–1167. https://doi.org/10.4315/0362-028X.JFP-12-481

- Landrum, M.A., Cox, N.A., Cosby, D.E., Berrang, M.E., Mize, S.C. & Jackson, J.S. 2017. 879 reduction of *Campylobacter* on chicken livers using a lowacid processing aid. *Journal of Animal Science*, 95(suppl\_4): 377–377. https://doi.org/10.2527/asasann.2017.879
- Lecompte, J.-Y., Collignan, A., Sarter, S., Cardinale, E. & Kondjoyan, A. 2009. Decontamination of chicken skin surfaces inoculated with *Listeria innocua*, *Salmonella enteritidis* and *Campylobacter jejuni* by contact with a concentrated lactic acid solution. *British Poultry Science*, 50(3): 307– 317. https://doi.org/10.1080/00071660902942742
- Lee, N.-K., Jung, B.S., Na, D.S., Yu, H.H., Kim, J.-S. & Paik, H.-D. 2016. The impact of antimicrobial effect of chestnut inner shell extracts against *Campylobacter jejuni* in chicken meat. *LWT - Food Science and Technology*, 65: 746–750. https://doi.org/10.1016/j.lwt.2015.09.004
- Liu, Y., Betti, M. & Gänzle, M.G. 2012. High pressure inactivation of *Escherichia* coli, Campylobacter jejuni, and spoilage microbiota on poultry meat. Journal of Food Protection, 75(3): 497–503. https://doi.org/10.4315/0362-028X.JFP-11-316
- Luning, P.A., Jacxsens, L., Rovira, J., Osés, S.M., Uyttendaele, M. & Marcelis, W.J. 2011. A concurrent diagnosis of microbiological food safety output and food safety management system performance: Cases from meat processing industries. *Food Control*, 22(3–4): 555–565. https://doi. org/10.1016/j.foodcont.2010.10.003
- McWhorter, A.R., Weerasooriya, G., Willson, N.-L. & Chousalkar, K.K. 2022. Peroxyacetic acid and acidified sodium chlorite reduce microbial contamination on whole chicken carcasses obtained from two processing points. *Food Microbiology*, 106: 104035. https://doi.org/10.1016/ j.fm.2022.104035
- Melero, B., Vinuesa, R., Diez, A.M., Jaime, I. & Rovira, J. 2013. Application of protective cultures against *Listeria monocytogenes* and *Campylobacter jejuni* in chicken products packaged under modified atmosphere. Poultry Science, 92(4): 1108–1116. https://doi.org/10.3382/ps.2012-02539
- Meredith, H., McDowell, D. & Bolton, D.J. 2013. An evaluation of trisodium phosphate, citric acid and lactic acid cloacal wash treatments to reduce *Campylobacter*, total viable counts (TVC) and total enterobacteriaceae counts (TEC) on broiler carcasses during processing. *Food Control*, 32(1): 149–152. https://doi.org/10.1016/j.foodcont.2012.11.026

96

- Meredith, H., Walsh, D., McDowell, D.A. & Bolton, D.J. 2013. An investigation of the immediate and storage effects of chemical treatments on *Campylobacter* and sensory characteristics of poultry meat. *International Journal of Food Microbiology*, 166(2): 309–315. https://doi.org/10.1016/ j.ijfoodmicro. 2013.07.005
- Moazzami, M., Bergenkvist, E., FernströM, L.-L., Rydén, J. & Hansson, I. 2021. Reducing *Campylobacter jejuni, Enterobacteriaceae, Escherichia coli,* and total aerobic bacteria on broiler carcasses using combined ultrasound and steam. *Journal of Food Protection*, 84(4): 572–578. https://doi. org/10.4315/JFP-20-395
- Moore, A., Nannapaneni, R., Kiess, A. & Sharma, C.S. 2017. Evaluation of USDA approved antimicrobials on the reduction of *Salmonella* and *Campylobacter* in ground chicken frames and their effect on meat quality. *Poultry Science*, 96(7): 2385–2392. https://doi.org/10.3382/ps/pew497
- Morgan, R.B., Sierra–Arguello, Y.M., Perdoncini, G., Borges, K.A., Furian, T.Q., Gomes, M.J.P., Lima, D. *et al.* 2022. Comparison of transport crates contamination with *Campylobacter* spp. before and after the cleaning and disinfection procedure in broiler slaughterhouses. *Poultry Science*, 101(7): 101909. https://doi.org/10.1016/j.psj.2022.101909
- Musavian, H.S., Krebs, N.H., Nonboe, U., Corry, J.E.L. & Purnell, G. 2014. Combined steam and ultrasound treatment of broilers at slaughter: A promising intervention to significantly reduce numbers of naturally occurring campylobacters on carcasses. *International Journal of Food Microbiology*, 176: 23–28. https://doi.org/10.1016/j.ijfoodmicro.2014.02.001
- Nagel, G.M., Bauermeister, L.J., Bratcher, C.L., Singh, M. & McKee, S.R. 2013. Salmonella and Campylobacter reduction and quality characteristics of poultry carcasses treated with various antimicrobials in a post-chill immersion tank. International Journal of Food Microbiology, 165(3): 281–286. https://doi.org/10.1016/j.ijfoodmicro.2013.05.016
- Nair, D.V.T., Nannapaneni, R., Kiess, A., Mahmoud, B. & Sharma, C.S. 2014. Antimicrobial efficacy of lauric arginate against *Campylobacter jejuni* and spoilage organisms on chicken breast fillets. *Poultry Science*, 93(10): 2636–2640. https://doi.org/10.3382/ps.2013-03858
- Nannapaneni, R., Chalova, V., Crandall, P., Ricke, S., Johnson, M. & Obryan, C. 2009. Campylobacter and Arcobacter species sensitivity to commercial orange oil fractions. International Journal of Food Microbiology, 129(1): 43–49. https://doi.org/10.1016/j.ijfoodmicro.2008.11.008

- Nauta, M., Andersen, J.K., Tuominen, P., Ranta, J. & Lindqvist, R. 2015. Riskbased microbiological criteria for *Campylobacter* in broiler meat: A comparison of two approaches. *Food Control*, 53: 177–184. https://doi. org/10.1016/j.foodcont.2015.01.019
- Nauta, M.J., Sanaa, M. & Havelaar, A.H. 2012. Risk based microbiological criteria for *Campylobacter* in broiler meat in the European Union. *International Journal of Food Microbiology*, 158(3): 209–217. https://doi. org/10.1016/j.ijfoodmicro.2012.07.018
- Northcutt, J.K., Cason, J.A., Ingram, K.D., Smith, D.P., Buhr, R.J. & Fletcher, D.L. 2008a. Recovery of bacteria from broiler carcasses after immersion chilling in different volumes of water, Part 2. *Poultry Science*, 87(3): 573–576. https://doi.org/10.3382/ps.2006-00444
- Northcutt, J.K., McNeal, W.D., Ingram, K.D., Buhr, R.J. & Fletcher, D.L. 2008b. Microbial recovery from genetically featherless broiler carcasses after forced cloacal fecal expulsion. *Poultry Science*, 87(11): 2377–2381. https:// doi.org/10.3382/ps.2007-00426
- Northcutt, J.K., Smith, D., Huezo, R.I. & Ingram, K.D. 2008c. Microbiology of broiler carcasses and chemistry of chiller water as affected by water reuse. *Poultry Science*, 87(7): 1458–1463. https://doi.org/10.3382/ps.2007-00480
- **Olaimat, A.N., Fang, Y. & Holley, R.A.** 2014. Inhibition of *Campylobacter jejuni* on fresh chicken breasts by κ-carrageenan/chitosan-based coatings containing allyl isothiocyanate or deodorized oriental mustard extract. *International Journal of Food Microbiology*, 187: 77–82. https://doi.org/10.1016/j.ijfoodmicro.2014.07.003
- Ortiz-Suárez, L.E., Redondo-Solano, M., Arias-Echandi, M.L., Valenzuela-Martínez, C. & Peña-Ramos, E.A. 2021. Optimization of the in vitro bactericidal effect of a mixture of chlorine and sodium gallate against *Campylobacter* spp. and *Arcobacter butzleri. Journal of Food Protection*, 84(7): 1127–1135. https://doi.org/10.4315/JFP-20-181
- Osiriphun, S., Iamtaweejaloen, P., Kooprasertying, P., Koetsinchai, W., Tuitemwong, K., Erickson, L.E. & Tuitemwong, P. 2011. Exposure assessment and process sensitivity analysis of the contamination of *Campylobacter* in poultry products. *Poultry Science*, 90(7): 1562–1573. https://doi.org/10.3382/ps.2009-00577

- Pacholewicz, E., Sura Barus, S.A., Swart, A., Havelaar, A.H., Lipman, L.J.A.
  & Luning, P.A. 2016. Influence of food handlers' compliance with procedures of poultry carcasses contamination: A case study concerning evisceration in broiler slaughterhouses. *Food Control*, 68: 367–378. https://doi.org/10.1016/j.foodcont.2016.04.009
- Purnell, G., James, C., James, S.J., Howell, M. & Corry, J.E.L. 2014. Comparison of acidified sodium chlorite, chlorine dioxide, peroxyacetic acid and tri-sodium phosphate spray washes for decontamination of chicken carcasses. *Food and Bioprocess Technology*, 7(7): 2093–2101. https://doi. org/10.1007/s11947-013-1211-8
- Rasschaert, G., Piessens, V., Scheldeman, P., Leleu, S., Stals, A., Herman, L., Heyndrickx, M. & Messens, W. 2013. Efficacy of electrolyzed oxidizing water and lactic acid on the reduction of *Campylobacter* on naturally contaminated broiler carcasses during processing. *Poultry Science*, 92(4): 1077–1084. https://doi.org/10.3382/ps.2012-02771
- Rattanachaikunsopon, P. & Phumkhachorn, P. 2010. Potential of coriander (*Coriandrum sativum*) oil as a natural antimicrobial compound in controlling *Campylobacter jejuni* in raw meat. *Bioscience, Biotechnology, and Biochemistry*, 74(1): 31–35. https://doi.org/10.1271/bbb.90409
- Reich, F., Atanassova, V., Haunhorst, E. & Klein, G. 2008. The effects of *Campylobacter* numbers in caeca on the contamination of broiler carcasses with *Campylobacter*. *International Journal of Food Microbiology*, 127(1–2): 116–120. https://doi.org/10.1016/j.ijfoodmicro.2008.06.018
- Ren, T., Hayden, M., Qiao, M., Huang, T.-S., Ren, X. & Weese, J. 2018. Absorbent pads containing N -Halamine compound for potential antimicrobial use for chicken breast and ground chicken. *Journal* of Agricultural and Food Chemistry, 66(8): 1941–1948. https://doi. org/10.1021/acs.jafc.7b05191
- Riedel, C.T., Brøndsted, L., Rosenquist, H., Haxgart, S.N. & Christensen, B.B. 2009. Chemical decontamination of *Campylobacter jejuni* on chicken skin and meat. *Journal of Food Protection*, 72(6): 1173–1180. https://doi. org/10.4315/0362-028X-72.6.1173
- Romero-Barrios, P., Hempen, M., Messens, W., Stella, P. & Hugas, M. 2013. Quantitative microbiological risk assessment (QMRA) of food-borne zoonoses at the European level. *Food Control*, 29(2): 343–349. https://doi. org/10.1016/j.foodcont.2012.05.043

- Rossow, M., Ludewig, M. & Braun, P.G. 2018. Effect of cold atmospheric pressure plasma treatment on inactivation of *Campylobacter jejuni* on chicken skin and breast fillet. *LWT*, 91: 265–270. https://doi.org/10.1016/j.lwt.2018.01.052
- Sahin, O., Kassem, I.I., Shen, Z., Lin, J., Rajashekara, G. & Zhang, Q. 2015. Campylobacter in Poultry: Ecology and Potential Interventions. Avian Diseases, 59(2): 185–200. https://doi.org/10.1637/11072-032315-Review
- Sampers, I., Habib, I., Berkvens, D., Dumoulin, A., Zutter, L.D. & Uyttendaele, M. 2008. Processing practices contributing to *Campylobacter* contamination in Belgian chicken meat preparations. *International Journal of Food Microbiology*, 128(2): 297–303. https://doi.org/10.1016/ j.ijfoodmicro.2008.08.024
- Sarjit, A. & Dykes, G.A. 2015. Trisodium phosphate and sodium hypochlorite are more effective as antimicrobials against *Campylobacter* and *Salmonella* on duck as compared to chicken meat. *International Journal of Food Microbiology*, 203: 63–69. https://doi.org/10.1016/ j.ijfoodmicro.2015.02.026
- Sasaki, Y., Maruyama, N., Zou, B., Haruna, M., Kusukawa, M., Murakami, M., Asai, T., Tsujiyama, Y. & Yamada, Y. 2013. Campylobacter crosscontamination of chicken products at an abattoir. Zoonoses and Public Health, 60(2): 134–140. https://doi.org/10.1111/j.1863-2378.2012.01509.x
- Schambach, B.T., Berrang, M.E., Harrison, M.A. & Meinersmann, R.J. 2014. Chemical additive to enhance antimicrobial efficacy of chlorine and control cross-contamination during immersion chill of broiler carcasses. *Journal of Food Protection*, 77(9): 1583–1587. https://doi. org/10.4315/0362-028X.JFP-14-092
- Schroeder, M.W., Eifert, J.D., Ponder, M.A. & Schmale, D.G. 2014. Association of *Campylobacter* spp. levels between chicken grow-out environmental samples and processed carcasses. *Poultry Science*, 93(3): 734–741. https:// doi.org/10.3382/ps.2013-03646
- Sears, A., Baker, M.G., Wilson, N., Marshall, J., Muellner, P., Campbell, D.M., Lake, R.J. & French, N.P. 2011. Marked Campylobacteriosis decline after interventions aimed at poultry, New Zealand. *Emerging Infectious Diseases*, 17(6): 1007–1015. https://doi.org/10.3201/eid/1706.101272
- Seliwiorstow, T., Baré, J., Berkvens, D., Van Damme, I., Uyttendaele, M. & De Zutter, L. 2016. Identification of risk factors for *Campylobacter* contamination levels on broiler carcasses during the slaughter process. *International Journal of Food Microbiology*, 226: 26–32. https://doi. org/10.1016/j.ijfoodmicro.2016.03.010

- Sevilla-Navarro, S., Marin, C., Cortés, V., García, C. & Catalá-Gregori, P. 2020. *Campylobacter* prevalence and risk factors associated with exceeding allowable limits in poultry slaughterhouses in Spain. *Veterinary Record*, 186(13): 415–415. https://doi.org/10.1136/vr.105558
- Sharma, C.S., Williams, S.K., Schneider, K.R., Schmidt, R.H. & Rodrick, G.E. 2012. Sodium metasilicate affects growth of *Campylobacter jejuni* in fresh, boneless, uncooked chicken breast fillets stored at 4 degrees Celsius for 7 days. *Poultry Science*, 91(9): 2324–2329. https://doi.org/10.3382/ ps.2011-01868
- Shin, D., Martin, B.C. & Sánchez-Plata, M.X. 2011. Pulsed electric field effects to reduce the level of *Campylobacter* spp. in scalder and chiller water during broiler chicken processing. *Asian-Australasian Journal of Animal Sciences*, 24(9): 1314–1317. https://doi.org/10.5713/ajas.2011.11075
- Shrestha, S., Wagle, B.R., Upadhyay, A., Arsi, K., Donoghue, D.J. & Donoghue, A.M. 2019. Carvacrol antimicrobial wash treatments reduce *Campylobacter jejuni* and aerobic bacteria on broiler chicken skin. *Poultry Science*, 98(9): 4073–4083. https://doi.org/10.3382/ps/pez198
- Shrestha, S., Wagle, B.R., Upadhyay, A., Arsi, K., Upadhyaya, I., Donoghue, D.J. & Donoghue, A.M. 2019. Edible coatings fortified with carvacrol reduce *Campylobacter jejuni* on chicken wingettes and modulate expression of select virulence genes. *Frontiers in Microbiology*, 10: 583. https://doi. org/10.3389/fmicb.2019.00583
- Šimunović, K., Zajkoska, S., Bezek, K., Klančnik, A., Barlič Maganja, D.
   & Smole Možina, S. 2020. Comparison of *Campylobacter jejuni* slaughterhouse and surface-water isolates indicates better adaptation of slaughterhouse isolates to the chicken host environment. *Microorganisms*, 8(11): 1693. https://doi.org/10.3390/microorganisms8111693
- Smith, J., Corkran, S., McKee, S.R., Bilgili, S.F. & Singh, M. 2015. Evaluation of post-chill applications of antimicrobials against *Campylobacter jejuni* on poultry carcasses. *Journal of Applied Poultry Research*, 24(4): 451–456. https://doi.org/10.3382/japr/pfv046
- Smulders, FJM., Gleisz, B., Sofka, D., Sacher, A., Omurtag, I., Paulsen, P. & Hilbert, F. 2011. Microbial ecology on poultry carcasses along the production line\* Mikrobielle Ökologie von Geflügelschlachtkörpern entlang der Fertigungslinie\*. Archiv für Lebensmittelhygiene (Archiv für Lebensmittelhygiene 62: 5, 170-174 (2011)): 170–174. https://doi. org/10.2376/0003-925X-62-170

- Soro, A.B., Whyte, P., Bolton, D.J. & Tiwari, B.K. 2020. Strategies and novel technologies to control *Campylobacter* in the poultry chain: a review. *Comprehensive Reviews in Food Science and Food Safety*, 19(4): 1353– 1377. https://doi.org/10.1111/1541-4337.12544
- Stella, S., Tirloni, E., Bernardi, C. & Grilli, G. 2021. Evaluation of effect of chilling steps during slaughtering on the *Campylobacter* sp. counts on broiler carcasses. *Poultry Science*, 100(3): 100866. https://doi.org/10.1016/ j.psj.2020.11.043
- Teh, A.H.T., Lee, S.M. & Dykes, G.A. 2019. Growth in the presence of specific antibiotics induces biofilm formation by a *Campylobacter jejuni* strain sensitive to them but not in resistant strains. *Journal of Global Antimicrobial Resistance*, 18: 55–58. https://doi.org/10.1016/ j.jgar.2019.05.020
- Thames, H.T., Fancher, C.A., Colvin, M.G., McAnally, M., Tucker, E., Zhang, L., Kiess, A.S., Dinh, T.T.N. & Sukumaran, A.T. 2022. The prevalence of *Salmonella* and *Campylobacter* on broiler meat at different stages of commercial poultry processing. *Animals*, 12(18): 2460. https://doi. org/10.3390/ani12182460
- Thanissery, R. & Smith, D.P. 2014. Marinade with thyme and orange oils reduces Salmonella Enteritidis and Campylobacter coli on inoculated broiler breast fillets and whole wings. Poultry Science, 93(5): 1258–1262. https:// doi.org/10.3382/ps.2013-03697
- Thormar, H., Hilmarsson, H., Thráinsson, J.H., Georgsson, F., Gunnarsson, E.
   & Dadadóttir, S. 2011. Treatment of fresh poultry carcases with emulsions of glycerol monocaprate (monocaprin) to reduce contamination with *Campylobacter* and psychrotrophic bacteria. *British Poultry Science*, 52(1): 11–19. https://doi.org/10.1080/00071668.2010.537308
- Vaddu, S., Kataria, J., Belem, T.S., Sidhu, G., Moller, A.E., Leone, C., Singh, M. & Thippareddi, H. 2021a. On-site generated peroxy acetic acid (PAA) technology reduces *Salmonella* and *Campylobacter* on chicken wings. *Poultry Science*, 100(7): 101206. https://doi.org/10.1016/j.psj.2021.101206
- Vaddu, S., Kataria, J., Rama, E.N., Moller, A.E., Gouru, A., Singh, M. & Thippareddi, H. 2021b. Impact of pH on efficacy of peroxy acetic acid against Salmonella, Campylobacter, and Escherichia coli on chicken wings. Poultry Science, 100(1): 256–262. https://doi.org/10.1016/ j.psj.2020.09.063

- Vetchapitak, T., Rana, M.S., Sasaki, S., Taniguchi, T., Sugiyama, S., Soejima, J., Luangtongkum, T., Yamaguchi, Y. & Misawa, N. 2021. A new disinfectant technique for *Campylobacter jejuni* and spoilage bacteria on chicken skin using a high-pressure pulsed jet spray apparatus. *Food Control*, 125: 107989. https://doi.org/10.1016/j.foodcont.2021.107989
- Vetchapitak, T., Shinki, T., Sasaki, S., Taniguchi, T., Luangtongkum, T. & Misawa, N. 2020. Evaluation of chemical treatment combined with vacuum and ultrasonication with a water resonance system for reducing *Campylobacter* on naturally contaminated chicken carcasses. *Food Control*, 112: 107087. https://doi.org/10.1016/j.foodcont.2020.107087
- Viglundsson, T. I., Agustsson, O. I., Sigurdsson, H., Kristjansson, B., Josafatsson, B., Hilmarsson, H., Palsdottir, E. O. & Kristanjsson, J.
  K. undated. New method of processing ice slurry for improved chilling of fresh foods. *International Institute of Refrigeration (IIR)*.[Cited 3 November 2023]. https://iifiir.org/datacite\_notices/144212
- Wagle, B.R., Arsi, K., Shrestha, S., Upadhyay, A., Upadhyaya, I., Bhargava, K., Donoghue, A. & Donoghue, D.J. 2019. Eugenol as an antimicrobial wash treatment reduces *Campylobacter jejuni* in postharvest poultry. *Journal of Food Safety*, 39(6): e12704. https://doi.org/10.1111/jfs.12704
- Wagle, B.R., Arsi, K., Upadhyay, A., Shrestha, S., Venkitanarayanan, K., Donoghue, A.M. & Donoghue, D.J. 2017. β-Resorcylic Acid, a phytophenolic compound, reduces *Campylobacter jejuni* in postharvest poultry. *Journal of Food Protection*, 80(8): 1243–1251. https://doi. org/10.4315/0362-028X.JFP-16-475
- Wagle, B.R., Shrestha, S., Arsi, K., Upadhyaya, I., Donoghue, A.M. & Donoghue, D.J. 2019a. Pectin or chitosan coating fortified with eugenol reduces *Campylobacter jejuni* on chicken wingettes and modulates expression of critical survival genes. *Poultry Science*, 98(3): 1461–1471. https://doi.org/10.3382/ps/pey505
- Wagle, B.R., Upadhyay, A., Upadhyaya, I., Shrestha, S., Arsi, K., Liyanage,
  R., Venkitanarayanan, K., Donoghue, D.J. & Donoghue, A.M.
  2019b. Trans-Cinnamaldehyde, Eugenol and Carvacrol reduce *Campylobacter jejuni* biofilms and modulate expression of select genes
  and proteins. *Frontiers in Microbiology*, 10: 1837. https://doi.org/10.3389/
  fmicb.2019.01837

- Weerasooriya, G., McWhorter, A.R., Khan, S. & Chousalkar, K.K. 2021. Transcriptomic response of *Campylobacter jejuni* following exposure to acidified sodium chlorite. *npj Science of Food*, 5(1): 23. https://doi. org/10.1038/s41538-021-00103-5
- Wieczorek, K. & Osek, J. 2015. Poultry flocks as a source of *Campylobacter* contamination of broiler carcasses. *Polish Journal of Veterinary Sciences*, 18(1): 101–106. https://doi.org/10.1515/pjvs-2015-0013
- Zampara, A., Sørensen, M.C.H., Gencay, Y.E., Grimon, D., Kristiansen, S.H., Jørgensen, L.S., Kristensen, J.R. et al. 2021. Developing innolysins against Campylobacter jejuni using a novel prophage receptor-binding protein. Frontiers in Microbiology, 12: 619028. https://doi.org/10.3389/ fmicb.2021.619028
- Zeiger, K., Popp, J., Becker, A., Hankel, J., Visscher, C., Klein, G. & Meemken, D. 2017. Lauric acid as feed additive – An approach to reducing *Campylobacter* spp. in broiler meat. *PLoS ONE*, 12(4): e0175693. https:// doi.org/10.1371/journal.pone.0175693
- Zhang, L., Garner, L.J., Mckee, S.R. & Bilgili, S.F. 2018. Effectiveness of several antimicrobials used in a postchill decontamination tank against *Salmonella* and *Campylobacter* on broiler carcass parts. *Journal of Food Protection*, 81(7): 1134–1141. https://doi.org/10.4315/0362-028X.JFP-17-507
- Zhang, L., Singh, P., Lee, H.C. & Kang, I. 2013. Effect of hot water spray on broiler carcasses for reduction of loosely attached, intermediately attached, and tightly attached pathogenic (*Salmonella* and *Campylobacter*) and mesophilic aerobic bacteria. *Poultry Science*, 92(3): 804–810. https:// doi.org/10.3382/ps.2012-02504
- Zhao, G., Huang, X., Zhao, J., Liu, N., Li, Y., Wang, L., Gao, Y. et al. 2020. Risk prevention and control points through quantitative evaluation of *Campylobacter* in a large broiler slaughterhouse. *Frontiers in Veterinary Science*, 7: 172. https://doi.org/10.3389/fvets.2020.00172
- Zweifel, C., Althaus, D. & Stephan, R. 2015. Effects of slaughter operations on the microbiological contamination of broiler carcasses in three abattoirs. *Food Control*, 51: 37–42. https://doi.org/10.1016/j.foodcont.2014.11.002

## A4.3 Post-processing

- Al-Qadiri, H.M., Ovissipour, M., Al-Alami, N., Govindan, B.N., Shiroodi,
   S.G. & Rasco, B. 2016. Efficacy of neutral electrolyzed water, quaternary ammonium and lactic acid-based solutions in controlling microbial contamination of food cutting boards using a manual spraying technique. *Journal of Food Science*, 81(5). https://doi.org/10.1111/1750-3841.13275
- Birk, T., Grønlund, A.C., Christensen, B.B., Knøchel, S., Lohse, K. & Rosenquist, H. 2010. Effect of organic acids and marination ingredients on the survival of *Campylobacter jejuni* on Meat. *Journal of Food Protection*, 73(2): 258–265. https://doi.org/10.4315/0362-028X-73.2.258
- Bolton, D., Meredith, H., Walsh, D. & McDowell, D. 2014. Poultry food safety control interventions in the domestic kitchen. *Journal of Food Safety*, 34(1): 34–41. https://doi.org/10.1111/jfs.12092
- De Jong, A.E.I., Verhoeff-Bakkenes, L., Nauta, M.J. & De Jonge, R. 2008. Crosscontamination in the kitchen: effect of hygiene measures. *Journal of Applied Microbiology*, 105(2): 615–624. https://doi.org/10.1111/j.1365-2672.2008.03778.x
- Eideh, A.M.F. & Al-Qadiri, H.M. 2011. Effect of refrigerated and frozen storage on the survival of *Campylobacter jejuni* in cooked chicken meat breast. *Journal of Food Science*, 76(1). https://doi.org/10.1111/j.1750-3841.2010.01924.x
- El-Shibiny, A., Connerton, P. & Connerton, I. 2009. Survival at refrigeration and freezing temperatures of *Campylobacter coli* and *Campylobacter jejuni* on chicken skin applied as axenic and mixed inoculums. *International Journal of Food Microbiology*, 131(2–3): 197–202. https://doi.org/10.1016/j.ijfoodmicro.2009.02.024
- Ivić-Kolevska, S., Miljković-Selimović, B. & Kocić, B. 2012. Survival of Campylobacter jejuni in chicken meat at frozen storage temperatures. Acta Microbiologica et Immunologica Hungarica, 59(2): 185–198. https:// doi.org/10.1556/amicr.59.2012.2.4
- Lahou, E., Wang, X., De Boeck, E., Verguldt, E., Geeraerd, A., Devlieghere, F. & Uyttendaele, M. 2015. Effectiveness of inactivation of foodborne pathogens during simulated home pan frying of steak, hamburger or meat strips. *International Journal of Food Microbiology*, 206: 118–129. https://doi.org/10.1016/j.ijfoodmicro.2015.04.014

- Lai, H., Tang, Y., Ren, F., Li, Z., Li, F., Cui, C., Jiao, X. & Huang, J. 2021. An investigation into the critical factors influencing the spread of *Campylobacter* during chicken handling in commercial kitchens in China. *Microorganisms*, 9(6): 1164. https://doi.org/10.3390/ microorganisms9061164
- Langsrud, S., Sørheim, O., Skuland, S.E., Almli, V.L., Jensen, M.R., Grøvlen, M.S., Ueland, Ø. & Møretrø, T. 2020. Cooking chicken at home: Common or recommended approaches to judge doneness may not assure sufficient inactivation of pathogens. *PLoS ONE*, 15(4): e0230928. https:// doi.org/10.1371/journal.pone.0230928
- Maziero, M.T. & Oliveira, T.C.R.M.D. 2010. Effect of refrigeration and frozen storage on the *Campylobacter jejuni* recovery from naturally contaminated broiler carcasses. *Brazilian Journal of Microbiology*, 41(2): 501–505. https://doi.org/10.1590/S1517-83822010000200034
- Mild, R.M., Joens, L.A., Friedman, M., Olsen, C.W., McHugh, T.H., Law, B. & Ravishankar, S. 2011. Antimicrobial edible apple films inactivate antibiotic resistant and susceptible *Campylobacter jejuni* strains on chicken breast. *Journal of Food Science*, 76(3). https://doi.org/10.1111/ j.1750-3841.2011.02065.x
- Nauta, M. J., Fischer, A. R., Van Asselt, E. D., De Jong, A. E., Frewer, L. J., & De Jonge, R. 2008. Food safety in the domestic environment: the effect of consumer risk information on human disease risks. *Risk Analysis: An International Journal*, 28(1): 179–192.10.1111/j.1539-6924.2008.01012.x
- Sampers, I., Habib, I., De Zutter, L., Dumoulin, A. & Uyttendaele, M. 2010. Survival of *Campylobacter* spp. in poultry meat preparations subjected to freezing, refrigeration, minor salt concentration, and heat treatment. *International Journal of Food Microbiology*, 137(2–3): 147–153. https:// doi.org/10.1016/j.ijfoodmicro.2009.11.013
- Trajkovic-Pavlovic, L., Novakovic, B., Martinov-Cvejin, M., Gusman, V., Bijelovic, S., Dragnic, N. & Balac, D. 2010. How a routine checking of *Escherichia coli* in retailed food of animal origin can protect consumers against exposition to *Campylobacter* spp. and *Listeria monocytogenes*? *Vojnosanitetski pregled*, 67(8): 627–633. https://doi.org/10.2298/ VSP1008627T

- Vaz, Vaz, C.S.L., Voss-Rech, D., Rebelatto, R., Duarte, S.C., Coldebella, A.
  & Bessa, M.C. 2021. Survival of *Campylobacter jejuni* in chicken at refrigeration and cooking temperatures. *Pesquisa Agropecuária Brasileira*, 56: e02405. https://doi.org/10.1590/s1678-3921.pab2021.v56.02405
- Verhoeff-Bakkenes, L., Beumer, R.R., De Jonge, R., Van Leusden, F.M. & De Jong, A.E.I. 2008. Quantification of *Campylobacter jejuni* cross-contamination via hands, cutlery, and cutting board during preparation of a chicken fruit salad. *Journal of Food Protection*, 71(5): 1018–1022. https://doi.org/10.4315/0362-028X-71.5.1018
- Zakarienė, G., Rokaitytė, A., Ramonaitė, S., Novoslavskij, A., Mulkytė, K., Zaborskienė, G. & Malakauskas, M. 2015. The antimicrobial effect of spice-based marinades against *Campylobacter jejuni* on contaminated fresh broiler wings. *Journal of Food Science*, 80(3). https://doi. org/10.1111/1750-3841.12788

## Microbiological Risk Assessment Series

- 1 FAO and WHO. 2002. Risk assessments of *Salmonella* in eggs and broiler chickens: interpretative summary.
- 2 FAO and WHO. 2022. Risk assessments of *Salmonella* in eggs and broiler chickens.
- 3 FAO and WHO. 2003. Hazard characterization for pathogens in food and water: guidelines.
- 4 FAO and WHO. 2004. Risk assessment of *Listeria monocytogenes* in ready-toeat foods: interpretative summary.
- 5 FAO and WHO. 2004. Risk assessment of *Listeria monocytogenes* in ready-toeat foods: technical report.
- 6 FAO and WHO. 2004. *Enterobacter sakazakii* and other microorganisms in powdered infant formula: meeting report.
- 7 FAO and WHO. 2008. Exposure assessment of microbiological hazards in food: guidelines.
- 8 FAO and WHO. 2005. Risk assessment of *Vibrio vulnificus* in raw oysters: interpretative summary and technical report.
- 9 FAO and WHO. 2005. Risk assessment of choleragenic *Vibrio cholerae* O1 and O139 in warm-water shrimp in international trade: interpretative summary and technical report.
- 10 FAO and WHO. 2006. *Enterobacter sakazakii* and *Salmonella* in powdered infant formula: meeting report.
- 11 FAO and WHO. 2008. Risk assessment of *Campylobacter* spp. in broiler chickens: interpretative summary.
- 12 FAO and WHO. 2008. Risk assessment of *Campylobacter* spp. in broiler chickens: technical report.
- 13 FAO and WHO. 2008. Viruses in food: scientific advice to support risk management activities: meeting report.
- 14 FAO and WHO. 2008. Microbiological hazards in fresh leafy vegetables and herbs: meeting report.
- 15 FAO and WHO. 2008. *Enterobacter sakazakii* (*Cronobacter* spp.) in powdered follow-up formula: meeting report.

- 16 FAO and WHO. 2011. Risk assessment of *Vibrio parahaemolyticus* in seafood: interpretative summary and technical report.
- 17 FAO and WHO. 2009. Risk characterization of microbiological hazards in food: guidelines.
- 18 FAO and WHO. 2010. Enterohaemorragic *Escherichia coli* in raw beef and beef products: approaches for the provision of scientific advice, meeting report.
- 19 FAO and WHO. 2009. *Salmonella* and *Campylobacter* in chicken meat: meeting report.
- 20 FAO and WHO. 2020. Risk assessment tools for *Vibrio parahaemolyticus* and *Vibrio vulnificus* associated with seafood: meeting report.
- 21 FAO and WHO. *Salmonella* spp. In bivalve molluscs: risk assessment and meeting report, in progress.
- 22 FAO and WHO. 2016. Selection and application of methods for the detection and enumeration of human-pathogenic halophilic *Vibrio* spp. in seafood: guidance.
- 23 FAO and WHO. 2014. Multicriteria-based ranking for risk management of foodborne parasites.
- 24 FAO and WHO. 2016. Statistical aspects of microbiological criteria related to foods: a risk managers guide.
- 25 FAO and WHO. 2020. Risk-based examples and approach for control of *Trichinella* spp. and *Taenia saginata* in meat: revised edition.
- 26 FAO and WHO. 2022. Ranking of low moisture foods in support of microbiological risk management: meeting report and systematic review.
- 27 FAO and WHO. 2022. Microbiological hazards in spices and dried aromatic herbs: meeting report.
- 28 FAO and WHO.2016. Microbial safety of lipid based ready-to-use foods for the management of moderate acute and severe acute malnutrition: first report.
- 29 FAO and WHO. 2021. Microbial safety of lipid based ready-to-use foods for the management of moderate acute and severe acute malnutrition: second report.
- 30 FAO and WHO. 2016. Interventions for the control of non-typhoidal *Salmonella* spp. in beef and pork: meeting report and systematic review.
- 31 FAO and WHO. 2018. Shiga toxin-producing *Escherichia coli* (STEC) and food: attribution, characterization, and monitoring.
- 32 FAO and WHO. 2019. Attributing illness caused by Shiga toxin-producing *Escherichia coli* (STEC) to specific foods: report.

- 33 FAO and WHO. 2019. Safety and quality of water used in food production and processing: meeting report.
- 34 FAO and WHO. 2019. Foodborne antimicrobial resistance: role of the environment, crops and biocides: meeting report.
- 35 FAO and WHO. 2021. Advances in science and risk assessment tools for *Vibrio parahaemolyticus* and *V. vulnificus* associated with seafood: meeting report.
- 36 FAO and WHO. 2021. Microbiological risk assessment guidance for food: guidance.
- 37 FAO and WHO. 2021. Safety and quality of water used with fresh fruits and vegetables.
- 38 FAO and WHO. 2022. *Listeria monocytogenes* in ready-to-eat (RTE) foods: attribution, characterization and monitoring, meeting report.
- 39 FAO and WHO. 2022. Control measures for Shiga toxin-producing *Escherichia coli* (STEC) associated with meat and dairy products, meeting report.
- 40 FAO and WHO. 2023. Safety and quality of water use and reuse in the production and processing of dairy products, meeting report.
- 41 FAO and WHO. 2023. Safety and quality of water used in the production and processing of fish and fishery products, meeting report.
- 42 FAO and WHO. 2023. Prevention and control of microbiological hazards in fresh fruits and vegetables. Parts 1 and 2: General principles, meeting report.
- 43 FAO and WHO. 2023. Prevention and control of microbiological hazards in fresh fruits and vegetables. Part 3: Sprouts, meeting report.
- 44 FAO and WHO. 2023. Prevention and control of microbiological hazards in fresh fruits and vegetables. Part 4: Specific commodities, meeting report.
- 45 FAO and WHO. 2023. Measures for the control of *Salmonella* spp. in poultry meat, meeting report.
- 46 FAO and WHO.2024. Measures for the control of *Campylobacter* spp. in chicken meat, meeting report.
- 47 FAO and WHO. Risk assessment of *Listeria monocytogenes* in foods, Part 1: Formal model, meeting report, in progress.
- 48 FAO and WHO. Risk assessment of *Listeria monocytogenes* in foods, Part 2: Risk Assessment, meeting report, in progress.
- 49 FAO and WHO. Risk assessment of viruses in foods. Part 1: Food attribution, analytical methods and indicator, meeting report, in progress.

To collate and assess the most recent scientific information relevant to the control of thermotolerant *Campylobacter* species in broiler production and chicken meat, the Joint FAO/WHO Expert Meeting on Microbiological Risk Assessment (JEMRA) convened a meeting in Rome, Italy in February 2023.

The expert committee reviewed the available data on *Campylobacter* control including scientific literature published from 2008 to October 2022 and data submitted in response to a call for data for this meeting. The experts: 1) determined the quality and quantity of evidence of control measures for *Campylobacter*, 2) evaluated the impact of measures to control *Campylobacter* in the broiler production chain, 3) determined which hazard-based interventions pertained specifically to *Campylobacter* and which were general to the control of foodborne pathogens in the pre- and post-harvest broiler production chain, and 4) reviewed and recommended revisions to the *Guidelines for the Control of Campylobacter and Salmonella in Chicken Meat*.

This report describes the output of this expert meeting and the advice herein is useful for both risk assessors and risk managers, at national and international levels and those in the food industry working to control the hazard in poultry.

Food Systems and Food Safety - Economic and Social Development jemra@fao.org http://www.fao.org/food-safety **Food and Agriculture Organization of the United Nations** Viale delle Terme di Caracalla 00153 Rome, Italy

Department of Nutrition and Food Safety jemra@who.int www.who.int/health-topics/food-safety/ **World Health Organization** 20 Avenue Appia 1211 Geneva 27, Switzerland



LODIO