

CURRENT STATUS AND OPTIONS FOR FOREST BIOTECHNOLOGIES IN DEVELOPING COUNTRIES

SUMMARY

The forestry sector differs from the crop or livestock sectors in a number of important ways. First, forest trees are highly heterozygous long-lived perennials with late sexual maturity and a lengthy regeneration cycle which places high priority on retaining genetic diversity as an insurance policy against rapid change. Second, most forest tree species have narrow regional adaptation so the numbers of species used for planting are much higher than for food crops. Third, forest trees serve as keystone species in dynamic ecosystems so managing against loss translates into more than tree survival. Fourth, forest trees are largely undomesticated although a few species have had some population-level improvement for one to four generations.

For management of naturally regenerated forests, DNA-based and biochemical markers are available for a growing number of tropical species. Today, findings are available to guide operational forest management plans including in developing countries, but only for a very limited number of the hundreds of tree species that are managed in naturally regenerated tropical forests. This area of forest biotechnology continues to expand, moving from tools development into more hypothesis-driven knowledge acquisition. Such research inquiry is a powerful source of pertinent knowledge for protecting tropical forests. This research is also moving from molecular markers into genomics. Biotechnology tools such as molecular markers and the field of genomics are therefore providing important knowledge about naturally regenerated tropical forests and important insights into the nature of the entire tropical forest ecosystems including the relationship between forest trees and the microbial communities with which they interact, which can influence the strategies employed for managing tropical forests.

For planted forests, although there is some overlap the range of biotechnologies used is generally quite different from that used for naturally regenerated forests. Plantations can have different types of management systems (e.g. intensive, semi-intensive) and use different types of genetic material (e.g. wild material, genetically improved trees). Depending on the level of management intensity and the genetic material used in the planted forest, different groups of biotechnologies can be used. For simplicity, three different groups of biotechnologies can be identified according to the type of planted forests, ranging from the least sophisticated to the most advanced.

A first group of biotechnologies is suitable for the least intensively managed planted forests, and includes a range of vegetative propagation methods (including micropropagation based on tissue culture), biofertilizers and genetic fingerprinting using molecular markers. It could also be complemented by conventional technologies, such as early-stage tree improvement programmes.

A second group of biotechnologies can be used for planted forests that provide industrial raw materials on a large planting scale. The single species used for plantations may be indigenous or exotic, but these plantations are intensively managed. This group of biotechnologies includes somatic embryogenesis (a tissue culture technique), molecular markers and quantitative trait locus (QTL) analyses, whole genome sequencing and functional genomics. A third and most sophisticated group of biotechnologies includes backward and reverse genomics approaches, whole-genome sequencing, low-cost vegetative propagation and genetic modification of forest trees. To date, the only report of commercial plantings with genetically modified (GM) trees is for poplar on 300 to 500 ha in China. However, most tree species used in planted forests have been successfully modified at the experimental level, and traits that have been the subject of extensive research include stem shape, herbicide resistance, flowering characteristics, lignin content, insect and fungal resistance.

Many developing countries currently have biosafety regulations for agricultural crops, including fruit-trees, although many others lack such frameworks and the capacity to implement them. There are no regulations, however, specific to the use of GM forest trees. Although policies and regulations adopted for agricultural crops are also likely to be used for forest trees, they present special challenges (long time frames and life spans, wild resource, major constituents of an ecosystem). Forests are not only trees, and forest ecosystems are more fragile, longer-lived and less closely controlled than crop fields. Decision-making is complicated by the fact that while agriculture is primarily viewed as a production system, forests are generally viewed as a natural system, important not only for the conservation of biodiversity but also for social and cultural values. Thus, the use of GM forest trees is viewed more as a political and environmental issue than as a technical or trade issue.

2.1 INTRODUCTION

In recent decades, forest biotechnology has grown into a dynamic portfolio of tools, moving beyond research into global trade and development. This portfolio concept is consistent with the *sensu lato* definition of biotechnology put forth in Article 2 of the UN Convention on Biological Diversity (CBD) “any technological application that uses biological systems, living organisms, or derivatives thereof, to make or modify products or processes for specific use.” This is the definition used in this and the other FAO background documents prepared for ABDC-10. The following is a brief description of the state of the world’s forests and some factors shaping forestry and forest biotechnology.

2.1.1 Forest and tree resources management - state of the world’s forests

Forests and other wooded areas perform key economic and ecological functions. Not only do they provide goods and livelihoods but they also protect soils, regulate water and absorb carbon. Forests also shelter much of the world’s biodiversity. FAO’s most recent review on the overall status of forest resources, the Global Forest Resources Assessment (FAO, 2006), indicate that the world has just under 4 billion hectares (ha) of forests, covering about 30 percent of the world’s land area. It also reveals that production of wood and non-wood forest products is the primary function for 34 percent of the world’s forests and that more than half of all forests are used for wood and non-wood production in combination with other functions such as soil and water protection, biodiversity conservation and recreation.

Only 5 percent of forests in the world are in plantations, with the balance found in natural or semi-natural, largely unmanaged and undomesticated forest stands. Planted forests are expanding and their contribution to global wood production is approaching 50 percent of the total. In 2004, the production of industrial roundwood was 1.6 billion cubic meters, representing some 45 percent of the global wood production, and forest products trade reached a total value of US\$327 billion. More than half the wood biomass consumed globally – and well over 80 percent in developing countries – is burned as fuel. About 1.6 billion people rely heavily on forest resources for their livelihoods (World Bank, 2001). Sixty million indigenous people living in the rain forests of Latin America, Southeast Asia and West Africa depend heavily on forests; 350 million people living in, or next to, dense forests rely on them for subsistence or income; and 1.2 billion people in developing countries use trees on farms to generate food and cash. Forest and tree resources are managed in different main types of systems, which are presented in Table 1. The intensity of management varies very much between primary natural forests and productive industrial plantations.

FAO (2006) indicates that the world's forested area is shrinking, particularly at tropical latitudes (Table 2). Only a few countries have seen a net increase in forested land area, and these include China, Vietnam, Cuba, Uruguay, Chile, United States and most of Europe, west and east. Forested land area is not increasing in tropical regions where biodiversity and growth rates per ha (not shown) are highest. This table points to a few of the factors shaping forest biotechnology opportunities.

TABLE 1

CHARACTERISTICS OF MAIN TYPES OF FOREST AND TREE RESOURCES MANAGEMENT

Naturally regenerated forests		Planted forests				Trees outside forests
Primary	Modified natural	Semi-natural		Plantations		
		Assisted natural regeneration	Planted component	Productive	Protective	
Forests of native species, where there are no clearly visible indications of human activities and the ecological processes are not significantly disturbed	Forests of naturally regenerated native species where there are clearly visible indications of human activities	Silvicultural practices by intensive management: <ul style="list-style-type: none"> • weeding by intensive • fertilizing • thinning • selective logging 	Forests of native species, established through planting or seeding intensively managed	Forests of introduced and/or native species, established through planting or seeding mainly for production of wood or non-wood goods	Forests of introduced and/or native species, established through planting or seeding mainly for provision of services	Stands smaller than 0.5 ha; tree cover in agricultural land (agroforestry systems, home gardens, orchards); trees in urban environments; and scattered along roads and in landscapes

TABLE 2

FORESTED AREAS AND ANNUAL RATES OF CHANGE FOR THE WORLD'S FORESTED LAND COVER BY REGION

Forest area	Land area (1 000 ha)	Land area (percent)	Annual change 2000-2005 (percent)	Forested countries with highest net increase
Africa	635 412	21.4	-0.62	Rwanda, Egypt
Asia and Pacific	734 243	25.8	+0.09	China, Vietnam, New Zealand
Europe	1 001 394	44.3	+0.07	Bulgaria, Spain
Latin America and Caribbean	859 925	47.3	-0.51	Uruguay, Chile, Cuba
North America	677 464	32.7	-0.01	United States
West and Central Asia	43 588	4.0	+0.03	Uzbekistan
World	3 952 025	30.3	-0.18	

Source: FAO (2006)

Rapid loss of forested areas is coming from changes in land use. In addition to deforestation, existing forests are being degraded by pathogens and pests, fire, atmospheric pollution, extreme weather events, climate change and unsustainable forest management practices.

2.1.2 Factors shaping forests, forestry and forest biotechnology

The following factors shape global opportunities, condition investment decisions and drive research priorities for forest biotechnologies. They also point to important differences in the use of biotechnologies compared with the crop or livestock sectors.

- Forest trees are highly heterozygous, long-lived perennials with late sexual maturity and a lengthy regeneration cycle which places high priority on retaining genetic diversity as an insurance policy against rapid change (Namkoong, Barnes and Burley, 1980).
- Most forest tree species have narrow, regional adaptation, so species numbers used for planting are orders of magnitude higher than those for food crops (Pautasso, 2009).
- Forest trees serve as keystone species in dynamic ecosystems, so managing against loss translates into more than tree survival (Whitham *et al.*, 2006). Survival for colonizing forest tree species often depends on the presence of specific symbiont microbial species (Bonfante and Anca, 2009).
- Forest trees are largely undomesticated although a few species have had some population-level improvement for one to four generations.
- Most of the world's forests have public ownership (Agrawal, Chatter and Hardin, 2008).
- A forest tree is utilized for multiple purposes, not a single product. A single log can be used for sawtimber, paper and pulp. Waste products from papermaking are sold in secondary markets. Pulping waste is a rich source of industrial solvents, livestock feed, lubricants and consumer products such as artificial vanillin and medication.

Against this context, the purpose of this Chapter is to review the state of biotechnology and its impact on forest activities. It addresses this first by looking at the past and then by looking forward. In looking at the past, Part 2.2 provides an overview of the history and status of application of conventional technologies in forestry with special attention to developing countries. Part 2.3 documents the current status of application of forest biotechnologies in developing countries. Part 2.4 provides an analysis of successes and failures of forest biotechnologies in developing countries, while Part 2.5 presents a small number of case studies. In looking forward, Part 2.6 addresses key issues in the sector where forest biotechnologies could be useful, Part 2.7 identifies options for developing countries and Part 2.8 presents priorities for action for the international community.

A. STOCKTAKING: LEARNING FROM THE PAST


2.2 OVERVIEW OF CONVENTIONAL TECHNOLOGIES IN DEVELOPING COUNTRIES

Oddly, planted forests are not domesticated forests. To explain this, consider the definition of domestication put forth by Allard (1960): “Domestication is the bringing of a wild species under the management of humans”. Another definition of domestication is when a plant or animal is modified for human use to the point where it relies solely on human intervention for its survival. Under either definition, even the most intensively managed forests are only semi-domesticated (Figure 1). This is seen as an opportunity by many authors, who advocate using advanced biotechnologies to accelerate domestication for the benefit of wood production (Robinson, 1999; Campbell *et al.*, 2003; Boerjan, 2005; Tuskan, 2007).

In any case, forest biotechnology applications have historically been developed for the benefit of planted forests. But today forests are still planted from undomesticated reproductive material, as explained below. Planted forests compose 5 percent of the world’s forested areas and a few forest tree species are in the early stages of domestication (Nelson and Johnsen, 2008) but even so, they are semi-domesticated at best (Figure 1). Forest biotechnology applications are specific to each type of forest.

FIGURE 1

TYPES OF PLANTED FORESTS SORTED BY DEGREE OF DOMESTICATION



A brief overview follows of some main applications of conventional technologies in the forestry sector in developing countries.

Forest tree improvement

Forest tree improvement spread as a concept in the twentieth century, well after the advent of quantitative genetics and World War II. The primary goal was to identify and select wild seed sources suitable for planted forests. Few recurrent breeding programmes developed from this. Breeding cycles were lengthy, in part because population level improvement was essential. Forest tree improvement proceeds along a separate trajectory from agriculture.

Namkoong, Barnes and Burley (1980) wrote on their opening page: “Tree breeding is now an accepted activity in approximately half of the countries of the world...the breeding strategy has stopped at the first generation concepts of selection, progeny tests and clonal or seedling seed orchards”. Three decades later, this still holds true. The decision to settle for a known seed or propagule source can be traced to shortfalls in the long-range stability, funding and continuity of efforts required to sustain any forest planting programme: political instability, policy shifts, timber surplus, land sales, warfare, famine, drought, extreme weather events, lost manufacturing capacity and shift in global markets. Rarely has the decision to halt a tree improvement programme rested on the choice of forest tree species, but shortfalls here include forest disease and pest outbreaks, poor wood quality and even a surfeit of seed production.

Tree improvement for indigenous species gained momentum after World War II, mostly in Europe, Canada, Australia, New Zealand and the United States when reliable and well-adapted seed sources were needed for massive planting programmes. The next step, making selections in natural stands that served as seed parents, was viewed as a radical practice that contrasted with natural regeneration, dysgenic logging and the occasional haphazard seed collection. These early programmes were government-led.

A few tree-improvement programmes matured into recurrent forest tree breeding programmes. Given large land and financial requirements, these became enduring public-private partnerships among governments, universities and timber companies. Vegetative propagation was used only in the early years, provided that the species could be propagated easily at a low cost. Whether seedlings or cuttings, the idea of a known/tested source of germplasm rapidly spread to Southern Hemisphere countries such as Argentina, Brazil, Chile, China, Colombia, the Congo, India, Malawi, South Africa and Zimbabwe, where introduced and indigenous forest species alike grew much faster than in the Northern Hemisphere. Notable among these were some of the world’s most successful exotic species today: *Pinus radiata*, *Eucalyptus* spp. and *Acacia* spp.

Southern Hemisphere tree improvement was founded on naturalized introductions, imported exotic species and a few indigenous species. Its link to markets and manufacturing grew with global trade. Multinational timber corporations could grow timber more cheaply in some Southern Hemisphere countries and this spurred closer connections between forest research initiatives in developed and developing countries. Planted forests and tree improvement programmes have reaped considerable benefits from globalization.

Recurrent tree breeding

Recurrent tree breeding refers here to the application of Mendelian genetics principles within a given silvicultural system for the purpose of improving the genetic quality of the forest. Its goal is to improve the genetic value of the population while maintaining genetic diversity. This advanced generation or recurrent breeding programme refers to population level improvement, not to the development of breeds or inbred lines. Few of the many forest tree species planted today have been subjected to even a single generation of population level improvement. This is a subtle but important point when comparing advances in forest biotechnology with advances in crop or livestock biotechnology. Forest tree breeders weigh the importance of genetic gain against the importance of sufficient genetic diversity, the avoidance of inbreeding depression and long-term uncertainty.

As such, the breeding programme requires highly skilled experts, considerable investment funds and continuity of effort, because it continuously provides the best individuals for planting with each new breeding-testing-selecting cycle (Balocchi, 1997; White and Carson, 2004). Selections are placed in a production population which can be a small indoor or outdoor orchard. For some programmes only a few seeds are needed for multiplying via vegetative propagation. Either seed or propagules may be sold or planted as “varietals” although each is highly heterozygous. To date, forest tree breeding programmes do not develop inbred lines or hybrid crosses as is the case with crop breeding.

In any breeding programme, the selection goal needs to be well defined. One important trait to consider in breeding programmes, be they conventional or biotechnology-based, is wood formation (Plomion *et al.*, 2000; Plomion, Pionneau and Baillères, 2003), which drives profit margin through age of harvest and product recovery.

For conifers, annual rings within a single tree generate differences in market value, and so much attention has centred on how to alter this aspect of wood formation. For example, the early corewood rings for *Pinus* spp. are less valued than the outer rings owing to their different warping and pulping qualities. Annual rings laid down at older ages compose so-called mature wood. Finding the genetic controls for wood quality at early and later

stages of development is a critical step for conifer plantations because higher quality wood in the early rings would lower the age of harvest or the “rotation age”. The rotation age is the earliest age in the tree’s lifespan at which harvest becomes profitable. Most forest trees can live decades or even centuries beyond the rotation age.

Even in the most intensively managed forest tree programmes, tree improvement has not followed the same path as the crop sector. Genetic gain is carefully balanced against genetic diversity. Unlike their crop counterparts, forest tree breeders attach great importance to maintaining genetic diversity for population level improvement. Genetic diversity is seen as an insurance policy against catastrophic loss beyond a single generation. Forest tree breeding programmes, so integral to molecular applications of forest biotechnology, work on long timelines as a biological necessity.

This biological imperative to balance genetic gain against genetic diversity has not only given rise to forest tree programmes that do not resemble those for crops or livestock, but also to novel solutions. One common approach in tree improvement programmes is to safeguard genetic diversity (Tanaka, Tsumura and Nakamura, 1999; FAO, 2001). Grafted archives are often established at multiple locations. Unlike agricultural crops, these are needed because there are no repositories to insure against the loss of indigenous forest tree species. The payoff for these backup collections often comes when these archives provide germplasm for disease resistance, catastrophic weather events or a change in market demands. Another and more cost-efficient method has been the multiple population breeding strategy which uses divergent selection and multiple populations for a 2-for-1 programme conserving genetic diversity at the same time as making genetic gain (Eriksson, Namkoong and Roberds, 1995; Williams, Hamrick and Lewis, 1995). In this respect too, the forest biotechnology portfolio follows a separate path from crops and livestock.

2.3 CURRENT STATUS OF APPLICATION OF FOREST BIOTECHNOLOGIES IN DEVELOPING COUNTRIES

Forest biotechnology can contribute to improving productivity and reducing vulnerability of forest ecosystems to disease, degradation and human disturbance. The challenge continues to be to ensure sufficient genetic gains while maintaining genetic diversity at the ecosystem and landscape levels. To date, forest biotechnology has provided knowledge on how to mitigate the effects of forest fragmentation on genetic diversity, and on how to promote gene flow by managing tropical forest ecosystems for pollination, seed dispersal and soil symbionts.

An overview now follows of applications of biotechnologies in naturally regenerated tropical forests and in planted forests. Some of the biotechnologies overlap, although the forest systems differ considerably.

2.3.1 Naturally regenerated tropical forests

Today, most molecular marker systems are DNA-based systems such as microsatellites (Brondani *et al.*, 1998; Yazdani *et al.*, 2003) or amplified fragment length polymorphisms (AFLPs) (Cervera *et al.*, 2000), although biochemical markers such as isozymes continue to provide important insights into tropical forest ecosystems (e.g. Brown and Moran, 1981; Hamrick, 2004). Molecular markers have been used for decades and are extensively reviewed in FAO (2007).

Molecular marker methods are available for a growing number of tropical hardwood species such as *Aucoumea klaineana*, *Bagassa guianensis*, *Entandrophragma cylindicum*, *Hopea odorata*, *Hymenea courbaril*, *Dryobalanops aromatica*, *Neobalanocarpus heimeii*, *Koompasia malaccense* and the endangered *Shorea lumutensi* (Born *et al.*, 2006, 2008; Garcia *et al.*, 2004; Hamrick and Murawski, 1990; Lacerda, Kanashiro and Sebbenn, 2008; Lee *et al.*, 2000, 2002, 2003, 2004a, 2004b, 2006; Lee and Krishnapillay, 2004; Lim *et al.*, 2002; Naito *et al.*, 2005; Ng, Lee and Koh, 2004; Ng *et al.*, 2006; Sebbenn *et al.*, 2008; Silva *et al.*, 2008). Today, findings are available to guide operational forest management plans in developing countries, but only for a very limited number of the hundreds of tree species that are managed in naturally regenerated tropical forests. This area of forest biotechnology continues to expand, moving from tools development into more hypothesis-driven knowledge acquisition (Table 3). Such research inquiry is a powerful source of pertinent knowledge for protecting tropical forests.

TABLE 3

HYPOTHESIS-DRIVEN MOLECULAR MARKER APPLICATIONS FOR INDIGENOUS TROPICAL FORESTS WHICH ARE NATURALLY REGENERATED

Topics	Region, biota or taxa	Reference
Life history and potential for resilience to climate change	Tropical forests	Hamrick, 2004
Phenology	Neotropics	Clark, 2004
Silvicultural diversity	Tropical forests	Finkeldey and Ziehe, 2004
Selective logging	<i>Shorea megistophylla</i>	Murawski, Dayanandan and Bawa, 1994
Organellar DNA diversity	<i>Cedrela odorata</i>	Cavers, Navarro and Lowe, 2003
Forest fragmentation	Many tree species	Nason and Hamrick, 1997; Young and Clarke, 2000
Genetic bottlenecks	<i>Pinus maximartinezii</i>	Ledig <i>et al.</i> , 1999
Reproductive biology	<i>Dunalia arborescens</i>	Cruz, 1981
Fitness by life cycle stage	<i>Platypodium elegans</i>	Hufford and Hamrick, 2003
Outcrossing rates	<i>Cordia alliodora</i>	Boshier <i>et al.</i> , 1995
Genetics of invasiveness	<i>Pinus</i> spp.	Richardson and Petit, 2006

The hypotheses were tested using DNA-based forest biotechnology tools in combination with other information sources such as meteorology, ecology and/or taxonomy

TABLE 4

EXAMPLES OF USE OF GENOMIC DATA IN FORESTRY

Research areas	Region, biota or taxa	Reference
Phylogeny, phylogeography, nuclear DNA diversity	Agroforestry <i>Leucaena</i> spp.	Hughes, Eastwood and Bailey, 2006
Speciation and the study of mechanisms which generate biodiversity	Neotropical forest genus <i>Inga</i>	Richardson <i>et al.</i> , 2001
Rapid species identification via DNA barcoding	Worldwide	CBOL Plant Working Group, 2009
DNA-based Phylogeny Tree of Life Project	Worldwide	Burleigh and Matthews, 2004

This research community is also moving from molecular markers into genomics. Genomics refers to sequencing DNA either from the nuclear genome or from plastid and mitochondrial organelles. Unlike other areas of forest biotechnology, genomics data are often found in the public domain, usually internet databases (see review in Dean, 2006) and this affords the opportunity for DNA-based computational biology research. This availability of DNA sequencing data brings a distinct advantage to worldwide research on tropical forests.

To date, genomics data are yielding new insights into comparative biology for tropical forests (Table 4). Perhaps the application of most immediate use is an international plant barcoding project under the Consortium for the Barcode of Life (CBOL) to identify genes that can be used to distinguish between plant species¹.

More recent applications from DNA sequencing are emerging for the study of naturally regenerated tropical forests. This emerging use of genomics has been applied to several areas of inquiry, including phylogeny, which refers to comparing two or more DNA sequences from related forest trees with their near relatives to infer past divergence and speciation events. DNA sequences can be assumed to diverge in a steady-state, linear manner such that they serve as a molecular clock (Table 3). A closely related area of inquiry is phylogeography, which refers to using DNA-based sequence data to infer the history and formation of one or more taxa (Table 4).

Genomics has yet to provide its full benefit: it is a growth area for the forest biotechnology portfolio. DNA sequencing can encompass well-characterized genes, entire chromosomes or even entire genomes. Not only are related taxa being compared but interrelationships among components of entire forest ecosystems can be studied. Taxonomy, complemented by phylogeny, has now given way to phylogeography and phylogenomics, where functional genes are compared across taxa (Eisen and Fraser, 2003; Burleigh and Matthews, 2004). DNA sequence data are available for comparative analyses via Internet databases (Table 5).

¹ http://barcoding.si.edu/plant_working_group.html

Biotechnology tools such as molecular markers and genomics can therefore provide important knowledge about naturally regenerated tropical forests and important insights into the nature of the entire tropical forest ecosystems, including the relationship between the forest trees and the microbial communities with which they interact, which can influence the strategies employed for managing tropical forests.

Mycorrhizae are symbiotic associations that form between the roots of plant species and fungi. The hyphae (thread-like structures that are part of the body of the fungi) spread through the soil, taking up nutrients such as phosphorus and absorbing water, and transporting them to the plant root. In return, the fungi receive sugars from the plant (FAO, 2008a). Trees colonized with fungal symbionts are therefore likely to be more resistant to microbial pathogens and less stressed by drought. These benefits hold particular relevance for tropical forest ecosystems, given that drought and pathogen increases are predicted under climate change.

Genomics-based research is elucidating how this symbiotic complex functions. First, not all fungal symbionts have the same mechanisms, as genomics knowledge is confirming. The two major types of associations are ectomycorrhizae (EM) and vesicular-arbuscular mycorrhizae (VAM). While both buffer the tree host against diseases and abiotic stress, EM is more desirable for slowing forest degradation (Connell and Lowman, 1989) and for hastening re-colonization of abandoned land (Viera, Holl and Peneireiro, 2009). To this end, Connell and Lowman (1989) hypothesized that EM would confer a greater advantage to their host species than VAM.

TABLE 5

SOME EXAMPLES OF RELEVANT DNA SEQUENCE DATABASES

Database	URL
NCBI Taxonomy Browser Entrez Site Map	www.ncbi.nlm.nih.gov/ www.ncbi.nlm.nih.gov/Taxonomy/taxonomyhome.html/ www.ncbi.nlm.nih.gov/Entrez/ www.ncbi.nlm.nih.gov/Sitemap/
EMBL-EBI UniProt Site Map EMBL-Heidelberg Bioinformatics Tools	www.ebi.ac.uk/ www.ebi.ac.uk/uniprot/index.html www.ebi.ac.uk/services/index.html www.embl.de/services/bioinformatics/index.php
DENDROME Treegenes	http://dendrome.ucdavis.edu/ http://dendrome.ucdavis.edu/treegenes/

Source: adapted from Dean (2006)

Analysis requires specialized software, also available at some of these sites

The symbiont complex is more than the forest tree's roots and the fungal symbiont. Symbiont EM fungi also have specific bacterial associates which together form complexes with the host tree's roots (Bonfante and Anca, 2009). Together, this fungal-bacterial complex with the tree's roots confers benefits within the roots and surrounding area. Genomic sequencing of some fungal symbionts has been completed (Kuhn, Hijri and Sanders, 2001) and this is leading the way towards an emerging field known as community genomics which uses DNA sequencing tools to unravel these and other complex interactions within an entire forest ecosystem (Whitham *et al.*, 2006). This plethora of DNA sequencing methods not only applies to a single species or its near relatives, but also can provide insights into a tropical forest ecosystem. Its potential is already emerging for testing ideas about paleoecology and community ecology.

2.3.2 Planted Forests

Although there is some overlap, the biotechnologies used for planted forests are generally quite different from those used for naturally regenerated forests. It is also important to emphasize that there are different kinds of planted forests. Plantations can have different types of management systems (e.g. intensive, semi-intensive) and use different types of genetic material (e.g. wild material, genetically improved trees). Depending on the level of management intensity and the genetic material used in the planted forest, different groups of biotechnologies can be used. For simplicity, three different groups of biotechnologies can be identified according to the type of planted forests, ranging from the least sophisticated to the most advanced.

2.3.2.1 Basic forest biotechnologies

This group of biotechnologies is suitable for the least intensively managed planted forests and includes a range of vegetative propagation methods such as tissue culture, biofertilizers and genetic fingerprinting using molecular markers. It can also be complemented by conventional technologies such as early-stage tree improvement programmes. For these least intensively managed planted forests, the tropical forest restoration staircase (Chazdon, 2008) is the example that illustrates this type. This starts with planting reliable and well-adapted seed or propagule sources for reforestation. Poorly adapted, dysgenic plantings cannot hope to achieve such outcomes as restoring soil fertility for crop or forestry use, payment for ecosystem services, timber production or biodiversity recovery (Quesada *et al.*, 2009). In this first stage of planted forests, forest biotechnologies contribute to the health and quality of indigenous tropical forests and of exotic species.

Vegetative propagation of forest tree species

This covers a wide range of techniques which are useful for the rapid multiplication of genotypes. This has been useful for species which produce few or recalcitrant seeds or seedlings and for multiplying selected genotypes in a short period of time. It is also among the most ancient of forest applications, dating back eight centuries in China (Minghe and Ritchie, 1999). In India, there are about 8.9 million ha of teak forest, much of which is propagated by tissue culture (Tiwari, Tiwari and Siril, 2002). The National Chemical Laboratory in Pune and the Tata Research Institute in Delhi produce up to a few million teak plantlets annually. Phytosanitary measures also require tissue culture when moving germplasm from one country to another. This reduces the spread of plant viruses. Some of the disadvantages are the high costs of maintaining a tissue culture laboratory and quality control. Without quality control, one often sees the occurrence of somaclonal variations and deformed plantlets.

Micropropagation is the development of clonal lines from small tissue samples such as buds, roots or embryos extracted from seeds (Yanchuk, 2001) and some examples are provided in Table 6. The principles and achievements relating to plant tissue culture and micropropagation have been well reviewed by FAO (1994, 2004) and Yanchuk (2001). Thorpe, Harry and Kumar (1991) listed over 70 angiosperm and 30 gymnosperm tree species for which successful methods for the production of plantlets have been reported. Almost two decades ago, Le Roux and van Staden (1991) listed over 25 species of *Eucalyptus* alone. This, therefore, is a maturing part of the forest biotechnology portfolio.

TABLE 6

A FEW OF THE MANY FOREST TREE PLANTATION SPECIES WHICH HAVE BEEN MULTIPLIED THROUGH TISSUE CULTURE ON A COMMERCIAL SCALE IN DEVELOPING COUNTRIES

Countries	Species
India	<i>Tectona grandis</i> , <i>Anogeissus latifolia</i> <i>Bamboo</i> spp.
Indonesia, Malaysia and Vietnam	<i>Acacia mangium</i> and <i>Acacia mangium</i> x <i>Acacia auriculiformis</i> hybrids
India, Vietnam and South America	<i>Eucalyptus</i> spp.
Chile	<i>Pinus radiata</i>
Brazil, Indonesia, Malaysia and Thailand	<i>Tectona grandis</i>

Between 2002 and 2004, FAO commissioned four studies to investigate the extent and patterns of research and application in biotechnologies in forest trees worldwide. Results from the studies indicate that Asia accounts for 38 percent of the activities in forest tree micropropagation, followed by 7 percent in South America, 3 percent in Africa and 2 percent in Oceania (FAO, 2004). As expected, micropropagation of tree species is active mostly in countries with significant tree planting programmes (Galiana *et al.*, 2003; Watt *et al.*, 2003; Goh and Monteuuis, 2005; Goh *et al.*, 2007). While a large number of tree species (78 to 80) have been used for vegetative propagation research, little of this effort continues. Most halt at the laboratory stages (94 percent), so few even get as far as the field-testing stage (5 percent). Less than just 1 percent of the species developed clonally and tested have reached the commercial application stage (FAO, 2004).

Biofertilizers

Soils are dynamic living systems that contain a variety of micro-organisms such as bacteria, fungi and algae. Maintaining a favourable population of useful microflora is important from a fertility standpoint. The most commonly exploited micro-organisms are those that help in fixing atmospheric nitrogen for plant uptake or in solubilizing/mobilizing soil nutrients such as unavailable phosphorus into plant available forms, in addition to secreting growth promoting substances for enhancing crop yield. As a group, such microbes are called biofertilizers or microbial inoculants.

The use of biofertilizers has yielded positive results for indigenous forest species in the eastern Madagascar littoral forests as well as for exotic forest species including eucalypts, acacia and cypress (Kisa *et al.*, 2007; Duponnois *et al.*, 2007, 2008; Ouahmane *et al.*, 2007; Remigi *et al.*, 2008). Other symbionts that are being considered include nitrogen-fixing bacteria such as *Rhizobium*, and *Azolla*, blue-green algae and mycorrhizal fungi (Caesar, 2009). In addition to the least intensively managed planted forests in developing countries, biofertilizers have also proved useful in forests under more intensive management.

Genetic fingerprinting with molecular markers

All types of molecular and biochemical markers have been used for decades in these early-stage tree improvement programmes. A few examples are as follows:

- measuring genetic diversity of breeding population accessions between indigenous provenances and naturalized landrace origins;
- testing paternity contributions to offspring grown in field tests;
- verifying genetic identity during vegetative propagation.

2.3.2.2 Intermediate forest biotechnologies

The second group of biotechnologies can be used for planted forests that provide industrial raw materials on a large planting scale. The single species used for plantations might be indigenous or exotic, but the plantations are intensively managed.

Somatic embryogenesis

Somatic embryogenesis (SE) is a tissue culture technique that can also be used for the micropropagation of forest trees, where a small group of vegetative cells which are stem-cell like, are induced on culture media to undergo tissue differentiation to form a somatic embryo. The somatic embryo goes through a maturing process before being “germinated” for planting (Tartorius, Fowke and Dunstan, 1991).

Regeneration through SE has been reported for over 50 woody species encompassing over 20 angiosperm families, and at least a dozen conifer species (Wann 1988; Attree and Fowke, 1991; Tartorius, Fowke and Dunstan, 1991; Watt *et al.*, 1991; Park, Barret and Bonga, 1998). Potential multiplication rates particularly from cell suspension cultures are very high. Additional advantages include the amenability of the process to handling in automated bioreactors and the possibility for mechanized delivery of the emblings (plants propagated from SE) through synthetic seed technology. SE is also ideally suited for efficient genetic transformation procedures because of the single cell origin of embryos. The advantages of SE in comparison with micropropagation by *in vitro* cuttings, especially with regards to multiplication rate and genetic modification, explain why large research investments have been made towards developing this technique. Although successes have been reported in some commercial species, there are still major obstacles to the large-scale operational application of the technique to forest trees. Most of the reports (Table 7) demonstrate that the results obtained are still in the experimental stages and are yet to reach the commercialization phase.

TABLE 7

SPECIES IN WHICH SOMATIC EMBRYOGENESIS IS AT THE EXPERIMENTAL STAGE

Country	Species reported	References cited
Chile	<i>P. radiata</i> , <i>P. taeda</i>	Park, 2002; Jones, 2002; Lelu-Walter and Harvengt, 2004
Brazil and India	<i>Eucalyptus globulus</i> , <i>E. grandis</i> and <i>E. dunnii</i>	Pinto <i>et al.</i> , 2002; Watt <i>et al.</i> , 2003
India	<i>Tectona grandis</i> <i>Bamboo</i> Sandal	Krishnadas and Muralidharan, 2008 Godbole <i>et al.</i> , 2004; Shali and Muralidharan, 2008 Rathore <i>et al.</i> , 2008
Bangladesh	<i>Gmelina arborea</i> , <i>Artocarpus chaplasha</i> , <i>A. heterophyllus</i> , <i>Azadirachta indica</i> and <i>Elaeocarpus robustus</i>	Sarker, Islam Rafiqul and Hoque, 1997; Roy, Islam and Hadiuzzaman, 1998

Moreover, SE is a costly high-precision operation usually funded through multinational timber companies which own or lease land. Only the most elite selections are propagated using SE, where the genetic gains from recurrent breeding are maximized through the high-volume propagation of a single genotype.


Molecular markers, maps and QTL analyses

As part of this high precision operation, molecular markers also take on new functions. Breeding and selection in the recurrent breeding programme can be optimized by localizing chromosomal regions which influence the trait of interest.

No longer used only for genetic fingerprinting, markers are now used to find associations between traits and chromosomal regions. Forest trees, as perennial plants, have an added temporal dimension which can be challenging (Gwaze *et al.*, 2003) even for a single pedigree. If a marker interval is found to change the trait value then it becomes known as a quantitative trait locus (QTL) or a QTL haplotype, delineated by the relative position of two molecular markers. A QTL haplotype is not a gene but a single chromosomal segment inherited from either the maternal or paternal parent. It can include one or many genes exerting some degree of influence over a phenotypic trait.

FIGURE 2

FORWARD GENOMIC METHODS



Source: modified from Grattapaglia and Kirst (2008)

Finding QTL for forest trees is more costly and more computationally demanding than for most crop and livestock species because forest tree pedigrees are outcrossing and highly heterozygous (Devey *et al.*, 2003; Williams and Reyes-Valdes, 2007; Williams, Reyes-Valdés and Huber, 2007). Large pedigrees are rare with few generations and the populations have no breed structure or strong degrees of differentiation. However, numerous reports have cited the identification of QTL for major traits ranging from growth to wood quality and disease resistance in both Northern and Southern Hemisphere countries for a cadre of forest tree species. Moving from the genetic map to the physical map is more feasible with small hardwood genomes such as poplar and eucalypts. For species with large genomes, molecular cytogenetics technology or placement of dye-tagged DNA segments from known genes on a fixed chromosome squash is proving to be a useful bridge technology (Doudrick *et al.*, 1995).

Translating QTL identification into marker-assisted selection (MAS) is moving into the realm of commercial applications. Choosing MAS requires a cost-benefit analysis as described for *Pinus radiata* (Wilcox *et al.*, 2001) and this is used for poplars, *Eucalyptus* spp., *Pinus radiata* and a few cases of temperate-zone *Pinus* spp. especially those planted as an exotic. MAS has led to some novel breeding strategies when applied to forest trees (e.g. El-Kassaby and Lstiburek, 2009).


The selection of QTL haplotypes is not straightforward because a given pedigree can be segregating for more than one or even two QTL haplotypes of interest, which can result in ambiguous genetic models for QTL inheritance. But MAS is operational, lending yet another level of forest biotechnology precision to plantation forestry. Figure 2 shows how large numbers of molecular markers are assayed on gels for segregation patterns, then placed on a genetic map. Each individual now has a known genetic fingerprint, a collection of marker intervals or haplotypes – and some trait measurements. QTL haplotypes can be identified from these elements.

However, as mentioned earlier, finding QTL in single pedigrees is an arduous process for forest trees. Other methods for identifying QTL have since been developed or borrowed from other biological systems. These include association genomics which was developed for humans and other mammals (reviewed by Darvasi and Shifman, 2005) but these methods are well suited for forest trees (Brown *et al.*, 2004). Association genomics is a population level QTL detection method that is only effective if enough gametic disequilibrium is present in the population. This has indeed been the case for several intensively managed forest trees in both Northern and Southern Hemisphere regions.

Trait measurements constitute the phenotypic value of an individual tree. Trait-based genomics approaches such as QTL mapping are known as forward genomics (Figure 2). The trait measurements for each individual can be compared with its marker haplotypes.

FIGURE 3

FORWARD AND REVERSE GENOMIC METHODS



Source: modified from Grattapaglia and Kirst (2008) and Grattapaglia et al. (2009)

This is known as co-segregation between linked molecular markers and a putative QTL haplotype. While forward genomics is trait-based, the reverse genomics approach (Figure 3) is gene-based. Reverse genomics identifies, tests and validates specific genes controlling the trait of interest. Together, they provide an integrated picture of which genes or chromosomal segments are influencing the trait of interest and the degree of independence among these genes. The most sophisticated forest biotechnology portfolios at this time use both forward and reverse approaches but these are limited to *Eucalyptus* spp, *Populus* spp. and *Pinus* spp.

Whole genome sequencing projects for forest tree species

For hardwood species such as *Eucalyptus* spp. or *Populus* spp., adding reverse genomics is rapid and feasible because genome sizes are in the same range as those for rice, tomato and *Arabidopsis* (Wakamiya et al., 1993). Conifer genomes, by contrast, are larger than any commodity species in agriculture. The poplar genome was the first forest tree species to be sequenced in its entirety (Tuskan et al., 2006). The *Eucalyptus* genome initiative for whole-genome sequencing is an even larger effort that is being coordinated between 130 scientists in 18 countries including Brazil and South Africa. Sequencing the pine genome has less momentum given its enormous size, almost seven times larger than the human genome. A number of groups are completing part of the pine genome at present. A total of 100 large chromosomal segments (also known as bacterial artificial chromosomes or BACs) from *Pinus taeda* are presently being sequenced at the Joint Genome Institute in the United States.

The impetus for sequencing these forest tree genomes tends to come from large-scale wood production in intensively managed plantations worldwide, but forest health has also provided an equally compelling case. One of the side benefits of a large-scale DNA sequencing project is a rich store of new molecular markers such as single nucleotide polymorphisms (SNPs).

Functional genomics

In recent years, sequencing entire genomes has shifted emphasis from analyzing sequence data to the elucidation of gene function, also referred to as functional genomics. Gene function is inferred by using sequence alignment-based comparisons, identifying homologues between and within organisms, transcript profiling to determine gene expression patterns for small numbers of transcripts and yeast two-hybrid interaction analysis for identifying metabolic pathways, gene networks and protein complexes.

It is often conducted using microarrays which refer to the parallel assessment of gene expression for tens of thousands of genes. It works on the principle of competitive hybridization between complementary DNA (cDNA) strands. This approach can identify candidate genes for quantitative traits in forestry, a form of reverse genomics. As an example, cDNA microarray technology generated a transcript-level profile of wood forming tissues (differentiating xylem) for a pedigree composed of individuals from a *Eucalyptus grandis* x *E. globulus* F₁ hybrid x *E. grandis* backcross population (Kirst *et al.*, 2004). Microarrays are information rich sources of information about genes controlling the trait of interest.

Proteomics

Just as a genome describes the genetic content of an organism, a proteome defines the protein complement of the genome. Proteomics includes the identification of proteins in cells or tissues and the characterization of their physio-chemical properties such as post-translational modifications, function and expression level. Proteomics is a powerful tool for studying proteins and their modifications under different developmental stages and/or in response to various environmental stimuli.

In the cell, proteins form transitory or stable complexes as part of pathways and act within protein networks. These protein-protein interactions can be used to unravel the various interactions. After processing and modifications, a single gene may express between one and a few dozen different protein products. A combination of methods is required to characterize expressed proteins (or proteomes) fully.

A standard procedure is two-dimensional gel electrophoresis as the separation method followed by mass spectrometry analysis of the separated and enzymatically digested proteins. The peptide mass fingerprints typically obtained by mass spectrometry are matched against

sequence databases using dedicated bioinformatics tools. The whole procedure can be automated and robotized for high throughput purposes. The aim of this technique is to evaluate the modifications of protein expression with respect to genetic, environmental and developmental factors. The question is which quantitative variation of proteins is responsible for which quantitative phenotypic variation. The application of two-dimensional gel electrophoresis coupled with mass spectrometry in forest tree genomics to map the expressed genome has been well reviewed by Plomion, Pionneau and Baillères (2003).

2.3.2.3 Advanced forest biotechnologies

This uses the most sophisticated forest biotechnology portfolio yet. It includes recurrent tree breeding, backward and reverse genomics approaches, whole genome sequencing, low cost vegetative propagation and genetic modification of forest trees. The latter is the focus of this part.

To date, the only report of commercial plantings with genetically modified (GM) trees is for poplar on 300-500 ha in China (Xiao-hua *et al.*, 2003; FAO, 2004). However, most tree species used in planted forests have been successfully transformed at the experimental level, and results have demonstrated the correct expression of new genes in these plants (Walter *et al.*, 1998; Bishop-Hurley *et al.*, 2001).

Benefits from genetic modification can arise in particular from the transfer of traits from species as wide apart as bacteria or other plants that are not readily available either in the breeding population or in the forest tree species as a whole. Traits that have been the subject of extensive research for genetic modification include stem shape (taper and “roundness”), herbicide resistance, flowering characteristics, lignin content, insect and fungal resistance (Li *et al.*, 2003; Grace *et al.*, 2005; Punja, 2001; Shin *et al.*, 1994; Tang and Tian, 2003).

The potential environmental benefits of such technology (Gianessi *et al.*, 2002) include new means to combat pathogen and pest outbreaks. Some intensively managed tree improvement programmes have been investigating the use of gene transfer methods for many years. An example is the introduction of the *Cry1Ac* gene from the bacteria *Bacillus thuriengensis* (Bt) into radiata pine, where the ultimate goal is to enhance resistance to the pine shoot tip moth (Grace *et al.*, 2005). Less attention has been given to other applications of GM forest trees such as environmental remediation, land reclamation and mercury sequestration. Forest trees (conifers and hardwoods) are useful for land reclamation purposes even without genetic modification, so the potential value is considerable.

As mentioned earlier, a small area of GM poplar is also planted on a commercial scale in China. China has a highly productive forestry plantation programme with six national forest planting programmes. To date, at least seven million ha have been planted with fast growing poplars (FAO, 2008b).

The first successful transformation was done on *Populus nigra* with the *Cry1Ac* gene in 1993. This tree was used in field testing as early as 1994 and was subsequently deployed in further pilot plantings. In 2000, the Chinese regulatory authority permitted the establishment of about one million trees on 300 ha (Hu *et al.*, 2001). This was followed by a smaller release with a hybrid poplar clone transformed with both *Cry1Ac* and *PI* genes (Xiao-hua *et al.*, 2003). The toxicity of this transformed clone was greatly enhanced as the GM plants contained two insect resistance genes. Subsequently, the transformation of poplars for disease resistance and tolerance to environmental stresses has been achieved, though these are still at the laboratory stage (Xiao-hua *et al.*, 2003).

Genetic modification is part of the reverse genomics approach that is used to evaluate gene function but its commercialization is shifting investment from the public domain into proprietary areas. As noted by FAO (2004), a notable trend is that the “numbers of publicly funded projects appear to be waning, while privately funded projects appear to be increasing, judging by field trials established in recent years”. This is a capital intensive effort requiring long-term continuity of funding, scientists and infrastructure. Developing a GM genotype on a commercial scale first requires a well established gene transfer technology (Walter *et al.*, 1998). Each GM genotype must then be vegetatively propagated on a large scale before shareholders can expect a return on the steep initial investment.

The issue of GMOs has received considerable attention over the last decade in scientific and non-scientific circles and from policy-makers worldwide. The focus of attention has been on the crop sector which is where most GMOs have been commercialized. In 2008, an estimated 125 million ha were cultivated with GM crops compared with just 400 ha of Bt poplars in China, with 20 000 seedlings prepared for planting in 2009 (James, 2008). If or when further GM forest trees are released commercially this situation may change. A regulatory framework to govern research and the applications of GM forest trees is essential. The issue goes beyond the country level because pollen flow and seed dispersal do not respect national boundaries. National and international regulatory systems should contain provisions for preliminary risk assessments, monitoring and control and for liability and redress.

Many countries currently have regulations for agricultural crops including fruit trees, although many developing countries lack such frameworks and the capacity to implement them. There are, however, no regulations specific to the use of GMOs in forestry. Although policies and regulations adopted for agricultural crops are also likely to be used for forest trees, forest trees present special challenges (they have long time frames and life spans, they form a wild resource and are major constituents of an ecosystem). Forests are not only trees, and forest ecosystems are more fragile, longer-lived and less closely controlled than crop fields. Decision making is complicated by the fact that while agriculture is primarily viewed as a production system, forests are generally viewed as natural systems, important

not only for the conservation of biodiversity but also for social and cultural values. Thus the use of GM forest trees is viewed more as a political and environmental issue than as a technical or trade one (El-Lakany, 2004).

2.3.3 Summary

Based on the current analysis, Table 8 attempts to summarize and compare the current status, key issues and future perspectives for a number of conventional approaches and biotechnologies in developing countries. The different technologies differ in respect of public acceptance, the technical capacity and infrastructure/materials required for their use and costs. For the near future, it is predicted that the potential impact is high for tree improvement, genomics, DNA barcoding and biofertilizers. To complement this information, Table 9 summarizes the anticipated contribution of forest biotechnology applications to natural and planted forests for developing and developed countries based on the worldwide survey commissioned by FAO (2004).

TABLE 8

CURRENT STATUS OF SOME CONVENTIONAL TECHNOLOGIES AND BIOTECHNOLOGIES, AND FACTORS INFLUENCING THEIR APPLICABILITY IN THE FORESTRY SECTOR IN DEVELOPING COUNTRIES

Emerging forest biotechnology applications	Extent of use	Public and government acceptance	Current technical capability for using technology	Current technical capability for adapting or developing new technology	Infrastructure and/or materials and tools available for using technology	Relative cost	Skills required for application	Potential for generating impact (time frame < 10 years)
Tree improvement	High	High	High	Low	Medium	Medium	Medium	High
Recurrent tree breeding	Low	High	Low	High	Medium	High	High	Medium
Molecular markers	Medium	High	Medium	Medium	Low	Medium	Medium	Medium
Genomics	Low	High	Low	Low	Low	Medium	High	High
Bioinformatics	Medium	High	Medium	Medium	Medium	Medium	High	Low
Genetic modification	Low	Low to medium	Low	Low	Low	High	High	Low
Biofertilizers	High	High	High	High	Low	Low	Low	High
Comparative phylogeny	Low	High	Medium	Medium	Low	Medium	High	Low
DNA barcoding	Low	High	Medium	Medium	Medium	Medium	Low	High

TABLE 9

ANTICIPATED CONTRIBUTION OF FOREST BIOTECHNOLOGY APPLICATIONS TO NATURAL AND PLANTED FORESTS WORLDWIDE

Applicable forestry component	Spatial scale	Development elements relevant to biotechnology	Broad technologies						
			Molecular applications						Regeneration
			Bioinformatics	Diversity measurement	Gene discovery	Genetic modification	Biosensors	Product verification	
Natural populations	Tree–population	Genetic resources characterization	X	X	X				
	Population	Mating system/gene flow		X					
	Population–landscape	Conserving diversity		X					
	Population–landscape	Silvicultural impact assessment		X					
Breeding populations	Tree	Selection	X	X	X				
	Tree–population	Mating designs		X					X
	Tree	Testing	X	X	X			X	X
	Population	Diversity management		X	X				
Production populations	Population	Mating system	X	X					X
	Population	Gene flow	X	X				X	X
	Population–landscape	Silvicultural impact		X			X		X
Regeneration	Stand	Natural		X	X			X	X
	Stand	Planted		X	X	X		X	X
Domestication	Population	Native species diversity	X	X	X		X	X	
	Population	Exotic species suitability	X	X	X		X	X	
Gene conservation	Population	Diversity assessment	X	X					
	Population	Gene flow/contamination		X			X		X
	Tree	Reproduction		X	X		X		X
Forest health	Tree–stand	Risk/hazard assessment	X	X			X		
	Tree	Resistance screening	X	X	X	X	X		X
Processing/ Value added	Logs	Pulp processing	X		X	X	X	X	
	Logs	Wood treatment	X		X	X	X	X	

Source: adapted from Table 2.4.2 in FAO (2004)

2.4 ANALYSIS OF SUCCESSES AND FAILURES OF FOREST BIOTECHNOLOGIES IN DEVELOPING COUNTRIES

To date, the use of biotechnologies has been beneficial only at very advanced stages of selection and improvement programmes. Unlike crops, where the number of species to choose from is relatively limited, an immense diversity at both the interspecific and intraspecific level is used in forestry. Thanks to this important diversity, the early stages of classical selection (exploration, collection, testing of genetic resources) provide important gains. By way of example at the species level, *Acacia crassiparpa*, which is currently the main plantation species in swamp areas was “unknown” as a plantation species only 20 years ago. At intraspecific level, coordinated, multilocational provenance trials have shown sometimes 200 percent variation in adaptive traits among populations across the natural range of distribution of the same species. Individual variation within populations is also very important, and selection at this level also yields important initial gains in particular through clonal development (the traditional rooting of cuttings).

For most species and forest tree management systems, advances registered in developing countries until now have been made without any incorporation of biotechnologies. There are very good examples of advanced tree breeding programmes using biotechnology tools in developing countries too, but they refer to a small part only of the forest area (although their share of timber production is relatively high).

One main reason for failure is an inadequate assessment of the real costs and benefits of using biotechnology tools in given conditions (the level of improvement and the intensity of management), often under pressure from providers. As a result, expectations are not met and unjustified costs are high. This is a common risk in the early stages of development of new technologies. The same problem occurred during the development of clonal forestry a few decades ago. The development of protocols for the mass vegetative propagation of eucalyptus (rooting cuttings) was a real breakthrough in the 1970s, making it possible to take advantage of outstanding individuals from highly heterogenous interspecific hybrid progenies (the genetic gain could not be captured otherwise). The first large-scale plantations and gains in the Congo and Brazil were very impressive. But a perverse side-effect was that insufficiently informed programmes (or projects that were under pressure from active clonal forestry promoters) overestimated the benefit from vegetative propagation and neglected all the necessary but time consuming and demanding basic work (systematic species and provenance exploration and testing, individual selection and breeding, etc). This resulted in disappointments and misconceived strategies in some cases.

Much still needs to be done along the lines of upgrading the skills of researchers by ensuring that they receive higher education or appropriate higher level training to

be able to plan, develop and execute proper tree improvement programmes. Sufficient financial resources also need to be committed at the national level to ensure that such programmes are carried out successfully with the final aim of producing improved and bred reproductive material.

In developed countries, the applications of advanced forest biotechnologies have developed faster than predicted by Robinson (1999) a decade ago. A shift in technology transfer models has contributed to this success. In particular, this has come from the engagement and contributions of many Southern Hemisphere governments and universities. These institutions have contributed funding, talent and impetus to virtual forest research consortia in areas such as whole genome sequencing and other genomic applications. This contrasts with the older technology transfer models which characterized early tree improvement programmes. Unlike tree improvements, the consortia are less formally structured within a government. These grassroots scientific exchanges often include one or more government partners, and they are hastening forest biotechnology in interesting ways that bear little resemblance to traditional models of technology transfer. The advances being made in developed countries are also relevant to the progress in the application of forest biotechnologies in developing countries.

An analysis of forest biotechnology successes and/or failures leads to the following seven observations:

Forest biotechnology applications are developing along a separate path from crop biotechnology

Policy-makers tend to put forest biotechnology and crop biotechnology on the same plane, but the benefits, goals, risks and deliverables are distinctly different. This points to an important knowledge deficit about forestry biotechnology that needs to be addressed.

Forest biotechnology is now expanding to a wide range of forest types

The forest biotechnology portfolio *sensu lato* appears to be growing beyond its utility to forest plantations. Tropical forest complexity, health and recovery are also benefiting from forest biotechnologies in the form of genomics and its panoply of related methods. This has new relevance for tropical forests given the major current focus on slowing climate change.

That forest biotechnology applications have rapidly expanded in the past 5–10 years is also apparent from FAO (2004) which indicates that 64 percent of research and application activities in forest biotechnologies worldwide were focused on only six genera (*Pinus*, *Populus*, *Eucalyptus*, *Picea*, *Quercus* and *Acacia*). As discussed earlier, this is no longer the case. In this respect, genomics can be seen as a technology spillover, no longer restricted to planted forests but being used also for the management of naturally regenerated tropical forests.

Forest biotechnology has advanced over the past decade during favourable economic conditions. This is expected to change. The next decade may see slower progress because the forestry industry itself has some inherent problems which may be accentuated by the global financial crisis. Like other research and development (R&D) areas, this downturn could reduce forest biotechnology investment at a critical time and shape how developing countries choose to invest in forest biotechnology. These problems are unique to the forestry context, as summarized by Robinson (1999).

Plantation forestry has less flexibility in tailoring its raw materials delivery due to long lead times

This means that tailoring raw materials for markets that are years or even decades into the future is a high-risk proposition. Historically, this is a point which has not been well understood by biotechnology leaders in the agricultural biotechnology or pharmaceutical industries (Robinson, 1999).

This suggests that it might be timely to re-examine the role of biotechnology within the wood manufacturing processes rather than modifying the raw material supply years in advance of market demand. This emerging field of science is known as molecular wood biotechnology. Biotechnology benefits to date have included energy savings, waste reduction, remediation of toxic chemicals (see reviews by Breen and Singleton, 1999; Mansfield and Esteghlalian, 2003; Ahuja, Gisela and Moreira, 2004). Perhaps the best known example is the use of microbial (fungal) enzymes that degrade lignin, a component of the plant cell wall. The use of microbial enzymes is a time-honoured method that has been applied in pulping processing since 1975. Economic feasibility studies have shown that recent microbial biotechnology applications can raise mill productivity by 30 percent (Mansfield and Esteghlalian, 2003). The genetic improvement of fungi, bacteria and other microbes is a faster way of improving the efficacy of pulping processes and degraded mill waste than attempting to modify the raw materials of forest trees.

The private forestry sector is cautious about investing in forests and forest biotechnologies on lands which are not wholly owned

This holds true both for forest biotechnology and for intensive plantation management. As noted earlier, most forests are not privately owned. A related issue is that for-profit licensing for genetically enhanced forest trees tends to have been a tricky business model in the past due to long timelines, low investment rates and public ownership of forests as a worldwide norm (Robinson, 1999). Thus forestry and its research, including tree breeding, are now more vulnerable to funding reductions and loss of continuity than before the financial crisis.

In most developing countries, the industrial sector is dominated by foreign firms that do not often solicit or require research input from local research establishments, as they

rely on research conducted in their countries of origin. The situation, however, is different for the agriculture, forestry and horticultural sectors, where research is heavily supported by public research institutions. This scenario does not bode well for the development of biotechnology because commercial biotechnology has its roots in academia. Researchers in universities and research institutes carry out nearly all the basic research from which biotechnologies and biotechnology processes are developed. Thus, private venture capital companies which could help supply equity capital in support for development of local biotechnology are lacking. Besides, local firms are very unlikely to invest their already limited financial resources in long-gestation projects when interest rates are often higher if the funds are kept in banks.

However, there are interesting experiences of genetic improvement cooperatives that have pooled the resources of various private companies and universities to establish a single genetic improvement programme to benefit all participants. This model was applied successfully in Chile where the state forest service, the main forest companies and a university created a cooperative. Most of the advances in biotechnologies both in pines and eucalypts were made by this consortium.

Costs and consequences of expanding the range of forest biotechnologies

To the above stresses and strains, one must add the cost of expanding the forest biotechnology portfolio itself. Such an expansion can also generate financial strain. Burdon (1992) considered how molecular-based forest biotechnology would fit with classic breeding programmes. He foresaw severe institutional strain "...without skilful and sensitive management various competitive forces can subvert the safe and successful application [of molecular-based methods]". He was also concerned that this internal tension could imperil the collaboration within and between organizations which had already brought so much success to tree breeding. That "proprietary technology is very tempting..." added yet another source of strain on forest tree breeding. Managing forest biotechnology appropriately requires strategic oversight.

Roller-coaster R&D funding

Another point is that forestry, and its R&D budget, have always had a "roller-coaster ride" (Robinson, 1999). Managing against the vagaries of a roller-coaster budget makes the success of forest biotechnology to date even more impressive. This is true also for government sponsored research. Unstable raw material costs, fixed labour and overhead costs coupled with small, uneven profit margins all mean that the scale of forestry operations tend to become ever larger to clear net profits. The roller-coaster ride is an external force which threatens the stability and continuity required for long term research on long lived forest species.

Infrastructure and capacity constraints

In most developing countries, the use of biotechnology has been mostly limited to using tissue culture for the multiplication of selected clones. While some form of master plan for the development of biotechnology is present, there have been no real efforts to popularize this technology in the countries, mainly because of socio-economic factors. The introduction of biotechnology to developing countries has been by means of multilateral or bilateral collaboration. Experts from the collaborating (often developed) countries have visited and worked in the developing countries as short- or long-term experts and counterparts. They have helped establish laboratories and equip them with the relevant facilities to carry out research. While work continues during the period of collaboration, it slows down considerably once the collaboration phase is over. There are several reasons for this:

- The local counterparts are not adequately trained to continue the work independently once the collaboration ends.
- Once the collaboration period is over and the experts have returned home, the work in the laboratories slows down considerably as a result of financial constraints or the lack of technical knowledge.
- When equipment breaks down, it takes a long time to be repaired or purchased due to lack of funds.
- The purchase of chemicals needed for the work can be delayed as a result of a shortage of funds or the need to wait for them to be imported into the country.

In spite of these shortcomings, countries such as Vietnam have successfully developed elite clonal hybrids of *Acacia mangium* x *Acacia auriculiformis* for planting programmes. There are currently 127 000 hectares under clonal acacia hybrids cultivation (van Bueren, 2005) and the planted area continues to increase each year. With the recent rapid growth of the economy in the country, the situation looks poised to change as the government commits more funds towards education, training, research, skilled manpower development and infrastructure development. The biotechnology agenda is also being given priority in the national development plans of countries like Vietnam, Laos and Cambodia.

2.5 CASE STUDIES OF APPLICATIONS OF FOREST BIOTECHNOLOGIES IN DEVELOPING COUNTRIES

2.5.1 Eucalyptus plantations in Brazil

More than 700 species of eucalyptus are found in Australia but their performance as a plantation species is far greater elsewhere (Borrvalho, 2001) and especially in Brazil.

The Aracruz Cellulose's plantations in Brazil are perhaps the most widely cited success story. The company won the coveted Wallenberg Prize for its intensification efforts. This operation comprises a total of 300 000 ha of eucalypts of which half are produced by low cost vegetative propagation (Aracruz Cellulose, 2008).

In support of this intensive plantation effort, Brazil has a large-scale eucalyptus genomic research initiative that is known as the Genolyptus Project or the Brazilian Network of Eucalyptus Genome Research (Grattapaglia, 2004). The Genolyptus Project builds on the international whole genome sequencing effort. It aims for a genome wide understanding of the molecular basis for wood formation in *Eucalyptus* and is coupled with ongoing tree breeding programmes.

The project's talented team of scientists is generating a suite of biological and computational resources to discover, sequence, map, validate and understand the underlying variation of genes and genomic regions of economic importance in *Eucalyptus* with a focus on wood formation and disease resistance (Grattapaglia *et al.*, 2009). The project is based on a partnership among agencies within the Brazilian federal government through the MCT-Fundo Verde Amarelo. The academic/research sector is represented by seven universities and the Brazilian Agricultural Research Corporation (EMBRAPA). Industry is represented by twelve forestry companies. The Genolyptus Project could be considered a good example of how genomics can be successfully integrated with traditional breeding programmes to return value in a reasonably short time. This project also represents a good model of how universities, government agencies and private enterprises can work together to benefit different categories of stakeholders.

2.5.2 Clonal propagation of teak in Malaysia

Teak, *Tectona grandis*, is widely planted in many countries in Asia, South and Central America and Africa. A fifteen-year collaboration between the Sabah Foundation Group (Malaysia) and the Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD, France), exploiting molecular markers and micropropagation, has led to the availability of superior quality planting material both for the local market as well as for export (Goh *et al.* 2007; Muralidharan, 2009a). The candidate trees for producing the superior clones were selected with reference to intrinsic wood qualities (e.g. natural durability, shrinkage, sapwood percentage, etc.). Simple sequence repeat markers were developed to determine the genetic background and diversity in order to reduce inbreeding and ensure the genetic fidelity of the clones mass produced by tissue culture. There is now widespread demand for these clones globally. In addition, a clone identification form provides detailed information on each clone, including the DNA fingerprinting profile.

2.5.3 Micropropagation applied to tree breeding of fast-growing forest tree species in Latin America

Biotechnology was introduced in Latin America in the 1980s. Networking has been very important for the research community there. By December 2008, there were 5 467 researchers in 738 agricultural biotechnology laboratories in 32 countries in the Technical Co-operation Network on Agricultural Biotechnology in Latin America and the Caribbean (REDBIO), based at the FAO Regional Office in Chile. The network has been in operation since 1991 to develop biotechnology for the sustainable use of regional genetic resources, promote the safe and responsible application of the technologies – especially in fragile environments, and enhance the regional development of new strategic technologies such as molecular genomics. It also encourages the application, whenever feasible, of advanced biotechnology tools in integrated crop management and sustainable production systems.

In terms of planted forests, the largest areas are in Brazil (7 million ha, 4.1 million of which are industrial man-made forests). Chile has 2.25 million ha of planted forest areas, practically all for industrial purposes; Uruguay has about 0.75 million ha; Argentina has 0.7 million ha; Venezuela, 0.5 million ha; Cuba, 0.4 million ha; Peru, 0.3 million ha; and Colombia and Mexico have about 0.2 million ha each. In the other countries of Latin America and the Caribbean the reforested area is less than 100 000 ha per country. The estimates of the current yearly forestation rate vary from 386 000 to 520 000 ha (FAO, 2006). Practically all the planted forests have been established on abandoned agricultural lands where erosion is prevalent, with the overwhelming majority being established with fast-growing exotic species in the *Eucalyptus* and *Pinus* genera. Many of these planted forests have been established by clonally propagated elite plants in the case of *Eucalyptus* or through somatic embryogenesis in the case of the *Pinus* species.

2.5.4 Bioprotection in Kerala, India

At the Forest Protection Division of the Kerala Forest Research Institute, India, investigations into control of a serious insect pest of teak viz. the teak defoliator (*Hyblaea parea*), have been carried out for several years. A *Hyblaea parea* nuclear polyhedrosis virus (HpNPV) isolated from natural populations of the insect larvae resulted eventually in a very effective biological control method. A permanent preservation plot where the pest outbreak was kept under control over several years with regular spraying of the HpNPV formulation clearly demonstrated the benefits in terms of increased volume of timber compared with the control plots. Research then went into the rearing of the insect larvae in the laboratory on an artificial diet and the mass multiplication of the virus, followed by the formulation of the pesticide incorporating UV protectants and other adjuvants, and finally the spraying technique in the planted forests. Almost two decades of research finally culminated in a

successful solution to a serious problem. Nevertheless, the technology has remained in the laboratory and there is no indication that it will make it to the standard package of practices of the teak plantings immediately. Since most teak in India today comes under the control of State Forest Departments, acceptability by the forestry professionals is important. Some farmers had shown interest and willingness to use the product in their plantations, and the initial response showed that the technology was effective. The case study demonstrates that research in forest biotechnology has a much better chance of producing results when conceived, developed and implemented in a broader framework that involves not only scientists and technologists but also at every stage the forestry professionals who work at the field level and, at some level, policy-makers who eventually have to give their approval (Muralidharan, 2009b).

B. LOOKING FORWARD: PREPARING FOR THE FUTURE


2.6 KEY ISSUES WHERE FOREST BIOTECHNOLOGIES COULD BE USEFUL

2.6.1 Adaptation to climate change

Forests, particularly tropical forests, play a central role in climate change and this is expected to shape the direction of forest biotechnology research in new ways. At the heart of the matter is how to ease forest adaptation. Forest adaptation is the foundation for all other forestry policy solutions aimed at slowing climate change (Hamrick, 2004; Millar, Stephenson and Stephens, 2007; Aitken *et al.*, 2008). In addition, all forestry policy solutions, i.e. reducing emissions from deforestation and degradation (REDD), forestry offsets, biofuels and biomass depend on the health and resilience of forests while adapting to climate change (Clark, 2004). Thus, forest adaptation deserves a closer look as predictive models for climate change effects become more regional. Already, predicted range shifts and assemblage mixing are being published to some degree but no two regions will experience climate change in the same way. Tropical forests are especially vulnerable to climate change (see Figure 4). Climate change related problem-solving will dominate forest R&D, particularly for tropical regions.

FIGURE 4

A SCHEMATIC DIAGRAM OF A FORESTED ECOSYSTEM WHICH IS ADJUSTING TO CLIMATE CHANGE



Thinking about forest tree adaptation under climate change has led to formulation of the concept of assisted migration (Aitken *et al.*, 2008; Marris, 2009). Assisted migration refers to the practice of matching seed source to location, assuming a different climate in the future. This is an important concept given that forest trees are so narrowly adapted.

As climate change models are becoming more regionally predictive, range shifts for indigenous forest trees are expected. Seed sources and provenances can be matched for optimum growth under these future site conditions (Aitken *et al.*, 2008). The issue is more critical at higher latitudes where forest rotation ages span 50–100 years, which is well within the realm of expected climate change by 2050.

With regard to biotic and abiotic stresses expected under climate change scenarios, the development of biotechnology tools for resistance to pests, tolerance to climatic extremes, bioremediation and carbon sequestration will be more relevant in the near future both for naturally regenerating and for planted forest tree species.

2.6.2 Sustainable management of forest genetic resources

Genetic diversity provides the fundamental basis for the evolution of forest tree species that has enabled forests and trees to adapt to changing conditions for thousands of years. Adaptation has resulted in a unique and irreplaceable portfolio of forest tree genetic resources. Fires, deforestation, new pests and diseases, and other factors are increasingly threatening forest genetic resources. The vast majority of forest genetic resources remain unknown and underutilized although the sustainable use of forest genetic diversity has great potential to contribute towards addressing new challenges and maintaining economic, social and cultural values, as well as providing environmental services and benefits. The field of forest genetic resources is undergoing significant changes. Traditionally, the sector has been concerned with technical issues of genetic conservation, tree improvement and seed supply for wood production. The scope of genetic management, however, is now expanding as the demand for products from forest species is increasing and diversifying (timber, fibre, fruits, resins and other non-wood products), which is contributing to food security and poverty reduction of rural populations. The emerging uses of forest genetic resources must be assessed to achieve sustainable use of these resources. Advances in biotechnology are rapidly enabling the improved use of genetic resources, and potentially greater economic and social contributions resulting from forest genetic resources. Biotechnology developments will also provide improved tools to enhance the effectiveness of conservation and development measures (knowledge about life-history traits and genetic diversity is lacking or inadequate for most tree species to define and implement conservation strategies).

2.7 IDENTIFYING OPTIONS FOR DEVELOPING COUNTRIES

Based on the stocktaking exercise carried out here, a number of specific options can be identified for developing countries to help them make informed decisions regarding the adoption of biotechnologies in the forestry sector in the future.

Biotechnologies should be integrated with conventional technologies

Forest biotechnology as a whole lacks strategic oversight to ensure the integration of its parts (Burdon, 1992). No policy exists that would ensure that tree breeding and molecular-based components of forest biotechnology work together in a complementary fashion. This is a problem because biotechnologies such as molecular markers and mass propagation methods can be useful only if stable conventional forest breeding programmes are in place.

Promote public-private partnership at national level

Effective public-private partnerships are a key factor in most successful cases of the development and implementation of forest biotechnologies, especially as regards industrial wood production and processing. It is therefore an important strategy to be considered by developing countries.

Improve information and communication strategies for biotechnologies

Public access to good and updated information on forest biotechnologies is very important in developing countries. Benefits from their use can be optimized if the end users know how to utilize them properly. Consolidated information and education mechanisms should be put in place to allow communication between the relevant sectors of society. Attention should be given to issues relating to the meaningful adoption of biotechnologies including socio-economic implications, efficiency, costs and benefits and environmental impacts.

2.8 IDENTIFYING PRIORITIES FOR ACTION FOR THE INTERNATIONAL COMMUNITY

The international community, including FAO and other UN organizations as well as NGOs, donors and development agencies, can play a key role in supporting developing countries by providing a framework for international cooperation as well as funding support for the generation, adaptation and adoption of appropriate biotechnologies. A set of Priorities for Action is given below that will help the international community fulfil this role.

Improve access to peer-reviewed scientific information about forest biotechnologies in developing countries

Even with Internet access, peer-reviewed journal and books continue to be central sources of information for scientists. Subscription costs for the best available scientific knowledge have risen exponentially, putting it out of the reach of many institutions even in the most science-literate developed countries. While open-source journals are a step in the right direction, publications in traditional forestry outlets such as proceedings, conferences and government printing office publications are declining. Today, forest biotechnology is adversely affected by barriers to knowledge acquisition. These barriers, when coupled with publication bias (defined as the propensity to forego publishing negative results), can only slow scientific progress.

The international community is already acting to reduce these barriers. For example, FAO is coordinating the Access to the Global Online Research in Agriculture (AGORA)² programme, providing free or low-cost access to scientific journals in the fields of food, agriculture (including forestry), environmental science and related social sciences. A sister programme called Online Access to Research in the Environment (OARE)³ also covers both forestry and biotechnology and its goal is to improve access to scientific research in developing countries by providing high quality, timely, relevant, environmental and related science journals and other scientific content for free or at nominal cost. OARE is an international public-private consortium coordinated by the United Nations Environment Programme (UNEP), Yale University and leading science and technology publishers. Such initiatives from the international community should be encouraged and strengthened.

Build capacity for understanding forest biotechnology issues at all levels

For most policy-makers, scientists and even students, forest biotechnology is a form of agriculture. As discussed earlier, however, this is not the case. At best, it is a tribute to those who have developed capacity for agricultural biotechnology, but applying agricultural biotechnology to trees will not optimize the benefits to be obtained from forestry resources. Agricultural biotechnology does not constitute the best form of knowledge for forests. Forest biotechnology is an area that is separate from crop and livestock biotechnology and requires its own capacity building. Capacity building initiatives in forest biotechnologies from the international community should be strengthened in view of this important observation. The capacity building initiatives should include training in emerging tools such as bioinformatics and computational biology for tropical forest studies. Intensive educational efforts for bioinformatics courses would benefit professionals and scientists in developing

² www.aginternetwork.org.

³ www.oaresciences.org.

countries. This skill set provides capacity for testing hypotheses using available information from DNA sequences and related databases. In addition to educational workshops, this action will also require upgrades of computing infrastructure and perhaps bandwidth in some cases. A wealth of data is being produced by whole genome DNA sequencing consortia, the Tree of Life project, the Consortium for the Barcode of Life and a host of other independent initiatives.

Review the status and potential of forest biotechnologies for developing countries

It is clear from this Chapter that the forestry sector in developing countries is in a very dynamic situation and facing a number of important challenges and opportunities for which biotechnologies can play a significant role. FAO commissioned a series of studies in 2002–2004 to investigate the extent and pattern of research and application of forest biotechnologies worldwide (FAO, 2004). These studies have informed and influenced policy-making in developing countries, providing good indicators and possible predictions of trends in forest biotechnologies around the world. Such global surveys are important, and the international community should continue to provide periodic reviews of the status and potential of forestry biotechnologies in developing countries. The reviews should cover synergies with other biotechnology sectors such as applications of biotechnology to micro-organisms to improve wood manufacturing processing, as well as with other fields of technology that may be useful such as nanotechnology, information technology and synthetic genomics that may converge to the benefit of wood products manufacturing. Another potential area for convergence is the combination of genomics tools with geographic information systems (GIS). Using GIS to track rare alleles, gene flow or expressed proteins is another area that deserves a closer look. Technology advances are delivering finer resolution at both the landscape and molecular ends of this molecule-to-landscape spectrum, and this will no doubt provide interesting ways to study all forest ecosystems.

Encourage North-South collaboration

As mentioned earlier in this Chapter, the application of forest biotechnologies has advanced faster in developed countries than originally predicted (Robinson, 1999). As much of the research refers to processes and/or tree species that are relevant to developing countries, these advances are of major potential relevance to developing countries as well. The international community should act to ensure that the results of research and application in forest biotechnologies in developed countries are made accessible to developing countries.

2.9 REFERENCES

- Agrawal, A., Chatter, A. & Hardin, R. 2008. Changing governance of the world's forests. *Science*, 320: 1460–1462.
- Ahuja, S.K., Gisela, M. & Moreira, A.R. 2004. Utilization of enzymes for environmental applications. *Crit. Rev. Biotechnol.*, 24: 125–154.
- Aitken, S.N., Yeaman S., Holliday, J.A., Wang, T. & Curtis-McLane, S. 2008. Adaptation, migration or extirpation: Climate change outcomes for tree populations. *Evol. Appl.*, 1: 95–111.
- Allard, R.W. 1960. *Principles of plant breeding*. New York, John Wiley & Sons.
- Aracruz Celulose. 2008. *Annual Report 2007*.
- Attree, S.M. & Fowke, L.C. 1991. Micropropagation through somatic embryogenesis in conifers. In Y.P.S. Bajaj, ed. *Biotechnology in agriculture and forestry 17. High-tech and micropropagation I*, pp. 53–70. Berlin, Heidelberg, New York, Springer-Verlag.
- Balocchi, C.E. 1997. Realised versus operational gain: The role of propagation strategies. In R.D. Burdon & J.M. Moore, eds., *Proceedings IUFRO'97 – Genetics of radiata pine*. FRI Bulletin 203, pp. 253–255.
- Bishop-Hurley, S.L., Zabkiewicz, R.J., Grace, L., Gardner, R.C., Wagner, A. & Walter, C. 2001. Conifer genetic engineering: Transgenic *Pinus radiata* (D. Don) and *Picea abies* (Karst) plants are resistant to the herbicide Buster. *Plant Cell Rep.*, 20: 235–243.
- Boerjan, W. 2005. Biotechnology and the domestication of forest trees. *Curr. Opin. Biotechnol.*, 16: 159–166.
- Bonfante, P. & Anca, I.-A. 2009. Plants, mycorrhizal fungi and bacteria: A network of interactions. *Ann. Rev. Microbiol.*, 63: 363–383.
- Born, C., Hardy, O.J., Chevallier, M.H., Ossari, S., Attéké, C. Wickings, E.J. & Hossaert-McKey, M. 2008. Small-scale spatial genetic structure in the Central African rainforest tree species *Aucoumea klaineana*: A stepwise approach to infer the impact of limited gene dispersal, population history and habitat fragmentation. *Mol. Ecol.*, 17: 2041–2050.
- Born, C., Vignes, H., Muloko, N., Wickings, E.J., Hossaert-McKey, M. & Chevallier, M.H. 2006. Isolation and characterization of polymorphic microsatellite loci from *Aucoumea klaineana* Pierre (Burseraceae), a tropical rainforest tree of Central Africa. *Mol. Ecol. Notes*, 6: 1054–1056.
- Borrhallo, N.M.G. 2001. The purpose of breeding is breeding for a purpose. *IUFRO Symposium: Developing the Eucalypt for the future*. Valdivia, Chile, 10–15 September 2001.
- Boshier, D. H., Chase, M.R. & Bawa, K.S. 1995. Population genetics of *Cordia alliodora* (Boraginaceae), a neotropical tree. 2. Mating system. *Am. J. Bot.*, 82: 476–483.
- Breen, A. & Singleton, F.L. 1999. Fungi in lignocellulose breakdown and biopulping. *Curr. Opin. Biotechnol.*, 10: 252–258.
- Brondani, R.P.V., Brondani, C., Tarchini, R. & Grattapaglia, D. 1998. Development and mapping of microsatellite based markers in *Eucalyptus*. *Theor. App. Genet.*, 97: 816–829.
- Brown, A.H.D. & Moran, G.F. 1981. Isozymes and the genetic resources of forest trees. In M.T. Conkle ed. *Proceedings symposium on isozymes of forest trees and forest insects*. USDA Forest Service Gen. Tech. Rep., pp. 1–10. PSW-48.
- Brown, G.R., Gill, G.P., Kuntz, R.J., Langley, C.H. & Neale, D.B. 2004. Nucleotide diversity and linkage disequilibrium in loblolly pine. *Proc. Natl. Acad. Sci. USA.*, 101: 15255–15260.

- Burdon, R.D. 1992. Tree breeding and the new biotechnologies: In damaging conflict or constructive synergism? *Paper to IUFRO S2.02-08 conference: Breeding tropical trees*, pp. 1–7. Cali, Colombia.
- Burleigh, J.G. & Matthews, L. 2004. Phylogenetic signal in nucleotide data from seed plants: implications for resolving the seed plant tree of life. *Am. J. Bot.*, 91: 1599–1613.
- Caesar, J. 2009. Experiences regarding agricultural biotechnology in Guyana. *Message 121 of the FAO e-mail conference on learning from the past: Successes and failures with agricultural biotechnologies in developing countries over the last 20 years*. (available at www.fao.org/biotech/logs/c16/090709.htm).
- Campbell, M., Brunner, A., Jones, H. & Strauss, S. 2003. Forestry's fertile crescent: The application of biotechnology to forest trees. *Plant Biotechnol. J.*, 1: 141–154.
- Cavers, S., Navarro, C. & Lowe, A.J. 2003. Chloroplast DNA phylogeography reveals colonization history of a Neotropical tree, *Cedrela odorata* L., in Mesoamerica. *Mol. Ecol.*, 12: 1451–1460.
- CBOL Plant Working Group. 2009. A DNA barcode for land plants. *Proc. Natl. Acad. Sci., USA*, 106: 12794–12797 (also available at www.barcoding.si.edu/plant_working_group.html).
- Cervera, M.T., Remington, D., Frigerio, J.M., Storme, V., Ivens, B., Boerjan, W. & Plomion, C. 2000. Improved AFLP analysis of tree species. *Can. J. For. Res.*, 30: 1608–1616.
- Chazdon, R.L. 2008. Beyond deforestation: Restoring forests and ecosystem services on degraded lands. *Science*, 320: 1458–1460.
- Clark, D.A. 2004. Tropical forests and global warming: Slowing it down or speeding it up? *Front. Ecol. Environ.*, 2: 73–80.
- Connell, J. H. & Lowman, M.D. 1989. Low-diversity tropical rain forests: Some possible mechanisms for their existence. *Am. Nat.*, 134: 88–119.
- Cruz, A. 1981. Bird activity and seed dispersal of a montane forest tree (*Dunalia arborescens*) in Jamaica. *Reprod. Bot.*, 34–44.
- Darvasi, A. & Shifman, S. 2005. The beauty of admixture. *Nat. Genet.*, 37: 118–119.
- Dean, J. 2006. Genomics resources for conifers. In C.G. Williams, ed. *Landscape, genomics and transgenic conifers*, pp. 55–74. New York, Springer.
- Devey, M., Carson, S.D., Nolan, M., Matheson, C., Te Riini, C. & Hohepa, J. 2003. QTL associations for density and diameter in *Pinus radiata* and the potential for marker-aided selection. *Theor. Appl. Genet.*, 108: 516–524.v
- Doudrick, R.L., Heslop-Harrison, J.S, Nelson, C.D., Schmidt, T., Nance, W.L. & Schwarzacher, T. 1995. Karyotype of slash pine (*Pinus elliottii* var. *elliottii*) using patterns of fluorescence *in situ* hybridization and fluorochrome banding. *J. Hered.*, 86: 289–296.
- Duponnois, R., Plenchette, C., Prin, Y., Ducouso, M., Kisa, M., Moustapha Bâ, A. & Galiana, A. 2007. Use of mycorrhizal inoculation to improve reforestation process with Australian Acacia in Sahelian ecozones. *Ecol. Eng.*, 29: 105–112.
- Duponnois, R., Kisa M., Prin Y., Ducouso, M., Plenchette, C., Lepage, M. & Galiana, A. 2008. Soil factors influencing the growth response of *Acacia holosericea* A. Cunn. ex G. Don to ectomycorrhizal inoculation. *New Forests*, 35: 105–117.
- Eisen, J. A. & Fraser, C.M. 2003. Phylogenomics: Intersection of evolution and genomics. *Science*, 300: 1706–1707.
- El-Kassaby, Y.A. & Lstiburek, M. 2009. Breeding without breeding. *Genet. Res. Camb.*, 91: 111–120.
- El-Lakany, M.H. 2004. Are genetically modified trees a threat to forests? *Unasylva*, 217: 45–47. (available at www.fao.org/docrep/007/y5507e/y5507e14.htm).
- Eriksson, G., Namkoong, G. & Roberds, J.H. 1995. Dynamic gene conservation for uncertain futures. *For. Ecol. Manage.*, 62: 15–37.

- FAO. 1994. *Biotechnology in forest tree improvement with special reference to developing countries*, by R. Haines. FAO Forestry Paper 118. Rome.
- FAO. 2001. *Forest genomics for conserving adaptive genetic diversity*, by K.V. Krutovskii & D.B. Neale. Forest Genetics Resources Working Paper FGR/3. Rome. (available at www.fao.org/DOCREP/003/X6884E/X6884E00.HTM).
- FAO. 2004. *Preliminary review of biotechnology in forestry, including genetic modification*. Forest Genetic Resources Working Paper FGR/59E. Rome. (available at www.fao.org/docrep/008/ae574e/ae574e00.htm).
- FAO. 2006. *Global forest resources assessment 2005 – progress towards sustainable forest management*. FAO Forestry Paper No. 147. Rome. (available at www.fao.org/docrep/008/a0400e/a0400e00.htm).
- FAO. 2007. *Marker-assisted selection: Current status and future perspectives in crops, livestock, forestry and fish*, by E.P. Guimaraes, J. Ruane, A. Sonnino, B.D. Scherf, & J.D. Dargie, eds. Rome. (also available at www.fao.org/docrep/010/a1120e/a1120e00.htm).
- FAO. 2008a. *Coping with water scarcity: What role for biotechnologies?*, by J. Ruane, A. Sonnino, P. Steduto & C. Deane. FAO Land and Water Discussion Paper 7. Rome. (also available at www.fao.org/docrep/011/i0487e/i0487e00.htm).
- FAO. 2008b. *Synthesis of country progress reports: Activities related to poplar and willow cultivation and utilization, 2004 through 2007*. FAO International Poplar Commission 23rd Session. Working Paper IPC/6E. (available at www.fao.org/docrep/011/k3380e/k3380e00.htm)
- Finkeldey, R. & Ziehe, M. 2004. Genetic implications of silvicultural regimes. *For. Ecol. Manage.*, 197: 231–244.
- Galiana, A., Goh, D., Chevallier, M.-H., Gidiman, J., Moo, H., Hattah, M. & Japarudin, Y. 2003. Micropropagation of *Acacia mangium* x *A. auriculiformis* hybrids in Sabah. *Bois et Forêts des Tropiques*, 275: 77–82.
- Garcia, F., Noyer, J.-L., Risterucci, A.-M. & Chevallier, M.H. 2004. Genotyping of mature trees of *Entandrophragma cylindricum* with microsatellites. *J. Hered.*, 95(5): 454–457.
- Gianessi, L., Silvers, C., Sankula, S. & Carpenter, J. 2002. *Plant biotechnology: Current and potential impact for improving pest management in U.S. agriculture: An analysis of 40 case studies* (executive summary). Washington, DC, National Center for Food and Agricultural Policy.
- Godbole, S., Anil Sood, A., Sharma, M., Nagar, P.K. & Ahuja, P.S. 2004. Starch deposition and amylase accumulation during somatic embryogenesis in bamboo (*Dendrocalamus hamiltonii*). *J. Plant Physiol.*, 161: 245–248.
- Goh, D.K.S. & Monteuis, O. 2005. Rationale for developing intensive teak clonal plantations, with special reference to Sabah. *Bois et Forêts des Tropiques*, 28: 5–15.
- Goh, D.K.S., Chaix, G., Baillères, H. & Monteuis, O. 2007. Mass production and quality control of teak clones for tropical plantations: The Yayasan Sabah Group and Forestry Department of CIRAD joint project as a case study. *Bois et Forêts des Tropiques*, 33: 6–9.
- Grace, L.J., Charity, J.A., Gresham, B., Kay, N. & Walter, C. 2005. Insect-resistant transgenic *Pinus radiata*. *Plant Cell Rep.*, 24(2): 103–111
- Grattapaglia, D. 2004. Genomics applied to Eucalyptus: The Genolyptus project. In R. Kellison, S. McCord & K.M.A. Gartland, eds. *Forest biotechnology in Latin America. Proceedings workshop biotecnologia forestal*. Global Biotechnology Forum 2–4 March, 2004, Concepcion, Chile.
- Grattapaglia, D. & Kirst, M. 2008. Eucalyptus applied genomics: From gene sequences to breeding tools. *New Phytol.*, 179: 911–929.
- Grattapaglia, D., Plomion, C., Kirst, M. & Sederoff, R. 2009. Genomics of growth traits in forest trees. *Curr. Opin. Plant Biol.*, 12: 148–156.

- Gwaze, D.P., Zhou, Y., Reyes-Valdés, M.H., Al-Rababah M.A. & Williams, C.G. 2003. Haplotypic QTL mapping in an outbred pedigree. *Genet. Res.*, 81: 43–50.
- Hamrick, J.L. 2004. Response of forest trees to global environmental changes. *For. Ecol. Manage.*, 197: 323–335.
- Hamrick, J.L. & Murawski, D.A. 1990. The breeding structure of tropical tree populations. *Plant Species Biol.*, 5: 157–165.
- Hu, J.J., Tian, Y.C., Han, Y.F., Li, L. & Zhang, B.E. 2001. Field evaluation of insect-resistant transgenic *Populus nigra* trees. *Euphytica*, 121: 123–127.
- Hufford, K.M. & Hamrick, J.L. 2003. Viability selection at three early-life stages of the tropical tree, *Platypodium elegans* (Fabaceae, Papilionoideae). *Evolution*, 57: 518–526.
- Hughes, C.E., Eastwood, R.J. & Bailey, C.D. 2006. From famine to feast? Selecting nuclear DNA sequence loci for plant species-level phylogeny reconstruction. *Philos. Trans. R. Soc. Lond. B Biol. Sci.*, 361: 211–225.
- James, C. 2008. *Global status of commercialized biotech/GM crops: 2008*. Brief No 39. New York, ISAAA.
- Jones, N. 2002. Somatic embryogenesis as a tool to capture genetic gain from tree breeding strategies: Risks and benefits. *S. Afr. For. J.*, 195: 93–101.
- Kirst, M., Myburg, A.A., Basten, C.J., Zeng, Z-B. & Sederoff, R.R. 2004. Transcript mapping by quantitative (QTL) analysis of microarrays in Eucalyptus. *Proceedings 12th plant & animal genome conference*, San Diego, CA. January 10–14.
- Kisa, M., Sanson A., Thioulouse J., Assigbetse K., Sylla S., Spichiger R., Dieng L., Berthelin J., Prin Y., Galiana A. & Lepage, M. 2007. Arbuscular mycorrhizal symbiosis can counterbalance the negative influence of the exotic tree species *Eucalyptus camaldulensis* on the structure and functioning of soil microbial communities in a sahelian soil. *FEMS Microbiol. Ecol.*, 62: 32–44.
- Krishnadas, K.R. & Muralidharan, E.M. 2008. Repetitive somatic embryogenesis from zygotic embryos of teak (*Tectona grandis* L.) cultured *in vitro*. In S.A. Ansari, C. Narayanan & A.K. Mandal, eds. *Forest biotechnology in India*, pp.149–155. Delhi, Satish Serial Publishing House.
- Kuhn, G., Hijri, M. & Sanders, I.R. 2001. Evidence for the evolution of multiple genomes in arbuscular mycorrhizal fungi. *Nature*, 414: 745–748
- Lacerda, A.E.B., Kanashiro, M. & Sebbenn, A.M. 2008. Effects of reduced impact logging on genetic diversity and special genetic structure of a *Hymenea courbaril* population in the Brazilian Amazon. *For. Ecol. Manage.*, 255: 1034–1043
- Le Roux, J.J. & van Staden, J. 1991. Micropropagation and tissue culture of Eucalyptus. A review. *Tree Physiol.*, 9: 435–477.
- Ledig, F.T., Conkle, M.T., Bermejo, B., Eguiluz, T., Hodgskiss, P., Johnson, D.R. & Dvorak, W.S. 1999. Evidence for an extreme bottleneck in a rare Mexican pinyon: genetic diversity, disequilibrium, and the mating system in *Pinus maximartinezii*. *Evolution*, 53(1): 91–99.
- Lee, C.T., Lee, S.L., Ng, K.K.S., Siti Salwana, H., Norwati, M. & Saw, L.G. 2003. *Effective population size of Koopassia malaccensis for conservation based on isozyme analysis*. In M.K. Thong, M.Y. Fong, M.E. Phipps, U.R. Kuppusamy, M. Ameen, M. Zulqarnain, K.A.R. Suzainur & M.N. Suzita, eds. *Proceedings 5th national congress of genetics – from peas to CHIPS: The globalization of genetics*, pp. 159–161. Genetic Society of Malaysia, Kuala Lumpur, Malaysia.
- Lee, S.L., Wickneswari, R., Mahani, M.C. & Zakri, A.H. 2000. Genetic diversity of *Shorea leprosula* Miq. (Dipterocarpaceae) in Malaysia: Implications for conservation of genetic resources and tree improvement. *Biotropica*, 32: 213–224.
- Lee, S.L., Ng, K.K.S., Saw, L.G., Norwati, A., Siti Salwana, M.H., Lee, C.T. & Norwati, M. 2002. Population genetics of *Intsia palembanica* (Leguminosae) and genetic conservation of virgin jungle reserves (VJRs) in Peninsular Malaysia. *Am. J. Bot.*, 89: 447–459.

- Lee, S.L. & Krishnapillay, B. 2004. *Status of forest genetic conservation and management in Malaysia*. In T. Luoma-aho, L.T. Hong, V. Ramanatha Rao & H.C. Sim, eds. *Proceedings Asia Pacific forest genetic resources programme (APFORGEN) inception workshop: Forest genetic resources conservation and management*, pp. 206–228. Serdang, Malaysia, IPGRI-APO.
- Lee, S.L., Tani, N., Ng, K.K.S. & Tsumura, Y. 2004a. Characterization of 15 polymorphic microsatellite loci in an endangered tropical tree *Hopea biltonensis* (Dipterocarpaceae) in Peninsular Malaysia. *Mol. Ecol. Notes*, 4: 147–149.
- Lee, S.L., Tani, N., Ng, K.K.S. & Tsumura, Y. 2004b. Isolation and characterization of 21 microsatellite loci in an important tropical tree *Shorea leprosula* and their applicability to *S. parvifolia*. *Mol. Ecol. Notes*, 4: 222–225.
- Lee, S.L., Ng, K.K.S., Saw, L.G., Lee, C.T., Norwati, M., Tani, N., Tsumura, Y. & Koskela, J. 2006. Linking the gaps between conservation research and conservation management of rare dipterocarps: A case study of *Shorea lumutensis*. *Biol. Conserv.*, 131: 72–92.
- Lelu-Walter, M.A. & Harvengt, L. 2004. L'embryogenèse somatique des conifères, état et perspectives. *Afocel Inf.-For.*, 694: 1–6. (available at www.fcbainfo.fr/pages/Archives/fif694.pdf).
- Li, L., Zhou, Y., Cheng, X., Sun, J., Marita, J.M., Ralph, J. & Chiang, V.L. 2003. Combinatorial modification of multiple lignin traits in trees through multigene co-transformation. *Proc. Natl. Acad. Sci. USA.*, 100: 4939–4944.
- Lim, L.S., Wickneswari, R., Lee, S.L. & Latiff, A. 2002. Genetic variation of *Dryobalanops aromatica* Gaertn. F. (Dipterocarpaceae) in Peninsular Malaysia using microsatellite DNA markers. *For. Genet.*, 9: 125–136.
- Mansfield, S.D. & Esteghlalian, A.R. 2003. Applications of biotechnology in the forest products industry. In S.D. Mansfield & J.M. Sadler, eds., *ACS Applications of enzymes to lignocellulosics*, pp. 2–29. Washington, DC, ACS Publications.
- Marris, E. 2009. Planting the forest of the future. *Nature*, 459: 906–908.
- Millar, C.M., Stephenson, N.L. & Stephens, S.L. 2007. Climate change and forests of the future: Managing in the face of uncertainty. *Ecol. Appl.*, 17: 2145–2151.
- Minghe, L. & Ritchie, G.A. 1999. Eight hundred years of clonal forestry in China: I. traditional afforestation with Chinese fir (*Cunninghamia lanceolata* (Lamb.) Hook.). *New Forest.*, 18: 131–142.
- Muralidharan, E.M. 2009a. Forestry – biotechnology – India. *Message 89 of the FAO e-mail conference on learning from the past: successes and failures with agricultural biotechnologies in developing countries over the last 20 years*. Rome. (available at www.fao.org/biotech/logs/c16/060709.htm).
- Muralidharan, E.M. 2009b. Biological control of forest pests – India. *Message 114 of the FAO e-mail conference on learning from the past: successes and failures with agricultural biotechnologies in developing countries over the last 20 years*. Rome (available at www.fao.org/biotech/logs/c16/080709.htm).
- Murawski, D.A., Dayanandan, B. & Bawa, K.S. 1994. Outcrossing rates of two endemic *Shorea* species from Sri Lankan tropical rain forests. *Biotropica*, 26: 23–29.
- Naito, Y., Konuma, A., Iwata, H., Suyama, Y., Seiwa, K., Okuda, T., Lee, S.L., Norwati, M. & Tsumura, Y. 2005. Selfing and inbreeding depression in seeds and seedlings of *Neobalanocarpus heimii* (Dipterocarpaceae). *J. Plant Res.*, 118: 423–430.
- Namkoong, G., Barnes, R.D. & Burley, J. 1980. *A philosophy of breeding strategy for tropical forest trees*. Oxford, University of Oxford, Department of Forestry, Commonwealth Forestry Institute.
- Nason, J. D. & Hamrick, J.L. 1997. Reproductive and genetic consequences of forest fragmentation: two case studies of neotropical canopy trees. *J. Hered.*, 88: 264–276.
- Nelson, C.D. & Johnsen, K.H. 2008. Genomic and physiological approaches to advancing forest tree improvement. *Tree Physiol.*, 28: 1135–1143.

- Ng, K.K.S., Lee, S.L. & Koh, C.L. 2004. Spatial structure and genetic diversity of two tropical tree species with contrasting breeding systems and different ploidy levels. *Mol. Ecol.*, 13: 657–669.
- Ng, K.K.S., Lee, S.L., Saw, L.G., Plotkin, J.B. & Koh, C.L. 2006. Spatial structure and genetic diversity of three tropical tree species with different habitat preferences within a natural forest. *Tree Genet. Genomes*, 2: 121–131.
- Ouahmane, L., Hafidi, M., Thioulouse, J., Ducousso, M., Kisa, M., Prin, Y., Galiana, A., Boumezzough, A. & Duponnois R. 2007. Improvement of *Cupressus atlantica* Gaussen growth by inoculation with native arbuscular mycorrhizal fungi. *J. Appl. Microbiol.*, 103: 683–690.
- Park, Y.S. 2002. Implementation of conifer somatic embryogenesis in clonal forestry: Technical requirements and deployment considerations. *Ann. For. Sci.*, 59: 651–656.
- Park, Y.S., Barret, J.D. & Bonga, J.M. 1998. Application of somatic embryogenesis in high-value clonal forestry: Deployment, genetic control, and stability of cryopreserved clones. *In Vitro Cell. Dev. Biol. Plant*, 34: 231–239.
- Pautasso, M. 2009. Geographical genetics and the conservation of forest trees. *Perspect. Plant Ecol. Evol. Syst.*, 11: 157–189.
- Pinto, G., Santos, C., Neves, L. & Araújo, C. 2002. Somatic embryogenesis and plant regeneration in *Eucalyptus globulus* Labill. *Plant Cell Rep.*, 21: 208–213.
- Plomion, C., Pionneau, C., Brach, J., Costa, P. & Baillères, H. 2000. Compression wood-responsive proteins in developing xylem of maritime pine (*Pinus pinaster* Ait.). *Plant Physiol.*, 123: 959–969.
- Plomion, C., Pionneau, C. & Baillères, H. 2003. Identification of tension-wood responsive proteins in the developing xylem of *Eucalyptus*. *Holzforschung*, 57: 353–358.
- Punja, Z.K. 2001. Genetic engineering of plants to enhance resistance to fungal pathogens – a review of progress and future prospects. *Can. J. Plant Path.*, 23: 216–235.
- Quesada, M., Sanchez-Azofeifa, G. A., Alvarez-Añorve, M., Stoner, K.E., Avila-Cabadilla, L., Calvo-Alvarado, J., Castillo, A., Espírito-Santo, M.M., Fagundes, M., Fernandes, G.W., Gamon, J., Lopezaraiza-Mikel, M., Lawrence, D., Cerdeira Morellato, L.P., Powers, J.S., de S. Neves, F., Rosas-Guerrero, V., Sayago, R. & Sanchez-Montoya, G. 2009. Succession and management of tropical dry forests in the Americas: review and new perspectives. *For. Ecol. Manage.*, 258(6): 1014–1024.
- Rathore, T.S., Rangaswamy, M., Goyal, B., Dubey, A.K. & Rao, P.S. 2008. Micropropagation through axillary shoot proliferation and somatic embryogenesis in Sandal wood (*Santalum album* L.) from mature trees. In S.A. Ansari, C. Narayanan & A.K. Mandal, eds. *Forest biotechnology in India*. Delhi, Satish Serial Publishing House.
- Remigi, P., Faye, A., Kane, A., Deruaz, M., Thioulouse, J., Cissoko, M., Prin, Y., Galiana, A., Dreyfus, B. & Duponnois, R. 2008. The exotic legume tree species *Acacia holosericea* alters microbial soil functionalities and the structure of the arbuscular mycorrhizal community. *Appl. Environ. Microbiol.*, 74 (5): 1485–1493.
- Richardson, D.M. & Petit, R.J. 2006. Pines and invasive aliens: Outlook on transgenic pine plantations in the southern hemisphere. In: C.G. Williams, ed. *Landscapes, genomics and transgenic conifer forests*. New York, Springer.
- Richardson, J.E., Pennington, R. T., Pennington, T.D. & Hollingsworth, P.M. 2001. Rapid diversification of a species-rich genus of neotropical rain forest trees. *Science*, 293: 2242–2245.
- Robinson, C. 1999. Making forest biotechnology a commercial reality. *Nat. Biotechnol.*, 17: 27–30.
- Roy, S., Islam, M. & Hadiuzzaman, S. 1998. Micropropagation of *Elaeocarpus robustus* Roxb. *Plant Cell Rep.*, 17: 810–813.
- Sarker, R.H., Islam Rafiqul, M. & Hoque, M.I. 1997. *In vitro* propagation of neem (*Azadirachta indica* A. Juss) plants from seedlings explants. *Plant Tissue Cult.*, 7: 125–133.

- Sebbenn, A.M., Degen, B., Azevedo, V.C.R., Silva, M.B., de Lacerda, A.E.B., Ciampi, A.Y., Kanashiro, M., Carneiro, F. da S., Thompson I. & Loveless, M.D. 2008. Modelling the long-term impacts of selective logging on genetic diversity and demographic structure of four tropical species in the Amazon. *For. Ecol. Manage.*, 254: 335–349
- Shali, T.P. & Muralidharan, E.M. 2008. Micropropagation of adult *Bambusa balcooa* Roxb. through multiple shoot formation and somatic embryogenesis. In S.A. Ansari, C. Narayanan, & A.K. Mandal, eds. *Forest biotechnology in India*. pp. 157–164. Delhi, Satish Serial Publishing House.
- Shin D.I., Podila, G.K., Huang, Y. & Karnosky, D.F. 1994. Transgenic larch expressing genes for herbicide and insect resistance. *Can. J. For. Res.*, 24: 2059–2067.
- Silva, M.B., Kanashiro, M., Ciampi, A. Y., Thompson, I. & Sebben, A.M. 2008. Genetic effects of selective logging and pollen gene flow in a low-density population of the dioecious tropical tree *Bagassa guianensis* in the Brazilian Amazon. *For. Ecol. Manage.*, 255: 1548–1558
- Tanaka, K., Tsumura, Y. & Nakamura, T. 1999. Development and polymorphism of microsatellite markers for *Fagus crenata* and the closely related species, *F. japonica*. *Theor. Appl. Genet.*, 99: 11–15.
- Tang, W. & Tian, Y. 2003. Transgenic loblolly pine (*Pinus taeda* L.) plants expressing a modified delta-endotoxin gene of *Bacillus thuringiensis* with enhanced resistance to *Dendrolimus punctatus* Walker and *Crypytothelea formosicola* Staud. *J. Expt. Bot.*, 54: 835–844.
- Tartorius, T.E., Fowke, L.C. & Dunstan, D.I. 1991. Somatic embryogenesis in conifers. *Can. J. Bot.*, 69: 1873–1899.
- Thorpe, T.A., Harry, I.S. & Kumar, P.P. 1991. *Application of micropropagation to forestry*. P.C. Debergh & R.H. Zimmerman, eds., pp. 311–336. Dordrecht (Netherlands), Kluwer Academic Publishers.
- Tiwari, S.K., Tiwari, K.P. & Siril, E.A. 2002. An improved micropropagation protocol for teak. *Plant Cell, Tissue Organ Cult.*, 71: 1–6.
- Tuskan, G.A. 2007. Bioenergy, genomics, and accelerated domestication: A U.S. example. *Paper prepared for the FAO seminar on the role of agricultural biotechnologies for production of bioenergy in developing countries*, Rome, 12 October 2007. (available at www.fao.org/biotech/docs/tuskan.pdf).
- Tuskan, G.A., Difazio, S., Jansson, S. et al. 2006. The genome of black cottonwood, *Populus trichocarpa* (Torr. & Gray) *Science*, 313: 1596–1604.
- Van Bueren, M. 2005. *Acacia hybrids in Vietnam. Impact assessment of ACIAR-funded project FST/1986/030*. Canberra, Australia, Australian Centre for International Agricultural Research.
- Viera, D.L.M., Holl, K.D. & Peneireiro, F.M. 2009. Agro-successional restoration as a strategy to facilitate tropical forest recovery. *Restor. Ecol.*, 17: 451–459.
- Wakamiya, I., Newton, R., Johnston, J.S. & Price, H.J. 1993. Genome size and environmental factors in the genus *Pinus*. *Am. J. Bot.*, 80: 1235–1241.
- Walter, C., Grace, L.J., Wagner, A., Walden, A.R., White, D.W.R., Donaldson, S.S., Hinton, H.H., Gardner, R.C. & Smith, D.R. 1998. Stable transformation and regeneration of transgenic plants of *Pinus radiata* D. Don. *Plant Cell Rep.*, 17: 460–468.
- Wann, S.R. 1988. Somatic embryogenesis in woody species. *Hortic. Rev.*, 10: 153–81.
- Watt, M.P., Blakeway, F., Cresswell, C.F. & Herman, B. 1991. Somatic embryogenesis in *Eucalyptus grandis*. *S. Afr. Forest. J.*, 157: 59–65.
- Watt, M.P., Blakeway, F.C., Mokotedi, M.E.O. & Jain, S.M. 2003. Micropropagation of *Eucalyptus*. In S.M. Jain & K. Ishii, eds. *Micropropagation of woody trees and fruits*, pp. 217–244. Dordrecht, Netherlands, Kluwer.
- White, T. L. & Carson, M.J. 2004. Breeding programmes of conifers. In: C. Walter & M. J. Carson, eds. *Plantation forest biotechnology for the 21st century*. Kerala, India, Research Signpost.

- Whitham, T.G., Bailey, J.K., Schweitzer, J.A., Shuster, S.M., Bangert, R.K., LeRoy, C.J., Lonsdorf, E., Allan, G.J., DiFazio, S.P., Potts, B.M., Fischer, D.G., Gehring, C.A., Lindroth, R.L., Marks, J., Hart, S.C., Wimp, G.M. & Wooley, S.C. 2006. A framework for community and ecosystem genetics: From genes to ecosystems. *Nat. Rev. Genet.*, 7: 510–523.
- Wilcox, P. L., Carson, S.D., Richardson, T.E., Ball, R.D., Horgan, G.P. & Carter, P. 2001. Cost–benefit analysis of marker based selection in seed orchard production populations of *Pinus radiata*. *Can. J. For. Sci.*, 31: 2213–2224.
- Williams, C.G. & Reyes-Valdes, M.H. 2007. Estimating a founder’s genomic proportion for each descendant in an outbred pedigree. *Genome*, 50: 289–296.
- Williams, C.G., Reyes-Valdés, M.H. & Huber, D.A. 2007. Validating a QTL region characterized by multiple haplotypes. *Theor. Appl. Genet.*, 116: 87–94.
- Williams, C.G., Hamrick, J.L. & Lewis, P.O. 1995. Multiple–population versus hierarchical conifer breeding programmes: A comparison of genetic diversity levels. *Theor. Appl. Genet.*, 90: 584–594.
- World Bank. 2001. *A revised forest strategy for the World Bank Group*. Draft. Washington, DC, World Bank.
- Xiao-hua, S., Bing-yu, Z., Qin-Jun, H., Lie-jian, H., & Xiang-hua, Z. 2003. Advances in tree genetic engineering in China. *Paper submitted to the XII world forestry congress, 2003*, Quebec City, Canada. (available at www.fao.org/docrep/article/wfc/xii/0280-b2.htm).
- Yanchuk, A.D. 2001. The role and implications of biotechnological tools in forestry. *Unasylva*, 204: 53–61. (also available at www.fao.org/docrep/003/x8820e/x8820e10.htm).
- Yazdani, R., Scotti, I., Jansson, G., Plomion, C. & Mathur, G. 2003. Inheritance and diversity of simple sequence repeat (SSR) microsatellite markers in various families of *Picea abies*. *Hereditas*, 138: 219–227.
- Young, A.G. & Clarke, G.M., eds. 2000. *Genetics, demography and viability of fragmented populations*. Cambridge, UK., Cambridge University Press.