


**LOBOTIDAE**

Vol. 5, p. 2942

**Tripletails**

To over 100 cm. In coastal waters and estuaries, some entering fresh water; also found offshore around floating objects. Three species in the area. Species of the genus *Coius* (formerly *Datniodes*) sometimes placed in a separate family, Coiidae or Datnioidae.





**GERREIDAE**

Vol. 5, p. 2946

**Silverbiddies, mojarras**

To about 35 cm. Near bottom in coastal and estuarine waters to depths of about 70 m; also entering fresh water. Fourteen species in the area.





**HAEMULIDAE**

Vol. 5, p. 2960

**Grunts, sweetlips, rubberlips**

To 120 cm. Near bottom in estuaries and in coastal waters on reefs or over soft bottom. Twenty-five species in the area.




**SPARIDAE**

Vol. 5, p. 2989


**Porgies**

To 75 cm. Coastal, demersal to a depth of about 150 m, occasionally entering estuaries. Twelve species in the area.

teeth canine-like, conical or incisor-like, molars often present, roof of mouth toothless




hind tip of premaxilla overlapping maxilla; maxilla in sheath when mouth closed

**LETHRINIDAE**

Vol. 5, p. 3002

**Emperors**

To 100 cm. Demersal in coastal waters, mostly over sandy, rocky, and coral reef bottom, to outer edge of continental shelf at depths to 180 m. Thirty-three species in the area.




no scales between eye and mouth but cheek scaled in Monotaxinae

*Monotaxinae (Gymnocranius)*

outer teeth canine-like or conical, molars often present, inner teeth villiform, roof of mouth toothless


single dorsal fin with X spines and 9 soft rays (10 in Monotaxinae)

*Lethrininae (Lethrinus)***NEMIPTERIDAE**

Vol. 5, p. 3049

**Threadfin breams**


To about 35 cm standard length. Demersal in coastal continental shelf waters, mostly over sand and mud bottom to a depth of about 300 m. Fifty-four species listed from the area.



suborbital spine in *Scolopsis*

*Scolopsis*

single dorsal fin with X spines and 9 soft rays


*Nemipterus*

**POLYNEMIDAE**

Vol. 5, p. 3088

**Threadfins**

To about 200 cm; marine coastal waters, estuaries, coastal lagoons and fresh water river mouths; benthic.





**SCIAENIDAE**

Vol. 5, p. 3115

**Croakers**

To 160 cm standard length. Demersal, coastal and continental shelves, mostly over soft bottoms; some also in estuaries and in fresh water. Around 50 species in the area.




**MULLIDAE**

Vol. 5, p. 3173

**Goatfishes**

To 50 cm. Bottom coastal fishes, mostly on soft bottoms or sandy areas around reefs. Twenty-nine species in the area.




**PEMPHERIDAE**

Vol. 5, p. 3199

**Sweepers**

To about 22 cm. Bottom coastal fishes, most common around coral reefs and down to a depth of around 100 m. Around 10 species in the area.




**GLAUCOSOMATIDAE**

Vol. 5, p. 3203

**Pearl perches**


To about 70 cm. Coastal on rocky or reef bottom in moderately deep water. Three species in the area.

**LEPTOBRAMIDAE**

Vol. 5, p. 3208

**Beachsalmon**


To about 43 cm. Coastal, entering brackish and fresh water. A single species in the family, also in the area.

**BATHYCLUPEIDAE**

Vol. 5, p. 3209

**Bathyclupeids**


To 30 cm. Deep-sea fishes on the continental slope and oceanic waters; meso-, bathy-, or benthopelagic. Three species in the area.

**TOXOTIDAE**

Vol. 5, p. 3210

**Archerfishes**

To about 50 cm. Mangroves and brackish waters, also entering fresh water. Two brackish-water species in the area.




**MONODACTYLIDAE**

Vol. 5, p. 3214

**Moonfishes**

To 25 cm. Coastal, estuaries and entering fresh water. Three species in the area.




**DREPANIDAE**

Vol. 5, p. 3219

**Sicklefishes**

To 50 cm. Coastal, mostly over sand or mud bottom, also on reefs and enters estuaries. Two species in the area.




**CHAETODONTIDAE**

Vol. 5, p. 3222

**Butterflyfishes**

To about 30 cm. Mostly on shallow coral reefs to a depth of 60 m; some species restricted to deep water below 60 m. Seventy species in the area.




**POMACANTHIDAE**

Vol. 5, p. 3264

**Angelfishes**


To about 50 cm. Most closely associated with coral reefs at depths between 5 and 60 m, some species restricted to greater depths. Around 44 species in the area.

**ENOPLOSIDAE**

Vol. 5, p. 3285

**Old wife**


To 31 cm. Coastal rocky reefs and seagrass beds from shallow water to a depth of 100 m. A single species in the area.

**PENTACEROTIDAE**

Vol. 5, p. 3286

**Armourheads**

To 35 cm. On hard bottom at depths from 40 to 400 m. A single species in the area.




**KYPHOSIDAE**

Vol. 5, p. 3288

**Seachubs**

To 60 cm. Occurs on rocky or coral reefs. Five species in the area.




**GIRELLIDAE**

Vol. 5, p. 3295

**Nibblers**

To 76 cm. Estuaries and around coastal rocky reefs. Two species marginally in the area.




**SCORPIDIDAE**

Vol. 5, p. 3297

**Halfmoons**

To 37 cm. Mostly on rocky reefs in estuaries and coastal areas. A single species marginally in the area.




**MICROCANTHIDAE**

Vol. 5, p. 3299

**Microcanthids**

To 25 cm. In coastal and estuarine waters, typically around rocky reefs. Two species marginally in the area.



**ARRIPIDAE**

Vol. 5, p. 3302

**Australian salmon**

To about 1 m. Shallow coastal waters and estuaries. A single species in the area.

terminal mouth,  
maxilla reaching  
below centre of eye

single dorsal fin with IX spines, the spinous  
portion noticeably higher than soft-rayed  
portion, and 15-18 soft rays

anal fin with III spines  
and 9-10 soft rays

**TERAPONTIDAE**

Vol. 5, p. 3303

**Terapon perches**

To about 35 cm. Coastal waters and estuaries, also entering fresh water. Nine species in the area.

opercle with 2 spines  
on posterior edge, the  
lower one stronger

single dorsal fin with XI-XIV spines  
and 8-14 soft rays

upper jaw not  
extending beyond  
centre of orbit

anal fin with III spines  
and 7-12 soft rays

caudal fin usually  
emarginate

**KUHLIIDAE**

Vol. 5, p. 3315

**Flagtails**

To 50 cm. Coastal waters and estuaries, also entering fresh water. Around 6 species in the area.

2 spines on  
opercle

single, deeply notched  
dorsal fin with X spines  
and 9-12 soft rays

well-developed scaly sheath at  
base of dorsal and anal fins

anal fin with III spines  
and 9-13 soft rays

**CIRRHITIDAE**

Vol. 5, p. 3319

**Hawkfishes**

To about 30 cm. On the bottom, mostly in shallow waters on coral or rocky reefs. Nineteen species in the area.

1 or more cirri on tips of  
membranes of spinous dorsal fin

single, notched dorsal  
fin with X spines and  
11-17 soft rays

fringe of cirri on  
posterior nostril

lower 5-7 rays of pectoral fin  
unbranched and usually enlarged

anal fin with III spines  
and 5-7 soft rays




CHEILODACTYLIDAE

Vol. 5, p. 3327

Morwongs

To around 45 cm. On the bottom, mostly inshore on rocky or coral reefs. Four species in the area.

mouth small, terminal to inferior, with thick lips  
teeth in jaws small, viliform, in bands; no teeth on vomer or palatines




CEPOLIDAE

Vol. 5, p. 3329

Bandfishes

To about 40 cm. In burrows in sand or mud bottom, from shallow waters to a depth of 475 m. Around 9 species in the area, although the taxonomy is very confused.




Order PERCIFORMES: LABROIDEI - Wrasses and allies

CICHLIDAE

Vol. 5, p. 3331

Cichlids

To about 60 cm. Usually fresh water, but entering brackish waters. Ten species tolerant to brackish water introduced into the area, but only 1 species regularly found in brackish water.




**POMACENTRIDAE**

Vol. 5, p. 3335

**Damselfishes**

To about 25 cm, but most species smaller than 5 cm. Very common on coral and rocky reefs and also occurring in other shallow habitats such as sand, rubble, silty, and seagrass bottoms. So far, 209 species recorded from the area.









**LABRIDAE**

Vol. 6, p. 3381

**Wrasses**

To about 200 cm (but most species less than 20 cm). Very common on coral reefs and also found on rocky, sand, and seaweed bottoms down to a depth of at least 100 m. So far, 183 species listed from the area.




diverse body shapes, a few representative forms shown here

**SCARIDAE**

Vol. 6, p. 3468

**Parrotfishes**

To 120 cm. Most common on coral reefs at depths shallower than 30 m. Around 44 species in the area.




**Order PERCIFORMES: ZOARCOIDEI - Eelpouts**

**ZOARCIDAE**

Vol. 6, p. 3493

**Eelpouts**

To around 20 cm. Deepsea pelagic. Three species in the area.



**Order PERCIFORMES: TRACHINOIDEI - Stargazers and allies**


A diverse assemblage of families difficult to characterize as a group; the composition of this suborder is provisional.

**CHIASMODONTIDAE**

Vol. 6, p. 3495

**Swallowers**

To about 26 cm. Meso- and bathypelagic. Twelve species in the area.



**CHAMPSODONTIDAE**

Vol. 6, p. 3497

mouth large, long  
depressible teeth2 dorsal fins, the 1<sup>st</sup> with IV-VI spines  
and the 2<sup>nd</sup> with 18-23 soft rays**Gapers**


To about 15 cm. Pelagic at depths from 34 to 1 100 m, reaching shallow depths only at night. A single species possibly in the area.

**PHOLIDICHTHYIDAE**

Vol. 6, p. 3500

**Convict fish**


To at least 34 cm. On rocky or coral reefs or over mixed soft and hard bottom. Two species in the area.

**PINGUIPEDIDAE**

Vol. 6, p. 3501

**Sandperches**


To about 30 cm. On sand, rubble, or mud bottoms in coastal waters. Twenty-two species in the area.

**TRICHONOTIDAE**

Vol. 6, p. 3511

**Sanddivers**


To about 16 cm standard length. Over sand bottom in shallow water to depths of about 50 m. Two described species from the area, at least 1 other undescribed species also in the area.

**CREEDIIDAE**

Vol. 6, p. 3513

**Sandburrowers**

To about 8 cm standard length. On sand or rubble bottom from shallow depths to around 150 m. Around 6 species in the area.




**PERCOPHIDAE**

Vol. 6, p. 3515

**Duckbills**

To about 30 cm standard length. Demersal, from shallow water to a depth of 600 m. About 10 species in the area.




**LEPTOSCOPIDAE**

Vol. 6, p. 3517

**Southern sandfishes**

To about 40 cm. Burrowing on sandy and muddy bottoms in coastal areas. A single species in the area.




**AMMODYTIDAE**

Vol. 6, p. 3518

**Sandlances**

To about 15 cm. On sandy coastal bottoms. Three species in the area.




**URANOSCOPIDAE**

Vol. 6, p. 3519

**Stargazers**

To about 65 cm. Burrowing in sand or mud bottom in coastal and estuarine areas. Sixteen species in the area.




**Order PERCIFORMES: BLENNIOIDEI - Blennies and allies**

**TRIPTERYGIIDAE**

Vol. 6, p. 3532

**Triplefins**

Most species less than 6 cm. Usually cryptic in shallow coastal rocky and coral habitats. Around 60 species in the area.




CLINIDAE

Vol. 6, p. 3536

Clinids

To about 60 cm, most smaller. Coastal inshore and estuaries, in rocky pools and among seaweeds. At least 4 species in the area.




BLENNIIDAE

Vol. 6, p. 3538

Blennies

Most less than 15 cm, 2 eel-like species reaching 53 cm. Coastal and estuarine, usually at depths less than 20 m, mostly on rocky and coral reefs, tidepools, and among seaweeds; a few in fresh water. About 195 species in the area.




Order PERCIFORMES: GOBIESOCOIDEI - Clingfishes

GOBIESOCIDAE

Vol. 6, p. 3547

Clingfishes

To about 7 cm. Coastal reef and seagrass habitats from shallow water to a depth of about 200 m. Nine species in the area.




Order PERCIFORMES: CALLIONYMOIDEI - Dragonets

CALLIONYMIDAE

Vol. 6, p. 3549

Dragonets


To about 45 cm. Demersal on sandy, muddy, rocky, or coral bottom, from the intertidal zone to a depth of 800 m. About 71 species in the area.



DRACONETTIDAE

Vol. 6, p. 3572

**Deepwater dragonets**  
To 14 cm. On sandy or muddy bottoms at depths between about 130 and 600 m. Five species listed from the area.




Order PERCIFORMES: GOBIOIDEI - Gobies and allies

ELEOTRIDAE

Vol. 6, p. 3574




**Sleepers**  
To about 60 cm but most do not exceed 20 cm. Generally fresh water, but entering brackish waters, some species found on coral reefs. Around 23 species in the area.



GOBIIDAE

Vol. 6, p. 3578

**Gobies**  
To about 30 cm, but most species less than 10 cm. Demersal in coastal waters; some in brackish waters and a few in fresh waters; usually in shallow depths, but a few species to below a 150 m; benthic; the largest family of fishes; of little interest to fisheries, but some species of interest to the aquarium trade. Around 500 species in the area.






**KRAEMERIIDAE**

Vol. 6, p. 3604

**Sandgobies**

To about 7 cm. Burrowing in sand or muddy sand in shallow coastal and estuarine bottoms. Six species in the area.




**XENISTHMIIDAE**

Vol. 6, p. 3605

**Wrigglers**

To 3.5 cm. Cryptic in coral, coral rubble, and sand habitats from the shoreline to a depth of around 30 m. Seven species in the area.




**MICRODESMIIDAE**

Vol. 6, p. 3607

**Wormfishes**

To almost 13 cm. Near bottom on coral reefs or sandy areas near coral reefs. Thirty-six species in the area.




**SCHINDLERIIDAE**

Vol. 6, p. 3609

**Schindler's fishes**

To around 2 cm. Pelagic, offshore or near coral reefs. Two species in the area.




**Order PERCIFORMES: KURTOIDEI - Nurseryfishes**

**KURTIDAE**

Vol. 6, p. 3610

**Nurseryfishes**

To about 36 cm. Brackish and fresh water. A single species.






Order PERCIFORMES: ACANTHUROIDEI - Surgeonfishes

EPHIPPIDAE

Vol. 6, p. 3611

Spadefishes

To 25 cm. Nearshore shallow water in estuaries, on coral reefs, sandy, or muddy bottom. Six species in the area.




SCATOPHAGIDAE

Vol. 6, p. 3623

Scats

To about 32 cm. Mostly in estuaries, but also entering fresh water. Two species currently recognized in the area.




SIGANIDAE

Vol. 6, p. 3627

Rabbitfishes

To 55 cm. Bottom dwelling on coral reefs, seagrass beds, estuaries, and magrove swamps, in shallow water to depths of around 50 m. Twenty-one species in the area.




**ZANCLIDAE**

Vol. 6, p. 3651

**Moorish idol**


To about 22 cm. On coral or rocky reefs from shallow water to a depth of 180 m.

**ACANTHURIDAE**

Vol. 6, p. 3653

**Surgeonfishes**


To about 75 cm. Near bottom on coral or rocky reefs, generally at depths less than 100 m. Usually at a depth of less than 50 m. Fifty-six species in the area.

**Order PERCIFORMES: SCOMBROLABRACOIDEI - Longfin escolar****SCOMBROLABRACIDAE**

Vol. 6, p. 3684

**Longfin escolar**

To 30 cm. On the continental shelf and slope between 100 and 900 m. A single species.



Order PERCIFORMES: SCOMBROIDEI - Tunas and allies

SPHYRAENIDAE


Vol. 6, p. 3685

Barracudas

To 170 cm. Nearshore pelagic and demersal from near surface to a depth of about 100 m. Around 9 species in the area.

UPPER JAW NOT PROTRUSIBLE, PREMAXILLA FIXED

PELVIC FINS WITH 0-1 SPINE AND 0-6 RAYS, OFTEN REDUCED OR ABSENT




GEMPYLIDAE


Vol. 6, p. 3698

Snake mackerels

To about 300 cm. Marine, from the surface (at night) to below a depth of 200 m; pelagic. Fourteen species listed from the area.



Gempylus




Lepidocybium

TRICHIURIDAE

Vol. 6, p. 3709

Cutlassfishes

To 225 cm. Benthopelagic on continental shelves and slopes from the surface to a depth of about 2 000 m. Sixteen species in the area.




SCOMBRIDAE


Vol. 6, p. 3721

Tunas

To over 5 m. Epipelagic nearshore and oceanic. Thirty species in the area.



Scomberomorus




Thunnus

**XIPHIIDAE**

Vol. 6, p. 3757

**Swordfishes**

To 450 cm. Pelagic, oceanic, from surface waters to a depth of 550 m. A single species.




**ISTIOPHORIDAE**


Vol. 6, p. 3759

**Billfishes**

To 5 m. Epipelagic oceanic, usually above the thermocline. Five species in the area.



*Makaira*



*Istiophorus*


**Order PERCIFORMES: STROMATEOIDEI - Butterfishes and allies**

**AMARSIPIDAE**

Vol. 6, p. 3765

**Amarsipas**

To 22 cm standard length. Epipelagic. A single species in the area.




**CENTROLOPHIDAE**

Vol. 6, p. 3767

**Medusafishes**

To 137 cm. Epi- and mesopelagic oceanic and nearshore demersal. Four species in the area.




**NOMEIDAE**

Vol. 6, p. 3771

**Driftfishes**

To 1 m. Epipelagic, oceanic. Around 10 species in the area.




**ARIOMMATIDAE**

Vol. 6, p. 3780

**Ariommas**

To 78 cm. Demersal over soft bottoms on the continental shelf and slope to a depth of at least 750 m. Three species in the area.




**TETRAGONURIDAE**

Vol. 6, p. 3784

**Squaretails**

To 50 cm. Oceanic epi- and mesopelagic. Two species in the area.




**STROMATEIDAE**

Vol. 6, p. 3786

**Butterfishes**

To about 60 cm. Coastal, usually at depths shallower than 100 m. Around 4 species in the area.




Order PLEURONECTIFORMES - Flatfishes

PSETTODIDAE

Vol. 6, p. 3792

Spiny turbot

To 60 cm. Demersal on sand and mud bottoms from shallow depths down to around 100 m. A single species in the area.




CITHARIDAE

Vol. 6, p. 3794

Largescale flounders

To 36 cm. Demersal, mostly on sandy bottoms. Three species in the area.




BOTHIDAE

Vol. 6, p. 3799

Lefteye flounders

To about 42 cm, most smaller than 20 cm. Demersal, from the coastline to a depth of about 980 m. Around 59 species in the area.




**PARALICHTHYIDAE**

Vol. 6, p. 3842

**Sand flounders**


To 40 cm. Demersal on shallow sandy and muddy bottoms; also entering brackish waters. Seventeen species in the area.



ventral view of pelvic fins

pelvic-fin bases  
nearly symmetrical  
in position

eyes on left  
side of body




elongate anterior dorsal-fin rays  
(present in 2 species of *Samaris*, but  
absent in all other species in the area)

**PLEURONECTIDAE**

Vol. 6, p. 3863

**Righteye flounders**

To about 22 cm. Demersal on mixed bottoms of mud, sand, silt, and crushed shells, mostly at depths of about 60 to 500 m; some species in shallow water on coral reefs. Twenty species in the area.




**SOLEIDAE**

Vol. 6, p. 3878

**Soles**

To about 32 cm. Demersal on shallow sandy and muddy bottoms; some species enter estuaries. Around 48 species in the area.




**CYNOGLOSSIDAE**

Vol. 6, p. 3890

**Tonguesoles**

To 40 cm. Demersal, mostly on muddy bottoms, from shallow coastal waters to depths of about 1 500 m. Around 35 species in the area.





## Order TETRAODONTIFORMES - Pufferfishes and allies

### TRIACANTHODIDAE

Vol. 6, p. 3902

#### Spikefishes

To 20 cm. Demersal, mostly at depths between 35 and 900 m. Thirteen species in the area.

A SMALL MOUTH WITH STRONG TEETH, FREQUENTLY COALESCED INTO A BITING PLATE

SKIN THICK OR ROUGH, SCALES USUALLY MODIFIED AS PRICKLES, SPINES, SHIELDS, OR PLATES

SMALL GILL OPENING


pelvic fins with I large spine and 1-2 rudimentary soft rays

VI visible dorsal-fin spines

caudal peduncle not distinctly tapering

caudal fin rounded to nearly truncate

PELVIC FINS ABSENT OR STRONGLY REDUCED



### TRIACANTHIDAE

Vol. 6, p. 3905

#### Tripodfishes, triplespines


To 30 cm. Mostly on flat, sandy, or seaweed bottoms. Seven species in the area.

pelvic fins with I large spine and no visible soft rays

V visible dorsal-fin spines

caudal peduncle distinctly tapering, narrow

caudal fin deeply forked



### BALISTIDAE

Vol. 6, p. 3911

#### Triggerfishes


To about 80 cm. Mostly bottom dwelling on coral or rocky reefs from the coastline to a depth of about 100 m; some species primarily pelagic in open water. Twenty-two species in the area.

pelvic fins rudimentary, only 4 enlarged scales encasing end of pelvis

III dorsal-fin spines visible

strong teeth

skin thick, with conspicuous plate-like scales



### MONACANTHIDAE

Vol. 6, p. 3929


#### Filefishes, leatherjackets

To 1 m. Mostly on coral and rocky reefs, seagrass beds, and sandy bottom at depths to over 200 m. Around 46 species in the area.

skin rough, with minute scales armed with spinules

II dorsal-fin spines, the 1<sup>st</sup> spine prominent, the 2<sup>nd</sup> very small

pelvic fins a bony rudiment fused to posterior end of pelvis




**OSTRACIIDAE**

Vol. 6, p. 3948

**Boxfishes**

To 40 cm. Bottom dwelling on rocky and coral reefs, and sandy bottoms, from the coastline to a depth of 100 m. Ten species in the area.




**ARACANIDAE**

Vol. 6, p. 3952

**Trunkfishes**

To about 40 cm. Bottom dwelling on rocky and coral reefs, and sandy bottom to a depth of about 200 m. A single species in the area.




**TRIODONTIDAE**

Vol. 6, p. 3953

**Threetooth puffers**

To 60 cm. In deep water to a depth of 300 m. A single species, also in the area.




**TETRAODONTIDAE**

Vol. 6, p. 3954

**Puffers**

To about 80 cm. Most in shallow, inshore water, entering estuaries and occasionally freshwater; some species pelagic. Fifty-one species in the area.




**DIODONTIDAE**

Vol. 6, p. 3958

**Porcupinefishes**

To 1 m. Mostly near bottom around coral or rocky reefs, sometimes over seaweed, sand, or mud bottoms, from the coastline to a depth of about 100 m. Eleven species in the area.




**MOLIDAE**

Vol. 6, p. 3966

**Molas**

To 300 cm or more. Open ocean pelagic, from the surface to about a depth of 360 m. Four species in the area.



**Class SARCOPTERYGII**

**Order COELACANTHIFORMES**

**LATIMERIIDAE**

Vol. 6, p. 3969

**Coelacanths**

To at least 1.24 m. Around volcanic islands at depths between 100 and 150 m. Probably spends the day inside caves and comes out at night to forage. A single representative recently discovered in the area and described as a new species. Recognized in the Coelacanthidae by some authors.

