ix

Tables

| · · · · · · · · · · · · · · · · · · · | |
|--|----|
| TABLE 1 | |
| Per caput DES by region and economic group, 1969-71, 1979-81 and 1990-92 | 11 |
| TABLE 2 | |
| Top five and bottom five developing countries in terms of per caput DES growth rates, 1969-71 to 1990-92 | 12 |
| TABLE 3 | |
| DES and population growth by region and economic group, 1969-71 to 1979-81 and 1979-81 to 1990-92 | 13 |
| TABLE 4 | |
| Distribution of 98 developing countries by per caput DES in 1969-71 and its growth from 1969-71 to 1990-92, and by population growth | 15 |
| TABLE 5 | |
| Total and animal protein supplies by region and economic group, 1969-71, 1979-81 and 1990-92 | 16 |
| TABLE 6 | |
| Total and animal fat supplies by region and economic group, 1969-71, 1979-81 and 1990-92 | 17 |
| TABLE 7 | |
| Share of major food groups in total DES by region and economic group, 1969-71 and 1990-92 | 23 |

| | TABLE 16 | |
|-----|---|----|
| | Average per caput energy consumption of undernourished population compared with minimum and average per caput energy requirements | 53 |
| | TABLE 17 | |
| | Relative inadequacy of food supply by developing region, 1969-71, 1979-81 and 1990-92 | 56 |
| | TABLE 18 | |
| | Distribution of 98 developing countries classified according to their relative inadequacy ratios, 1969-71 and 1990-92 | 57 |
| Н., | TABLE 19 | |
| | Classification of 98 developing countries into four categories based on roles of per caput DES growth and redistribution in eliminating food inadequacy | 60 |
| | TABLE 20 | |
| | Comparison of actual and required levels of per caput DES, 1990-92 | 61 |
| | TABLE 21 | |
| | Estimates of prevalence and numbers of wasted, stunted and underweight children in developing countries, 1990 | 69 |
| | TABLE 22 | |
| | Change over time in the prevalence of underweight children (under 5 years) in selected developing countries | 72 |

| TABLE 23 | |
|---|----|
| Prevalence of stunting indicated by national height surveys among schoolchildren in Latin America and the Caribbean | 75 |
| TABLE 24 | |
| Prevalence of stunting and thinness among adolescent males and females in selected countries | 76 |
| TABLE 25 | |
| Percentage of adults in different BMI classes based on available country surveys | 77 |