

Annexure 6

ANALYSIS OF SOME MORPHOMETRIC AND MERISTIC CHARACTERS OF *HILSA ILISHA* OF BANGLADESH WATERS

by N.N. Das, T. Nishida, S.A. Azad, M.S. Islam, M. Hossain and Q.M. Huq*

1. INTRODUCTION

The Hilsa Shad, *Hilsa ilisha* (Hamilton), is one of the most important species of food fish caught in Bangladesh and is a major source of protein in the people's diet and for providing employment. Over the past four decades or so, a number of reports have dealt with morphometric and meristic differences in *Hilsa ilisha* in the Indo-Bangladesh region. The important work on Bangladesh hilsa is that of Quddus et al. (1984), besides brief studies of Quereshi (1968) and Shafi, Quddus and Hossain (1977). However, all these investigations had covered inland waters only — riverine and estuarine. From the review attempted by Raja (1985), it seems that these investigations have resulted in establishing individually different stocks in each of the major river systems in the Indo-Bangladesh region. In addition he had also drawn attention to the report of segregation of stocks into at least two varieties, a broad and slender one, in India, Bangladesh and Burma. It was also pointed out by him that no studies have been made on the composition, continuity, independence or interdependence of the stocks of the marine sector. Provisionally, based purely on the evidence of available literature, he suggested that there may be at least four stocks in the region, three anadromous and one marine. The three anadromous stocks may be distributed thus: one in the Indo-Bangladesh area, one in eastern Bangladesh and one in the central and southern regions of Burma. The marine stock may be contributing to the fishery of south-eastern Bangladesh coast contiguous with the Arakan coast of Burma. These are supposed to be in addition to the purely fluvial stocks in the upper reaches of the five major river systems in India, Bangladesh and perhaps in Burma.

Because of the total lack of any racial studies on the hilsa of the marine environment, the investigations launched by the UNDP/FAO project "Marine Fishery Resources Management in the Bay of Bengal" included the aspect of racial studies in the programme to cover the hilsa from riverine, estuarine and marine environments,

All approaches were considered -biometric, biochemical, cytogenetic and tagging. It was also originally planned to subcontract the work to a university. Financial demands for undertaking biochemical/cytogenetic studies were too high to be accommodated within the project's budget. Tagging experiments are a still more expensive undertaking. Hence it was decided to limit the approach to the conventional one, biometrics. In its attempt at exploring the possibilities of subcontracting the studies to the university, the project lost considerable time. Hence the studies could be initiated only during the middle of the project period, September 1985, thus limiting the study period to about 6-7 months.

* T. Nishida is from the BOBP, the other authors are from the Directorate of Fisheries, Bangladesh.

2. MATERIALS AND METHODS

The samples were collected from September/October 1985 to March 1986 from the experimental and commercial catches from Cox's Bazar, Chittagong, Khepupara and Chandpur.

From Chandpur, an adequate sample size could not be obtained from experimental fishing; hence a part of the samples of Chandpur was procured either from the fishing ground or from the landing centre. For Khepupara, all the samples were procured from commercial catches since no experimental fishing could be conducted in that area. The number of fish examined from different stations by month and sex is shown in Table I.

Originally the plan was to examine the fish the same day they were procured in respect of Chandpur, Khepupara and Chittagong stations. Only with regard to Cox's Bazar would there be some delay in the arrival of iced samples at Chittagong, but they were to be examined immediately after. For this purpose the senior author was expected to be at Chittagong when the samples arrived but unfortunately he could not adhere to this plan. Consequently, the samples caught during experimental fishing were preserved in ice on board, and transferred to deep freeze after landing until they were taken up for examination. Except the Khepupara samples which were examined fresh, the other samples from the fishing grounds or from the landing centre were also kept in deep freeze for later examination. The period that elapsed between the time of procurement and the time of examination is shown in Table II.

The morphometric and meristic characters chosen were those which had in the past been found to have exhibited significant differences. Such selected morphometric characters were the total length (TL), depth at dorsal origin (DP), head length (HL), thickness of body (TH), eye diameter (ED) and caudal peduncle length (CPL). The meristic characters selected are post-pelvic scutes (PS) and pectoral fin rays (PF).

Total length was measured with the help of a metre scale to the nearest millimeter, while body depth, caudal peduncle length, head length, thickness of body and eye diameter were measured with the help of a slide caliper. The accuracy of the slide caliper was 0.5 mm. All meristic counts and morphometric measurements were made following Pillay (1957).

The following definitions were applied to the characteristics measured or counted.

Total length (TL) is the distance from the tip of snout, when the mouth is closed, to the tip of the lower lobe of the caudal fin, when stretched out.

Depth at dorsal origin (DP) is the distance between the dorsal and ventral edges of the body as measured from the origin of dorsal fin.

Head length (HL) is the distance from the tip of snout, when mouth is closed, to the posterior-most point of the operculum.

Thickness of body (TH) is measured across the thickest part, usually at the 5th-6th lateral line scale of each side.

Eye diameter (ED) is measured longitudinally across the left eye.

Caudal peduncle length (CPL) is the distance from the posterior edge of the base of the anal fin to the base of caudal fin.

Pectoral fin rays (PF) is the number of the fin rays in the pectoral fin.

Post-pelvic scutes (PS) is the number of scutes posterior to the origin of the pelvic fin at the anal opening.

In addition, the individual weights of fish and stages of maturity were also recorded.

The data were subjected to regression analysis and analysis of covariance with the help of the Microstat Programme on an Apple IIe microcomputer.

The descriptive statistics of morphometric characters and those of meristic characters are shown in Table III and IV.

3. RESULTS

(i) Morphometric characters

Among all the morphometric characters, the estimated regression statistics between TL vs DP, TL vs HL, TL vs TH, TL vs CPL and HL vs ED between months and stations are shown in Table V to IX. It may be seen from these tables that only for the relationship between TL and HL (Table VI) was the r^2 value more or less uniformly high, at the level of 85 to 100% at all the stations, for the months of November to February. The data were therefore subjected to analysis of covariance. Since the r^2 values of other morphometric characters were very erratic and were not of a significant order, analysis of combinations of these variables, month-wise and station-wise, could not be justified. The erratic and/or low value of r^2 is attributed to the data being affected by a certain factor or factors. Hence no further treatment of data was taken up for other morphometric characters.

Total length and head length

The scatter diagrams for the variables is shown in Fig. 1, and the estimated parameters of the regression equation for the pooled data of each station in Table X. The analysis of covariance for the relationship between total length and head length showed that there were no significant differences in the slopes of the regression lines between stations, but the intercepts exhibited significant differences. The test for the intercepts indicated that those relating to Chittagong and Cox's Bazar were not different from each other and the same was the case between Chandpur and Khepupara. It was hence concluded that some factor could be responsible for this separation of Chittagong/Cox's Bazar fish on the one hand and Chandpur/Khepupara fish on the other.

It may be recalled that the frozen periods for different stations were different (Table II). In order to see whether differences in the frozen period could have affected the intercept, a dozen Hilsa from Chandpur were collected on 12-6-1986 and frozen till they were analysed on 1-7-1986.

Fig. 2 shows the relationship between the number of days of preservation and the values of intercept in the regression equation. It may be seen that the values of intercept increase as the days of preservation increased. After analysis it was found that while the value of the slope remained the same, i.e., 0.23, the value of intercept, 3.0278, falls on the line for the values obtained earlier on the basis of which Fig. 2 was drawn.

Since in the case of samples from Chittagong and Cox's Bazar the preservation days were more than double that of Chandpur, it is possible that the high values of intercept obtained for Chittagong and Cox's Bazar was due to the longer period of preservation. If this difference were to be rejected on the basis of this causal factor, the conclusion is that between TL and HL there is no significant difference between the samples of the four stations.

(ii) Meristic characters

Post-pelvic scutes: Analysis of variance to test the significance of averages among four stations (Table XI) showed that there was no significant difference in the number of post-pelvic scutes among four stations, a conclusion which is completely different from earlier investigations on this character (Pillay, 1957 ; Rao, 1969 ; Quddus *et al.*, 1984).

Pectoral fin rays: The results of analysis (Table XII) indicated the same conclusions as for the post-pelvic scutes, namely, no significant difference between the stations, which again is a departure from some of the earlier works (Rao, 1969, Quddus *et al.*, 1984).

In general, the results from these two characters did not indicate the existence of more than one race of *Hilsa ilisha* at the stations investigated, and there was no evidence or basis for an unbiased separation of samples for any comparative study.

4. SUMMARY

Nearly 500 specimens were measured from four stations representing marine, estuarine and riverine sectors. The morphometric characters selected for the study were total length, depth at dorsal origin, thickness of body, caudal peduncle length, head length and eye diameter; the meristic characters were post-pelvic scutes and pectoral fin rays.

Among the five different pairs of non-meristic characters, the correlation between four of them was found to be erratic and not significant. Although in the case of total length versus head length, the correlation was of high order and there were no significant differences in the regression coefficient, the differences in the values of intercept were traced to the differences in the period of storage under frozen conditions. Thus the results obtained on the morphometric data have to be totally rejected for reasons of defective methodology in preservation of fish.

Analysis of variance on the two meristic characters showed that there were no significant differences in the mean values between stations.

5. LITERATURE CITED

- Pillay, T.V.R., 1957. A morphometric study of the population of hilsa, *Hilsa ilisha* (Hamilton) of the river Hooghly and of the Chilka lake. *Indian J. Fish.*, **4** (2) : 344-386.
- Quddus, M.M.A., Shimizu and Y. Nose, 1984. Meristic and morphometric differences between two types of *Hilsa ilisha* in Bangladesh waters. *Bull. Jap. Soc. Sci. Fish.*, **50** (1) : 43-49.
- Quereshi, M.R., 1968. Hilsa fishery in East Pakistan *Pak. J. Sci. Ind. Res.*, **11** : 95-103.
- Raja, B.T., Antony, 1985. A review of the biology and fisheries of *Hilsa ilisha* in the upper Bay of Bengal. Marine Fishery Resources Management in the Bay of Bengal. BOBP/WP/37 66 PP.
- Rao, Babu. M., 1969. Some observations on the juveniles of *Hilsa ilisha* (Hamilton) (Pisces: Clupeidae) from Godavari estuary. *J. Bombay Nat. His. Soc.*, **66** (1) : 116-131.
- Shafi, M., M.M.A. Quddus, H. Hossain, 1977. A morphometric study of the population of *Hilsa ilisha* (Hamilton-Buchanan) from the river Meghna. *Proc. Second Bangladesh Sci Conf.*, Abstracts : A-40.

Annexure 6




Fig. 1 Scatter diagram of Total Length (TL) versus Head Length (HL) of *Hilsa ilisha*, at Cox's Bazar Chittagong, Khepupara and Chandpur.




Fig. 2 Relationship between period of preservation of *Hilsa ilisha* in deep-freezer and intercept values obtained from regression analysis of "total length versus head length".

Table I
Number of fish examined by month, sex and station

Month	Cox's Bazar			Chittagong			Chandpur			Khepupara		
	Female	Male	Total	Female	Male	Total	Female	Male	Total	Female	Male	Total
Sep. 1985	—	—	—	—	—	—	5	21	26	—	—	—
Oct. 1985	14	7	21	18	5	23	10	10	20	—	—	—
Nov. 1985	15	3	18	9	12	21	12	12	24	8	12	20
Dec. 1985	14	10	24	11	12	23	4	15	19	2	18	20
Jan. 1986	12	7	19	—	—	—	7	8	15	14	10	24
Feb. 1986	3	15	18	7	11	18	7	8	15	10	14	24
March 1986	13	10	23	15	10	25	10	14	24	6	18	24
Total	71	52	123	60	50	110	55	88	143	40	72	112

Table II
Interval between collection of samples and their examination

Station	Date of collection	Date of examination	Period under frozen condition	Average period of preservation
Chittagong	Sept. 15	Oct. 10	25	21 days
	Oct. 14	Oct. 29	15	
	Nov. 14	Nov. 27	13	
	Dec. 23	Jan. 25	33	
	Feb. 20	Mar. 21	29	
	Mar. 15	Mar. 26	11	
Chandpur	Sept. 25	Oct. 7	12	10 days
	Oct. 28	Nov. 12	15	
	Nov. 28	Dec. 1	4	
	Dec. 26	Dec. 27	1	
	Jan. 26	Jan. 28	2	
	Feb. 27	Mar. 23	24	
Cox's Bazar	Sept. 7	Oct. 2	25	23.6 days
	Oct. 7	Oct. 30	23	
	Nov. 7	Nov. 28	21	
	Dec. 27	Jan. 25	29	
	Jan. 7	Jan. 26	19	
	Feb. 20	Mar. 21	29	
Khepupara	Mar. 7	Mar. 26	19	0 days
	Nov. 19	Nov. 19	0	
	Dec. 23	Dec. 23	0	
	Jan. 19	Jan. 19	0	
	Feb. 19	Feb. 19	0	
	Mar. 31	Mar. 31	0	

Table III
Minimum and maximum values of morphometric characters and weight of fish at different sampling stations

		Chittagong	Chandpur	Cox's Bazar	Khepupara
Total length (TL)	Min	288	262	287	279
	Max	488	529	488	488
Depth at dorsal origin (DP)	Min	72.3	44.4	58.5	75.8
	Max	123.4	138.6	135.3	136.8
Head length (HL)	Min	69.85	40.6	68.2	63.4
	Max	117.1	122.3	114.4	115.6
Caudal peduncle length (CPL)	Min	19.5	10.8	19.6	20.2
	Max	39.1	51.2	35.45	35.2
Eye diameter (ED)	Min	9.8	10	10.4	10.08
	Max	15.8	20.3	15.4	18.2
Weight (WT)	Min	190	140	220	260
	Max	1330	1600	1420	1320

Table IV
Descriptive statistics for meristic character examined

Meristic character		Chittagong	Chandpur	Cox's Bazar	Khepupara
		n=136	n=121	n=130	n=113
Post-pelvic scutes (PS)	Min	12	11	10	12
	Max	15	15	14	15
	Mean	13.1324	13.1135	12.9813	13.1681
	Std. Dev.	0.541723	0.631246	0.507452	0.532894
			n=142	n=111	n=133
Pectoral fin rays (PF)	Min	14	13	14	13
	Max	16	15	16	16
	Mean	14.7817	14.5594	14.7929	14.5664
	Std. Dev.	0.49273	0.51214	0.533137	0.564992

Table V

Regression statistics on the relationship between depth (DP) at dorsal origin and total length (TL) with sex combined and separated by station and by month

(CTG=-Chittagong; CHA=Chandpur; CB=Cox's Bazar; KHE=Khepupara)

Month	Station	Sex Slope	Combined Intercept	n	r ²	n	Male r ²	Female r ²	n
October	CTG	0.27	0.97	23	0.42	5	0.8243	0.4049	18
	CHA	0.19	30.36	20	0.51	10	0.8029	0.2752	10
	CB	0.28	-4.97	21	0.76	7	0.9111	0.4311	14
	KHE	—	—	—	—	—	—	—	—
November	CTG	0.21	17.90	21	0.91	12	0.9258	0.9696	9
	CHA	0.26	4.00	24	0.88	12	0.9604	0.5636	12
	CB	0.19	33.18	18	0.62	3	0.6955	0.7889	15
	KHE	0.21	21.89	20	0.76	12	0.7059	0.9643	8
December	CTG	0.27	-4.66	23	0.91	12	0.9567	0.9228	11
	CHA	0.24	10.47	19	0.95	15	0.8684	0.9821	4
	CB	0.22	13.85	24	0.82	10	0.7764	0.8427	14
	KHE	0.18	30.64	20	0.70	18	0.7599	—	2
January	CTG	—	—	—	—	—	—	—	—
	CHA	0.27	-1.05	15	0.92	8	0.6872	0.8966	7
	CB	0.25	-3.55	19	0.90	7	0.9072	0.8847	12
	KHE	0.27	1.51	24	0.70	10	0.5584	0.5751	14
February	CTG	0.18	35.36	18	0.44	11	0.5728	0.0781	7
	CHA	0.32	-19.52	15	0.96	8	0.9188	0.8325	7
	CB	0.22	10.94	18	0.30	15	0.2871	0.9996	3
	KHE	0.23	13.13	24	0.84	14	0.8327	0.6341	10
March	CTG	0.23	8.50	25	0.74	10	0.0317	0.7120	15
	CHA	0.17	31.41	24	0.57	14	0.4678	0.7294	10
	CB	0.26	-4.36	23	0.80	10	0.8199	0.7670	13
	KHE	0.30	-16.83	24	0.88	18	0.8567	0.8417	6

Table VI

Regression statistics on the relationship between head length (HL) and total length (TL) by month and station

Month	Station	Slope	Intercept	r ²	n
October	CTG	0.23	5.28	0.86	23
	CHA	0.22	9.64	0.80	20
	CB	0.21	12.88	0.93	21
	KHE	—	—	—	—
November	CTG	0.23	2.56	1.00	21
	CHA	0.23	1.17	0.94	24
	CB	0.20	16.16	0.94	18
	KHE	—	—	—	—
December	CTG	0.22	6.01	0.97	23
	CHA	0.22	3.49	0.99	19
	CB	0.22	6.29	0.96	24
	KHE	0.25	-7.28	0.97	20
January	CTG	—	—	—	—
	CHA	0.23	0.82	0.95	15
	CB	0.22	4.85	0.99	19
	KHE	0.23	-0.68	0.93	24
February	CTG	0.26	-9.67	0.75	18
	CHA	0.24	-2.64	0.99	15
	CB	0.27	-8.89	0.95	18
	KHE	0.22	6.81	0.92	24
March	CTG	0.25	-7.59	0.85	25
	CHA	0.13	31.89	0.58	24
	CB	0.21	12.86	0.89	23
	KHE	0.24	-2.92	0.92	24

Table VII

Regression statistics on the relationship **between** thickness of body (TH) and total length (TL) by sex, month and station

Month	Sex	Station	Slope	Intercept	r ²	n
October	Female	CTG	0.0769	13.6226	0.2989	18
		CHA	0.0509	26.4186	0.4301	10
		CB	0.0961	8.7043	0.5021	14
		KHE	—	—	—	—
October	Male	CTG	0.2410	-1.3907	0.9558	5
		CHA	0.0646	18.0475	0.4192	10
		CB	0.1362	-9.0618	0.8224	7
		KHE	—	—	—	—
November	Female	CTG	0.1019	3.8713	0.9598	9
		CHA	0.0836	14.3803	0.4943	12
		CB	0.0802	16.2136	0.3195	15
		KHE	0.0867	11.0026	0.8268	8
November	Male	CTG	0.2140	7.8892	0.9917	12
		CHA	0.1122	6.6294	0.7296	12
		CB	-0.1036	82.0281	0.2379	3
		KHE	0.0727	13.7824	0.3010	12
December	Female	CTG	0.1385	-7.5309	0.7289	11
		CHA	0.0471	30.3289	0.6222	4
		CB	0.1369	-8.7068	0.9298	14
		KHE	—	—	—	—
December	Male	CTG	0.1049	1.8133	0.7180	12
		CHA	0.1063	7.3638	0.7477	15
		CB	0.0869	9.5123	0.7121	10
		KHE	0.0620	21.3347	0.6524	18
January	Female	CTG	—	—	—	—
		CHA	0.0907	10.4579	0.8761	7
		CB	0.1540	-16.7977	0.8970	12
		KHE	0.0727	17.7859	0.6830	14
January	Male	CTG	—	—	—	—
		CHA	0.0014	-8.1417	0.5233	8
		CB	0.1131	0.3027	0.7894	7
		KHE	0.0616	20.9932	0.4488	10
February	Female	CTG	-0.0253	60.5406	0.0220	7
		CHA	0.1789	-31.3052	0.6197	7
		CB	0.2275	-35.4532	0.8620	3
		KHE	0.0446	27.7230	0.4253	10
February	Male	CTG	0.0545	24.8258	0.1570	11
		CHA	0.1378	-8.5301	0.6512	8
		CB	0.1279	-4.4318	0.7527	15
		KHE	0.0923	9.1503	0.6555	14
March	Female	CTG	0.0412	29.5690	0.2502	15
		CHA	0.0699	21.4957	0.3496	10
		CB	0.1070	0.3080	0.7781	13
		KHE	0.0961	4.4403	0.7327	6
March	Male	CTG	0.1524	-17.4828	0.3301	10
		CHA	0.0492	17.3203	0.3831	14
		CB	0.1138	0.7238	0.7183	10
		KHE	0.1325	-7.3213	0.6199	18

Table VIII

Regression statistics on the relationship between caudal peduncle length (CPL) and total length (TL) by month and station (sex combined)

Month	Station	Slope	Intercept	r ²	n
October	CTG	0.0930	-8.5158	0.5074	23
	CHA	0.0704	0.7623	0.7683	20
	CB	0.0625	4.1248	0.6015	21
	KHE	—	—	—	—
November	CTG	0.0885	-5.0843	0.9471	21
	CHA	0.1065	-12.6249	0.7679	24
	CB	0.0651	3.1477	0.6448	18
	KHE	0.0685	1.4547	0.6703	20
December	CTG	0.0712	0.4181	0.8285	23
	CHA	0.0843	-4.8500	0.8812	19
	CB	0.0641	2.5350	0.8901	24
	KHE	0.0759	-0.9208	0.8158	20
January	CTG	—	—	—	—
	CHA	0.0493	6.8991	0.8750	15
	CB	0.0499	8.1233	0.7994	19
	KHE	0.0551	5.6477	0.4954	24
February	CTG	0.0510	8.3137	0.1712	18
	CHA	0.0503	9.4707	0.4312	15
	CB	0.0607	4.6054	0.3470	18
	KHE	0.0601	3.4518	0.6703	24
March	CTG	0.0669	1.3451	0.2788	25
	CHA	0.0426	9.4204	0.5868	24
	CB	0.0410	13.9919	0.3975	23
	KHE	0.0263	16.8228	0.2331	24

Table IX

Regression statistics on the relationship between Head Length (HL) and Eye Diameter (ED) by month and by station (sex combined)

Month	Station	Slope	Intercept	r ²	n
October	CTG	0.0738	6.0068	0.3623	23
	CHA	0.1439	-1.1495	0.3835	20
	CB	0.0749	5.7675	0.6068	21
	KHE	—	—	—	—
November	CTG	0.0922	4.2445	0.9042	21
	CHA	0.0823	5.4694	0.8512	24
	CB	0.1180	1.6617	0.7776	18
	KHE	—	—	—	—
December	CTG	0.0936	4.0142	0.8201	23
	CHA	0.0787	5.5047	0.7247	19
	CB	0.0782	5.8367	0.6687	24
	KHE	0.1119	3.3214	0.8114	20
January	CTG	—	—	—	—
	CHA	0.0623	7.2982	0.4714	15
	CB	0.0820	5.2841	0.7601	19
	KHE	0.0821	6.0749	0.2793	24
February	CTG	0.0475	8.7574	0.1699	18
	CHA	0.0794	5.0583	0.8404	15
	CB	0.0701	6.2683	0.4071	18
	KHE	0.0478	9.1752	0.3277	24
March	CTG	0.0371	9.7882	0.0800	25
	CHA	0.0774	5.8876	0.8432	24
	CB	0.0493	8.2810	0.0821	23
	KHE	0.0864	5.1045	0.6961	24

Table X

Regression statistics of the posted data on the relationship between Head Length (HL) and Total Length (TL) by stations

Station	N	Pooled months	r ²	Slope	Intercept	TL (Max; Min) in mm	Average number of days after freezing
Chittagong	44	Nov. to Feb.	0.9890	0.2268	4.5702	487; 288	21.0
Chandpur ¹	99	Nov. to Feb.	0.9732	0.2351	-0.2862	529; 278	10.0
Cox's Bazar	79	Nov. to Feb.	0.9688	0.2245	5.6538	473; 287	23.6
Khepupara ²	68	Nov. to Feb.	0.9517	0.2365	-1.2254	488; 279	0

¹Sept. data included

²No data for November

Table XI**Analysis of variance on mean values of post-pelvic scutes for the four sampling stations**Analysis of variance
One-way anova

Station	Range	Mean	In
Chittagong	12-15	13.132	136
Chandpur	11-15	13.116	121
Cox's Bazar	10-14	12.992	130
Khepupara	12-15	13.168	113
Pooled	10-15	13.100	500

Analysis of variance

Source	Sum of squares	D.F.	Mean square	F Ratio
Between	2.205	3	0.735	2.267
Within	160.795	496	0.324	
Total	163.000	499		

Table XII**Analysis of variance on mean values of pectoral fin rays for the four sampling stations**Analysis of variance
One-way anova

Station	Range	Mean	In
Chittagong	14-16	14.782	142
Chandpur	13-15	14.613	111
Cox's Bazar	14-16	14.722	1333
Khepupara	13-16	14.676	111
Pooled	13-16	14.704	497

Analysis of variance

Source	Sum of squares	D.F.	Mean square	F Ratio
Between	1.915	3	0.638	2.321
Within	135.606	493	0.275	
Total	137.52 1	496		