

42

2008

ISSN 1014-2339

**ANIMAL GENETIC
RESOURCES
INFORMATION**

**Special issue:
Scientific forum of Interlaken Conference**

**BULLETIN
D'INFORMATION
SUR LES RESSOURCES
GÉNÉTIQUES ANIMALES**

**Numéro spécial:
Forum scientifique de la Conférence d'Interlaken**

**BOLETÍN
DE INFORMACIÓN
SOBRE RECURSOS
GENÉTICOS ANIMALES**

**Número especial:
Foro científico de la Conferencia de Interlaken**



Editors - Editeurs - Editores:
S. Galal & I. Hoffmann

Viale delle Terme di Caracalla,
00153 Rome, Italy

Animal Genetic Resources Information is published under the auspices of the Food and Agriculture Organization of the United Nations (FAO). It is edited in the Animal Genetic Resources Group of the Animal Production and Health Division of FAO. It is available direct from FAO or through FAO sales agents.

ANIMAL GENETIC RESOURCES INFORMATION will be sent free of charge to those concerned with the sustainable development conservation of domestic livestock. Anyone wishing to receive it regularly should send their name and address to the Editor, at the address shown above.

AGRI can also be found in the "Library" of DAD-IS at www.fao.org/dad-is.

Le *Bulletin d'information sur les ressources génétiques animales* est publié sous les auspices de l'Organisation des Nations Unies pour l'alimentation et l'agriculture (FAO). Cette publication est éditée par le Groupe des ressources génétiques de la Division de la production et de la santé animales de la FAO. On peut se le procurer directement au siège de la FAO ou auprès des dépositaires et agents de vente des publications de l'Organisation.

LE BULLETIN D'INFORMATION SUR LES RESSOURCES GÉNÉTIQUES ANIMALES sera envoyé gratuitement aux personnes intéressées par le développement durable et la conservation du cheptel national. Les personnes souhaitant recevoir cette publication régulièrement voudront bien faire parvenir leurs nom et adresse à l'éditeur, à l'adresse susmentionnée.

AGRI peut être consulté également dans la "Bibliothèque" de DAD: www.fao.org/dad-is.

El *Boletín de información sobre recursos genéticos animales* se publica bajo los auspicios de la Organización de las Naciones Unidas para la Agricultura y la Alimentación (FAO). Se edita en el Grupo de Recursos Zoogenéticos de la Dirección de Producción y Sanidad Animal de la FAO. Se puede obtener directamente de la FAO o a través de sus agentes de venta.

EL BOLETÍN DE INFORMACIÓN SOBRE RECURSOS GENÉTICOS ANIMALES será enviado gratuitamente a quienes estén interesados en el desarrollo sostenible y la conservación del ganado doméstico. Si se desea recibirlo regularmente, se ruega comunicar nombre, apellido y dirección al editor a la dirección arriba indicada.

AGRI puede consultarse también en la "Biblioteca" de DAD-IS en: www.fao.org/dad-is.

**ANIMAL GENETIC
RESOURCES INFORMATION**


**BULLETIN
D'INFORMATION
SUR LES RESSOURCES
GÉNÉTIQUES ANIMALES**

**BOLETÍN DE
INFORMACIÓN SOBRE RECURSOS
GENÉTICOS ANIMALES**

CONTENTS

Page

Editorial	I
Introduction	1
Dynamics of livestock production systems, drivers of change and prospects for animal genetic resources	3
<i>C. Seré, A. van der Zijpp, G. Persley & E. Rege</i>	
Panellists' comments and discussion	25
Inventory, characterization and monitoring	29
<i>M. Tixier-Boichard, W. Ayalew & H. Jianlin</i>	
Panellists' comments and discussion	45
Sustainable use and genetic improvement	49
<i>C. Nimbkar, J. Gibson, M. Okeyo, P. Boettcher & J. Soelkner</i>	
Panellists' comments and discussion	66
Conservation of animal genetic resources: approaches and technologies for <i>in situ</i> and <i>ex situ</i> conservation	71
<i>J.A. Woolliams, O. Matika & J. Pattison</i>	
Panellists' comments and discussion	86
Recent publications	91
Editorial policies and procedures	97



Editorial Advisory Board (EAB) of Animal Genetic Information (AGRI)

- Editor-in-Chief: I. Hoffmann, Chief Animal Production Service, FAO
- Editor: S. Galal
- Technical Editor: C. Mosconi
- Editorial Board: L. Ollivier, Chairperson
L. Alderson
J.S. Barker
P. Bhat
J.V. Delgado Bermejo
M. Djemali
J. Hodges
K. Ramsay
E. Rege
A. Tewolde

The following is the address for each of the members of the Editorial Advisory Board.

- Irene Hoffmann, FAO, Viale delle Terme di Caracalla 1, 00153 Rome, Italy
Tel.: +33 06 5705 52796, irene.hoffmann@fao.org
- Salah Galal, Animal Production Department, Faculty of Agriculture, University of Ain Shams, PO Box 68, Hadaeq Shubra 11241, Cairo, Egypt
Tel.: +20 2 634408, sgalal@tedata.net.eg
- Louis Ollivier, INRA-Station de Génétique Quantitative et Appliquée, CRJ, Domaine de Vilvert, 78352 Jouy-en-Josas Cédex, France
Tel.: +33 1 34652190, ugenlol@dga2.jouy.inra.fr
- Lawrence Alderson, Countrywide Livestock Ltd, 6 Harnage, SY5 6EJ Shrewsbury, Shropshire, UK
Tel.: +44 195 2510030, alderson@clltd.demon.co.uk
- J. Stuart Barker, Dept. of Animal Science, University of New England, NSW 2351 Armidale, Australia
Tel.: +61 7 54358365, sbarker@une.edu.au
- Pushkar Nath Bhat, Indian Association for Animal Production, World Buffalo Trust, Flat N° 205 N° F64-C/9 Sector 40, 201 303 Noida (UP), India
Tel.: +91 11 91579627, pnbhat@bol.net.in
- Juan Vicente Delgado Bermejo, Departamento de Genética, Universidad de Córdoba, Campus de Rabanales Edificio C-5 (Gregor Mendel), 14071 Córdoba, Spain
Tel.: +34 957 218661, id1debej@lucano.uco.es
- Mnouer Djemali, Institut National Agronomique de Tunis, 43, avenue Charles Nicole, Tunis, Tunisia
Tel.: +216 1 289683/289431, djemali.mnaouer@inat.agrinet.tn
- John Hodges, Lofererfeld 16, 5730 Mittersill, Austria,
Tel.: +43 6562 5481, hodgesjohn@compuserve.com
- Keith Ramsay, Senior Livestock Specialist, Private Bag X 138, 0001 Pretoria, Gauteng, South Africa
Tel.: +27 12 3197448, KeithR@nda.agric.za
- Ed Rege, International Livestock Research Institute (ILRI), P.O. Box 30709, 00100 Nairobi, Kenya
Tel.: +254 20 4223201, e.rege@cgiar.org
- Assefaw Tewolde, Inovacion y Tecnologia/ Biotecnologia, Instituto Interamerican de Cooperacion para la Agricultura (IICA), Sede Central, Coronado, San Jose, Costa Rica, atewolde@uamac.uat.mx
- Cesare Mosconi, EAAP, Via G. Tomassetti 3, 00161 Rome, Italy
Tel.: +39 06 44202639, mosconi@eaap.org

Editorial - The *Global Plan of Action*

The International Technical Conference on Animal Genetic Resources for Food and Agriculture, which took place in Interlaken, Switzerland, from 3 to 7 September 2007, was attended by delegations from 109 countries and 42 intergovernmental and non-governmental organizations. The Conference was a resounding success. It welcomed *The State of the World's Animal Genetic Resources for Food and Agriculture*, and formally adopted the *Global Plan of Action for Animal Genetic Resources* through the *Interlaken Declaration on Animal Genetic Resources*. The *Global Plan of Action* is reviewed in the Recent Publications section of this issue.

In addition to the negotiations, the Conference provided opportunity for learning and networking, and brought together technical people and diplomats. At the beginning of the Conference, there was an all-day Scientific Forum, highlighting and discussing the importance of use, development, characterization, monitoring and conservation of animal genetic resources (AnGR). The Scientific Forum, and the many well attended side events, contributed to the positive spirit among the delegates and the eventual adoption of the *Global Plan of Action*. The accompanying technical events also set the stage for the further elaboration of the technical programme of work to support the implementation of the *Global Plan of Action*.

The *Global Plan of Action* includes several Strategic Priorities aiming to address the current low level of capacity in all fields of AnGR management, through partnerships and cooperation for enhanced educational and training opportunities, technical cooperation, technology transfer and exchange of experience. The scientific and donor communities have important roles to play in this regard. Although research on AnGR management has increased in the past few years, linkages between groups and collaborative networks of researchers and stakeholders need to be strengthened, and standardization of approaches

needs further enhancement. Specifically, there is urgent need to strengthen research and development of methods for characterization, and breed evaluation, valuation and comparison, as well as *in situ* and *ex situ* conservation methods and technologies. Technical standards, guidelines and protocols for phenotypic and molecular characterization, particularly for the assessment of comparative breed performance in different production environments, and for *in situ* and *ex situ* conservation, should be developed. Cooperation and coordination in the fields of conservation, sustainable use and development of AnGR should be strengthened at national, regional and international levels, including through *ex situ* backup systems for protection against emergencies or disasters. Because of countries' shared responsibilities for transboundary breeds, several actions in the *Global Plan of Action* refer to the international collaboration needed for the characterization, use and development, and conservation of these breeds.

Research outcomes should assist National Coordinators for AnGR to update their breed data in DAD-IS and provide them with relevant information. The importance given to research as a basis for policy development and for decision-making in the field of AnGR conservation implies an important role for the scientific community, in particular the Consultative Group on International Agricultural Research (CGIAR) network.

Governments must now demonstrate the sustained political will to implement the *Global Plan of Action*, and resources will have to be found, nationally and internationally. Success will depend on farsighted cooperation among many stakeholders. Governments, international organizations, the scientific community, donors, civil society organizations and the private sector all have important roles to play.

The Editors

Éditorial - Le *Plan d'Action Mondial*

La Conférence Technique Internationale sur les Ressources Zoogénétiques pour l'Alimentation et l'Agriculture a eu lieu à Interlaken, en Suisse, du 3 au 7 septembre 2007 et a réuni des délégations de 109 pays et 42 organisations intergouvernementales et non-gouvernementales. La conférence a eu un succès retentissant et a donné la bienvenue à *L'État des Ressources Zoogénétiques pour l'Alimentation et l'Agriculture dans le Monde* en adoptant formellement le *Plan d'Action Mondial pour les Ressources Zoogénétiques* à travers la *Déclaration d'Interlaken sur les Ressources Zoogénétiques*. Le *Plan d'Action Mondial* se trouve aussi dans la section consacrée aux Publications Récentes de ce numéro.

Autre aux négociations, la Conférence a offert la possibilité d'apprendre et de mettre en place des réseaux, en réunissant des techniciens et diplomates. Au début de la Conférence, un Forum Scientifique d'une journée a été organisé pour discuter de l'importance de l'utilisation, du développement, de la caractérisation, du suivi et de la conservation des ressources génétiques animales (AnGR). Le Forum Scientifique, ainsi que les différentes réunions organisées en parallèle, ont contribué à la formation d'un esprit positif parmi les délégués pour l'adoption éventuelle du *Plan d'Action Mondial*.

Le *Plan d'Action Mondial* inclut différentes Stratégies Prioritaires qui ont comme objectif de résoudre le manque de capacité de gestion de AnGR actuellement dans tous les domaines, à travers des associations pour améliorer les opportunités de formation et éducation ainsi que la coopération technique, le transfert de technologie et l'échange d'expérience. La Communauté scientifique et les donateurs jouent un rôle important dans ce sens. Malgré la recherche sur la gestion des AnGR ait augmenté au cours des dernières années, les relations entre groupes et réseaux de collaboration des chercheurs et éleveurs doivent être renforcées, tout comme il est nécessaire d'améliorer ultérieurement la normalisation des différentes approches.

En particulier, il existe le besoin urgent de renforcer la recherche et le développement des

méthodes de caractérisation et d'évaluation, de valorisation et comparaison des races, ainsi que les méthodes et technologies de conservation *in situ* et *ex situ*. On devra développer des standards techniques, des normes et des protocoles pour la caractérisation phénotypes et moléculaire, et en particulier pour la valorisation comparative des rendements des races dans les différents milieux de production, et pour la conservation *in situ* et *ex situ*. La coopération et la coordination dans le domaine de la conservation, l'utilisation durable et le développement de AnGR devront être renforcés au niveau nationale, régionale et internationale, y compris les systèmes de soutien *ex situ* pour la protection en cas d'émergence ou désastre. Etant donné les responsabilités conjointes entre les pays dans les cas des races transfrontalières, une série d'actions ont été mises au point dans le *Plan d'Action Mondial* soulignant la nécessité d'une collaboration internationale pour la caractérisation, l'utilisation et le développement.

Les résultats de la recherche devront aider les Coordinateurs Nationaux de AnGR dans la mise à jour des données sur les races dans DAD-IS, tout en fournissant une information importante. L'importance donnée à la recherche comme base pour la politique de développement et la législation en matière de conservation de AnGR, comporte un rôle important de la communauté scientifique, surtout du réseau du Groupe Consultatif sur la Recherche Agricole Internationale (CGIAR).

Les gouvernements devront maintenant démontrer leur politique durable pour réaliser le *Plan d'Action Mondial* et pour cela il sera nécessaire de trouver les ressources financières au niveau national et international. Le succès dépendra de la vision de future entre éleveurs pour la coopération. Les gouvernements, les organisations internationales, la communauté scientifique, les donateurs, les organisations civiles et le secteur privé devront jouer un rôle important.

Les Editeurs

Editorial - El Plan de Acción Mundial

La Conferencia Técnica Internacional sobre los Recursos Zoogenéticos para la Agricultura y la Alimentación que tuvo lugar en Interlaken, Suiza, del 3 al 7 de septiembre de 2007, reunió delegaciones de 109 países y 42 organizaciones intergubernamentales y no gubernamentales. La Conferencia tuvo un éxito resonante. Dio la bienvenida a *La Situación de los Recursos Zoogenéticos Mundiales para la Alimentación y la Agricultura* y adoptó formalmente el *Plan de Acción Mundial para los Recursos Zoogenéticos* a través de la *Declaración de Interlaken sobre Recursos Zoogenéticos*. El *Plan de Acción Mundial* queda también reflejado en la sección dedicada a Publicaciones Recientes de este número.

Además de la negociación, la Conferencia ofreció la oportunidad de aprender y establecer redes, al poner en contacto técnicos y diplomáticos. Al principio de la Conferencia hubo un Forum Científico de un día en el que se subrayaron y discutieron la importancia del uso, desarrollo, caracterización, seguimiento y conservación de los recursos zoogenéticos (AnGR). El Forum Científico, así como los distintos eventos organizados paralelamente, contribuyeron a la formación de un espíritu positivo entre los delegados en vistas a la adopción del *Plan de Acción Mundial*. Los eventos técnicos paralelos también contribuyeron a poner las bases para una futura elaboración del programa técnico de trabajo para ayudar en la realización del *Plan de Acción Mundial*.

El *Plan de Acción Mundial* incluye diversas Estrategias Prioritarias que tienen como objetivo resolver el escaso nivel actual de capacidad de gestión de los AnGR en todos los terrenos, a través de la asociación y cooperación para mejorar las oportunidades de formación y educación así como la cooperación técnica, la transferencia de tecnología y el intercambio de experiencias. La comunidad científica y los donantes desempeñan un papel importante en este sentido. A pesar de que la investigación sobre la gestión de los AnGR ha aumentado en los últimos años, las relaciones entre grupos y redes de colaboración de investigadores y ganaderos necesitan ser reforzadas, así como es

necesario mejorar ulteriormente la normalización de los distintos enfoques. En particular, existe la necesidad urgente de reforzar la investigación y el desarrollo de los métodos de caracterización y de evaluación, valoración y comparación de las razas, así como los métodos y tecnologías de conservación *in situ* como *ex situ*. Se deberán desarrollar estándares técnicos, normas y protocolos para la caracterización fenotípica y molecular, y en particular para la valoración comparativa de los rendimientos de las razas en distintos medios de producción y para la conservación *in situ* y *ex situ*. La cooperación y coordinación en el campo de la conservación, el uso sostenible y el desarrollo de AnGR deberán ser reforzadas a nivel nacional, regional e internacional, incluyendo los sistemas de apoyo *ex situ* para la protección en caso de emergencias o desastres. Dadas las responsabilidades compartidas entre países en los casos de razas transfronterizas, se han determinado una serie de acciones en el *Plan de Acción Mundial* referidas a la necesidad de una colaboración internacional para la caracterización, el uso y el desarrollo y conservación de estas razas.

Los resultados de la investigación deberán ayudar a los Coordinadores Nacionales de AnGR en la actualización de datos sobre sus razas en DAD-IS, además de proporcionar información importante. La importancia dada a la investigación como base para la política de desarrollo y para la legislación en materia de conservación de AnGR, implica un papel importante de la comunidad científica, en particular la red del Grupo de Consulta sobre Investigación Agrícola Internacional (CGIAR).

Los gobiernos deberán ahora demostrar su política sostenible para llevar a cabo el *Plan de Acción Mundial* y, para ello, será necesario encontrar recursos a nivel nacional e internacional. El éxito dependerá de la visión de futuro de la cooperación entre ganaderos. Los gobiernos, organizaciones internacionales, la comunidad científica, los donantes, las organizaciones civiles y el sector privado, tendrán que jugar un papel importante.

Los Editores

Introduction

The Scientific Forum on Animal Genetic Resources, organized as part of the FAO International Technical Conference on Animal Genetic Resources for Food and Agriculture, was held in Interlaken, Switzerland, on 3 September 2007. The Forum was chaired by Mr Fritz Schneider of the Swiss College of Agriculture.

The Forum provided an opportunity for informal discussion and clarification of key issues in advance of the Conference negotiating sessions. The Forum was structured around the presentation of four papers prepared by teams of leading experts, which highlighted the latest scientific developments in the management of animal genetic resources, following the main themes dealt with in Part 4 (the state of the art in the management of animal genetic resources) of *The State of the World's Animal Genetic Resources for Food and Agriculture*:

- The dynamics of livestock production systems, and the drivers of change.
- Inventory, characterization and monitoring.
- Sustainable use and genetic improvement.
- Conservation of animal genetic resources.

Each of the papers was presented by the lead author. Each presentation was followed by a panel discussion involving a range of stakeholders, after which the meeting was opened for general discussion and interventions from the floor.

The Forum concluded with a round-up session entitled "*Animal science meets society*" which sought to address the question: how can the wider society be involved in the implementation of the *Global Plan of Action*, and in research for development? During the ensuing discussion it was recognized that scientific developments had to be complemented by the mobilization of financial resources and the development of policy and legal frameworks that support the sustainable management of animal genetic resources. The need for greater cooperation at all levels and between different disciplines was emphasized.

Dynamics of livestock production systems, drivers of change and prospects for animal genetic resources

C. Seré¹, A. van der Zijpp², G. Persley¹ & E. Rege¹

¹International Livestock Research Institute, P.O. Box 30709, Nairobi 00100, Kenya

²Animal Production Systems Group, Department of Animal Sciences, Wageningen University, P.O. Box 338, 6700 AH Wageningen, The Netherlands

Summary

This overview analyses the key drivers of change in the global livestock sector and assesses how they are influencing current trends and future prospects in the world's diverse livestock production systems and market chains; and what are their consequent impacts on the management of animal genetic resources for food and agriculture. The trends are occurring in both developing and industrialized countries, but the responses are different. In the developing world, the trends are affecting the ability of livestock to contribute to improving livelihoods and reducing poverty as well as the use of natural resources. In the industrialized world, the narrowing animal genetic resource base in industrial livestock production systems raises the need to maintain a broader range of animal genetic resources to be able to deal with future uncertainties, such as climate change and zoonotic diseases.

This chapter discusses:

- What are the global drivers of change for livestock systems? Economic development and globalization; changing market demands and the "livestock revolution"; environmental impacts including climate change; and science and technology trends.
- How are the livestock production systems responding to the global drivers of change? Trends in the three main livestock production systems (industrial, crop-livestock and pastoral systems); the range and rate of changes occurring in different systems and how these affect animal genetic resources. The implications are that breeds cannot adapt in time to meet new circumstances. Hence new strategies and interventions are necessary to improve the management of animal genetic resources in situations where these genetic resources are most at risk.

- What are the implications for animal genetic resources diversity and for future prospects of their use? - Industrial livestock production systems are expected to have a limited demand for biodiversity, while crop-livestock and pastoral systems will rely on biodiversity to produce genotypes of improved productivity under changing environmental and socio-economic conditions. All systems will rely on biodiversity, albeit to varying degrees, to cope with expected climate change.
- What immediate steps are possible to improve animal genetic resources characterization, use and conservation? Appropriate institutional and policy frameworks are required to improve animal genetic resources management and these issues are being addressed at national and intergovernmental levels, in a process led by FAO to promote greater international collaboration on animal genetic resources. Based on an analysis of the current situation, the continuing loss of indigenous breeds and new developments in science and technology, there are several complementary actions that can begin to improve the management of animal genetic resources and maintain future options in an uncertain world.

These are summarized here as:

- a. "Keep it on the hoof" - Encouraging the continuing sustainable use of traditional breeds and *in situ* conservation by providing market-driven incentives, public policy and

This paper has benefited from inputs from several reviewers and other contributors, and we thank all for their thoughtful insights. We acknowledge the contributions of our colleagues at FAO, particularly Irene Hoffmann, Dafydd Pilling and Henning Steinfeld, and at the International Livestock Research Institute (ILRI): Ade Freeman, Mario Herrero, Olivier Hanotte, Steve Kemp, Sandy McClintock, Sara McClintock, Margaret MacDonald-Levy, Susan MacMillan, Grace Ndungu, An Notenbaert, Mwai Okeyo and Robin Reid.

other support to enable livestock keepers to maintain genetic diversity in their livestock populations.

- b. "Move it or lose it" – Enabling access to and the safe movement of animal genetic resources within and between countries, regions and continents is a key factor in use, development and conservation of animal genetic resources globally.
- c. "Match breeds to environments" – Understanding the match between livestock populations, breeds and genes with the physical, biological and economic landscape. This "landscape livestock genomics" approach offers the means to predict the genotypes most appropriate to a given environment and, in the longer term, to understand the genetic basis of adaptation of the genotype to the environment.
- d. "Put some in the bank" -- New technologies make *ex situ*, *in vitro* conservation of animal genetic resources feasible for critical situations and are a way to provide long-term insurance against future shocks.

The multiple values, functions and consequences of livestock production systems and their rapid rate of change lead to divergent interests within and between countries. Conversely, the uncertainty about the implications of rapid, multifaceted global change for each livestock production system and the resulting future changes in the required genetic make-up of animal genetic resources make collective action to tackle conservation of animal genetic resources a long-term, global public good. Conserving animal genetic resources will not by itself solve these problems, but it is an important first step towards maintaining future options.

Advances in science and the technology, in areas such as reproductive technology, genomics and spatial analysis, as well as progress in conceptualization of global public good production for the future management of animal genetic resources, should enable the international community to address both the short- and long-term challenges in innovative ways.

Résumé

Ce résumé analyse les facteurs clés qui ont subi des changements dans le secteur élevage et propose une évaluation de l'influence qu'ils ont eu sur la situation actuelle et les perspectives futures dans les différents systèmes d'élevage et de marché au

niveau mondial. On analyse également les impacts sur la gestion des ressources génétiques animales pour l'alimentation et l'agriculture. Cette tendance se retrouve aussi bien dans les pays industrialisés que dans ceux en développement, mais les réponses sont différentes. Dans les pays en développement ces tendances ont une influence directe sur la capacité que présente l'élevage à contribuer à l'amélioration de la qualité de vie et à la réduction de la pauvreté, ainsi qu'à l'utilisation des ressources naturelles. Dans le monde industrialisé la proximité de la base des ressources génétiques animales avec les systèmes de production d'élevage au niveau industriel ont porté au besoin de conserver une plus grande gamme des ressources génétiques animales pour faire face aux incertitudes futures telles que le changement climatique et les zoonoses.

Dans l'article on discute de:

- Quels sont les principaux facteurs de changement dans les systèmes d'élevage?
- Comment répondent les systèmes de production d'élevage aux facteurs de changement au niveau mondial?
- Quelles sont les implications sur la diversité des ressources génétiques animales et pour les perspectives d'utilisation futures?
- Quels sont les démarches immédiates qui permettront une amélioration de la caractérisation des ressources génétiques animales, leur utilisation et conservation?

D'après une récente analyse de la situation actuelle, de la perte continue de races indigènes et du nouveau développement de la science et de la technologie, il existe différentes actions complémentaires qui pourraient aider à améliorer la gestion des ressources génétiques animales et conserver des options pour le futur dans un monde plein d'incertitude.

Ces actions peuvent se résumer comme il suit:

- Encourager l'utilisation durable des races traditionnelles.
- Permettre l'accès et la vente de ressources génétiques animales dans et entre pays.
- Compréhension du rapport entre élevage, races et gènes avec le milieu physique, biologique et économique.
- La formation de stock comme assurance future.

L'incertitude sur les implications des changements rapides sur chacun des systèmes de production animale et les changements futurs que cela entraîne en terme de demande de ressources génétiques animales, requière d'une action collective pour faire face à la conservation des ressources génétiques animales en tant que bien

public. La conservation des ressources génétiques animales en soi ne résoudra pas les problèmes mais il s'agit d'un pas important pour conserver les options futures.

Resumen

Este resumen analiza los factores clave que han cambiado en el sector ganadero y hace una evaluación de cómo han influenciado la corriente actual y las perspectivas futuras en los distintos sistemas de producción ganadera y mercados en el mundo. También se analizan los consiguientes impactos sobre la gestión de los recursos zoogenéticos para la alimentación y la agricultura. La tendencia se da tanto en países industrializados como en vía de desarrollo pero las respuestas son distintas. En los países en vía de desarrollo estas tendencias están afectando la capacidad ganadera para contribuir a la mejora de la calidad de la vida y reducción de la pobreza, así como la utilización de los recursos naturales. En el mundo industrializado la proximidad de la base de recursos zoogenéticos con los sistemas de producción ganadera industrial plantean la necesidad de mantener un mayor rango de recursos zoogenéticos para hacer frente a las incertidumbres futuras, tales como el cambio climático y las zoonosis.

En este capítulo se discute:

- ¿Cuáles son los principales factores de cambio en los sistemas ganaderos?
- ¿Cómo responden los sistemas de producción ganadera a los factores de cambio a nivel mundial?
- ¿Cuáles son las implicaciones para la diversidad de recursos zoogenéticos y para las perspectivas futuras de su utilización?
- ¿Cuáles son los pasos inmediatos que puedan permitir la mejora de la caracterización de los recursos zoogenéticos, su utilización y conservación?

Sobre la base de un reciente análisis de la situación actual, la pérdida de razas indígenas y el nuevo desarrollo de la ciencia y la tecnología, existen distintas acciones complementarias que pueden empezar a ayudar a mejorar la gestión de los recursos zoogenéticos y mantener opciones futuras en un mundo lleno de incertidumbres.

Tales acciones se resumen así:

- Fomentar la continua utilización sostenible de razas tradicionales.
- Permitir el acceso y movimiento para venta de recursos zoogenéticos dentro y entre países.

- Conocer la relación entre poblaciones ganaderas, razas y genes con el entorno físico, biológico y económico.
- Conservar stocks para hacer frente a incertidumbres futuras.

La incertidumbre sobre las implicaciones de cambios rápidos, multifacéticos y globales para cada sistema de producción ganadera y los consiguientes cambios futuros en la demanda de recursos zoogenéticos requieren una acción colectiva para hacer frente a la conservación de recursos zoogenéticos a largo plazo como bien público mundial. La conservación de los recursos zoogenéticos por sí sola no resolverá los problemas pero es un paso importante para mantener las opciones futuras.

Keywords: *Global livestock sector, Livestock production systems, Market chains, Environmental effects, Climate change, Management, Sustainable use.*

Introduction

This overview paper analyses the key drivers of change in the global livestock sector and assesses how they are influencing current trends and future prospects in the world's diverse livestock production systems and market chains; and what are their consequent impacts on the management of animal genetic resources for food and agriculture. The trends are occurring in both developing and industrialized countries, but the responses are different. In the developing world, the trends are affecting the ability of livestock to contribute to improving livelihoods and reducing poverty as well as the use of natural resources. In the industrialized world, the narrowing animal genetic resource base in industrial livestock production systems raises the need to maintain a broader range of animal genetic resources to be able to deal with future uncertainties, such as climate change and zoonotic diseases.

The range of livestock covered here are domesticated species, particularly the five major economic species (cattle, sheep, goats, chickens and pigs). There are no detailed figures yet to link specific breeds with specific production systems. We are tackling the problems from a production system angle. Throughout the paper, and based on the findings of *The State of the World's Animal Genetic Resources for Food and Agriculture*, we use the approximation that commercial breeds, as a subgroup of international transboundary breeds, are used in intensive, high-external input livestock

production systems (termed “*industrial systems*”), and that local breeds are the basis in most extensive and low-external input systems. These are called here “*pastoral systems*” and “*crop-livestock systems*”, respectively.

This paper covers four main areas:

- What are the global drivers of change for livestock systems?
- How are the three main livestock production systems (industrial, crop-livestock and pastoral systems) responding to the global drivers of change, and what are the implications of the range and rate of changes for the management of animal genetic resources in these systems?
- What are the implications for animal genetic resources diversity and future prospects of their use?
- What immediate steps are possible to improve animal genetic resources characterization, use and conservation?

Drivers of change in global livestock systems

Economic development and globalization

Livestock production is a complex and heterogeneous part of global agriculture. It ranges from highly automated, intensive large-scale production of pigs and poultry and, to a lesser degree, cattle, to small-scale, largely scavenging production of backyard pigs and chicken. Domestication of livestock started several millennia ago and humans have shaped the genetic make-up of domesticated animals to respond to human needs in different production environments.

This genetic make-up of livestock that resulted from this long-term process has been put under stress by fast-paced changes over the past few decades, across the entire range of biophysical, social and economic contexts in which humans keep animals. These changes can be subsumed under terms of economic development and globalization. These are themselves largely driven by technical progress, plus the global exchange of knowledge and products. These trends are also characterized by unequal access to natural resources, financing, markets, technology and personal mobility.

Since 1945, the world has seen an unprecedented economic growth, starting in the industrialized economies (countries of the

Organisation for Economic Co-operation and Development [OECD]) and expanding into the rest of the world over the past two decades. The latter is epitomized by the economic growth path of China. A number of developing countries, mainly in Asia and Latin America, have undergone major transformations associated with significant growth in their economies and increases in per capita incomes.

The socio-economic indicators for selected countries are given in table 1. The following inferences can be drawn from the data:

- The contribution of livestock to agricultural gross domestic product (GDP) demonstrates the significance of the livestock sector in many economies (providing value addition); this occurs even in countries that are experiencing rapid economic growth (India and China) and/or have a growing share of industrial livestock systems (China, Brazil and Argentina).
- The key demand drivers of GDP growth and urbanization point towards growing demand for livestock products across all regions in the developing world. This “*livestock revolution*” is discussed further below.
- The trends in foreign direct investment (FDI) show that increases in FDI are concentrated in a few countries (China and India). These countries are ones in which the industrialization of livestock production has been rising sharply. Some other countries in Africa (e.g. Kenya and Botswana) have also recorded significant increases in FDI over the past decade, although from a lower base.

Economic development has led to important changes in the spatial distribution of the world’s population, leading to a rapid process of urbanization in the developing world. At the same time, breakthroughs in medical research and their applications have led to dramatic increases of the human population in developing countries. In the industrialized world, population growth rates have declined in the last decades as social security, female employment in labour-scarce economies and cultural/social changes have led to declining birth rates and gradually aging populations. In terms of consumer demand, there is more demand for “*fast food*” and processed animal products. Food safety requirements are becoming increasingly stringent, due to disease problems such as bovine spongiform encephalopathy (BSE) associated with processed animal products. A similar trend is occurring in developing countries, although currently limited to the affluent urban class.

Table 1. Socio-economic indicators for selected countries.

	Contribution of livestock to agricultural GDP (in %)		GDP growth (annual change) ^a - (in %)				Urban population ^b			FDI ^c		
	1990-1995 average	2000-2005 average	1990	1995	2000	2005	1990	2004	1990-2004 average annual growth (in %)	1997-1999	2000-2002	2003-2005
							population (in %)	population (in %)		Annual average in US\$ million		
Sub-Saharan Africa												
Botswana	85.0	82.1	6.8	4.5	8.3	6.2	42	52	3.0	77	161	363
Kenya	42.5	44.5	4.1	4.3	0.6	5.8	25	40	6.1	15	48	50
South Africa	46.1	44.0	-0.3	3.1	4.2	5.1	49	57	3.0	1 955	2 991	2 581
Latin America and the Caribbean												
Argentina	45.9	36.5	-1.3	-2.8	-0.8	9.2	87	90	1.4	13 480	4 911	3 552
Brazil	41.8	44.4	-4.2	4.2	4.3	2.9	75	84	2.3	26 713	23 942	14 501
Peru	36.0	33.1	-5.1	8.6	3.0	6.4	69	74	2.2	1 908	1 370	1 890
East Asia and Pacific												
Cambodia	20.5	20.1	1.1	6.5	8.4	13.4	13	19	5.5	226	148	198
China	26.9	24.6	3.8	10.9	8.4	10.4	27	40	3.6	42 247	43 983	60 380
Viet Nam	16.7	18.0	5.0	9.5	6.8	8.4	20	26	3.4	1 768	1 333	1 671
South Asia												
India	26.51	30.75	6.0	7.6	5.3	9.2	26	29	2.5	2 794	4 894	5 552
Pakistan	49.1	53.5	4.5	5.0	4.3	8.0	31	34	3.3	585	505	

Sources:

^aIMF (2007).^bWorld Bank (2006).^cUnited Nations (2007).

Another key driver of change that is leading towards larger-scale, cereal-based animal production systems around the world has been the rise in labour costs in the industrialized economies and in some parts of the developing world, as a result of economic growth and rising incomes.

Changing economic policy associated with rapid economic growth in parts of the developing world (e.g. Asian “tiger” economies) has changed the investment climate in emerging economies and led to massive inflows of FDI. Similarly, labour migration from developing to industrialized economies has generated capital flows back to developing countries, which are often larger than official development assistance. Capital investments from outside the farming community, for example in the feed industry and livestock production chains in Southeast Asia, are also influencing changes in livestock production systems.

The effects of globalization and growing incomes have by no means been evenly distributed within or between countries. In the context of rapid population growth, many countries and social and ethnic groups within countries have not participated in the growth process. Large numbers of poor people, particularly in rural areas, have been left behind or adversely affected by the changes. For example, such communities may actually suffer from loss of access to natural resources, bear the brunt of environmental impacts and be characterized by the breakdown of traditional social and economic ties and values, without a better (or at least viable) alternative. Also, local breeds of animals are often not competitive in this changing world.

These inequalities pose a major challenge for the global community, which has responded by setting the Millennium Development Goals (MDGs), a UN-driven process to address several core problems facing the world. The MDGs include a commitment to halve the numbers of people living in poverty by 2015, as well as setting several other key development targets, including protecting the environment and conserving biodiversity. The sustainable use and conservation of the world’s animal genetic resources for food and agriculture supports the Millennium Development Goals 1 and 7, and is also covered by the Convention on Biological Diversity (CBD).

Market demand for livestock products – the “livestock revolution”

Growing demand for animal products – as well as higher standards to improve the quality and safety of the products – and more processed animal products have substantial consequences for the evolution of livestock production systems. Overall, the processes of economic development, population growth, urbanization and changing patterns of consumption have led to a dramatic increase in the consumption of animal products in the developing world, a process that has been termed the “livestock revolution”. FAO data suggest that this trend is expected to continue for several decades because of the strong direct correlation between rising income and increasing animal product consumption.

Figure 1 shows the expected percentage changes in per capita consumption of selected food commodities in developing and industrialized countries between 2001 and 2030, providing evidence of the “livestock revolution” occurring in the developing world. There are large differences between the projected per capita growth rates in consumption of livestock products (meat and milk) between developing and industrialized countries. There are also marked differences in the per capita growth rates of the different products in developing countries, with meat and milk being the highest, followed by oil seeds. Growth rates for cereal consumption as human food are stagnating everywhere, but increasing for other uses, especially for animal feed and biofuels.

The consumption of milk and meat per capita are shown in figures 2 and 3, respectively. These data illustrate substantial differences in current consumption of meat and milk between industrialized and developing countries; the rates of growth in consumption are higher in the developing world. This trend is part of the “livestock revolution” and is the result of increased demand and increased incomes, economic growth and urbanization in developing countries. Consumption per capita of milk and meat is currently between two and four times higher in industrialized countries than in the developing world but, in absolute terms, demand is higher in the developing world.

The growing demand for animal products in the developing world is associated with the changes in production location, facilitated by the increasing ease of transporting feed and animal products around the world. Animal products were previously produced close to where the consumers live. Increasingly, livestock production now takes




Figure 1. Expected percentage changes in per capita consumption of selected food commodities in developing and industrialized countries, 2001-2030.

Source: adapted from IAASTD (2007).




Figure 2. Milk consumption per capita to 2050 (kg/person).

Source: adapted from IAASTD (2007).




Figure 3. Meat consumption per capita to 2050 (kg/person).
Source: adapted from IAASTD (2007).

place close to the locations with good access to feed, either in feed production areas or ports. The animal products are then transported to markets. This trend is changing the competitiveness of diverse livestock production systems worldwide, with more animal products being produced in lower cost economies (mainly in industrial and crop-livestock systems) and traded in domestic, regional and international markets.

At the same time, large numbers of poor people depend on livestock production for their livelihoods and, for some of them, livestock offer a pathway out of poverty. These smallholders and pastoralists frequently compete for markets with the commercial sector, which is producing animal products in industrial systems worldwide. Smallholders and pastoralists together with their traditional breeds are increasingly being pushed out by the industrial systems coming into the developing world. Hence there is pressure for smallholders and pastoralists to replace their traditional breeds with more productive but less resilient breeds in order to be able to compete in the expanding livestock markets in the developing world.

Technological developments associated with international transport, partially related to the increased access to capital and the opening of many economies, have dramatically increased the role of

international trade in animal products. The expansion of international trade in animal products has brought to the fore the need to establish more stringent animal health and food safety standards, in order to manage the risks to the domestic sector of individual countries and to protect consumers. These health and food safety requirements have been driven by the growing problems of animal diseases, including zoonoses. These disease risks are linked to a number of factors including increasing stock numbers, the intimate cohabitation of poor families with their animals and the increased global movement of animals and animal products.

Domestic markets, including the informal livestock product markets, handle the largest share of the livestock products consumed in developing countries. However, in urban areas, the modern food retail sector is also growing rapidly, and imposing specific requirements in terms of quality assurance and homogeneity of the products (of national and international origin). The term "supermarket revolution" has been coined for these processes. These two marketing systems require markedly different food safety and biosecurity standards, affecting livestock production systems supplying these markets.

Table 2 shows that the share of supermarkets in food retailing has been increasing over the past two decades in much of the developing world. If current trends in expanding urban populations continue, the share of supermarkets in the urban food retail sector in the developing world will increase to levels that they are now in the industrialized economies (i.e. about 80 percent of the total food retail sector). The changing set of actors implied by the supermarket revolution and the growing importance of agribusiness in food retailing will have important implications for poor farmers.

The coexistence of three markets for animal products in the developing world (the traditional, frequently informal markets, the growing formal (super)markets for the urban middle classes and the regional/international export markets) poses particularly daunting challenges for policy-makers in pursuing mutually compatible policies of:

1. protecting livelihoods among the smallholder livestock keepers and pastoralists;
2. supporting efficient markets for the urban population; and

3. encouraging active engagement of livestock producers and their traditional breeds in the regional and global livestock markets.

The livestock product markets in industrialized countries are evolving along quite different paths. Besides consuming relatively inexpensive livestock products from large-scale industrial systems, there is increasing demand for niche products, frequently linked with certification of origin, often produced in traditional ways or with specific breeds, by “*organic agriculture*”, and/or with particular concern for animal welfare.

Animal welfare is an increasing area of concern, especially in markets in industrialized countries. These concerns include caring for animals in all types of production systems. There is particular criticism of intensive housing systems for animals (e.g. chickens, pigs, dairy cows). This is leading to more animal friendly housing systems such as group housing of sows; and free range hens as alternatives for the caging for laying hens. Some consumers in industrialized countries are prepared to pay a premium for animal products coming from

Table 2. Trends in share of supermarkets in total food retail for selected countries.

Waves of diffusion and average market share	Country	Year	Supermarket share in food retail (%)
Industrialized country example	United States of America	2005	80
First wave of developing countries (10–20% market share around 1990)	Argentina	2002	60
	Brazil	2002	75
	Taiwan Province of China	2003	55
	Czech Republic	2003	55
	Costa Rica	2001	50
	Chile	2001	50
	Republic of Korea	2003	50
	Philippines	2003	50
	Thailand	2003	50
Second wave of developing countries (5–10% market share around 1990)	South Africa	2001	55
	Mexico	2003	56
	Ecuador	2003	40
	Colombia	2003	47
	Guatemala	2002	36
Third wave of developing countries (Virtually zero market share around 1990)	Indonesia	2001	30
	Bulgaria	2003	25
	Kenya ^a	2004	20
	Nicaragua	2006	20
	China*	2004	30
	India	2007	9

^aShare of urban food retail.

Source: Reardon, Henson and Berdegué (2007).

such production systems that take account of animal welfare concerns. Animal welfare concerns are highly culture-specific and, while important in some societies, others consider them to be non-tariff trade barriers. Some of these trends will dictate breeds and breeding practices – for example, performance under range conditions and “broodiness” of hens will be important attributes for the niche markets.

In the industrialized countries, hobby farming has become a popular activity, using relatively small land areas for limited numbers of livestock such as sheep, goats, horses and cattle. For *in situ* conservation of species and breeds within species, these part-time farmers are important contributors.

Environmental effects of livestock production

The rapid population growth and the growing consumption of goods and services by people whose incomes are growing puts pressure on natural resources and the environment. Livestock production, under certain conditions, is driving degradation processes and is at the same time affected by them. Increasing land use for food crops and crops for biofuels is increasing the pressure on rangelands and other open access or community managed resources. This affects the viability of the low-input production systems, the sustainable use of traditional breeds and thus the livelihoods of pastoralists and smallholders.

At the same time, the rapid growth of large-scale, intensive animal production units puts a serious constraint on the capacity of the environment to deal with carbon dioxide and methane output, nutrient loading in certain areas, effluent into rivers and seas, loss of biodiversity because of land clearing to grow feeds (for example, soybeans in Latin America) and other environmental impacts.

The recent FAO (2006) report *Livestock's long shadow: environmental issues and options* focused on the effects of livestock on the environment. The “long shadow” refers to the negative effects of the livestock food chain on almost all aspects of the environment; livestock production is associated with carbon dioxide, methane and nitrous oxide emissions, water depletion, soil erosion, soil fertility, damage to plants, loss of biodiversity and competition with wildlife.

As population and living standards grow, natural resources become a limiting factor. Particularly in marginal zones for rangeland-based animal production (pastoral systems), alternative

land uses such as provision of opportunities for carbon sequestration through trees or wildlife conservation may become increasingly competitive with livestock production. On the other hand, livestock production in pastoral systems can be complementary to other services – for example, livestock production provides a means to maintain shrub/rangeland systems, with grazing reducing the risk of fire in extensive rangelands and providing other ecological services.

Climate change effects

The relationship between livestock production and climate change works in both directions. On the one hand, livestock contributes significantly to climate change via carbon dioxide, methane and nitrous oxide production (calculated in FAO (2006) at 18 percent of the total global greenhouse gas emissions from human sources). On the other hand, climate change will have important effects on farming systems and on the role of livestock, both directly and indirectly.

For example, large parts of Africa and Central Asia are likely to experience reductions in the length of growing period as a result of increased temperatures and lower rainfall. This is likely to lead to lower crop yields and reduced rangeland productivity, thus affecting the provision of feeds for animals. Climate change is also likely to change the distribution of animal diseases and their vectors. Large parts of South and Southeast Asia are likely to experience increases in rainfall and in the number of extreme climatic events (e.g. cyclones). This could lead to increased exposure of livestock to diseases, such as those caused by helminths. Crop losses due to extremes in climate could result in less animal feed being available, especially in crop-livestock and pastoral systems.

Science and technology drivers of change: general aspects and in relation to animal breeding and genetics

Science and technology have had a major influence on the transformation of animal production in industrialized economies and increasingly in developing countries. With increasing labour scarcity, larger, high-output and more productive animals were bred. From multipurpose breeds, highly specialized breeds were developed. Generally, disease resistance was sacrificed for

higher output, taking into account that through capital investments it became possible to adapt the environment to the existing animals in ways that had not been possible in the past. Research into housing and mechanization allowed significant labour productivity increases. These advances occurred in many species but particularly in short-cycled monogastric species such as poultry and pigs.

Animal nutrition research, linked with breeding, has made major contributions to improving feed efficiency and shortening production cycles and thereby reducing maintenance feed requirements and allowing a more efficient use of the capital investments and natural resources.

In the developing world, the impact of modern livestock science and technology has been uneven. Industrial livestock production systems (mainly for chickens) with limited links to the local resource base have been developed in some locations close to urban demand and/or to ports, given their frequent dependence on imported feed. Smallholder crop-livestock systems are much more reliant on locally available feed and traditional breeds. These crop-livestock systems are highly complex, delivering multiple products and services. Progress in improving the sustainable productivity of these systems has been much more limited and is a significant research challenge. System-based research is required to help these systems change in line with the changing social, economic and environmental context in which they operate. Currently, the speed of change of animal production systems and market chains is very high in some locations/regions, and is accompanied by loss of animal genetic resources. (This is discussed further below.)

Science and the management of animal genetic resources

The science related to the management of animal genetic resources has made significant progress, based mainly on advances in molecular biology and genetics as well as new developments in information and communications technology (ICT). The main advances are summarized in this paper and are discussed in more detail in the following papers. The advances include:

- Technologies are increasingly available for characterizing animal genetic resources. Molecular characterization is providing a better understanding of the genetic diversity in global livestock populations. Functional genomics is

also making it possible for genomes to be characterized, specific genomic regions and genes identified and gene functions elucidated. These technologies are based on a combination of genetic analysis and bioinformatics.

- New technologies are becoming increasingly available for utilizing animal genetic resources better, to meet changing needs, threats and opportunities. New genetic technologies enable the better characterization of breeds and populations. Other technologies, such as geographic information systems (GIS), enable the better characterization of the environment. Linking this knowledge will enable making a better fit between a genotype and an environment and, in the longer term, understanding the genetic basis of genotype x environment interaction. In this way, we can begin to identify appropriate genotypes for fast-changing environments. For example, there are increasing threats from drier climates that increase the need for hardier animals, tolerant to drought and disease. Animal reproduction technologies such as sexed semen and *in vitro* fertilization of embryos will enable the rapid development of new populations and faster distribution of superior animal genetics. These technologies are not yet widely used in developing countries, but offer future options in areas where a genetic solution is possible.
- Technologies are increasingly available for conserving animal genetic resources. New technologies are available for improved cryopreservation of embryos and semen that are applicable in more species. These technologies lead to new options for *ex situ*, *in vitro* conservation of animal genetic resources. For example, use of testes and ovaries obtained from livestock as sources of frozen semen and *in vitro* fertilization (IVF) embryos for long-term cryopreservation of animal genetic resources in gene banks.
- ICTs enable more precise linkage of genotypes and locations/production environments. New developments in ICTs also have implications for animal genetic resources characterization and conservation. These developments are linked to improvement of infrastructure and communication systems, such as the widespread use of mobile phones. ICTs also allow georeferencing to link particular genotypes with specific geographic locations. This knowledge provides the scientific underpinning of *in situ* conservation practices.

In order to take full advantage of the opportunities presented by advances in ICT, it is necessary to develop common standards for characterizing animal genetic resources, in terms of their genetics, phenotype and production system, so that knowledge can be shared among different communities and countries. Given such systematic and standardized descriptions of livestock, the intersection between new ICTs and modern genetics, through genomics and bioinformatics, presents opportunities to examine genome function by integration of these rich data sets.

Current status and trends in livestock production systems

In the light of the above drivers of change, this section discusses:

- The relative importance of the three main livestock systems worldwide (industrial, crop-livestock and pastoral) and the breeds they harbour.
- The implications of global drivers of change for the different livestock production systems.
- The implications for livelihoods.
- The implications of the scope and rate of changes in the main livestock production systems for current and future animal genetic resources management.

Livestock species by region

The geographic distribution of the major livestock species worldwide is given in table 3. This table shows that for all species the majority of animals are in the developing world. It also shows the importance of different species by region. For example, ruminants are most important in sub-Saharan Africa (SSA) and Latin America (LAC), both continents with vast areas of savannah and relatively low population densities. Poultry is most important in East Asia and the Pacific and LAC, regions of either high economic growth or with middle-income countries with high degrees of urbanization and adequate market infrastructure.

Livestock production systems by region

Three major types of livestock production systems can be identified worldwide – industrial livestock systems (IS); crop/livestock systems, mainly in high potential areas (CLS); and pastoral systems, mainly in marginal areas (PS).

The share of livestock in each of these systems in different geographic regions is shown in table 4. These data show that most livestock are located in crop-livestock systems. The proportion of livestock in industrial systems by region is mainly a function of economic status and rate of growth (e.g. higher proportions of industrial systems in the industrialized world and Asia).

Implications of global drivers of change for livestock production systems

Current status of livestock production systems

Each of the three main livestock production systems responds differently to the effects of the global drivers of change, and therefore has different development and investment needs. The overarching trends are increasing intensification in both industrial systems and in crop-livestock systems in order to meet increasing demand for animal products and consumer preferences for higher-quality products that meet stringent food safety standards.

- Intensification and scaling up trends in industrial and crop-livestock production systems.
The demand for livestock products has been met by intensification of livestock production systems in both developing and industrialized countries. Among other factors, this intensification has been based on using cereal grains as livestock feed. For example, in OECD countries, livestock feeding in intensive systems accounts for two-thirds of the average per capita grain consumption. In contrast, crop-livestock systems in sub-Saharan Africa and India use less than 10 percent of grains as feeds as they rely mostly on crop-residues (40–70 percent of feed), grazing and planted fodders.
- Market characteristics and demand.
The trend towards intensification of industrial systems and crop-livestock systems is largely driven by consumer demands for livestock products, both fresh and processed. The market characteristics are increasing demand for animal

Table 3. Geographic distribution of livestock (millions of head).

	Cattle	Sheep and goats	Pigs	Poultry
Sub-Saharan Africa ^a	219	365	22	865
Near East and North Africa ^a	23	205	0	868
Latin America and Caribbean ^a	370	112	70	2 343
North America ^a	110	10	74	2 107
East Europe and Central Asia ^a	84	121	72	1 160
West Europe ^a	83	119	125	1 072
East Asia and Pacific ^a	184	514	543	7 168
South Asia ^a	244	303	15	777
Industrial world ^b	318	390	284	4 663
Developing world ^b	1 046	1 460	659	12 735

^aAverage 2000–2005 number.

^bReported number for 2004.

Source: FAOSTAT (2007).

Table 4. Share of livestock (total livestock units [TLU]: cattle, goats, sheep, pigs and poultry) per livestock production system for selected regions and countries.

	TLU shares (%)		
	Livestock production system		
	PS	CLS	IS
Sub-Saharan Africa			
Botswana	80	19	0.14
Kenya	34	50	14
Mali	47	51	0.9
South Africa	55	36	8
Latin America and Caribbean			
Argentina	42	40	16
Brazil	18	63	17
Peru	44	21	33
East Asia and Pacific			
Cambodia	6	73	20
China	9	70	19
Viet Nam	0.75	82	16
South Asia			
India	2	82	15
Pakistan	25	63	10
Developed World			
European Union	9	67	22
Russian Federation	16	50	32

Source: FAO (2004).

products in developing countries, plus quality preferences and food safety requirements in all markets. Public-private partnerships that provide services and market opportunities also play a key role in intensifying industrial and crop-livestock systems.

Future trends in livestock production systems

Intensive systems. Intensive systems are facing increasing restrictions, owing to their associated negative environmental effects, such as problems of waste disposal and water contamination. Demand

for cereals is also increasing for other purposes (e.g. biofuels) and this is driving up the price of cereals, and subsequently the price of livestock products coming from intensive systems.

Crop-livestock systems. Crop-livestock systems in developing countries are constrained by farm size and lack of access to inputs and services. These constraints affect soil fertility, crop yields, income generation and ultimately livestock production through the limited provision of high-quality feeds. There is also increasing competition for land and associated opportunity costs.

Pastoral systems. The remoteness and the limited agricultural potential of pastoral systems in marginal areas of the developing world create difficulties for these systems to integrate into the expanding markets for livestock products. This poses a set of different needs related to adaptation of systems to reduce the vulnerability of livestock keepers and their animals and expanding access to markets.

A major driver of change in pastoral systems over the past decades has been the widespread policy to settle pastoralists and allocate them individual land rights. This approach and the increasing encroachment of crop production have seriously affected the viability of these systems by reducing the mobility of livestock and access to feed resources. Although the negative aspects of these policies are increasingly acknowledged, they will continue to shape political processes in many developing countries.

Future implications of structural changes in livestock production systems

In the industrial and mixed crop-livestock systems, rising demand for livestock products will continue to drive structural changes in these livestock production systems and markets. Market transformation, particularly in urban markets, will lead to the increasing importance of supermarkets, large livestock processors and transformation of wholesale livestock markets. Much of this transformation has taken place in the industrialized countries. This pattern is expected to increase in the developing world with a growing share of industrial livestock systems.

Farmers in intensifying crop-livestock systems will diversify their production into dairy and other livestock products even more in response to market opportunities arising from rising demand for high-value foods. Similarly, income growth and urbanization will increase diversification of

consumer diets and the share of livestock products in diets.

The major changes in livestock markets are going to take place in domestic markets. The relative importance of domestic markets versus trade in the future will reflect past trends in which domestic market dynamics were far more important than trade. For example, in 1980 and 2001, meat exports and imports were approximately four percent of output and consumption in the developing world. In contrast, the share of domestic urban markets in total livestock consumption has been increasing over the past 25 years.

The growing importance of domestic urban markets as opposed to international trade implies changes of actors in domestic livestock industries, particularly in agribusiness in wholesale markets, livestock processing and the retail industry, with more fresh and processed animal products being sold through supermarkets.

These structural changes in markets, transformation in urban markets, and in retail and distribution sectors in the livestock industry will have profound impacts for the future of smallholders and poor livestock keepers in competing with intensifying industrial and crop-livestock systems in high potential areas. Empirical evidence from Asia shows that smallholder farmers provide up to half of the share of production in dairy and meat markets. Undercapitalized small producers are likely to be squeezed out of dynamic domestic livestock markets. Policy action that supports small producers who can be helped to become competitive will have substantial equity pay-offs. In the absence of such pro-poor policies in the livestock sector, market changes and the entry of new actors in livestock processing, distribution chains and the retail sector can marginalize poor people who depend on livestock for their livelihoods.

High transaction costs and limited access to markets will lead to a dramatic decline of share of livestock production from pastoral systems in marginal areas. Without significant public investments in infrastructure and services, poor producers in these areas will become increasingly marginalized and many will have to leave livestock production as a source of income. Livestock will continue to be important in traditional pastoral systems as sources of food and fulfil multiple other uses, providing traction, transport, skins and hides for shelter.

Implications for livelihoods

In terms of livelihood impacts, the above changes will lead to changes in the role of animal genetic resources for livelihoods in two divergent ways: in intensive systems livelihoods will have a weak link to genetic resources, which will play very specialized production roles. The major livelihood impacts will be through employment. Frequently this will be limited direct employment in large-scale operations but some increased employment will be expected along the value chain. Consumer livelihoods will be affected in terms of impact of prices and of changed attributes of the animal products coming from these intensive systems. Society-wide, there may be negative impacts on livelihoods of traditional smallholders displaced from markets by industrially produced animal products. The net effects will depend significantly on the policy environment and the extent of substitution between animal products produced by industrial systems and smallholder systems.

In crop-livestock systems, livelihoods will be affected by the pressures to intensify and specialize production. Systems may change from grazing to zero-grazed systems, increasing milk production while reducing animal traction. This will imply changes in the labour patterns and possibly gender distribution of work and benefits from animal production. More intensively kept animals will require higher levels of management and external inputs. Increasing livelihood opportunities can be expected to develop in these forward and backward linkages associated with these commodity chains.

Pastoral systems in developing countries tend to have very strong linkages to diverse species and breeds of animals, which allow them to adapt to the exploitation of natural resources with very unique attributes and generally very limited alternative uses. Livelihoods are intimately linked to the animal genetic resources under these conditions. Risk is a major issue and the management of multiple species and multiple outputs is a key way of coping. Increasing competition for the resources, as well as policy orientations towards settling pastoralists, significantly affect these peoples' livelihoods.

In the industrialized world, highly specialized pastoral production systems rely heavily on their animal genetic resources – normally a narrow genetic base comprising one or two commercial breeds of one or two species or a defined crossbred animal population. In relation to pastoral and smallholder systems in developing countries, these systems do not involve much labour. Therefore, the

livelihoods of fewer people are generally involved in these production systems.

Implications of the scope and rate of changes in livestock production systems for animal genetic resources management

The drivers of change and the evolution of the farming systems that they induce will have important effects on livestock biodiversity and its use. This in turn implies that needs and opportunities for human intervention will vary.

In industrial systems, where it is largely possible to adapt the environment to the needs of the animals, highly productive commercial breeds and hybrids are going to be the main genetic pillar. Genetic resources are handled by the specialized private sector firms and traded internationally. Their interest in hardiness or disease-resistance traits will be limited unless diseases emerge for which no alternative control strategies are available or policies require important changes in the management systems, e.g. free-ranging instead of caged laying hens.

In crop-livestock systems, pressure to intensify will be a major force shaping the production system and the genetic resources underpinning it. Significant increases in productivity will be required to meet demand and these will be achieved by simultaneously improving the conditions (feed, health, etc) and adapting the genetic resources. Given the heterogeneous environments, many different breeds will be required. In higher potential areas with good market access this specialization will increasingly involve crossbreeding with exotic breeds. Given the relatively small numbers of animals of each breed required in these niches, these genetic materials will not be produced by private multinational companies but will require active engagement of farmers, public sector and non-governmental organizations (NGOs). These systems will continue to be an important source of genetic diversity and will also demand a range of solutions to fit their specific conditions. As science improves its capacity to understand the role of specific genes and their interaction with environmental factors triggering their expression, the value of local breeds in targeted breeding programmes for these systems will increase. These systems will naturally use a diverse genetic base and will be amenable to engage with *in situ* conservation. Supportive institutional arrangements will be key to driving such efforts.

In pastoral systems in developing countries, high levels of diversity can be encountered and traits of disease-resistance and tolerance of harsh environments are widely present. These systems are frequently declining in livestock numbers and in particular small endemic populations are at risk. In these settings, conservation will require public action because of the limited resources of the generally poor pastoralists. This will be an area where NGOs can be expected to play a key role in assisting in *in situ* conservation.

Given the fragility of institutional arrangements in many developing country contexts and their exposure to natural and human-induced crises, there is merit in designing *ex situ*, *in vitro* conservation strategies as a back up and long-term insurance against loss of diversity in the field. These conservation strategies will need to be coordinated at national and regional/international levels to be efficient and cost-effective.

Climate change considerations add an important dimension to the discussion of livestock biodiversity. Different systems will be affected in different and highly uncertain ways, but access to genetic resources could be a critical ingredient for most adaptation responses in the medium to long term. Table 5 summarizes major trends in livestock system evolution and their implications for the management of animal genetic resources.

Conclusions and next steps

What immediate steps are possible to improve animal genetic resources characterization, use and conservation?

Appropriate institutional and policy frameworks are required to improve animal genetic resources management and these issues are being addressed at national and intergovernmental levels, in a process led by FAO to promote greater international collaboration. Based on an analysis of the current situation, the continuing loss of indigenous breeds of farm animals, new developments in science and technology, and the strategies suggested for the future management of animal genetic resources (as summarized in table 5), there are several complementary actions that can begin to improve the management of animal genetic resources and maintain future options in an uncertain world. The scientific basis that underpins these proposed actions is discussed in more detail in subsequent papers. Four areas for action to improve the

sustainable use and *in situ* conservation, characterization and long-term *ex situ* conservation of animal genetic resources are summarized here, and are addressed in further detail in the companion papers:

Sustainable use and *in situ* conservation of animal genetic resources

*“Keep it on the hoof” – Encouraging the continuing sustainable use of traditional breeds and *in situ* conservation of animal genetic resources, by providing market-driven incentives, public policy and other support to enable livestock keepers to maintain genetic diversity in their livestock populations.*

In this context, *sustainable use* refers to the continuing use of traditional breeds by livestock keepers, as a result of market-driven incentives. *In situ* conservation refers to animal genetic resources conservation measures supported by public policy and, on occasion, public investments to support *in situ* conservation of traditional breeds by livestock keepers.

In regard to encouraging the sustainable use of animal genetic resources, market-driven incentives applicable in developing countries include facilitating access to markets for livestock products coming from traditional breeds. This may include identifying niche markets for traditional products and providing infrastructure (such as transport) to help livestock keepers to get their products to market.

Increasing the productivity of traditional breeds through breeding is also an incentive for livestock keepers to retain these breeds. (The companion paper discusses the role of breeding in more detail.) These breed improvement strategies could also make more use of the widespread crossing that has occurred in traditional populations over time, as livestock keepers seek to improve their breeds.

In regard to encouraging *in situ* conservation of particular breeds, especially in the diversity-rich crop-livestock and pastoral systems in developing countries, the incentives include having public policies that support the conservation of traditional breeds and providing public services (e.g. human and livestock health services, schools, roads) to support communities in livestock producing areas. Such services may encourage people to stay with their animals in rural areas rather than migrate to urban areas where more services are available.

In situ conservation makes use of local and indigenous knowledge, which can also be validated scientifically. For example, some farmers have

realized that by crossbreeding part of their herd to an exotic breed, they can make more profit during the good times but avoid the risk of losing all their animals when conditions are bad. Exotic animals tend to be poorly adapted to harsh conditions and tend to die during droughts, for example. Thus genetic variability reduces vulnerability to sudden changes and shocks in the system.

The concept of *in situ* conservation also extends to conserving livestock as part of the landscape, within an overall biodiversity conservation strategy, as a long-term global public good.

“Move it or lose it” – Enabling access and safe movement of animal genetic resources within and between countries, regions and continents.

Maintaining mobility of animal breeds, populations and genes within and between countries, regions and continents is one of the key actions for facilitating the sustainable use and thereby the conservation of animal genetic

resources. Safe movement of animal genetic resources enables their access, use and conservation for mutual benefit by livestock keepers worldwide. *Mobility* here refers to facilitating informed access to genetic diversity, based on systematic breed evaluations and analysing the potential usefulness of various breeds in different environments.

There are benefits and risks in increasing the mobility of animal genetic resources. The benefit is that, in a fast-changing, unpredictable world, mobility of animal genetic resources enables flexibility in response to changing climate, disasters, civil strife, etc. For example, when civil strife has occurred in some part of Africa, animals are moved across borders to avoid their unintended death in conflicts. One risk of increased mobility is that animals moving to different environments may not be adapted to their new environment, livestock system or social system. There are also animal health risks, in terms of the possible spread of disease, or by animals not being tolerant to the diseases prevalent in a new environment. For

Table 5. Trends in livestock system evolution and their implications for the management of animal genetic resources.

Livestock production system: description and trends	AnGR – current status in system	AnGR management: future strategy for each livestock production system
Industrial systems (IS)		
Industrial systems changing quickly, expanding globally.	Breeding by private sector, with narrow genetic base in pigs, poultry, cattle.	Commercial systems will continue to adapt environment to suit genetics (IS prefer to use most productive breeds and manage other production issues by non-genetic means).
Controlled system, almost “landless” environment, able to adapt environment to genetics.	High-value genetic stock protected by know-how and traded internationally.	IS need to be able to respond to future shocks (e.g. identify tolerance to zoonotic diseases such as avian influenza and also identify more disease-resistant breeds able cope with diseases of intensification without antibiotics).
Systems changing to reduce negative environmental impacts, meet market demands and consumer preferences, and address new issues (e.g. animal health and welfare).	Limited interest or incentive for private firms in conserving species/breed biodiversity.	Conserving AnGR of main industrial species (pigs, poultry, cattle) to maintain biodiversity is a long term, public (and private) good to enable IS to deal with future options and new shocks .
Changing systems require broader genetic base to address new issues and future shocks.		

(... continued)

Livestock production system: description and trends	AnGR – current status in system	AnGR management: future strategy for each livestock production system
<p>Crop-livestock systems (CLS) Diverse systems with broader genetic base, in industrialized and developing countries.</p> <p>CLS dependent on natural resource (NR) base.</p> <p>CLS less in control of environment than IS.</p> <p>Future of CLS affected by market demands, NR availability, climate change, land-use options.</p> <p>CLS changing and intensifying production, especially in developing countries; but rate of change less than for IS.</p> <p>Intensification options – better feed, land, water use, genetic improvement.</p>	<p>Developing and conserving AnGR by use in CLS (<i>in situ</i>).</p> <p>Genetic base more diverse than IS, as animals need to be in balance with system and co-evolve with natural resource base.</p> <p>Sustainable delivery of genetic material occurring in some CLS.</p>	<p>Need to adapt animal genetics to changing environment.</p> <p>CLS need to be able to respond to changing environment, climate change effects, other drivers of change; conserving diverse AnGR in CLS is a public good.</p> <p>Sustainable use of AnGR will help CLS maintain diversity and ability to respond to future drivers of change.</p> <p>Smallholders may require incentives to continue to conserve AnGR <i>in situ</i> with changing, more productive CLS (e.g. foster niche markets to encourage farmers to keep traditional breeds, for short- and long-term value).</p> <p>Mobility of AnGR critical to maintain future options as CLS change in response to global drivers (mobility favours sustainable use of AnGR).</p> <p>Example of moving adapted AnGR to new areas when climate change affects system, such as moving hardier animals to areas more prone to drought.</p> <p>Institutional development to support sustainable AnGR management in CLS (e.g. farmers associations, environmental, food safety and animal health regulations).</p>

(... continued)

Livestock production system: description and trends	AnGR – current status in system	AnGR management: future strategy for each livestock production system
Pastoral systems (PS) in marginal areas		
PS comprise rangelands in industrial and developing countries.	PS in industrial countries have narrow genetic base.	Need to adapt animal genetics to marginal environment.
Systems determined by NR base, usually in marginal environments.	PS in developing countries have diverse AnGR, conserved through sustainable use.	Maintaining diverse AnGR is desirable to reduce vulnerability of livestock keepers.
Multiple value and uses of animals in traditional PS in developing countries.	Traditional AnGR conservation <i>in situ</i> by livestock keepers, linked with indigenous knowledge of animals and land.	Future need to improve productivity of PS, maintain livelihoods, with less people likely to be living in marginal lands (e.g. animal health interventions).
PS changing more slowly than IS or CLS as least likely to be influenced by global drivers of change.		Genetic solutions through hardier animals, able to adapt to harsher environments, with few interventions.
Some PS changing more quickly (e.g. in parts of India where there is competition for pastoral land for alternative uses).		Incentives to maintain <i>in situ</i> conservation practices and promote sustainable use (e.g. improve market access through better infrastructure; foster niche markets for traditional animal products).
PS closely related to traditional (cultural) practices and institutions for the management of natural resources and traditional knowledge.		Risk mitigation (e.g. better forecasting and strategies for handling risks in PS, such as droughts).
		Payments for environmental services may mean alternative land-use options that complement or compete with livestock production; requires adaptation of PS and related AnGR, depending on the nature of the environmental service.
		Institutional development to support policies and practices for grazing, water and land-use rights.

transboundary movements, these risks as well as the benefit should be identified and shared with stakeholders prior to importation, and risk mitigation steps taken before importing semen, embryos or live animals into a country.

Characterizing animal genetic resources

“Match breeds to environments” – Understanding the match between livestock breeds, populations and genes and the physical, biological and economic landscape. This “landscape livestock genomics” approach offers the means to predict the genotypes most appropriate to a given environment and, in the longer term, to understand the genetic basis of adaptation of the genotype to the environment.

In regard to the long-term prospects for this research, the advances in our ability to describe the genome of an animal in unprecedented detail, coupled with our ability (through spatial analysis) to describe the landscape in which it resides – a landscape description that includes biotic, abiotic, human and market influences – are beginning to provide an opportunity to probe genome function in a unique way. This is an approach already used to study the distribution of particular alleles in livestock and to probe the human genome for disease-causing genes. Its potential for understanding the fit between livestock genotype and landscape is significant, and it depends on sophisticated data-management tools. It also offers the opportunity not only to understand the function of the genome, but also to predict the genotype most appropriate to a given environment.

This is a long-term research objective that can be linked with existing data-gathering exercises to add to their value. For example, building in systematic sampling of DNA of livestock breeds in combination with a careful description of the systems under which each population presently functions, and georeferencing the data, will add greatly to our ability to understand and utilize animal genetic resources. For example, we can begin to ask *“what combination of genotypes is appropriate for a milking cow under a given management regime, under a given range of disease pressures and under a given set of physical stresses?”* Knowing this will enhance the value of genotypes *“in the bank”* or *“on the hoof”* and will provide the tools we need to identify intelligently appropriate genotypes for specific agro-ecological niches. (Approaches to

characterizing AnGR are discussed further in the companion paper.)

Ex situ conservation of animal genetic resources in gene banks

“Put some in the bank” – New technologies make ex situ, in vitro conservation of animal genetic resources feasible for critical situations and a way to provide long-term insurance against future shocks in all livestock production systems.

Improving technology (e.g. cryopreservation) is making long-term, *ex situ*, *in vitro* conservation of semen and embryos more feasible, affordable and applicable to a wider range of species. The challenge is to decide which animal genetic resources to conserve; how to collect them; where to store them; when and how to characterize them; and who can access, use and benefit from them in the future. It is particularly important to collect the rich diversity of traditional livestock breeds in crop-livestock and pastoral systems in developing countries before it is lost forever.

A risk is that *ex situ*, *in vitro* gene banks can become *“stamp collections”*, put away in the deep freeze and never characterized. Another potential risk is that this approach may be a disincentive to *in situ* conservation through sustainable use, where the genetic resources are more accessible in the short to medium term and where not only the genetic resources but also the traditional knowledge associated with them are conserved. In fact, *in situ* and *ex situ* conservation approaches are complementary rather than competing approaches, serving short- and long-term needs. *Ex situ*, *in vitro* animal genetic resources conservation is a long-term insurance policy and an important first step in conserving animal genetic resources for future generations. (Further details on conservation approaches are given in the companion paper.)

Closing remarks

Several important drivers of change are leading to rapid changes in the livestock production sector that have implications for the future management of animal genetic resources. The multiple values, functions and consequences of livestock production systems and their rapid rate of change lead to divergent interests within and between countries. Conversely, the uncertainty about the implications

of rapid, multifaceted global change for each livestock production system and the resulting future changes in the required genetic make-up of the animals makes collective action to tackle conservation of animal genetic resources a long-term, global public good. Developing and conserving animal genetic resources will not by themselves solve all these problems, but are important first steps towards maintaining future options.

Advances in science and technology, in areas such as reproductive technology, genomics and spatial analysis, as well as progress in conceptualization of global public good production for the future management of animal genetic resources, should enable the international community to address both the short- and long-term challenges in innovative ways.

List of references

- FAO.** 2004. Working files produced by Environmental Research Group, Oxford, UK, unpublished.
- FAO.** 2006. Livestock's long shadow – environmental issues and options, by H. Steinfeld, P. Gerber, T. Wassenaar, V. Castel, M. Rosales & C. de Haan. Rome.
- FAO.** 2007. FAOSTAT statistical database. Rome (available at <http://faostat.fao.org>).
- IAASTD.** 2007. International Assessment of Agriculture Science and Technology for Development. World Bank, Washington, DC (in press).
- IMF (International Monetary Fund).** 2007. World economic outlook database. Washington, DC.
- Reardon, T. & Timmer, P.C.** 2005. Transformation of markets for agricultural output in developing countries since 1950: how has thinking changed? In R. Evenson, P. Pingali & T.P. Schultz, eds. Handbook of agricultural economics, Vol. 3A. Amsterdam, Elsevier.
- Reardon, T., Henson, S. & Berdegue, J.** 2007. Proactive fast-tracking' diffusion of supermarkets in developing countries: implications for market institutions and trade. *Journal of Economic Geography*, 7: 399–432.
- United Nations.** 2007. Common database. New York, USA.
- World Bank.** 2006. World development indicators. Washington, DC.

Bibliography for further reading

De Fraiture, C., Wichelns, D., Rockstrom, J., Kemp-Benedict, E., Eriyagama, N., Gordon, L., Hanjra, M.A., Hoogenveen, J., Huber-Lee, A. & Karlberg, L. 2007. Looking ahead to 2050: scenarios of alternative investment approaches. Water for food, water for life. In: A comprehensive assessment of water in agriculture. London, Earthscan, and Colombo, International Water Management Institute, pp. 9–145.

Delgado, C., Rosegrant, M., Steinfeld, H., Ehui, S. & Courbois, C. 1999. Livestock to 2020: the next food revolution, Food, Agriculture, and the Environment Discussion Paper 28. Washington DC., International food Policy Research Institute.

Fisher, P., Hedeler, C., Wolstencroft, K., Hulme, H., Noyes, H., Kemp, S., Stevens, R.T. & Brass, A. 2007. A systematic strategy for large-scale analysis of genotype-phenotype correlations: identification of candidate genes involved in African trypanosomiasis. *Nucleic Acids Research* (in press).

Gibson, J., Gamage, S., Hanotte, O., Iñiguez, L., Maillard, J.C., Rischkowsky, B., Semambo, D. & Toll, J. 2006. Options and strategies for the conservation of farm animal genetic resources. Report of an international workshop, 7-10 November 2005, Montpellier, France. Rome, CGIAR System-wide Genetic Resources Programme (SGRP)/Bioversity International, pp. 53.

ILRI (International Livestock Research Institute). 2002. Livestock, a pathway out of poverty: ILRI's strategy to 2010. Nairobi.

ILRI/FAO. 2006. The future of livestock in developing countries to 2030, Workshop Report on Future of Livestock in Developing Countries, 13-15 February 2006. Nairobi, ILRI.

.....

Kierstein, S., Noyes, H., Niessens, J., Nakamura, Y., Pritchard, C., Gibson, J., Kemp, S. & Brass, A. 2006. Gene expression profiling in a mouse model for African trypanosomiasis. *Genes Immun.*, 7: 667–679.

Lockhart, D.J. & Winzeler, E.A. 2000. Genomics, gene expression and DNA arrays. *Nature*, 405: 827–836.

MacKenzie, A.A. ed. 2005. Biotechnology applications in animal health and production. *Scientific and Technical Review*, 24(1), April 2005.

Pittroff, W., Cartwright, T.C. & Kothmann, M.M. 2002. Perspectives for livestock on grazinglands. *Archivos Latinoamericanos de Produccion Animal*, 10(2): 133–143.

.....

Panellists' comments and discussion

Mr Ken Laughlin, European Forum of Farm Animal Breeders (EFFAB), the Netherlands

Firstly I would like to thank FAO for the opportunity to participate in this meeting and panel discussion.

This paper recognizes the need for action and action plans to ensure the retention of biodiversity and not just to conserve the *status quo* for another day.

Further it recognizes the need for market driven incentives as part of the process for using and developing traditional breeds at local, national and international levels where this might be possible. The paper suggests having public policies that support the conservation of traditional breeds and it may also be valuable to have discussion on the perceived positive and negative opportunities of some system of "*animal breeder's rights*" in this regard.

Critically in the paper there is an appreciation of the need to focus on understanding the real values of these breeds, which can be validated scientifically following the generation of a sufficient number of animals with accurate trait measurements. This will be a necessary basis for understanding the match between breeds, populations and genes with the physical biological and economic landscape. Fortunately this does not require high tech science but does require diligence and attention to detail. There is a vast amount of available experience amongst existing breeders in this area of trait definition and collection.

There is a clear opportunity to engage with the breeders in the "*industrial system*" who have seen many of the opportunities and pitfalls of the last century of "*informed animal breeding*". Specifically there is an opportunity to learn from the experience and data of poultry breeding which has probably evolved further than any other species in terms of balanced breeding.

Interestingly, the importance of focusing on reducing and halting the loss of indigenous breeds of farm animals is strongly supported by a recent genomic study of poultry which has concluded that modern agricultural practices were not the major source of allele loss. The majority of alleles were lost prior to the formation of the current intensive industry.

The breeders in industrial systems have further experience of the opportunities and threats of breeding stock movements where the OIE systems of disease recognition and subsequent movement control are followed.

Finally it is important that during this forum we try to identify what the real drivers are in reducing biodiversity and protecting biodiversity. These may not simply be opposites.

Mr Fernando Madalena, Brazil

I would like to compliment Dr. Seré and co-authors on their comprehensive paper. I have the following comments, not so much to the paper itself, but rather to the general atmosphere related to the management of animal genetic resources. It would seem that we keep changing our flags over time before the goals are reached. In the sixties we talked about development, then -before- development was accomplished in most countries- we started talking about sustainable development, and again before development, let alone sustainable, was accomplished, we now talk about alleviating poverty ... of the future generations. Incidentally, poverty should be eliminated, not just alleviated, and the proper utilization of animal genetic resources may contribute some to that end, by the widespread use of the more economic stocks, breeds or crosses. Less productive resources need then special preservation programmes. Using new genomic technologies seems particularly attractive to researchers in developed and developing countries. However, this comes at a price, because of patenting and problems of accessing knowledge.

While in the past emphasis was in the utilization of genetic resources, I have the impression that we are now focusing too much on characterization and conservation, and a lot of technical work and funding are dedicated to these two. However, I must confess I fail to grasp the logic of caring only of preserving resources for the possible use by future generations when we do not use them properly now to help meeting the so many urgent human needs. Some "*stakeholders*" would agree with me, as the duck in this quoted cartoon, who is telling its chick "*Humans are odd. They save animals but let their own kids die of hunger*". I therefore

would like to emphasize the need for a renewed and stronger focus on the utilization of animal genetic resources to meet current human needs and not only on conservation for potential future needs. In order to achieve this, we must go beyond characterization to evaluation. Only by comparatively evaluating the specific features of our breeds in given environmental and economic circumstances, we will be in a position to better utilize them.

Ms Ilse Köhler-Rollefson, League for Pastoral Peoples and Endogenous Livestock Development, Germany/India

I agree with many of the observations and conclusions made in the paper, however I feel the analysis remains on a superficial level, and that there is not enough critical reflection on why we have gotten where we are and whether the rapid erosion of animal genetic resources that we have witnessed in the last several decades has been as inevitable as it is depicted here.

The paper zeros in on four drivers of change, i.e. economics and globalization, poverty, environmental concerns, and science and technology. This is the way in which the situation may present itself to scientists and is a correct analysis on some level. However for an organization that is working with poor and marginal livestock keepers, especially pastoralists, on the ground, the scenario looks quite different.

The paper does not mention that the poor livestock keepers that have stewarded diversity have been progressively losing their rights, and that industrialized systems and the introduction of exotic breeds have been heavily subsidized. The example of how exotic pig breeds in Viet Nam have been promoted through a variety of subsidies is now well known, but it is only one case that happens to have been scrutinized. In fact, in the last several decades, livestock development in developing countries everywhere has been practically synonymous with the replacement or “upgrading” of local breeds with exotic ones. By contrast, very little endeavour has been made with respect to endogenous development and taking local resources, knowledge, and institutions as a starting point for livestock-related interventions. There have been no efforts to strengthen the livestock keepers themselves, by assisting them to develop their own breeds, by supporting them organizationally, by involving them in policy development, and by integrating them into research

projects. In order to protect industrial production, we are now making it almost impossible for diversity-conserving livestock keepers to continue keeping animals.

In the wake of avian influenza, some countries have prohibited the keeping of smallholder poultry in the vicinity of industrial holdings. In Germany, there has been a close to 50 percent decrease in the number of smallholder poultry because their keepers could not comply or cope with regulations to confine their animals.

For us, the drivers of change are thus the progressive loss of rights of smallholder livestock keepers and of pastoralists – such as the right to keep animals, in order to protect the interests of industrial producers, and the loss of customary grazing rights and access to land.

We are also concerned that farmers are increasingly losing their breeding function, becoming mere raisers of animals that are dependent on a handful of companies for all their inputs which have often been produced at the other end of the world and leave a huge carbon footprint.

Cryoconservation as a backup is ok, but in the final analysis it can not be expected to be of much relevance, if the pressure on small-scale diversity-conserving livestock keepers continues and they are forced to, or choose to, give up keeping animals. Banking on cryoconservation basically means taking the easy way out, and is once again a way of strengthening scientists and government bureaucracies, rather than livestock keepers.

We must be clear that even the best science will not be able to move things in a better direction and to save diversity – it can only play a supporting role. The only real care takers of animal genetic resources are farmers and pastoralists. If we want to save animal genetic resources, then we need to save these livestock keepers first, by creating appropriate policies that strengthen them and that reward them for their crucial role in sustaining diversity. Such policies need to be combined with the monitoring and regulation of industrial production systems. Instead of subsidizing them and providing them with unfair advantages, these systems should be taxed.

My organization, the League for Pastoral Peoples will continue to contribute to the implementation of the *Global Plan of Action* on two levels:

- Globally and nationally, we will support pastoralist and small-scale farming communities to advocate for their rights and to achieve recognition of their role in conserving breeds and as custodians of diversity.

- Equally important, we will continue to work on the ground through our partners in the LIFE-Network to help pastoralists and small farmers to get organized and develop the skills and knowledge that they will need to remain competitive in a globalized economy.
- Maintaining animal genetic resources as an integral part of production systems offers opportunities to address conservation, and food and livelihood security in a complementary way.
- Animal genetic resources should be matched to the production environment. Exotic breeds and cross-breeding programmes offer opportunities to meet increased demand and support livelihoods where conditions are appropriate. However, in some production systems, the productivity of local breeds may be as good as, or better than, potential replacements if all outputs and inputs are taken into consideration – non-market roles of livestock, including insurance and financing functions are important here, but are often receive insufficient attention from researchers and policy-makers.

Summary of plenary discussion

The meeting was then opened for general discussion and interventions from the floor. Issues raised during this discussion included:

- The importance of non-market livestock functions and the question of their inclusion in economic statistics.
- The need to assess consumer demographics and trends with respect to their effects on animal genetic resources.
- The importance of networking and capacity-building.
- The impact of natural disasters, wars and unrest, and global warming on animal genetic resources.
- The important role of small-scale production.
- The potential conflict between maintaining genetic diversity and meeting increased demand for animal products in developing countries.
- The nature of the key issues that need to be addressed with respect to the rights of livestock keepers and communities, and differences compared to the plant sector.
- The important contribution of “minor” species (e.g. guinea pigs, snakes, insects).
- The influence of religious and cultural factors on the development of animal genetic resources.
- The need to consider possible trade-distorting effects of providing subsidies to livestock keepers.
- There is a need to identify institutional arrangements that can offer smallholders the opportunity to successfully market their products. The roles of “minor” species also require greater research attention as these are often important to livelihood and food security.
- Addressing all these issues, and providing the basis for informed decision-making, is a challenge to the scientific community - including the social sciences.
- With respect to the rights of farmers and communities, it is important to recognize that issues are not identical to those in the plant sector. In the case of livestock, the replacement of local breeds with exotic animal genetic resources rather than the expropriation of local genetic resources by outside interests is a key issue, although the latter also needs to be addressed. An equitable framework for access and benefit sharing is needed.
- In the case of acute threats such as wars, pre-emptive actions, for example to ensure that appropriate animal genetic resources are available for restocking programmes, are needed.

The authors’ responses and concluding remarks included the following points:

Inventory, characterization and monitoring

M. Tixier-Boichard^{1,2}, W. Ayalew³ & H. Jianlin⁴

¹Institut National de la Recherche Agronomique (INRA), AgroParisTech, UMR1236 Génétique et Diversité Animales, F-78350 Jouy-en-Josas, France

²Département sectoriel A4, "Biotechnologies - Ressources - Agronomie", Direction Générale de la Recherche et de l'Innovation, Ministère de l'Enseignement Supérieur et de la Recherche, 1 rue Descartes, F-75231 Paris, France

³National Agricultural Research Institute (NARI), P.O. Box 1639, Lae 411, Morobe Province, Papua New Guinea

⁴CAAS-ILRI Joint Laboratory on Livestock and Forage Genetic Resources, Institute of Animal Science, Chinese Academy of Agricultural Sciences (CAAS), No. 2, Yuan Min Yuan Xi Lu, Haidian District, Beijing 100094, P.R. China

Summary

Inventory of species and breeds, their population sizes, geographic distribution and possibly their genetic diversity is generally undertaken as a first step in any national programme for the management of animal genetic resources for food and agriculture. The primary purpose of such an assessment is to document the current state of knowledge in terms of a population's ability to survive, reproduce, produce and provide services to farmers. Starting an inventory requires some knowledge of the inventory items and their characteristic attributes. Inventory and characterization are, therefore, complementary processes, in which the characterization step provides the baseline information as well as the criteria that will be used to establish and update the inventory. Characterization provides data on present and potential future uses of the animal genetic resources under consideration, and establishes their current state as distinct breed populations and their risk status. As use and management of animal genetic resources are dynamic processes, monitoring the status of a population has to be done on a regular basis. Thus, risk status indicators for use during the monitoring process need to be defined following the inventory and characterization steps.

This paper discusses methods and criteria currently available, from research and past experience, for inventory, characterization and monitoring of animal genetic resources, with the view to assist in the development of a more comprehensive framework. Particular consideration is given to emerging tools and technologies. The scope of the review includes all livestock species and their wild ancestors and wild related species. Examples focus on cattle, sheep, goats, pigs and chickens.

Résumé

L'inventaire des espèces et des races, la taille des populations, la distribution géographique et si possible leur diversité génétique, est en général le premier pas à accomplir dans un programme national pour la gestion des ressources génétiques animales pour l'alimentation et l'agriculture. Le principal objectif de ce genre d'évaluation est de documenter la situation actuelle en termes de connaissances sur la capacité de survivre et de se reproduire d'une population, ainsi que d'offrir des services aux éleveurs. Pour initier un inventaire il est nécessaire de disposer de certaines connaissances sur les points principaux et sur les attributions des caractéristiques. En outre, inventaire et caractérisation sont des procédures complémentaires étant donné que la caractérisation fournit l'information de base et les critères qui s'utiliseront pour établir et mettre à jour l'inventaire. Tenant compte que l'utilisation et la gestion des ressources génétiques animales sont des procédures dynamiques, le suivi d'une population doit être réalisé sur des bases concrètes. Pour cette raison, il est nécessaire de définir les indicateurs pour les situations de risque qui seront utilisés pendant le suivi tenant compte des différents points de l'inventaire et de la caractérisation.

L'article présente les méthodes et critères disponibles actuellement à partir de la recherche et des expériences passées pour classer, caractériser et suivre les ressources génétiques animales dans le but d'aider au développement d'un réseau plus efficace. On souligne en particulier les nouveaux outils et technologies. L'objectif de cette révision comprend toutes les espèces d'élevage ainsi que leurs ancêtres sauvages et les espèces sauvages voisines. Certains exemples se sont centrés sur les bovins, ovins, caprins, porcins et volailles.

Resumen

El inventario de las especies y razas, el tamaño de sus poblaciones, su distribución geográfica y posiblemente su diversidad genética es en general lo que se hace como primer paso en un programa nacional para la gestión de los recursos zoogenéticos para la alimentación y la agricultura. El principal propósito de este tipo de evaluación es documentar la situación actual en términos de conocimientos sobre la capacidad de sobrevivir y de reproducirse de una población, y de proveer y producir servicios para los ganaderos. Iniciar un inventario requiere algunos conocimientos sobre los puntos de un inventario y la atribución de sus características. Por lo tanto, inventario y caracterización son procesos complementarios en los que el paso de la caracterización proporciona la información de base así como los criterios que se utilizaran para establecer y poner al día el inventario. La caracterización proporciona datos sobre el uso actual y potencial futuro de los recursos zoogenéticos en estudio, y establece cual es el estado actual de cada población de razas y situación de riesgo. Teniendo en cuenta que la utilización y gestión de los recursos zoogenéticos son procesos dinámicos, el seguimiento de la situación de una población debe llevarse a cabo sobre bases regulares. Por lo tanto, se necesitan definir indicadores sobre situaciones de riesgo para su utilización durante el seguimiento en base a los puntos del inventario y de la caracterización.

El artículo discute métodos y criterios disponibles actualmente provenientes de la investigación y de experiencias pasadas para inventariar, caracterizar y monitorear los recursos zoogenéticos, con vistas a asistir al desarrollo de una red más efectiva. Se da particular consideración a las nuevas herramientas y tecnologías. El objetivo de esta revisión incluye todas las especies ganaderas y sus antepasados salvajes así como especies salvajes relacionadas. Algunos ejemplos se han centrado en bovinos, ovinos, caprinos, porcinos y especies avícolas.

Keywords: *Descriptors, Relevant scales, Inventory, Characterization, Production systems, Phenotypic characterization, Molecular characterization, Genetic diversity.*

Conceptual framework

Genetic diversity within a livestock species is reflected in the range of breeds and populations and in the variation present within each.

The concept of the breed

The commonly used unit of reference of animal genetic diversity is the breed. Although the term "breed" is generally defined in terms of morphological, geographic, utility and genetic criteria, it is difficult to establish a definition that can be universally applied in both developed and developing countries. Definition of breed identities and characteristics requires at least a preliminary characterization of the breeds that are known to exist within a country. However, using the breed concept may lead to the exclusion of local populations that are not well described or not identified as breeds by the national authorities. In order to avoid missing data relevant to the efficient design of strategies for the management of animal genetic resources, it is useful to recall the different types of populations that are covered by the broad concept of the breed and that should be included in the inventory.

Traditional populations are mainly local and are considered to be adapted to their environment. They often exhibit a large phenotypic diversity (particularly for coat or plumage colour). They are managed by the farmers with low selection intensity, and are also affected by natural selection. Their genetic structure is mainly influenced by migration events and mutations, which would generally be counter selected in the wild. Population size is generally large.

Standardized breeds are selected on the basis of morphological traits, with a recognized "standard" breed descriptor, generally established by a community of breeders. They derive from traditional populations, but exhibit less phenotypic diversity as they are selected to meet minimum standards of phenotype. Their genetic structure may be influenced by important founder effects. Total population size may be very variable, depending on history and breeders' organization.

Selected breeds or commercial lines are characterized by an economic selection objective and the use of quantitative genetics methods. Molecular markers are often used, for instance for parentage testing. These populations derive from standardized breed or from traditional populations. Breeders are organized for pedigree and

performance recording. Total population size is generally large.

Derived lines arise from the use of specific breeding methods. Close inbreeding leads to highly specialized lines which exhibit low genetic variability. Conversely, composite breeds are derived from crosses between standardized breeds or selected lines, and exhibit a high level of genetic variability. Experimental selected lines used for research are part of this group, as they are generally derived from known breeds and selected for very specific traits. Transgenic lines would also belong to this group. Total population size is generally limited, except for composite breeds, which can form the basis of a new selection programme.

These different types of population may be easily identifiable in highly commercialized species such as cattle, pigs or chickens in Europe or Asia, for instance. The classification may not apply directly to other species such as camelids or geese, but can be considered a general framework for all types of domesticated populations.

In addition to these categories, wild ancestors and wild related species are also relevant for inventories. Indeed, spontaneous cross-breeding may still take place between wild relatives and livestock in interface areas. For example, the mountainous regions of north Viet Nam provide permanent contact between wild species and domesticated chicken populations. This “*free breeding*” increases introgression from wild genomes and plays an important role in maintaining a high genetic diversity and adaptation to particular conditions. Thus, these local populations should undoubtedly be considered in any inventory.

Descriptors (items) for inventory, characterization and monitoring

Primary indicators of animal genetic diversity should address both between-breed and within-breed components. Using breeds as the main indicator of total animal genetic diversity would miss out the important contribution of within-breed diversity. National authorities need to recognize the limitations of the breed concept and ensure that as much intraspecific genetic diversity as possible is accounted for in strategies for inventory, characterization and monitoring.

Typically, inventory, characterization and monitoring efforts will start by itemizing genetically distinct populations or “*breeds*”, the number of animals per population, and the number of farms

that keep these resources. As stated in *The State of the World's Animal Genetic Resources for Food and Agriculture* (SoW-AnGR), inventory, characterization and monitoring should include the identification, quantitative and qualitative description, and documentation of breed populations and the natural habitats and production systems in which they are embedded. Traits such as adaptation to a harsh environment, disease resistance, provision of environmental services, and product quality may receive specific attention depending on the context. Thus, it is necessary to describe the economic, social and environmental context in which the breeds are used, including cultural aspects of peoples' livelihoods. Furthermore, as socio-economic and environmental contexts evolve, criteria for evaluating breeds and their traits will also have to evolve.

Relevant scales

In principle, the strategy for inventory, characterization and monitoring should canvass all breeding populations across relevant production systems within a country, and include the sampling of representative animals to generate population descriptor data.

However, depending on the geographical distribution of the breeding population, the population size, breed risk status and economic significance, actions may be undertaken at different scales. For endangered and at-risk populations, they may be carried out at the level of individual animals, or populations of breeding animals in farms or stations. In the case of transboundary breeds, the exercise may involve intercountry collaboration, as in the case of commercial dairy and beef breeds included in multicountry breed evaluation programmes.

Inventory

A nationally mandated institution for inventory and monitoring is needed. At least in developing countries, this institution should set up a national mechanism to verify whether a particular breed or population represents a distinct unit of animal genetic diversity in the country, and as such needs to be included in the primary inventory.

In any country, it will be necessary to identify the number of farmers or communities that keep a particular population that is registered in the

national primary inventory. The national institution in charge of the inventory will collect data from government extension services, as well as from farmers' organizations – at any level from local communities to commercial companies. Involving livestock keepers and breeding organizations in the process has the added value of raising awareness about the value of the breeds in question. Bottom-up approaches also exist, in which a community describes a breed and brings it to the attention of the authorities. Confidentiality issues may affect inventories of commercial lines; breeding companies do not always agree to divulge numbers for the nucleus lines under selection.

In countries or areas where neither extension services nor breeding organizations can be identified to provide census data, on-field counting and systematic georeferencing may be set up as a special effort to improve inventory. Georeferencing will provide very useful information, as it allows geographical and climatic data to be linked to the distribution of breeds within a country.

Characterization

The first step of characterization is the primary assessment or baseline survey, which should include collection of data on population size and structure, geographical distribution, production systems in which the breed is found, phenotypic attributes (physical features, performance levels and any unique features), historical development of the breed through exchange, upgrading and selection, and the genetic connectedness of populations when these are found in more than one country (e.g. the N'Dama cattle breed of West Africa). The within-population genetic diversity is measured both at the phenotypic level (phenotypic breed diversity) and at molecular level; the two are complementary. All these data are needed to inform decisions on the utilization, improvement and conservation of the population.

Production systems and social organizations

As noted in the SoW-AnGR, the term “breed” is often accepted as a cultural rather than a biological or technical term. Hence, in order to depict direct and indirect use values of breeds, they need to be characterized in the context of the production systems and social structures in which they are

used. The objective is to allow comprehensive input/output analysis of the genetic resources in the context of the agro-ecosystems of which they form a part. The environmental impact of a breeding population should also be considered as part of the characterization of the production system. Such data can be collected by survey. FAO has already developed simplified formats for data collection for mammals and poultry. The cost and time needed for such surveys should not be underestimated, but they could benefit from being linked to training programmes, e.g. for MSc and PhD students.

Surveys will be organized differently depending on the institutional background. In developed countries, where commercial and conservation farms keep registers of individual animals and their pedigrees, structured surveys can be used to collect information on production systems and the environment. The procedure should take advantage of current data collection systems and additional costs should be quite limited. Yet, measurements related to environmental impact of breeds and their production systems are generally not included in routine procedures and specific actions are needed to collect such information.

In countries where such data are not regularly recorded, specific surveys need to be set up. For traditional communities in pastoral and farming production systems, participatory surveys and structured interviews can be used to generate data on breeding objectives, breed and trait preferences and production system constraints. In the context of traditional breeds, these descriptions give insights into the multitude of functions and services that breeds provide for their keepers. Statistical sampling procedures can be applied to study localities, farms and individual animals once the sampling framework is defined.

In situations where limited documented information on breed identification and characteristics is available, extensive exploratory surveys may be necessary. However, exploratory surveys have limitations; the facts generated are highly subject to the biases of questionnaire respondents. Thus, steps need to be taken to ground-truth and cross-check findings using complementary procedures such as key-informant interviews, focus-group discussions and reporting-back sessions with respondent communities. Consequently, these surveys become demanding in terms of time, skilled personnel and financial resources. This has been observed, for example, in livestock breed surveys in Zimbabwe and Ethiopia.

Phenotypic characterization

The different phases of characterization involve morphological attributes, biometrical indices, production levels (growth, reproduction, milk, egg, fibre, traction) and specific adaptations, including survival. Morphological variants may be associated with known genes (coat colour, morphological mutations) and will benefit from their molecular characterization.

It is important that phenotypic measurements (biometrics and performance) should not focus on means or averages alone, but also account for variations. It is the variation that provides the basis for conservation and for present as well as future utilization. For this reason, a large proportion of the population should be included in the assessment of performance.

Performance may be assessed either by direct recording of the animals or by exploiting information that is available in published literature, extension service field reports and reports of breeding units and organizations. Performance testing may be done either on-farm or in testing stations.

On-farm testing

When genetic evaluation is performed utilizing national records from on-farm testing, the associated data can be made available for characterization, and breeding values should be incorporated. However, this is not feasible for pig or chickens breeding schemes run by companies which will not share their data.

For species or countries where there is no national on-farm testing, specific action to collect

on-farm data is required. Technicians should be trained to collect morphological data. Pictures should be taken utilizing a tape measure to document phenotypic variability as thoroughly as possible. In traditional communities, indigenous knowledge and practices associated with breed identity and unique utility should also be compiled along with population performance descriptors. A variety of relevant participatory methods exist, including methods that allow livestock keepers to rank breed and trait preferences, including traits with non-market values. Simple criteria such as sales and survival rates provide valuable information.

When georeferencing of phenotypic data is available, further biophysical data from the environment (climate, soil, vegetation cover, water availability, type and level of disease challenges) can be overlaid, and joint analysis in GIS (geographical information system) will help to assess adaptability traits.

On-station testing

On-station characterization makes it possible to evaluate breed performance and potential in a relatively defined and controlled production environment. The limitations are that animals may not necessarily be adapted to the controlled environment and that some traits such as grazing behaviour and response to environmental stressors cannot be measured. Thus, the specific advantages of a local population may not be recognized. Indeed, it is currently difficult to find objective criteria to describe the adaptation of local populations to specific climatic or feed conditions. Research is needed in this field – identifying morphological and

Box 1. The Management of Farm Animal Genetic Resources in the SADC Sub-Region project

The implementation of the animal genetic resources characterization project for the Southern African Development Community between 2000 and 2004 demonstrated that the human, financial and networking resources of public institutions and international research and development organizations can be harnessed to run large-scale exploratory surveys. In this particular case, the United Nations Development Programme provided funding; FAO and the International Livestock Research Institute provided expert advice and guidance in the design, execution and evaluation of breed characterization surveys.

physiological predictors for heat tolerance or walking ability, for instance. Moreover, such unknown adaptive traits are usually not captured in a standardized research protocol; new protocols need to be developed. Conversely, a controlled environment allows more precise measurement of individual performance, pedigree recording and estimation of genetic parameters, and provides opportunities to undertake multiple comparisons (breeds and production environments) across stations, so as to assess genotype by environment interactions. A positive aspect of on-station characterization is that it may contribute to the establishment of a nucleus population and contribute to the conservation of the resource being characterized.

Advanced phenotyping

Product quality is generally considered by breeding organizations using precise descriptors, which are defined according to the destination of the product, taking into account indications from nutritionists and food processors. For instance, fat percentage in milk is analysed in terms of fatty-acid composition, and protein percentage can be detailed according to the different types of caseins. Furthermore, systems have been set up in Europe to associate a product with a certificate of origin, such as Protected Geographic Indication¹ and Protected Designation of Origin², which generally include the breed of origin of the product (Box 2). The same concept is applied for goat meat in Argentina (Box 3). In many African and Asian countries, specific products are also associated with local breeds, and accurate description of the product should be undertaken in order to better define it and, consequently, characterize the breed. This requires capacity-building for the definition of product quality requirements, and the establishment of an official system for certifying that the product and production methods meet these requirements.

Disease resistance is a high priority for several reasons: local breeds survive in harsh environments and this needs to be better understood; epidemics are major threats for all animal genetic resources across the world; climatic change is likely to increase the spread of tropical diseases to temperate areas. In addition to claims that local breeds are adapted and resistant, scientific evidence has been obtained in several instances (examples are reported in the SoW-AnGR). The effect is particularly well documented for parasitic diseases,

which are very prevalent in tropical areas, with local breeds maintaining a better performance in the presence of parasites and/or exhibiting lower levels of parasite infestation. Generally, this condition is better described as tolerance, a typical example being trypanotolerance in cattle. Generally, more data are needed on exposure and response of animal populations to parasites, viruses and bacteria. One delicate question involves possible confusion between resistance and a healthy-carrier state for a given pathogen. True resistance, in which the host does not allow the pathogen to disseminate, is the objective of most research studies in developed countries. This is consistent with the assumption that it will be possible to eradicate the pathogen. However, this seems unrealistic for tropical parasites. Thus, research is focusing on defence mechanisms, in order to better understand the permanent race between hosts and pathogens. Furthermore, epidemiological studies suggest that pathogens may adapt more easily to uniform genotypes, and that genetic variation of the host is one key to limiting pathogen expansion. Thus, cooperation between Northern and Southern countries is needed to better characterize the potential usefulness of animal genetic resources for disease control. This may benefit from progress in genomics and the identification of genes for resistance to major diseases, as well as in the understanding of general immune response.

Molecular characterization

The impressive development of molecular tools in the past 20 years benefits the characterization of animal genetic resources in many ways – which are already well documented in the SoW-AnGR. It is important that countries are aware of what questions molecular tools can or cannot answer at

¹Protected geographical indication: the name of a region, specific place or country describing a product originating in that region, specific place or country and possessing a quality or reputation which may be attributed to the geographical environment with its inherent natural and/or human components.

²Protected designation of origin: the name of a region, specific place or country referring to a product originating in that region, specific place or country and whose quality or other characteristics are essentially or exclusively due to a particular geographical environment.

Box 2. Differentiation in chicken meat production in France

The French production of chicken meat is differentiated into several categories: standard broiler (SB), label chicken (LB), certified chicken (CF), organic chicken, and *Appellation d'Origine Contrôlée* (AOC = Protected Designation of Origin) for the Bresse breed only. Whereas LB production represented nearly 100 million chickens in 2002 (www.synalaf.fr), the Bresse AOC represented 1.4 million chickens raised only in the Bresse geographical area as defined by law. The LB category was created in 1965, to promote product quality throughout the production process. The LB and CF legal definitions do not require reference to a particular breed, but only slow-growing lines are eligible. These slow-growing lines are generally characterized by a coloured phenotype, easy to distinguish from the white plumage of SB. The philosophy of AOC is quite different since it defines a geographical district which is characterized by specific features of the natural conditions and production system. For the Bresse, the district was defined as early as 1936 and the protection of the name "*Volaille de Bresse*" was enshrined in law number 57-866 on August 1, 1957. The Bresse breed standard includes white plumage and blue shanks, which is a rare association among French poultry breeds. A fixed set of growing conditions (density, open-air access, type of feed) must be applied for at least 9 weeks, starting from 5 weeks of age. Then, the finishing period, slaughtering conditions and carcass processing are strictly regulated. The minimal age at slaughter for the Bresse is 112 days, whereas it is 84 days for LB and 39 days for SB. Tasting panels are regularly organized to check the meat quality. The selection procedure for the Bresse breed has also been strictly regulated and is managed by a selection centre (CSB) which is working in close collaboration with farmers. The Bresse breed is the only local French chicken breed the population size of which has not decreased since 50 years, and credit must be given to the AOC for this success (Verrier *et al.*, 2005).

Box 3. Differentiation of goat meat in Argentina

The traditional goat production system from North Neuquén (Patagonia, Argentina), developed by transhumant goat keepers is a marginal system with low economic input and fragile environment but with a high cultural capital, an adapted genetic resource and a product with high reputation but not differentiated. To overcome this situation the application of a Geographical Indication was developed. This process was based on the organization of the local goat meat marketing chain and the description of technological properties of the product of the Neuquén Criollo breed. The chain actors developed a common vision about the system and its identity, which is reflected in the Protocol of the Designation of Origin of the "*Criollo Kid of North Neuquén*". A study on the product's typical characteristics and quality has contributed to define quality indicators and traceability of the product. As a result, the goat keepers' organizations have been empowered, a common ground of communication has been established enhancing the understanding level among local actors, which was previously not existent. This has reinforced regional development and given projection to sustainability of the system and genetic resource (Pérez Centeno *et al.*, 2007)

present, and how this may change in the future. It is also important to consider that the broad array of tools that is available in the case of the "*big five*" species is not available for species with a more limited geographic distribution, but which should not be neglected.

Some practical considerations

The first step is to collect samples of sufficient quality from representative animals of the population to be described – either a well-known breed or a non-described population (FAO, 1993).

The general recommendation is to sample 30 to 50 unrelated individuals, in flocks or herds covering a wide geographical area, taking into consideration historical exchange of breeding stocks, and agro-ecological zones as possible barriers to gene flow between populations. These are minimum numbers. Ideally, half the sample should be females and half males. A clear description of the sampling procedure is needed, both for immediate use of the samples and to allow the samples to be used for future studies. Ideally, the animals sampled should also have been subject to phenotypic characterization.

The required DNA quality depends on the intended future use. Several protocols are available, and good quality should be the aim. Blood or ear-tissue samples are ideal for typing nuclear and mitochondrial DNA (mtDNA) markers, but such sampling is not always accepted by the farmer. It is possible to extract sufficient DNA from hair bulbs to allow the typing of microsatellite DNA markers, but such samples are not easy to work with in the case of mtDNA and other markers. Extraction kits are expensive, but should provide repeatable quality. Manual extraction needs trained personnel. Whatever the protocol, DNA quality should be checked before samples are used or sent for genotyping.

Molecular markers involve genomic and mtDNA loci. Microsatellite markers are most commonly used because they are multi-allelic and numerous, and can be genotyped on automatic machines. New microsatellite marker sets of 20 to 30 loci per species recommended by the International Society of Animal Genetics (ISAG)/FAO Standing Committee are available for most species (FAO/ISAG, 2004). It is highly recommended that a core set of a minimum of 15 markers be included so as to allow comparative studies across countries. Merging genotype data sets produced in different laboratories has proven to be possible though quite challenging. Exchange of reference samples between laboratories is mandatory, and training of technicians to score the genotypes following the same procedure is necessary. Statistical methods for meta-analysis are also under development to make the best possible use of available data in order to merge all information and facilitate international comparisons. The problem of standardizing microsatellite typing is not encountered in the case of typing single nucleotide polymorphisms (SNPs) because technologies are available to provide standardized reading of SNPs and to produce data that can be merged between laboratories. SNPs are discussed in more details in Box 7.

Assessment of genetic diversity with anonymous DNA markers

The first question that anonymous DNA markers can answer relates to the diversity level within a population, which can be described by number of alleles, number of private alleles, or observed and expected heterozygosity. Generally, the diversity level of domestic breeds/populations has been found to be lower than that of wild relatives and ancestors. Diversity can be expected to have gradually declined during the dispersal of livestock populations from their centres for domestication or origin to their current locations, mainly as a result of random genetic drift. However, this pattern may be distorted by the introduction of exotic breeds, cross-breeding between populations, admixture of populations from different centres of domestication and human selection. Thus a careful examination of the population's history is warranted. It is also well known that heterozygosity estimates are not so sensitive to the change of number of alleles, particularly in the case of multi-allelic microsatellite markers. Therefore, the adjusted mean number of alleles according to sample sizes could be a better parameter to measure genetic diversity within breeds or populations.

Methods have been proposed, and are still under development, to estimate the effective population size of a breed or population from molecular data, particularly from linked markers. It is also possible to detect departure from the equilibrium state either due to excessive inbreeding or to population fragmentation in subgroups that have few or no exchanges between them. Thus, DNA markers can be used for monitoring conservation programmes aimed at avoiding inbreeding, genetic bottlenecks and fragmentation. Furthermore, they can be used to identify "livestock biodiversity hotspots" as priority areas for conservation of indigenous livestock populations. Typically, populations containing large variation at anonymous loci are expected also to exhibit large variation for functional traits. Thus, DNA markers could be most useful in cases where little information on population history is available. However, anonymous markers do not at present provide a reliable prediction of phenotype; they do not replace performance measurements and should not be used alone to make conservation decisions.

The second area in which DNA markers provide useful answers includes questions of relatedness between populations, detection of admixture, introgressions and breed identity. Between-breed variation may be described by genetic differentiation indices, such as F_{ST} for which

statistical significance can be calculated in order to conclude whether or not genetic differentiation takes place between pairs of populations. Allelic frequencies for molecular loci also provide the basis on which to calculate genetic distances. As mentioned in the SoW-AnGR, phylogenetic reconstruction of the evolution of breeds or populations is not well adapted to the dynamics of domesticated populations, which do not diverge strictly from a common ancestor and may include cross-breeding, admixture and introgression events in their histories (Box 4). In the case of selected lines derived from the same breed, phylogenetic reconstruction with neighbour-joining tree can reveal clustering (Box 5).

Multivariate methods offer a different approach, which unlike phylogenetic trees, does not rely on any evaluative assumption. Bayesian clustering has been shown to be very efficient for the assignment of individuals to breeds or populations and as a means to detect population structure and admixture without any prior information on population ancestry. Recent results obtained in chicken populations, both traditional and commercial lines, showed that more than 90 percent of individuals could be assigned to their true breed of origin according to their genotypes for microsatellite markers (Box 6).

Thus, DNA markers allow the definition of the genetic entity behind the breed. This can clarify the procedure of inventories and identify the base population for conservation programmes. Knowledge of the molecular identity of certain breeds or populations may also be used to establish biological identification systems for certification and traceability of living animals and derived products.

In addition to nuclear markers, both mtDNA and markers from the Y chromosome of mammals provide additional information on the history of domestication and introgression events. Very interesting results have been obtained for ruminants in this respect. These data may also be useful because they shed light on peculiar adaptive traits that these populations may have accumulated over time.

Known genes and functional diversity

Progress in genome annotation and quantitative trait loci (QTL) programmes has led to the identification of many candidate genes that are likely to influence traits of interest. QTL programmes and genome databases are available for the “big five” species. Comparative genomics may also facilitate the assessment of functional diversity by transferring knowledge between species. Significant progress has been made in the molecular identification of genetic abnormalities as well as major genes affecting meat quality or muscular growth. Some causal mutations, as well as diagnostic methods for these mutations, have been patented, and new alleles may be present in some indigenous populations. Therefore, the issue of intellectual property (IP) arising from the discovery of functional diversity and exclusive or non-exclusive use of this IP has to be addressed. As far as QTL are concerned, finding genes responsible for the quantitative effect on the performance is still rare. Furthermore, the effect of a QTL region may depend on the genomic background: epistatic interactions are known to take place, so that a given QTL region identified in one population may not be

Box 4. Sheep biodiversity

The ECONOGENE project combined a molecular analysis of biodiversity, socio-economics and geostatistics to address the diversity and conservation of small ruminants in marginal agro-ecosystems. The population structure and genetic diversity of 57 European and Middle Eastern sheep breeds from 15 countries were analysed by typing 31 microsatellite markers, thereby extending the available knowledge of sheep diversity at the molecular level. The domestication centre for sheep lies in the Near and Middle East, and the results showed high levels of genetic variation among Middle-Eastern and South-eastern European breeds. The analysis of markers and of the spatial distribution revealed the occurrence of two clusters, one with north-western European breeds and the other with Middle-Eastern/southeastern European breeds.

Source: Peter *et al.* (2007)

Box 5. Pig biodiversity

The European PigBiodiv project used 50 microsatellite markers to assess the between- and the within-breed genetic diversity for a set of 59 pig breeds. The resulting structure of eight groups (bootstrap) showed within-breed clustering of pig lines. The national populations of major breeds and the commercial lines were clustered around their breeds of reference (Duroc, Hampshire, Landrace, Large White and Pietrain) in most cases. The Meishan breed represented a specific outgroup. Local breeds did not group into one cluster and appeared to be scattered within the global frame. Using only 18 markers decreased the reliability of the clustering, particularly for the Landrace breed.

Source: San Cristobal *et al.*, (2006).

Box 6. Breed assignment with anonymous markers

The AvianDiv project used 27 microsatellite markers to genotype 30 animals from 20 chicken populations, ranging from the wild ancestor to highly selected commercial lines. After an analysis with the “*Structure*” software, it was possible to assign birds to their correct breeds with 90% efficiency using 12 markers. After 24 markers, efficiency remained close to 97%. Correct assignment of commercial birds to their true line of origin was the most difficult and required all markers.




Figure adapted from Rosenberg *et al.* (2001).

relevant for another population. An integrated strategy using molecular markers would be to map the genetic diversity among indigenous livestock breeds/populations to test hypotheses about which of them may carry unique QTL for disease resistance.

The transcriptomics approach has enabled the exploration of gene expression patterns for thousands of genes simultaneously. But this approach has not been used to a large extent for diversity studies. It raises a number of questions, regarding the tissue to be sampled, the stage of sampling, and very often requires animals to be slaughtered. The best examples deal with the study of disease resistance, where multigenic expression patterns can efficiently describe the mechanisms involved in defence responses, and can identify relevant differences between breeds. Thus, more experimental data are needed before gene expression patterns are incorporated into characterization.

The final effector molecules are proteins. Proteomics has also made significant progress, although it raises delicate methodological issues and has not yet been applied to the characterization of genetic diversity. Research is needed to improve this approach which may open the way to intensive phenotyping.

Prospects with high-density single nucleotide polymorphisms

The full genome sequence is or will soon be available for chickens, cattle, pigs and sheep (and goats) and large numbers of *single nucleotide polymorphisms* (SNPs) are becoming available (Box 7). As compared to microsatellites, mtDNA and polymorphisms of known genes, the use of high-density SNP markers offers quite new perspectives: these markers are so numerous that they may unravel the fine structure of the genome and identify chromosomal segments showing selection signatures. This will greatly improve our knowledge of population genetic make-up. Large-scale SNP typing has already started in selection programmes for cattle (dairy and beef) and chickens. Performance recording is still necessary at crucial steps of characterization programmes, to define the association between genotypes and desired phenotypes. Alleles, haplotypes or quantitative trait nucleotides (QTN) could then be used to estimate a breeding value genome-wide. This represents one step forward from the current marker-assisted selection programmes, which track

a limited number of QTL regions to whole genome selection. Thus, the whole organization of data collection may change in the coming years. FAO's information system will have to be updated to take into account these trends.

Advanced inventory and monitoring

All countries need an active inventory and monitoring strategy for their animal genetic resources – to better understand, use, develop, maintain, conserve and access these resources. The *Global Plan of Action for Animal Genetic Resources* recognizes the need to have a country-based strategy so that activities for inventory and monitoring can be linked and coordinated with relevant country-level action plans such as agricultural censuses or livestock population surveys. Indicators are needed for population trends, breed risk status and changes in the production environment. Apart from the opportunity of carrying out meta-analysis of nationwide data to establish trends and information gaps, country-based strategies also encourage the establishment of information databases of animal genetic resource inventory which can provide a comprehensive source of information for research, development of breeding strategies, conservation programmes, policy frameworks and even training.

Monitoring driving forces and describing production environments

Production environments are dynamic, albeit at different scales and rates. As discussed in the introductory paper to this series, the major drivers of change that are of relevance to the management of animal genetic resource diversity are population growth, urbanization, and the associated changes in the structure and volume of demand for livestock products, globalization, climate change and global health hazards such as avian influenza. All of these drivers should be monitored to predict future scenarios and allow improved preparedness to meet future challenges.

Indicators related to production environment were elaborated at an FAO expert consultation which met in Armidale, Australia in 1998. Five main criteria (climate; terrain; disease, disease

Box 7. A new approach of genome diversity with SNPs

Large numbers of SNPs have been or will be generated as companion programmes of the genome sequencing efforts undertaken for the “big five” species. SNPs are mainly bi-allelic due to the low frequency of mutations. Therefore, only a higher number of SNPs can achieve information content comparable to that obtained using a given number of microsatellite markers. Characterization of the same set of ten chicken breeds using 29 microsatellite markers and 145 SNPs confirmed that increasing the number of SNPs had a higher impact on the reliability of the results than increasing the sample size (Hillel *et al.*, 2007). Heterozygosity and allelic-richness estimates obtained for SNP markers exhibit a lower order of magnitude as compared to microsatellite markers, with values in the range of 0.34 and 1.94, respectively, across a set of Holstein-Friesian bulls (Zenger *et al.*, 2007). It is likely that systematic molecular studies of animal genomes will use SNPs and handle questions of selection and management of genetic diversity at the same time. Cost of typing SNPs is steadily decreasing, but SNPs are valuable only when they are very numerous (e.g. more than 3 000). Therefore, the absolute cost of typing is still a matter to be considered.

complexes and parasites; resource availability; and management) were identified as the basis for the characterization of production environments for all livestock species, with three to seven indicators for each criterion (FAO, 1998). The framework is demanding in terms of resource requirements and needs to be operationalized, but can be used to select priority criteria and indicators that better meet specific needs.

The application of georeferencing tools can make a major contribution to improving the scope and scale of advanced inventory and monitoring both at country and global levels.

Monitoring animal populations

Through their ratification of the Convention on Biological Diversity, countries are committed to inventory and monitoring of the status of their animal genetic resources. However, country reports prepared during the SoW-AnGR reporting process show that national inventories have either not been carried out or are still incomplete.

Monitoring requires regular checking of population status, and the evaluation of trends in the size and structure of breed/populations, their geographical distribution, risk status and genetic diversity. If breeders' associations or other groups interested in breed maintenance and promotion exist, it may be possible to update the inventory annually. In the absence of such groups, the mandated national institutions must ensure that

periodic assessment of breed status are carried out ideally on annual or biennial basis, or at least at intervals of one generation for the species in question. This would require comprehensive updating at intervals of about eight years for horses and donkeys, five years for cattle, buffaloes, sheep and goats, three years for pigs and two years for chickens. Once a breed has been identified as at risk, a more intensive monitoring programme is needed on an ongoing basis.

As noted in the SoW-AnGR, monitoring can be an extremely expensive aspect of the management and should take as much advantage as possible of existing resources and activities.

Defining indicators for animal genetic diversity

A compromise has to be found between the ideal list of indicators needed to provide accurate information, and the cost of collection and ease of interpretation. As stated by OECD (2001), four main criteria may be used to assess the value of indicators: policy relevance, analytical soundness, measurability and interpretation. In general, a small number of indicators is preferable in terms of measurability and communication, but relevant information needs to be captured in order to support sound decisions.

The existing FAO definitions of breed risk status (extinct, critical, endangered and not at risk) are

based on numbers of breeding females and males, but do not relate to how matings are handled (e.g. random or high selection intensity within breeds, use of crossbreeding). Major drivers of change can lead to rapid changes in the population size and structures of locally adapted breeds. Regular monitoring is therefore required, at least for those breeds classified as critical or endangered. At present, most national livestock censuses do not contain breed-level data; therefore, regular reporting of breed population numbers does not usually take place. In addition to population size, the number of farms and number of breeding organizations could be considered. The number of breeding males should be made available. Such a monitoring scheme can serve as the basis for national early warning, so that timely management interventions can be planned. Monitoring programmes need to ensure that feedback is provided to farmers, researchers and other stakeholders.

Recent research suggests that several issues need to be taken into account for the development of indicators for animal genetic diversity:

- the concept of the breed as a genetic entity for measuring diversity would benefit from the use of molecular markers for the assignment of individuals to breeds .
- the assessment of breed risk status should not rely on population size alone, but would benefit from more accurate parameters calculated on the basis of extensive pedigree analysis, such as inbreeding coefficients of current breeding animals, or the number of ancestors with a cumulated contribution of 50 percent of the total gene pool.
- in the absence of pedigree recording, loss in diversity may be monitored using molecular markers, particularly on the basis of the adjusted mean number of alleles calculated for reference sets of microsatellite markers.
- occurrence of introgressions or fragmentations may be monitored with molecular markers, combining nuclear markers and mtDNA, provided that reference data sets for a range of populations are available for comparative analysis within a country or region.

Target values for country-based early warning tools are yet to be developed. It is essential to establish both baseline (inventory) and follow-up (monitoring) assessments to effectively inform decision-making in the management and utilization of animal genetic diversity. Monitoring of diversity should address both the level of between-breeds

diversity, with setting up conservation programmes for endangered populations, and the level of within-breed diversity with updating rules for the genetic management of the population (Fikse & Philippson, 2007).

Conclusions and recommendations

Inventory and characterization of animal genetic resources should be an iterative process. Regular updates are necessary, because animal genetic resources are exposed to strong driving forces, both from the viewpoint of production systems and emerging technologies.

Data from all types of populations are relevant for the Domestic Animal Diversity Information System (DAD-IS) managed by FAO. In order to minimize information gaps, the concept of the breed should be understood in a broad sense. Inventory should include criteria to assess within-breed diversity. National databases have to be set up and should be coordinated at a regional level and with FAO, in order to facilitate the comparisons and the updating of information.

A comprehensive description of production environments is needed in order to better understand the comparative adaptive fitness of specific animal genetic resources. It will also help to identify threats and options for the management of these resources.

On-field and on-station phenotypic characterization are complementary. Performance data should focus on variability as much as possible and not only include means. Defence mechanisms against pathogens should be a priority, given the significance of the threats posed by epidemics and climate change.

It is likely that microsatellite markers will remain the first choice for the analysis of genetic diversity in many domestic species in the near future. Steps should be taken to support comprehensive multicountry studies, and to facilitate meta-analysis. On the technical side, this requires improved exchange of reference samples and standardization of genotyping procedures. From the methodological perspective, appropriate models need to be developed and tested.

Anonymous markers provide a range of information, from population history to breed identity. However, the number of markers which is sufficient for population genetics studies does not allow any prediction of performance. Thus,

available molecular genetic markers should be used together with phenotypic data.

Recent technologies for large-scale gene expression studies and high-throughput SNP genotyping are likely to greatly modify characterization tools, with the prospect of better connecting phenotypes to genotypes. Costs are still too high for these procedures to be used in systematic surveys of genetic diversity, but in species such as cattle and chickens in which the genome has been sequenced, these technologies are likely to rapidly prove their usefulness in achieving a comprehensive approach to the assessment of genetic diversity.

Data on production systems, phenotypes and molecular markers should be used altogether in an integrated approach to characterization. Decisions regarding conservation should incorporate all descriptors. Conserving without documenting would be useless. National authorities should be aware that sharing information and data is essential to support cost-effective decisions in the management of animal genetic resources.

List of references

- FAO. 1993. Secondary Guidelines for Development of National Farm Animal Genetic Resources Management Plans. Measurement of Domestic Animal Diversity (MoDAD): Original Working Group Report. Rome.
- FAO. 1998. Report: Working group on production environment descriptors for farm animal genetic resources. Report of a Working Group, held in Armidale, Australia, 19–21 January 1998. Rome.
- FAO. 2007. The State of the World's Animal Genetic Resources for Food and Agriculture, edited by B. Rischkowsky & D. Pilling. Rome, (also available at <ftp://ftp.fao.org/docrep/fao/010/a1250e/a1250e.pdf>), pp. 511.
- FAO/ISAG. 2004. Secondary Guidelines. Measurement of Domestic Animal Diversity (MoDAD): New recommended microsatellite markers. (also available at <http://dad.fao.org/cgi-bin/getblob.cgi?sid=-1,50005882>).
- Fikse, W.F. & Philipsson, J. 2007. Development of international genetic evaluations of dairy cattle for sustainable breeding programs. *Animal Genetic Resources Information Bulletin*, 41, 29–44
- Hillel, J., Granevitze, Z., Twito, T., Ben-Avraham, D., Blum, S., Lavi, U., David, L., Feldman, M.W., Cheng, H. & Weigend, S. 2007. Molecular markers for the assessment of chicken biodiversity. *World Poultry Science Journal*, 63: 33–45.
- International Chicken Polymorphism Map Consortium. 2004. A genetic variation map for chicken with 2.8 million single-nucleotide polymorphisms. *Nature*, 432(7018): 717–722.
- OECD. 2001. OECD Expert meeting on agri-biodiversity indicators. Summary and recommendations. 5–8 November 2001, Zurich, Switzerland. (also available at ftp://ftp.fao.org/agl/agll/ladadocs/oecd_zurich.doc).
- Oldenbroek, K. (Ed.). 2007. Utilization and conservation of farm animal genetic resources. Wageningen, the Netherlands, Wageningen Academic Publishers, pp. 103–130.
- Pérez Centeno, M., Lanari, M.R., Romero, P., Monacci, L., Zimerman, M., Barrionuevo, M., Vázquez, A., Champredonde, M., Rocca, J., López Raggi F. & Domingo, E. 2007. Puesta en valor de un sistema tradicional y de sus recursos genéticos mediante una Indicación Geográfica: El proceso de la Carne Caprina del Norte Neuquino en la Patagonia Argentina. *Animal Genetic Resources Information Bulletin*, 41, 17–24
- Peter, C., Bruford, M., Perez, T., Dalamitra, S., Hewitt, G., Erhardt, G. & The Econogene Consortium, 2007. Genetic diversity and subdivision of 57 European and Middle-Eastern sheep breeds *Animal Genetics*, 38: 37–44
- Rosenberg, N.A., Burke, T., Elo, K., Feldman, M.W., Freidlin, P.J., Groenen, M.A.M., Hillel, J., Maki-Tanila, A., Tixier-Boichard, M., Vignal, A., Wimmers, K., Weigend, S. 2001. Empirical evaluation of genetic clustering methods using multilocus genotypes from 20 chicken breeds. *Genetics*, 159, 699–713.
- SanCristobal, M., Chevalet, C., Haley, C.S., Joosten, R., Rattink, A.P., Harlizius, B., Groenen, M.A.M., Amigues, Y., Boscher, M.Y., Russell, G., Law, A., Davoli, R., Russo, V., Désautés, C., Alderson, L., Fimland, E., Bagga, M., Delgado, J.V., Vega-Pla, J.L., Martinez, A.M., Ramos, M., Glodek, P., Meyer, J.N., Gandini, G.C., Matassino, D., Plastow, G.S., Siggens, K.W., Laval, G., Archibald, A.L., Milan, D., Hammond, K.

& **Cardellino, R.** 2006. Genetic diversity within and between European pig breeds using microsatellite markers. *Animal Genetics*, 37: 189–198.

Verrier, E., Tixier-Boichard, M., Bernigaud, R. & Naves, M. 2005. Conservation and value of local livestock breeds: usefulness of niche products and/or adaptation to specific environments. *Animal Genetic Resources Information Bulletin*, 36, 21–32

Zenger, K.R., Khatkar, M.S., Cavanagh, J.A.L., Hawken, R.J. & Raadsma, H.W. 2007. Genome-wide genetic diversity of Holstein Friesian cattle reveals new insights into Australian and global population variability, including impact of selection. *Animal Genetics*, 38: 7–14.

Bibliography for further reading

Ayalew, W., van Dorland, A. & Rowlands, J. (Eds). 2004. Design, execution and analysis of the livestock breed survey in Oromia Regional State, Ethiopia. Nairobi, OADB (Oromia Agricultural Development Bureau), Addis Ababa, and ILRI (International Livestock Research Institute), pp. 260.

Berthier, D., Chantal, I., Thévenon, S., Marti, J., Piouemal, D. & Maillard, J.C. 2006. Bovine transcriptome analysis by SAGE technology during an experimental *Trypanosoma congolense* infection. *The Annals of the New York Academy of Sciences*, 1081: 286–299.

Cunningham, E.P. & Meghen, C.M. 2001. Biological identification systems: genetic markers. *Revue scientifique et technique de l'Office internationale des epizooties*, 20: 491–499.

Doherty, M.K., McLean, L. & Beynon, R.J. 2007. Avian proteomics: advances, challenges and new technologies. *Cytogenetics and Genome Research*, 117: 358–369.

Drucker, A., Gomez, V. & Anderson, S. 2001. Economic valuation of farm animal genetic resources: a survey of available methods. *Ecological Economics*, 36 (1): 1–18.

Eaton, D., Windid J., Hiemstra, S.J., van Heller, M., Trach, N.X., Hao P.X., Doan, B.H. & Hu, R., 2006. Indicators for livestock and crop biodiversity. Centre for Genetic Resources, CGN/DLO foundation, report 2006/05, Wageningen, the Netherlands.

FAO. 2006. A System of Integrated Agricultural Censuses and Surveys, Volume 1, World Programme for the Census of Agriculture 2010, (SDS No. 11). (also available at www.fao.org/ess/ess/census/default.asp).

FAO. 2006. The role of biotechnology in exploring and protecting agricultural genetic resources, edited by J. Ruane & A. Sonnino. Rome. (also available at www.fao.org/docrep/009/a0399e/a0399e00.htm).

Freeman, A.R., Bradley, D.G., Nagda, S., Gibson, J.P. & Hanotte, O. 2006. Combination of multiple microsatellite datasets to investigate genetic diversity and admixture of domestic cattle. *Animal Genetics*, 37: 1–9.

Gandini, G.C., Ollivier, L., Danell, B., Distl, O., Georgoudis, A., Groeneveld, E., Martyniuk, E., van Arendonk, J.A.M., & Woolliams, J.A. 2004. Criteria to assess the degree of endangerment of livestock breeds in Europe. *Livestock Production Science*, 91:173–182.

Gibson, J.P. & Bishop, S.C. 2005. Use of molecular markers to enhance resistance of livestock to disease: a global approach. *Revue scientifique et technique de l'Office internationale des epizooties*, 24: 343–353.

Gibson, J.P., Ayalew, W. & Hanotte, O. 2007. Measures of diversity as inputs for decisions in conservation of livestock genetic resources. In: D.I. Jarvis, C. Padoch & H. D. Cooper (Eds). *Managing biodiversity in agricultural ecosystems*. New York, Columbia University Press, pp. 117–140.

Hanotte, O., Bradley, D.G., Ochieng, J., Verjee, Y., Hill, E.W. & Rege, J.E.O. 2002. African pastoralism: genetic imprints of origins and migrations. *Science*, 296(5566): 336–339.

Kadarmideen, H.N., von Rohr, P. & Janss, L.L.G. 2006. From genetical genomics to systems genetics: potential applications in quantitative genomics and animal breeding. *Mammalian Genome*, 17: 548–564.

Moazami-Goudarzi, K. & Laloe, D. 2002. Is a multivariate consensus representation of genetic relationships among populations always meaningful? *Genetics*, 162: 473–484.

.....

Mburu, D.N., Ochieng, J.W., Kuria, S.G., Jianlin, H., Kaufmann, B., Rege, J.E.O. & Hanotte, O. 2003. Genetic diversity and relationships of indigenous Kenyan camel (*Camelus dromedarius*) populations: implications for their classification. *Animal Genetics*, 34: 26–32.

Muchadeyi, F.C., Eding, H., Wollny, C.B.A., Groeneveld, E., Makuza, S.M., Shamseldin, R., Simianer, H. & Weigend, S. 2007. Absence of population substructuring in Zimbabwe chicken ecotypes inferred using microsatellite analysis. *Animal Genetics*, 38: 332–339.

Pritchard, J.K., Stephens, M. & Donnelly, P. 2000. Inference of population structure using multilocus genotype data. *Genetics*, 155: 945–959.

Tano, K., Kamuanga, M., Faminow M.D. & Swallow, B. 2003. Using conjoint analysis to estimate farmers' preferences for cattle traits in West Africa. *Ecological Economics*, 45(3): 393–408.

Tuggle, C.K., Wang Y.F. & Couture, O. 2007. Advances in swine transcriptomics. *International Journal of Biological Sciences*, 3: 132–152.

Vignal, A., Milan, D., SanCristobal, M. & Eggen, A. 2002. A review on SNP and other types of molecular markers and their use in animal genetics. *Genetics Selection Evolution*, 34: 275–305.

Panellists' comments and discussion

Mr Richard Clarke, Rare Breeds Survival Trust (RBST), United Kingdom

As the first NGO interested in the conservation of farm animal genetic resources we fully support the approach presented in this paper. Inventory, characterization, monitoring and utilization are the core principles that have driven the work of the RBST since its creation in 1973.

Since 1973, the RBST followed these principles for over 70 breeds of large livestock native to the United Kingdom. Since 1976, RBST has sought to share its awareness of breeds at risk through the publication of an annual watchlist. Eligibility is based on numerical, distribution and genetic factors. Over the past thirty years RBST has overseen the movement of 11 breeds from an "at risk" classification to mainstream.

Initially the Trust was independent of governmental agencies, but now is working more closely with them. Surveys of endangered breeds have been undertaken by RBST since 1976, but since 1994 the RBST, in close association with government agencies, has undertaken surveys of all livestock breeds in the United Kingdom. Surveys have been every 4-5 years.

In the past twelve months much of the work of the RBST has been encapsulated in the DEFRA (Department of Environment, Food and Rural Affairs) *UK National Action Plan on Farm Animal Genetic Resources*.

Finally, I would like to make a couple of observations concerning issues raised in this paper.

From the RBST's years of experience in considering breeds at risk of extinction, a preoccupation on numbers at the cost of an awareness of within-breed diversity, and geographical distribution can be dangerous.

This paper clearly demonstrates how technology is rapidly developing to allow us to better connect phenotype to genotype, but with it comes the threat that these technologies will allow more rapid selection for a specific phenotype, thus threatening genetic diversity.

Mr Milan Zjalic, International Committee for Animal Recording (ICAR), Italy
Animal identification and recording as tools in characterization of animal genetic resources

"Characterization of AnGR encompasses all activities associated with the identification, quantitative and qualitative description, and the documentation of breed population..." (State of the World report, Part 4 - Section B, Methods for characterization).

Identification and registration (I&R) are important tools at all stages of an animal's life, as well as in any part of the food production process such as:

- Farm and herd management.
- Animal recording.
- Animal breeding.
- Animal health and disease surveillance.
- Trade.

Animal marking is associated to the domestication of different animal species by humans. Identification techniques are classified according to characters used and to their permanence on the animal. Main artificial permanent systems are branding (hot-iron and freezing), tattooing, ear notching, ear tagging (metal and plastic) and electronic identification (injectable, ear tags and bolus), but natural systems are also used (mainly retinal imaging and molecular markers). Recent experience has shown that electronic identification in spite of its high-tech nature is the most suitable method of identification, also for developing countries. In this respect, the use of boluses for ruminants and injectables for monogastric animals are recommended. Molecular markers are routinely used in parentage registration.

Description and documentation of phenotypic characteristics of the breed population includes recording all traits of economic importance such as product yield, quality of products, longevity and reproduction traits. For decision-making on conservation strategies and on developments of breeds, it is necessary also to collect data on birth weight, age at sexual maturity, daily gain and others. Data should be collected, stored and processed in a uniform and standardized manner, so that they can be retrieved and compared with future data as well with data related to other local or mainstream breeds.

The International Committee for Animal Recording (ICAR) is composed of national and local organizations involved in animal identification, recording and genetic evaluation. The object of ICAR is to promote the development and improvement of performance recording and evaluation of farm animals through establishing definitions and standards for measuring animal characteristics having economic importance. Working together, Members of ICAR establish rules, standards and guidelines for the purpose of identifying animals, the registration of their parentage, recording their performance, evaluating their genetics and publication of such.

The present structure of ICAR as a registered non-profit INGO provides for full participation of its members in developing guidelines and recommendations on the basis of the sound scientific evidence. Guidelines represent minimum requirements set up to ensure a satisfactory degree of uniformity of recording among member countries and a maximum flexibility in the choice of methods. ICAR's logo – an antique Roman scale and inscription in Latin "*Quod scriptum est manet*" (what is written remains) symbolizes the message that one can measure animal traits using available tools and that data should be registered for future use. However, only standardized methods ensure uniformity of data and adequate evaluation of traits.

Animal identification, performance recording and genetic evaluation are of particular importance for the sustainable utilization of animal genetic resources and development of animal production in developing countries. For this reason, ICAR encourages and supports experts from countries with prevalently low to medium input production systems in developing standards and guidelines suitable for their specific conditions. The ICAR Task Force Developing Countries promotes participation of experts from developing countries in ICAR research and training activities and promotes the establishment of country-specific identification and recording systems. Success stories, such as guidelines for buffalo milk recording, recommendations for animal identification and breeding programmes in low to medium input systems and activities of ICAR Members from developing countries in Africa, Asia and Latin America, indicate that this type of international cooperation is necessary also in support of conservation and sustainable utilization of local and endangered breeds.

Mr Jacob Wanyama, VETAID
Mozambique/LIFE Network, Mozambique

The paper covers different levels and methods of characterization, and highlights the importance of covering more than only production. It provides details on the role of emerging technologies for molecular characterization. It provides a useful basis for conducting the inventory and monitoring animal genetic resources.

However, from the points of view of the livestock keepers:

- The breed definition does not give justice to the fact that many local breeds are products of deliberate manipulations by their keepers. It does not capture the selection criteria of breeders in the communities.
- Documentation methods should put more emphasis on the participation of stakeholders in the communities.
- If collecting materials for molecular characterization, those doing the inventories should obtain the prior informed consent of the owners of the animals!

The paper outlines what needs to be done. But it lacks information on transparent procedures on how the information is stored and used. It should be clear:

- Who has access to the data.
- Who can use them.
- How the data will be used.

The monitoring describes an early warning system. But again, it does not recognize the important role that livestock keepers can play in this process.

The future:

- NGOs have the direct contacts to livestock keepers in communities and therefore have the ability to aid communities in the inventory, characterization and monitoring of their breeds.
- The information collected should be stored in community-based and driven documentation centres. NGOs have the capacity to facilitate such process. They can build the capacity of communities to document their own breeds.

Summary of plenary discussion

The meeting was then opened for general discussion and interventions from the floor. Issues raised during this discussion included:

- The practical steps needed to improve inventory and monitoring.
- The need to ensure that results of characterization processes are widely disseminated.
- The need to consider legal issues – such as those related to intellectual property rights associated with methods for characterization and the animal genetic resources themselves.

The authors' responses and concluding comments included the following points:

- The need for a multi-disciplinary approach to characterization and for the appropriate allocation of roles in the characterization process.
- The importance of encouraging participation of breeders and livestock keepers and their organizations.
- The need for clear explanation of how the information gathered will be used, and for those that provide the information to be involved in the analysis and use of this information.

Sustainable use and genetic improvement

C. Nimbkar¹, J. Gibson², M. Okeyo³, P. Boettcher⁴ & J. Soelkner⁵

¹Animal Husbandry Division, Nimbkar Agricultural Research Institute, P.O. Box 23, Phaltan 415 523, Maharashtra, India

²The Institute for Genetics and Bioinformatics, C.J. Hawkins Homestead, University of New England, Armidale, NSW 2351, Australia

³International Livestock Research Institute, P.O. Box 30709, Nairobi 00100, Kenya

⁴Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, International Atomic Energy Agency, P.O. Box 100, A-1400 Vienna, Austria

⁵University of Natural Resources and Applied Life Sciences, Vienna, Austria

Summary

Sustainable use of animal genetic resources for agriculture and food production is proposed as the best strategy for maintaining their diversity. Achievement of sustainable use would continue to support livelihoods and minimize the long-term risk for survival of animal populations. The concept of sustainable use has economic, environmental and socio-cultural dimensions. Sustainable use of animal genetic resources also contributes to food security, rural development, increasing employment opportunities and improving standards of living of keepers of breeds. Supporting the rearing of breeds through better infrastructure, services, animal health care, marketing opportunities and other interventions would make a significant contribution to the sustainable use of animal genetic resources.

Sustainable use envisages the use and improvement of breeds that possess high levels of adaptive fitness to the prevailing environment. It also encompasses the deployment of sound genetic principles for sustainable development of the breeds and the sustainable intensification of the production systems themselves. Sustainable use and genetic improvement rely on access to a wide pool of genetic resources.

Genetic improvement programmes need to be considered in terms of national agriculture and livestock development objectives, suitability to local conditions and livelihood security as well as environmental sustainability. Genetic improvement can involve choice of appropriate breeds, choice of a suitable pure breeding or crossbreeding system and application of within-breed genetic improvement. The choice of appropriate breeds and crossbreeding systems in developed countries has been a major contributor to the large increases in productivity,

and has benefited greatly from the fact that developed country animal genetic resources are well characterized and relatively freely exchanged. Where proper steps have been followed by careful assessment of demand, execution, delivery, impact and cost-benefit analyses, successful within-breed improvement has been realized within indigenous populations in developing countries. Breeding objectives and programmes for subsistence oriented and pastoralist systems are likely to be entirely different from conventional programmes. Crossbreeding has been most successful where it is followed by a rigorous selection programme involving livestock owners' participation and substantial public sector investment in the form of technical support. In any genetic improvement programme, inbreeding needs to be monitored and controlled.

Within-breed genetic improvement is normal practice in the developed world, and has become a highly technical enterprise, involving a range of reproduction, recording, computing and genomic technologies. Emerging genomic technologies promise the ability to identify better, use and improve developing world animal genetic resources in the foreseeable future. Useful systems can, however, be established without the need for application of advanced technology or processes.

Résumé

On propose une utilisation durable des ressources génétiques animales pour l'agriculture et l'alimentation comme meilleure stratégie pour la conservation de la diversité. Atteindre l'utilisation durable permettra d'améliorer la qualité de vie et diminuera le risque à long terme de la survie des populations animales. Le concept d'utilisation

durable entraîne des mesures économiques, environnementales et socioculturelles. L'utilisation durable des ressources génétiques animales contribue aussi à la sécurité alimentaire, au développement rural, à l'augmentation des opportunités d'emploi et à l'amélioration des standards de vie des éleveurs. Soutenir l'amélioration des races à travers une meilleure infrastructure de services, de santé animale, d'opportunités de marché et d'autres interventions pourrait aider de façon significative à l'utilisation durable des ressources génétiques animales. L'utilisation durable comporte l'utilisation et amélioration des races qui possèdent des hauts niveaux d'adaptation physique aux principaux milieux. Cela comporte aussi l'application de principes génétiques adéquats au développement durable des races et à l'intensification durable des systèmes de production en soi. L'utilisation durable et l'amélioration génétique se basent sur l'accès à une large gamme de ressources génétiques.

Les programmes d'amélioration génétique doivent être considérés en termes d'agriculture nationale et développement des objectifs d'élevage, ainsi que compatible avec les conditions locales de moyens d'existence et d'environnement durable. L'amélioration génétique peut entraîner le choix de races plus appropriées, races plus pures adaptées ou un système de croisement de races et l'application de l'amélioration génétique à l'intérieur de la race elle-même. Le choix de la race et des systèmes de croisement de races dans les pays en développement a été un des facteurs qui a influencé le plus l'augmentation de la productivité et a bénéficié largement le fait que dans les pays développés les ressources génétiques animales soient bien caractérisées et puissent bénéficier d'un mouvement relativement libre. Là où les démarches appropriées ont été suivies à travers des évaluations correctes sur la demande, l'exécution, la remise, l'impact et l'analyse de coût-bénéfice, le succès de l'amélioration à l'intérieur de la race a tout de suite été atteint avec les populations indigènes dans les pays en développement. Les objectifs d'amélioration et les programmes pour la subsistance et les systèmes de pâturage seront différents des programmes conventionnels. Les croisements de races ont eu plus de succès lorsqu'un programme de sélection rigoureux a été suivi et quand la participation des éleveurs et une partie du secteur public a été présente en forme d'investissement et appui technique. Dans tout programme d'amélioration génétique il est nécessaire de contrôler et faire un suivi de la consanguinité.

L'amélioration génétique de la race est une pratique normale dans le monde développé et est devenue une entreprise hautement technique qui met ensemble les domaines de la reproduction, le contrôle, l'identification et technologies du génome. Les nouvelles technologies du génome promettent dans un futur proche une meilleure capacité d'identification et l'utilisation et amélioration des ressources génétiques animales dans le monde en développement. Des systèmes utiles peuvent cependant être établis sans la nécessité d'appliquer des procédures ou des technologies à l'avant-garde.

Resumen

Se propone una utilización sostenible de los recursos zoogenéticos para la agricultura y la alimentación como mejor estrategia para el mantenimiento de su diversidad. Alcanzar el uso sostenible contribuirá a la mejora de la calidad de vida y minimizará el riesgo a largo plazo de la supervivencia de las poblaciones animales. El concepto de utilización sostenible conlleva dimensiones económicas, ambientales y socioculturales. La utilización sostenible de los recursos zoogenéticos también contribuye a la seguridad alimentaria, el desarrollo rural, el aumento de oportunidades de empleo y la mejora de los estándares de vida de los ganaderos. Apoyar la cría de razas a través de una mejor infraestructura, servicios, cuidados sanitarios de los animales, oportunidades de mercado y otras intervenciones contribuiría de forma significativa a la utilización sostenible de los recursos zoogenéticos.

La utilización sostenible comporta el uso y mejora de las razas que poseen altos niveles de adaptación de su forma física a los principales ambientes. También conlleva el despliegue de principios genéticos adecuados para el desarrollo sostenible de las razas y la intensificación sostenible de los sistemas de producción en sí mismos. La utilización sostenible y la mejora genética se basan en el acceso a una amplia gama de recursos genéticos.

Los programas de mejora genética necesitan ser considerados en términos de agricultura nacional y desarrollo de objetivos ganaderos, así como compatibilidad con las condiciones locales y seguridad de sustento y sostenibilidad ambientales. La mejora genética puede implicar la elección de las razas más apropiadas, la raza más pura adecuada o un sistema de cruce de razas y la aplicación de mejora genética dentro de la raza. La elección de la raza adecuada y de los sistemas de cruces de razas

en los países en vía de desarrollo ha sido uno de los factores que más ha influido en el incremento de la productividad, y se ha beneficiado ampliamente del hecho que en los países desarrollados los recursos zoogenéticos están bien caracterizados y gozan de un intercambio relativamente libre. Donde se han seguido los pasos adecuados con evaluaciones correctas sobre la demanda, ejecución, consigna, impacto y análisis de costo-beneficio, el éxito de la mejora dentro de la raza ha sido alcanzado con poblaciones indígenas en países en vía de desarrollo. Los objetivos de mejora y los programas para la subsistencia y sistemas pastorales serán mayormente distintos de los programas convencionales. Los cruces de razas han tenido mayor éxito donde se ha seguido un programa de selección riguroso que implique la participación de ganaderos y parte substancial del sector público en forma de inversión y soporte técnico. En todo programa de mejora genética es necesario controlar y monitorear la consanguinidad.

La mejora genética dentro de la raza es una práctica normal en el mundo desarrollado y se ha convertido una empresa altamente técnica que cubre los campos de reproducción, control, computo y tecnologías de genoma. Las nuevas tecnologías de genoma prometen la capacidad para identificar mejor y la utilización y mejora de los recursos zoogenéticos del mundo en vía de desarrollo en un futuro próximo. Los sistemas útiles pueden sin embargo ser establecidos sin la necesidad de aplicar procedimientos o tecnologías de vanguardia.

Keywords: *Targeting breeds, Production systems, Market access, Adding value, Dissemination, Sustainable breeding programmes, Technology, Intellectual property.*

Introduction

Sustainable use “*is the use of components of biological diversity in a way and at a rate that does not lead to the long-term decline of biological diversity, thereby maintaining its potential to meet the needs and aspirations of present and future generations*”. This is the definition of “*sustainable use*” proposed in Article 2 of the Convention on Biodiversity (CBD). *The State of the World’s Animal Genetic Resources for Food and Agriculture (SoW-AnGR)* (FAO, 2007) identified key elements of this concept as it applies to animal genetic resources. It reviewed existing concepts but did not attempt a comprehensive description of the state of the art. The general conclusions of the SOW-AnGR were that there is a

need for the concept of “*sustainable use*” to be “*interpreted in the context of agricultural biodiversity, and for concrete management strategies to be developed for AnGR*”. After the drafting of SOW-AnGR, FAO held an expert meeting that identified the guiding principles of sustainable use, made specific recommendations addressing relevant aspects of the concept and focused on work required to clarify and develop the concept further (FAO, 2006a; FAO, 2008). This paper describes the state of the art of scientific thinking on the key technical issues, options and opportunities in relation to sustainable use of AnGR.

Animals are reared in production systems, each with its unique geographical, environmental, cultural and socio-economic context. Sustainable use of animals for agriculture and food production in robust, ecologically compatible production systems is widely accepted to be the best strategy to maintain their diversity. Continued use of animal genetic resources within the environment in which they were developed provides a number of advantages, including maintenance of local knowledge about how best to manage the animal, maintenance of the production environment, and continued opportunities for the livestock to adapt to local production conditions and the needs of the society (FAO, 2006a). However, allowing movement of animal genetic resources to new locations and production and market systems is also a way of promoting their sustainability. Use of animal genetic resources inevitably includes development. Animal genetic resources are dynamic resources, changing with each generation in interaction with the physical environment and according to the selection criteria of their keepers (Wurzinger *et al.*, 2006). The concept of sustainable use therefore encompasses genetic improvement.

Even in the most rapidly developing countries, there are striking inequities in access to the benefits of economic development. Many families continue to keep a few animals of traditional breeds, often with very low use of external inputs, to provide a wide variety of products and services for household consumption and for sale in local markets. The development of opportunities for most of these families to intensify production and participate in national or niche markets or to find more lucrative non-agricultural employment is not likely to occur in the near future. In the meantime, continued access to well-adapted, local animal genetic resources will remain important for them (FAO, 2006b). Support for sustainable use of animals in developing countries will thus contribute to the broader socio-economic goals of livelihood security

and rural development, increasing employment opportunities and standards of living in rural areas and reducing migration to cities. Animal genetic resources play an important role in maintaining vital rural areas in developed countries also. In addition, in both developing and developed countries, animal genetic resources supply nutritious, protein-rich foods to people.

This paper discusses opportunities to enhance sustainable use of animal genetic resources, given the identified drivers of change articulated in the first paper in this series, mainly in agropastoral systems in marginal areas and crop-livestock systems in high potential areas. This focus is justified because industrial production systems using commercial breeds are already well developed, supported by heavy investment of capital, other resources and knowledge, and have efficient monitoring and corrective mechanisms in place if needed. The paper also presents the current scientific understanding in the area of genetic improvement and sustainable breeding programmes for development of animal genetic resources. The impact of revolutionary technologies in the field of genetic improvement and issues related to intellectual property rights are also discussed.

Facilitating sustainable use within production systems

Animal genetic resources form the basis of the livelihood and the cultural identity of a large number of farming and pastoral groups. Livestock have a critical role in maintaining sustainable agricultural systems, assuring food security and alleviating poverty. This role is especially important given the prospect of climate change or emerging diseases and the unpredicted rate and consequences of such change. It is expected that sustainable use would lead to the maintenance of vibrant and vigorous populations of breeds in their appropriate production systems. Increasing the profitability of rearing animals, particularly by increasing their market value, as well as enhancing their non-market values can maximize the probability of their continued use in the long term. Adaptive fitness and increased productivity can be achieved and maintained more effectively by improving inputs, environmental conditions and genetic resources concurrently. There is a range of alternatives and opportunities available for such facilitation including institutional strengthening.

There are, however, also many examples where opportunities have been wasted or inadequately exploited due to inappropriate policies and lack of support in critical areas (Philipsson, Rege and Okeyo, 2006).

Targeting breeds that require interventions

In general, more effort to promote sustainable use needs to be directed to those breeds that are likely to become threatened without support. Another factor to be considered in the targeting of interventions is the specific characteristics of breeds that make them unique – for example, adaptive traits such as disease and heat-resistance or specific feeding behaviour. Other criteria might include a focus on breeds that are specific to restricted regions or are unique in terms of their genetic, morphological, functional or cultural characteristics or the products that they produce. Development of a breed is likely to be more successful where there is a local community that highly values the breed in question and has a long history of local knowledge and experience of working with the animals. Continuous monitoring of the status of breeds by periodic breed surveys and censuses would help to provide information on population trends and impending threats. Such data can inform decision-making and help in formulating sound development schemes. This aspect is dealt with more comprehensively in the companion paper on characterization.

Strengthening production systems

Breeds fit into specific production systems and agricultural landscapes. If particular production systems disappear, the associated animal populations may no longer continue to be used sustainably. Strengthening these production systems so that they are robust in the face of changing circumstances would support the sustainable use of animal genetic resources. Various ways of strengthening these systems are elaborated below.

- Opportunities for small changes in farming systems. Small changes in farming systems, designed according to the prevailing climate, resource profile and agricultural practices, can make livestock rearing more profitable and beneficial to the farming system and thus more

sustainable. One example is to use novel ways of integrating crop farming and livestock rearing such as ley farming. Another example of an alternative model of farming using livestock is growing grass/leguminous forage on marginal, rainfed lands and rearing livestock instead of sowing grain crops that usually do not yield any grain because of inadequate rainfall.

- Provision of technical services. In some cases, technical improvements to animal nutrition, management and health may improve the economic viability of animal populations. The sustainability of animal genetic resources in existing production systems could be improved substantially by provision of basic veterinary services, including disease prevention measures such as vaccination. Improvements in management and genetics go together in reality as changes in one create new opportunities for the other. Provision of credit to purchase animals and for capital expenditure and a reliable supply of feed resources can provide significant impetus to the rearing of endangered breeds. These services may have to be tailored to specific needs – for example, they need to be mobile for nomadic herding. Other improvements to rural and agricultural infrastructure would also encourage livestock rearing in addition to other general benefits, for example by improving market access through provision of market information and objective pricing structures.
- Ensuring continued resource availability to livestock keepers. Sustainable use of animal genetic resources is closely linked to the continued availability of adequate grazing and water. Pastoralist production systems are increasingly under threat worldwide. The reasons for this are numerous:
 - deterioration of natural pastures as a result of droughts, inappropriate management of grazing and soil erosion;
 - curtailed access of livestock to common property resources;
 - diversion of grazing lands to other uses such as irrigated crop-farming, establishment of industries, urbanization or creation of national parks;
 - increasing difficulties in migration owing to increased cross-border disease-related controls, and traffic and highway codes that restrict livestock movement along and across major highways.

There are also other increasing demands, such as for biofuel production, on common property

resources and government lands in almost all countries. A pragmatic approach would be to take into account the vital role of animal genetic resources in diverse spheres, from production of much-demanded animal protein to maintenance of fertility of farmlands and creation of space for animal genetic resources in land-use plans (Köhler-Rollefson and LIFE Network, 2007).

- Capacity building. Training will help to inform livestock keepers of the latest scientific developments applicable to their livestock, such as availability of new vaccines, and will help to protect them from inappropriate advice (Malmfors *et al.*, 2002). Training should build upon existing local knowledge of the production system and enable livestock keepers to make informed decisions.
- Improving the status of animal genetic resources by raising awareness among policy- and decision-makers. Sustainable use of animal genetic resources has not achieved a high priority in the strategies of many governments or national and international funding agencies. In the Consultative Group on International Agricultural Research (CGIAR), institutional capacity and availability of funds are generally skewed heavily towards the plant sector (FAO, 2006b). Animal husbandry usually gets a raw deal compared to crop farming in governmental financial allocations because of inadequate awareness of policy-makers of the importance of livestock. It is therefore necessary to raise the awareness of the contribution of livestock to national economies and to the well-being of large numbers of families to give a higher profile and status to livestock rearing. Raising awareness will help in encouraging policy-makers to develop sound policies that are beneficial for sustainable use of animal genetic resources rather than policies that may have an adverse impact on livestock rearing. For example, supportive public policy and long-term technical support systems are largely responsible for the success of the dairy subsector in India (Kumar, Birthal and Joshi, 2003).
- Promoting “organizations” of livestock keepers. A key aspect to promoting sustainable use is creating or strengthening structures to organize the keepers of animals and help motivate communal efforts (Kosgey & Okeyo, 2007). Organizations are stronger than individuals and can safeguard group interests better, by advocacy with authorities. In the longer term, building these structures may serve a capacity-building role – allowing livestock

keepers better access to information, strengthening their position in relation to extension services, facilitating the organization of training and improving bargaining power when marketing products. In Europe, there are strong farmer cooperatives and breeding organizations that go back a century and have also received much public support over the years.

Improving market access and promoting novel uses of animal genetic resources

Developing markets for livestock breeds, their products and services

The value of animal production can be increased by marketing products more effectively. Ease of marketing and lucrative prices for animals and their products can provide the biggest boost for continued use of animal genetic resources (Boxes 1 and 2).

Development of niche markets is also important from the perspective of promoting sustainable utilization. Niche markets rely on creating perceived value regarding the conditions of production, product quality or a combination of these. Consumers that particularly value food quality or specific production methods are the most likely to purchase specialized niche products such as Parmigiano Reggiano cheese produced from Regianna cattle in Italy, high-value cured pork products from Iberian pigs reared in oak-forest production environments in Spain and meat from the black boned chicken breed in Viet Nam, known for its medicinal value. One of the ways to create demand for products of breeds reared in pastoralist systems with no chemical inputs is to market them as “range-fed” or “fed on natural vegetation”. Such products could also benefit from “geographical indication” recognition.

In almost all areas of India, a niche market for local breed chickens and eggs, perceived to be “high-quality” and therefore more expensive, exists side by side with broiler chicken and commercial layer hen eggs. Similarly, in Malaysia, meat from the Kampong chicken is considered to be better tasting than the commercial breeds. In the United Kingdom, a ready market was developed for beef from Angus cattle as high-quality beef (with high marbling), which served to increase the Angus population. The measures adopted for this included promoting

Angus beef through a restaurant chain. The fragility of some such niche markets is, however, demonstrated by the collapse of the restaurant chain following the outbreak of mad cow disease.

Novel uses for animals and animal products

New uses have been developed for animals and their products with desirable consequences for continued maintenance of animal genetic resources. The unique immune system enhancement properties of Panchagavya (a mixture of milk, curd, ghee, urine and dung of indigenous cows prepared according to a recipe from ancient Ayurveda [Sushruta Samhita, 1985]), identified by new research (Chauhan *et al.*, 2004) have led to new marketing possibilities in India and Sri Lanka. A non-governmental organization (NGO) in Rajasthan, India, has successfully introduced camel milk ice cream (desert dessert) as part of a comprehensive strategy to make camel rearing more profitable. Research in the United States of America on “aversive conditioning” using boluses with lithium chloride (Mueller, Poore and Skroch, 1999) shows that sheep can be trained to bypass the tender shoots of grapevines and trees for the weeds sprouting underneath.

Promoting use of animals in landscape conservation

Use of traditional grazing livestock for landscape heritage and biodiversity maintenance and for nurturing more complete ecosystems is a growing management practice in many developed countries. In the United Kingdom and Europe, and specifically the Balkans, the role of grazing livestock has been recognized as critical in the maintenance of wildlife and native plant biodiversity in many high nature value ecosystems. In the Mediterranean, grazing for shrub control helps to reduce forest fires. Cultural tourism associated with the unique culture of rearing local breeds has been expanding rapidly in Europe and also in South America where camelids are great attractions at parks and tourist sites. Similar approaches are needed in other developing countries, since here too particular breeds have shaped certain landscapes. Functioning pastoralist systems also have value as tourist attractions, besides contributing to ecosystem health.

Box 1. Adding value to Nguni cattle

The Nguni of South Africa is an African taurine breed with a slight admixture of zebu blood that reached the region together with southward migrating pastoralists in about 300 AD. After white settlers arrived with exotic cattle, the Nguni cattle were long perceived as inferior because of smaller carcass size, non-uniform colour pattern and lack of information on their production potential. Even the people who had originally kept this breed started crossbreeding or keeping exotic cattle. Research in the 1980s then revealed that the Nguni breed is very tick-tolerant, can maintain its condition during seasonal food shortages, can obtain optimal nutritional value from the available forage, is a good walker and very docile. Its adaptation to harsh extensive production systems offers many advantages to smallholders. The Animal Improvement Institute has therefore initiated a project to supply selected Nguni bulls to smallholders together with training and infrastructural support. Nguni cattle have a wide range of colours. The colour variation indicates the cultural heritage of the breed, which has been raised by African stockmen for centuries. Colour variation frequently had a ceremonial and symbolic importance. The colourful Nguni hides are much in demand these days for pelts that are tanned with the hair on, for use as rugs, clothing and home furnishings. Being able to predict and generate specific colours has taken on a new economic aspect as these uses have recently increased. In addition, certain colours or pigmentation patterns (such as pigmented skin beneath white hair) can be helpful in adaptation of animals to harsh conditions of high solar radiation. All three of these factors (tradition, utility, adaptation) combine to make colour important for Nguni breeders, and unravelling the details of colour genetics can be useful for them (Köhler-Rollefson, 2004; Spönenberg, 2007).

Box 2. Value-adding to peri-urban dairy farming in Latin America

Straddling the border of Peru and Bolivia, the Altiplano – a high-altitude plain at 4 000m above sea level – is one of the poorest regions in the world. At such high altitudes, the environment is unforgiving: drought and extreme cold are common. The region supports six million people, who mostly depend on agriculture. Potato is the staple but crop failure is a regular occurrence and many families live in extreme poverty. However, for some Altiplano farming families living close to urban centres, nutritional and income stability is not completely unattainable. Milk production is growing in importance in the region and a pilot project, under the ALTAGRO initiative of the International Potato Center (CIP) and its partners, has created a market in several large towns for cheese made from local cows' milk. Most households in the area earn around US\$1 per day. With this initiative, dairy producers have more than doubled their income, with some now earning up to US\$850 per year. The ALTAGRO project, financed by the Canadian Government, has supported the construction of two small dairies in Atuncolla-Illpa, a Peruvian town with a population of 10 000 people. A training plant set up at the experimental station of the Instituto Nacional de Investigación Agraria (INIA, Peru's National Agricultural Research Institute) is providing technical assistance to farmers and processors in how to transport the milk and process it into cheese (www.new-ag.info/07/04/focuson/focuson1.php).

Research and dissemination of research results

Public-funded applied research needs to focus on improving livestock rearing as an integral part of production systems and finding innovative solutions to real problems rather than on obscure theoretical topics. Successes as well as failures need to be published in order to capitalize on experiences. Research on the beneficial interrelationships between livestock and their environment and the necessity of livestock to maintain the sustainability of the landscapes they use is likely to provide enlightening results (Lewis, 2003). It is important to publish research results in accessible sources to ensure wider dissemination.

Promoting sustainability as the main objective

The supporting interventions should be such that they create an enabling environment to make livestock rearing self-sustained in the long term rather than dependent on outside support. If support is withdrawn due to a change in the macroeconomic situation or in the government, the livestock rearing system it has strengthened should not collapse. In fact, consequences of interventions could be tested against the potentiality that the support may be terminated.

Appropriate strategies for sustainable use will differ from country to country or among groups of countries because of the large differences among areas of the world, especially in terms of gross national product and available technology (Gandini and Oldenbroek, 2007).

Genetic improvement and sustainable breeding programmes

Introduction

The concept of sustainable use encompasses the development of animal genetic resources, ensuring that they remain a functional part of production systems, and the sustainable intensification of these production systems. Genetic improvement is the systematic exploitation of genetic variation in important traits among individuals within or

between breeds. Breeding programmes for animal genetic resources are generally undertaken in order to improve their productivity and the quality of food and products derived from them and to ensure the availability of such food/products at affordable prices. Genetic improvement of livestock has made and will continue to make major contributions to agricultural development, food security, sustainability and livelihoods. In high-input production systems, which are common in the developed world, modern chicken and pig hybrids consume less than half the feed per kilogram of meat produced than the strains of 50 years ago. Such genotypes cannot, however, stand the harsh rigours (disease challenge, poor-quality feed, high temperature) of the low-input, livelihood-focused systems in most of the developing countries. The high feed conversion efficiency has allowed the demand for meat of affluent societies to be met from a greatly reduced land area, thus releasing large areas of agricultural land that would otherwise have been required to produce poultry and pig feed. The importation of these improved genetics along with their associated production systems into developing countries has benefited consumers through availability of cheap broiler meat and pork and has also brought profits to farmers, although some other farmers were crowded out of markets because of these developments. There are other examples of benefits (with some qualifications) in the developing world. For example, the use of improved dairy genotypes has allowed the development of a large informal milk market that has dramatically improved smallholder livelihoods and human nutrition in the densely populated highlands of Kenya. A recent study has, however, shown that these animals are of higher milk potential than tropical climates and feed resources can support. In some situations, this resulted in drastic reductions in farmers' profits (King *et al.*, 2006).

Genetic improvement can take many forms, but generally and logically follows an ordered hierarchy of events. This starts with understanding of the production and marketing systems, choice of appropriate breeds or strains (sometimes resulting in replacement of existing breeds), establishment of an effective pure breeding or crossbreeding system, and then further improvement through selection of superior genotypes within populations that best suit the production and market conditions. The past 50 years have seen a drastic change in breed use. As a consequence, genetic improvement in the developed world is now primarily based on a few breeds and within-breed improvement. Almost all

pigs in developed country markets are, however, crossbred and some strategic crossbreeding is being undertaken increasingly in cattle and sheep. In the developing world, most genetic change is taking place through change of breeds via crossbreeding programmes aimed at “grading up” of indigenous breeds towards exotics from the developed world. Systematic within-breed improvement is much less prevalent, although livestock keepers themselves continuously make decisions to keep and cull animals according to criteria they consider important. However, apart from a few cases, most of the structured breeding programmes have seen limited success, mainly because of inadequate understanding of the prevailing agro-ecological and marketing conditions.

Within-breed improvement

Within-breed genetic improvement programmes are routine for all the breeds and strains of livestock used in the dominant livestock production systems of the developed world. The genetic improvement typically accounts for 40 to 60 percent of the annual productivity gains in these systems. In the developing world, however, within-breed improvement to improve productivity is not common and has not often been sustainable. The relative lack of effort is partly due to the perception that greater genetic change is possible through the choice of specialized and improved exotic breeds and strains and crossbreeding systems. However, inadequately planned crossbreeding programmes have seen as much failure, if not more, as within-breed improvement programmes. Lack of suitable infrastructure, expertise and sustained government support has also hampered the establishment of within-breed improvement programmes in developing countries. Many factors have contributed to the lack of success in existing programmes – inadequate initial characterization of local populations, lack of participation of smallholder beneficiaries, inadequate dissemination mechanisms, inadequate or unsustainable infrastructure and expertise and/or rapid evolution of production systems (such as breed replacement), apparently eliminating the need or demand for the improved stock. Successful application of within-breed improvement is undoubtedly attainable in the developing world, but requires more careful assessment of demand, execution, delivery, impact and cost-benefit analyses.

Within-breed improvement presents a particular challenge in subsistence-oriented systems. It has to be based on adequate knowledge of the breeds in question and of the production system. Serious consideration has to be given to social, economic and environmental sustainability in this situation. Potential strategies for breed development appropriate to the local conditions and in keeping with the country’s overall livestock development objectives should then be identified, assessed and prioritized (Box 3).

Generally, breeding objectives have focused on increasing productivity, often measured at the individual animal level. However, breed improvement should take into account the full range of attributes that make production systems sustainable. Selective breeding efforts can vary in scope from highly organized breeding programmes through to simple culling decisions based on individual phenotypic information under less controlled environments. The choice of methods will depend on the objectives of the breeding programme, their acceptability to the whole spectrum of stakeholders, access to improved genetic resources and the technology and infrastructure available.

In harsh mountainous or arid rangelands and pastoral systems where the environment and markets are unlikely to change in the medium to long term and where existing genotypes are well adapted, simple within-breed selection programmes focusing on as few traits as possible provide the best approach. The traits to be included need to be easily recorded for the animals to be selected and depend on the primary use of a breed. They will be multiple for multipurpose breeds. While natural selection will take care of many adaptive traits, fertility of male animals needs to be considered based on the results of a first mating season. When the environment and market requirements are changing, then more planning, better designs and institutional integration/coordination are required.

Where proper steps have been followed, successful within-breed improvement has been realized, even within indigenous populations in developing countries. The improved Boran cattle in Kenya, the Nguni cattle in South Africa, the Tuli cattle of Zimbabwe and the Murrah buffalo programme of India (with some limitations) are success stories in regions where many programmes have failed. What is unique about all these examples is that the production, policy and market environments were well understood, the locally available genetic resources/breeds were well

Box 3. Community sheep breeding programme in Peru

In the highlands of the Sierra Central in Peru (an isolated high mountain range environment at an altitude of about 4 000 metres above sea level), dual purpose Corriedale sheep and native-type sheep with different levels of exotic upgrading are kept in an extensive pastoral system. A survey conducted in 1996 identified three types of sheep production systems: individual family flocks, communal flocks belonging to villages and multicommunal flocks managed by cooperatives often involving several villages in a region. The survey identified two major requests of farmers related to breeding: the need for suitable rams and the need for training in breeding techniques. After extensive discussions, an interesting breeding structure based on the open nucleus concept was established and made functional. The land and labour necessary to run the nucleus were provided by the communities based on a series of arrangements and technical support was provided by the university. The nucleus was established by mating imported and locally produced top rams with 50 "best" females of each of nine communal and multicommunal flocks. Half of the ewes were returned pregnant to the suppliers and the other half were used for starting a central nucleus providing improved rams to communal and regional flocks, which in turn also provided rams to family flocks. Incidentally, the progeny of local rams proved to be better suited to the local market conditions than the progeny of imported rams. Farmer organization and farmer training are the backbone of this successful community-based sheep breeding programme, which is still in operation (Mueller, Flores and Gutierrez, 2002).

evaluated and simple selection criteria agreed upon and implemented.

Intensive selective breeding will inevitably result in some reduction in genetic diversity within the breed. Systems for allocating breeding males to females based on the relative genetic contributions of parents have been developed to optimize genetic improvement while minimizing the rate of inbreeding (e.g. Sonesson and Meuwissen, 2000). These are used in commercial breeding and can be applied to local breeds if animals are appropriately identified and their pedigrees recorded accurately.

Choice of breeds and cross-breeding

The matching of appropriate breeds to evolving production systems has been a major contributor to growth in productivity and improvement of product quality in the developed world. This has been possible because developed world breeds and

strains are relatively well characterized and are easily accessible through established processes such as genetic evaluation rankings and semen and breeding male distribution schemes. In the developing world, most animal genetic resources are inadequately characterized and access to animal genetic resources from other developing countries is often difficult or impossible. In fact it is ironic that recently developed well-intentioned instruments such as the Convention on Biodiversity may hamper the sharing of breeds across countries even if it appears to be the most technically logical option and would actually contribute to the maintenance of agricultural biodiversity. Unless livestock genetic resources of the developing world are better characterized and made more accessible, it is inevitable that the choice of breeds and strains for breed replacement and crossbreeding will be dominated by those of the developed world. This is evident, given the strong marketing strategies of the improved livestock genetics companies from the

high-input systems of some developed countries. This may severely restrict the options of developing countries to develop their local breeds to meet goals for agricultural and economic development, sustainability and improvement of livelihoods.

With now widely predicted climate changes through direct and indirect effects (i.e. reduced number of growing days, hence herbage yields, increased disease outbreaks and challenges), the developing regions of the world's production systems are likely to be severely affected. Therefore, the need to source appropriate (those that best match the predicted future scenarios) breeds and genes from one developing country to another would be the most logical option. For example, if, as a result of global climate change, most of the sub-Saharan regions receive less rainfall and have hotter climates than currently is the case, then instead of embarking on long-term within-breed improvement of local breeds to match the predicted future environments in the affected areas, it would be better to access and move breeds. For example, Kenana and Butana cattle breeds of the Sudan that are already naturally adapted and reasonably productive under a harsh environment could be moved to those areas where harsher conditions are expected in future. Such realities add a new dimension to the potential utility of indigenous breeds.

In pastoral systems, and when market opportunities for improved milk and meat production exist but where large erratic environmental changes such as droughts are common, livestock keepers may maintain a range of diverse genotypes, some of which can survive drought conditions. Traditionally, pastoralists may keep a mix of species and breeds in their herds to maximize the advantages of good seasons and to reduce risk during bad seasons. For example, crossbred animals generated by crossing locally adapted females to an improved "exotic" breed male may be more profitable than their local purebred mothers when conditions are good, but may be the first to die when there is a bad drought. Farmers may use some indigenous breed sires and some exotic sires on parts of their herds/flocks while practising within-breed selection in part of the herd/flock. A good example is the Ankole cattle breed in the African Great Lakes region, where many keepers of large herds adopt a strategy of splitting their herds in this manner (Wurzinger *et al.*, 2006). Better planning is then necessary to find a balance between high-profit/high-risk and low-profit/low-risk and to ensure a good bio-economic balance.

The use of crossbreeding has also made major contributions to productivity and product quality in the developed world. Structured cross-breeding systems, such as "terminal crossing" where first generation cross-bred (F1) animals are slaughtered or where specialized crossbred dam lines are used, are common. Cross-breeding may also be used for gradual breed replacement with upgrading or the controlled maintenance of various proportions of exotics leading to formation of composites. The need to maintain pure breeds for the production of crossbred animals or commercial production is either managed by farmers or by commercial companies. Farmers have had extensive support and training and now understand the need to maintain a balance of breeds to make the system sustainable in the long term.

There are also examples of successful crossbreeding programmes in developing countries. In some situations, carefully conceived and executed crossbreeding programmes have merit as a rapid method of introducing desirable traits into local well-adapted breeds. The development of the Dorper sheep is one of the most successful programmes of composite breed development for a low-input production environment (de Waal and Combrinck, 2000). The breed was developed in South Africa by crossing Dorset Horn sheep with the fat-rumped, black-headed Persian sheep, a local Somali breed. Other successful crossbreeding programmes include the formation of the Sunandini synthetic dairy cattle in Kerala State, India (Box 4), the Boer goat of South Africa (Malan, 2000) and the Brazilian Milking Hybrid (MLB) cattle (Madalena, 2005). Crossbreeding has been most successful where it was followed by a rigorous selection programme involving livestock owners' participation and substantial public sector investment in the form of technical support.

However, very often, crossbreeding has been indiscriminate and the local breeds that underpin the crossbreeding programme have been lost because of a lack of understanding by the authorities, companies and/or farmers involved that these pure breeds must be maintained to support the system. The strategic use of crossbreeding as a way out of a narrowed genetic base in commercial breeds is also considered important. It is gaining acceptance, for example, for fixing the increasing adverse trends in reproductive traits in commercial dairy cattle in North America. Such strategic crossbreeding is desirable to prevent long-term reduction of genetic diversity.

Finally, it should be recognized that large, highly variable and rich genetic pools of crosses

Box 4. The Sunandini cow in Kerala, India

Conditions in the State of Kerala in southern India are generally not conducive to classical dairy farming. These conditions are: the year round hot and humid climate, relentless pressure on land for human needs, acute scarcity of fodder, high rainfall and consequent mineral depletion of the soil. However, the Kerala dairy development programme, implemented over four decades (1964–65 to 2000–01), increased the State's average yield per cow per day from less than a litre to nearly 7 litres and milk production from 200 000 to 2.6 million tonnes per year. It has provided livelihood support to over one million smallholder households. The phenomenal growth in milk production can be attributed to a planned effort to develop the cattle genetically for milk production, supported by an extension programme for fodder development and a well-organized milk collection, processing and marketing system. A new composite breed, called "*Sunandini*" has been established by crossbreeding local cattle and further selection among the crosses. During the process, however, almost 80 percent of the local cattle have been converted to Sunandini and the local Vechur breed of cattle has almost been lost. The composite has a wide genetic base of exotic donor breeds – Brown Swiss, Jersey and Holstein Friesian and, to a lesser extent, the Indian donor breeds Sahiwal, Gir, Rathi and Kankrej. The Sunandini breed combines the positive qualities of local cattle such as adaptability, resistance to disease and strong hooves with the high production potential of the exotics. The level of exotic inheritance is limited to 50 percent. Its overall average lactation milk yield is 3 400 kg with a milk fat percentage of 4.0 (KLDB, 2004).

between exotic and indigenous breeds exist in developing countries today. Such populations would serve as a quick foundation for synthetic breed formation; especially given the surviving individuals have the combination of genes that best fit the prevailing environments. Strategic use of such crosses to develop breeds for specific production systems is prudent and timely. For example, in trypanosomiasis endemic areas, it would make good sense to combine N'dama crosses that have survived and are productive with purebreds or crosses of equally tolerant cattle breeds such as the Orma Boran of Kenya and Sheko of Ethiopia (which is at risk). This underlines the importance of sorting out the problem of cross-country access to such genetics.

Applications of technology in genetic improvement

Current use of technology

Breeding programmes in the industrial production systems are complex and have evolved over many years of technical inputs in terms of design, determination of breeding objectives, calculation of

economic weights, genetic evaluation methods, breeding strategies and delivery of services, as well as structures and techniques for dissemination of improved genetics. They involve the extensive use of technologies for data recording and storage, advanced computing and statistical analysis, reproduction, genetics and genomics. For example, dairy cattle improvement generally involves automatic milk recording of several hundred thousand cows each year, compositional quality assessment, data download to a central database, large computers and advanced computer algorithms that estimate the genetic merit of millions of animals simultaneously, artificial insemination of millions of cows and embryo transfer of several thousand cows, laboratory assays to determine parentage and, increasingly, molecular genetic testing to determine which animals carry particularly desirable sets of genes.

In the developing world, advanced technologies are more difficult to implement because of high cost, lack of expertise and infrastructure and are consequently not widely used. A contrasting situation, however, exists in some developing countries (such as India) where several top research institutes pursue the use of mainly molecular technologies for their glamour rather than for supporting a practical breeding programme.

Research involving use of technologies is preferred over more tedious field research, which is perceived to be less rewarding. It is therefore necessary to ensure that simple breeding programmes based on proven genetic principles are not abandoned in favour of molecular genetic technologies that, in turn, need the existence of sound breeding programmes to be used effectively (see Box 5 for an example of effective use of advanced technology in a breeding programme in a developing country.) An example of effective use is that reproductive technologies, such as frozen semen or embryos, are used in several species to transfer germplasm between countries and sometimes to expand and/or disseminate rapidly an imported population. In addition to greater efficiency and reduced cost achieved, the use of such technologies greatly reduces the risk of disease transmission compared with importation of live animals.

Progress with simple technology

The low level of use of advanced technologies in most aspects of genetic improvement in the developing world need not prevent effective improvement being achieved (Box 6). For example, a well designed improvement programme, based on selection of the best animals assessed on their own performance, with no other information or analysis, can achieve from 40 to 70 percent of the maximum

possible rate of genetic improvement when compared with the use of all advanced technologies. The use of advanced technologies in the developed world is driven by the intense competition among breeding groups or companies and the desire to improve characteristics that are not easily or accurately recorded for every animal. In the absence of such intense competition in developing countries, there is no immediate need to introduce expensive, advanced technologies. A lower rate of genetic progress using simple cost-effective techniques is preferable and certainly better than no selection.

The level of sophistication in terms of breeding strategies to be adopted in order to ensure sustainability and effectiveness needs to be carefully considered. It will depend on the state of the local infrastructure, the product market and available supportive technical expertise and institutional arrangements. An example of unsustainable levels of sophistication is the Kenyan National Dairy Cattle Breeding Programme, with sophisticated progeny testing comprising multibreed centralized milk and butter fat recording and data processing systems involving several institutions. The programme was modelled along a European type of system without considering the local infrastructure and institutional limitations. The result is an ineffective system in which an unacceptably low (five or less) number of bulls per breed are recruited each year, with up to 11 years before the test results are completed, leading to a

Box 5. Marker-assisted introgression/gene introduction in India

A good example of a clear gene effect successfully implemented in a marker-assisted introgression (MAI) programme is found in India. The Booroola gene is being introgressed from the small Garole breed into the local Deccani breed that is suitable for meat production but has a limited reproductive performance. The Booroola gene has tremendous economic effects in this production system, increasing the weaning rate by nearly 50 percent. The breeding programme is undertaken by a research institute, but there are clear strategies and activities to ensure that the improved stock finds its way to shepherd flocks. Evaluation of the results in these shepherd flocks is an explicit part of the project, and initial results look very promising. Long-term impact, however, needs to be assessed. Early results also indicate that the litter size of Booroola carriers has a direct correlation with feed availability during mating/pregnancy. This means shepherds would be able to reap the benefits of the higher litter size during "good" years while the flock's average litter size would not be unsustainably high during "bad" years. Shepherds may also like to keep a mixed flock of Booroola carrier and non-carrier animals as a risk insurance. MAI should not be ruled out for breeding programmes in developing countries, but should be assessed based on the merit of each case (van der Werf, 2007).

near-zero genetic gain. In this situation, a simpler nucleus-herd-based young bull scheme would have been more effective and sustainable, given the very limited number of herds actually contributing to genetic improvement. Location and management of the nucleus and recorded herds should ensure that production conditions in such herds match or mimic those of the smallholder and/or commercial farms under which most of the progenies of the bulls are raised.

Emerging technology applications

Reproduction, data and statistical analysis technologies continue to show regular incremental improvements and are expected to benefit but not fundamentally change the current design and operation of genetic improvement programmes. After decades of research and development, sexed semen has recently become available on a commercial basis (Johnson *et al.*, 1987, Weigel, 2004). The use of sexed semen could be especially beneficial in countries such as India where religious beliefs preclude the consumption of beef. In such countries, the male animals are neglected and are a wasted resource. Technologies for management of female reproduction, such as synchronization of oestrus and (non) pregnancy diagnosis, can contribute to faster genetic improvement by decreasing the intervals between successive

parturitions and increasing the number of candidates for selection.

Some technologies, such as the Livestock Identification and Trace-Back System (LITS) implemented in Botswana as a deterrent to cattle thefts (http://practicalaction.org/?id=peace5_cattle_tracking_botswana), could have huge potential for a genetic improvement programme where lack of individual identification is one of the main hurdles. The digital identification system uses radio frequency identification technology, is safe, environmentally friendly and tamperproof, and is used to identify individual livestock throughout the country. Other than managing cattle records and deterring cattle thefts, the system would also potentially open up access to important livestock markets such as the European Union (EU). The EU beef market regulation requires that imported beef be traceable from the export slaughter facilities to the individual animal that the meat came from.

Genomic technologies that have emerged from the human genome project are rapidly being developed for livestock. For example, in the past two years the ability to detect variations in the genetic code of individual cattle has risen from testing two or three variations in a single test to 50 000 variations in a single test, and the cost of testing has dropped more than a hundredfold (see companion paper on characterization). Such technology developments are truly revolutionary and provide prospects for radical changes in

Box 6. Simplifying phenotypic measurement of performance

The marginal gain obtained by increasing the precision of information on phenotypic traits is subject to the law of diminishing returns. For this reason, developing countries that are attempting to implement an open nucleus breeding scheme may be advised to begin by collecting “*low tech*”, simple measurements of phenotypes from more animals and farms, rather than asking a few farmers to record complicated measures. For example, recording of milk yield could be bi-monthly or quarterly, rather than monthly. Lactation milk yield estimates based on only two test-days have been shown in some studies to have a correlation of greater than 0.85 with estimates based on ten test-days (Vasconcelos *et al.*, 2004). Measurements of heart girth can serve as a proxy for body weight when scales are not available, as the traits are both highly heritable and highly correlated genetically (Janssens and Vandepitte, 2004). For traits such as overall likeability, temperament and general disease resistance that would be difficult or expensive to measure objectively, farmers can be asked to assign simple, ordered categorical scores for phenotypes.

genetic characterization and improvement. Several groups have already demonstrated that using such tests it is possible to determine the genetic merit of individual animals for most commercially important characteristics, without the need for any prior phenotypic information on the animal (Meuwissen, Hayes and Goddard, 2001). Huge quantities of molecular level data are, however, needed. The ramifications of this are still being explored, but it is clear that radical changes in design and operation of genetic improvement in the developed world could emerge. The ability to apply such technologies for routine genetic improvement in the developing world will require substantial reductions in cost, which seem likely to be achieved but cannot be guaranteed. However, it is already clear that these new technologies can be applied to achieve a much greater understanding of the functional genetic variation of developing and developed world animal genetic resources, which can then be used indirectly to better target genetic improvement globally.

Intellectual property issues

Virtually all the processes of relevant reproduction, data capture, statistical analysis and computing technology are in the public domain. Proprietary software is either readily available at reasonable cost or can easily be duplicated without infringement of proprietary rights. A small number of commercially valuable molecular genetic tests have been patented. In most cases these patents have not been registered in developing countries and therefore provide little or no restriction on use in developing countries. Coupled with the fact that such existing patents are for inventions with little practical value in the developing world, the willingness of patent owners to provide free or low-cost access to the developing world does not appear to have been tested. The recent development of high-throughput tests for genetic variants has led to several applications for patents for simultaneous use of large numbers of genetic polymorphisms. It is understood that in recent months the United States Patent Office has ruled that such patents are not valid and that the test for each polymorphism must be patented separately. The most likely consequence of this is that inventors will seek to protect such intellectual property (IP) by maintaining commercial secrecy rather than applying for thousands of separate patents. This will mean that the technology will be available to competing companies or countries, but the exchange of

information will be hindered. It may also mean more difficulties for inventors to share IP with others, even where no commercial competition exists. This is because of the risk that key information might be leaked, thereby devaluing the IP. This situation is likely to be more damaging to technology use in the developing world where resources are less likely to be available to duplicate discoveries that have been protected by commercial secrecy.

Conclusions

Enhanced use and development of animal genetic resources in all relevant production systems play key roles in achieving food security and alleviating poverty. Ongoing utilization is also regarded as an effective means of maintaining diversity and ensuring the availability of resources for the future. Utilization is likely to continue if the breeds are perceived to provide genuine benefits – whether these are private benefits for the livestock keeper or public benefits for which society is willing to pay.

Continued increases in animal production and productivity will be necessary to enhance food security and provide critical income, products and services to hundreds of millions of poor families. Strategies involving incremental improvements in the production potential and productivity of traditional breeds, and corresponding gradual improvements in management and access to veterinary services, supplemental feeds and markets, appear most promising. The continued use of traditional breeds is likely to remain the most effective strategy for resource-poor farmers in many of the least-developed countries. However, opportunities may exist both to improve local breeds and for carefully managed and limited introductions of exotic breeds in areas of greatest production potential. These opportunities must be seized when genuinely available.

List of references

- Chauhan, R.S., Singh D.D., Singhal, L.K. & Kumar, R.** 2004. Effect of cow urine on interleukin-1 and 2. *Journal of Immunology and Immunopathology*, 6(1): 38–39.

- de Waal, H.O. & Combrinck, W.J.** 2000. The development of the Dorper, its nutrition and a perspective of the grazing ruminant on veld. *Small Ruminant Research*, 36(2): 103–117.
- FAO.** 2006a. Report on an expert meeting on sustainable utilization. Commission on Genetic Resources for Food and Agriculture. Intergovernmental Technical Working Group on Animal Genetic Resources for Food and Agriculture. Fourth session. 13–15 December 2006. Rome.
- FAO.** 2006b. A strategic approach for conservation and continued use of animal genetic resources. Commission on Genetic Resources for Food and Agriculture. Intergovernmental Technical Working Group on Animal Genetic Resources for Food and Agriculture. Fourth session. 13–15 December 2006. Rome.
- FAO.** 2007. The State of the World's Animal Genetic Resources for Food and Agriculture, edited by B. Rischkowsky & D. Pilling. Rome.
- FAO.** 2008. Report of the FAO/WAAP Expert Meeting on Sustainable Utilization of Animal Genetic Resources, Ferentillo, Italy, 2–4 July 2006, by D. Weary, D. Pilling & B. Rischkowsky. Rome. (Forthcoming – full proceedings including presented papers).
- Gandini, G. & Oldenbroek, K.** 2007. Strategies for moving from conservation to utilization. In K. Oldenbroek, (Ed.). *Utilization and conservation of farm animal genetic resources*, Wageningen, Netherlands, Wageningen Academic Publishers, pp. 29–54.
- Janssens, S. & Vandepitte, W.** 2004. Genetic parameters for body measurements and linear type traits in Belgian Bleu du Maine, Suffolk and Texel sheep. *Small Ruminant Research*, 54: 13–24.
- Johnson, L.A., Flook, J.P., Look, M.V. & Pinkel, D.** 1987. Flow sorting of X and Y chromosome-bearing spermatozoa into two populations. *Gamete Research*, 16: 1–9.
- King, J.M., Parsons, D.J., Turnpenny, J.R., Nyangaga, J., Bakari, P. & Wathes, C.M.** 2006. Modelling energy metabolism of Friesians in Kenya smallholdings shows how heat stress and energy deficit constrain milk yield and cow replacement rate. *Animal Science*, 82(5): 705–716.
- KLDB (Kerala Livestock Development Board).** 2004. Capitalization of experiences in Kerala Livestock Development Board. Kerala, India. pp. 158.
- Köhler-Rollefson, I.** 2004. Farm animal genetic resources. Safeguarding national assets for food security and trade. *GTZ, FAO, CTA*, pp. 60.
- Köhler-Rollefson, I. & LIFE Network.** 2007. Keepers of genes: the interdependence between pastoralists, breeds, access to the commons, and livelihoods. Sadri, Rajasthan, India, LIFE Network. pp. 57.
- Kosgey, I.S. & Okeyo, A.M.** 2007. Genetic improvement of small ruminants in low input, smallholder production systems: technical and infrastructural issues. *Small Ruminant Research*, 70: 76–88.
- Kumar, A., Birthal, P.S. & Joshi, P.K.** 2003. Research on crossbreeding in cattle: an analysis of its economic and social impact in India. *Agricultural Economics Research Review*, 16(2): 91–102.
- Lewis, M.** 2003. Cattle and conservation at Bharatpur: a case study in science and advocacy. *Conservation and Society*, 1: 1–21.
- Madalena, F.E.** 2005. Considerations on the management of animal genetic resources in Latin America. Proceedings of EAAP/SLU/FAO/ICAR Workshop on Sustainable Management of Animal Genetic Resources: Linking perspectives globally. 2 June 2005, Uppsala, Sweden, pp. 10.
- Malan, S.W.** 2000. The improved Boer goat. *Small Ruminant Research*, 36(2): 165–170.
- Malmfors, B.M., Smalley, M., Philipsson, J., Ibrahim, H., Anderson-Eklund, L., Mwai, O., Mpofu, N. & Rege, J.E.O.** 2002. Capacity building for sustainable use of animal genetic resources in developing countries- A new approach. 7th World Congress on Genetics Applied to Livestock Production. 19–23 August 2002, Montpellier, France. CD-ROM Communication No.29–04.
- Meuwissen, T.H.E., Hayes, B.J. & Goddard, M.E.** 2001. Prediction of total genetic value using genome wide dense marker maps. *Genetics*, 157: 1819–1829.
- Mueller, J.P., Flores, E.R. & Gutierrez, G.A.** 2002. Experiences with a large scale sheep genetic improvement project in the Peruvian highlands.

7th World Congress on Genetics Applied to Livestock Production. 19–23 August 2002, Montpellier, France. CD-ROM Communication No. 25-12.

Mueller, J.P., Poore, M.H. & Skroch, W.A. 1999. Damage assessment in Christmas tree plantations following vegetation control with sheep and geese. *Southern Journal of Applied Forestry*, 23: 11–15.

Philipsson, J., Rege, J.E.O & Okeyo, A.M. 2006. Sustainable breeding programs for tropical farming systems. In J.M. Ojango, B. Malmfors, & A.M. Okeyo, (Eds). *Animal genetics training resources*. CD-ROM, Version 2. Nairobi, International Livestock Research Institute and Uppsala, Sweden, Swedish University of Agricultural Sciences.

Sonesson, A.K. & Meuwissen, T.H. 2000. Mating schemes for optimum contribution selection with constrained rates of inbreeding. *Genetics Selection Evolution*, 32: 231–248.

Sponenberg, D. P. 2007. Rainbow livestock: Nguni cattle, Damara sheep, and indigenous goat types. *South African Studbreeder*. Issue 17. South African Studbook Authority.

Sushruta Samhita. 1985. The medical science of ancient Aryans, 2nd ed., edited and translated by A.C. Bandyopadhyaya. Calcutta, India.

van der Werf, J.H.J. 2007. Marker-assisted selection in sheep and goats. In E. Guimaraes, J. Ruane, B. Scherf, A. Sonnino & J. Dargie (Eds). *Marker-assisted selection: current status and future perspectives in crops, livestock, forestry and fish*. Rome, FAO.

Vasconcelos, J., Martins, A., Petim-Batista, M.F., Colaço, J., Blake, R.W. & Carvalho, J. 2004. Prediction of daily and lactation yields of milk, fat, and protein using an autoregressive repeatability test day model. *Journal of Dairy Science*, 87: 2591–2598.

Weigel, K.A. 2004. Exploring the role of sexed semen in dairy production systems. *Journal of Dairy Science*, 87: E120–E130.

Wurzinger, M., Ndumu, D., Baumung, R., Drucker, A., Okeyo, A.M., Semambo, D.K., Byamungu, N. & Sölkner, J. 2006. Comparison of production systems and selection criteria of Ankole cattle by breeders in Burundi, Rwanda, Tanzania and Uganda. *Tropical Animal Health and Production*, 38: 571–581.

Panellists' comments and discussion

Mr Jan Philipsson, Interbull Centre, Sweden

This paper essentially concludes that the best strategy to maintain animal genetic diversity is based on the sustainable use of the AnGR. This is a pro-active way of considering the conservation issue and it includes economic as well as natural resource aspects, and has environmental and sociocultural dimensions. Interesting examples with diverse use of the AnGR are given as illustrations, primarily from developing countries.

In reviewing the options for the sustainable use of AnGR these may indeed vary considerably between developed and developing countries, due to differences in climatic challenges, resources and infrastructure. It is underlined that genetic improvement of livestock productivity in most cases is a prerequisite for sustainability and that production systems also will be intensified. It is an important task to investigate how this can best be achieved in different regions and production systems considering possible environmental impacts. However, intensification by such interventions leading to increased fertility, survival rate and health of animal populations would under many circumstances mean considerably higher productivity and less environmental stress.

The authors emphasize the need to make use of both traditional as well as new, molecular genetic tools to catch the opportunities for genetic improvement of livestock populations. However, they also point out that considerable improvements can be achieved already with traditional knowledge and rather simple techniques. They point for example to both the use of nucleus herd selection schemes and cross-breeding for the formation of composite breeds to improve productivity while maintaining adaptation to the environment. Effective use is thereby made of genes from different populations, a method that historically always has been practised in the dynamic forming of breeds. The proposed wider use geographically of tropically adapted breeds is important and should be further explored.

Obstacles for sustainable use of AnGR which I think need to be further addressed are:

1. Improper definitions of breeding objectives to be sustainable in relation to both market conditions and a broad range of traits in animals. This is still lacking in many populations of the

developed world, and may lead to serious biological problems with the animals (e.g. declining fertility and disease resistance). In developing countries the choice of exotic breeds for use in pure-breeding and cross-breeding has too often not been properly assessed before embarking on breeding systems, that later proved to be non-sustainable, partly because of a lack of long-term breeding strategies.

2. The role of farmer involvement needs to be emphasized even more. There are neither short- nor long-term benefits of knowledge in genetics or animal breeding if this knowledge cannot be transferred to the farming community for their action. More research on this technology transfer issue and its proper consideration in development projects are needed.
3. There are generally too few trained animal breeding scientists and extension staff in most developing countries to effectively support implementation and running of sustainable breeding programmes. Also in the developed countries there is a shortage of quantitative geneticists needed for development of AnGR. Increased research supporting the sustainable use of AnGR and intensified capacity-building is therefore necessary.

In conclusion, this paper emphasizes the sustainable use of AnGR as the best method to maintain necessary genetic diversity. For that to happen, I would like to emphasize that genetic improvement programmes are greatly dependent not only on genetics, but also on policy, organizational and infrastructural issues to be sustainable.

The issue of short-term vs. long-term benefits of genetic improvement programmes was raised. It was stated that it simply takes too long for the benefits to be visible.

My answer to this is that the time it takes to see the results of genetic improvement programmes depends very much on the species, breeding strategies chosen and the production system. For species with short generation intervals, such as poultry and pigs, results of both pure-breeding and cross-breeding are realized within one to two years, whereas effects of within-breed selection of dairy cattle takes five to ten years if germplasm from other more productive strains of the same breed cannot be used. On the other hand, all genetic effects are

accumulated for each generation of selection. The later you start a selection programme with a breed, the more it will fall behind other breeds under continuous selection. However, genetic improvement can be achieved much faster through cross-breeding with a superior breed assessed to be suitable for the environment in question, and provided that a long-term strategy has been worked out and considered feasible. In many cases such a strategy ends up with a new synthetic breed as the authors of this paper have indicated.

The other point I would like to raise on this issue, and which has an impact on the outcome, is that genetic improvement programmes should not be seen in isolation. First of all they must be applicable to the prevailing environment and production systems. Secondly, improvements in management and feeding must also be made, and some of these improvements may be realized even before the genetic ones. A holistic view of an improvement programme must be aimed for, where the genetics is just one component, but an important one. Whatever the strategy chosen, a long-term perspective must always be taken.

Mr Raúl Perezgrovas, Instituto de Estudios Indígenas, Chiapas, México

I would like to thank FAO for the invitation to be a panellist during the Scientific Forum, and for the opportunity to share some comments on the paper on "*Sustainable use and genetic improvement of Animal Genetic Resources*". I would like to congratulate Chanda and the co-authors of the paper for the comprehensive coverage of such a wide number of topics regarding the sustainable use of AnGR.

I have been requested to present some reactions, and, as an animal scientist I can easily relate to all of the strategies and technical approaches presented by the different authors. It was possible to identify the key words, as they were constantly mentioned throughout the document, and some of them were:

- Animal genetic resources.
- Farm animal breeds.
- Livestock.
- Breed improvement.
- Genomic technologies.
- Animal productivity; to mention just a few.

However, as a social scientist, I noticed the low frequency in the use of "*other*" key words, such as:

- Livestock keepers.
- Small-scale farmers.
- Livelihoods.

- Local knowledge.

And this is very important when the topic under discussion is precisely the "*sustainable use and genetic improvement of AnGR*", because we know that sustainability has a social, an economic and a cultural component.

So, I present here a couple of suggestions to be considered by the agricultural technicians, the extension workers, the university professors and researchers, and the policy-makers.

First, we all need to take a more humble approach and recognize that livestock keepers can teach us a few things regarding the sustainable utilization of farm animals; it is they who have conserved and improved many indigenous or local animal breeds, making them a vital part of their livelihoods.

And second, there are many participatory methodologies that will allow the livestock keepers and the researchers, and the policy-makers to undertake a fruitful learning experience for all.

There is in the paper a very good example of how to integrate the local peoples' expertise. When talking about "*capacity-building and training*", I came across the following statement, and I quote:

"...training should build upon existing knowledge of the production system, and enable livestock keepers to make informed decisions..."

I would suggest that the same spirit is considered also when we speak of genetic improvement, promoting novel uses of animal genetic resources, and the design of sustainable breeding programmes.

Earlier today, the issue of Livestock Keepers' Rights was raised as necessary in a global action plan. I suggest that the issue of "*Listening to Livestock Keepers' Voices*" needs also to be discussed.

Ms Xuan Li, South Centre, Switzerland

First of all, I would like to extend my appreciation to FAO for inviting me to be a panellist today. I represent the South Centre, an intergovernmental organization and think tank of the developing countries, with its headquarter in Geneva, Switzerland. The member states of the South Centre comprise G-77 and China. Our mandate is to provide strategic vision and technical support for the developing countries, particularly in the field of intellectual property, trade and global governance.

Coming to the paper, sustainable use and genetic improvement is considered one of four strategic components of the *Global Plan of Action for Animal Genetic Resources*. The paper properly demonstrates that there is no inherent incompatibility between conservation and utilization if animal genetic resources are properly managed. The immediate questions arising are, what are the challenges and what management strategy will work?

Before addressing the challenges, I would like to give a brief picture on what is happening in major international fora in the field of animal genetic resources in Geneva. In WIPO-IGC (World Intellectual Property Organization Intergovernmental Committee on Intellectual Property and Genetic Resources, Traditional Knowledge and Folklore), the issue is rather limited to the discussion of plant genetic resources, and no binding international agreement has been achieved. In WTO (World Trade Organization), the matter is overshadowed by the debate on “*disclosure of source*” and “*prior informed consent*”. In WHO (World Health Organization), emphasis has been made to urge member states to establish mechanisms that ensure the routine and timely sharing of biological materials and isolates from both humans and animals, and no proper benefit-sharing scheme is available. In short, neither effective incentive to promote sustainable use of animal genetic resources nor proper international regulation on genetic resource protection has been provided.

Reading from the paper, in my view, there are at least four challenges that we are facing to ensure a sustainable use of animal genetic resources. The first challenge is to re-think livestock sector policies that “*distort the playing field*” on which indigenous breeds compete. As we have seen, farmers are often disadvantaged by subsidies on feed, artificial insemination and other inputs that tend to favour exotic breeds. From a policy perspective, there is a need to conduct some systematic policy analysis to assess the implications of the existing livestock policies, and respond accordingly.

The second challenge is, from an economic perspective, that appropriate incentive and funding mechanisms to foster innovation is crucial. Choice needs to be made between patent and alternative mechanisms, whichever is more cost-effective to foster sustainable use and innovation.

The third challenge is that the interface between animal genetic resources and intellectual property (IP) seems to be underestimated. The statement “*all the processes of relevant reproduction, data capture, statistics analysis, etc are in the public domain*” is

inaccurate. In addition, a range of rapidly developing molecular and reproductive biotechnologies also has important implications for animal genetic resources management. In order to ensure a sustainable use of animal genetic resources, efforts must be made to carefully examine the interface between animal genetic resources and IP.

The fourth challenge is that a proper scheme to ensure access to animal genetic resources and technology transfer as well as benefit-sharing is necessary. Specifically, there is a strong need to establish an international binding treaty to stimulate the sustainable conservation and use of animal genetic resources. Such a treaty should cover “*any genetic material of animal origin of actual or potential value for food and agriculture*”. The objective should target the conservation and sustainable use of animal genetic resources for food and agriculture and the fair and equitable sharing of benefits derived from their use, in harmony with the Convention on Biological Diversity, for sustainable agriculture and food security. The International Treaty on Plant Genetic Resources for Food and Agriculture serves as an excellent analogy. Considering the slow movement in WIPO, WTO and WHO on genetic resource discussion, Member states are encouraged to work towards the establishment of an efficient, effective and transparent multilateral system at FAO, to facilitate access to animal genetic resources for food and agriculture, and to share the benefits in a fair and equitable way.

To conclude, if properly managed, a global action plan for animal genetic resources will be a major contribution to the implementation of the Convention on Biological Diversity in the field of food and agriculture. Efforts should be made to establish a series of coherent policy and legal regimes to achieve the purpose above.

Summary of plenary discussion

The meeting was then opened for general discussion and interventions from the floor. Key issues raised during this discussion included:

- The need to reinforce research, human resources and institutions, particularly in the developing world.
- The significance of buffaloes in the context of South Asia and the limited attention that this species attracts from international donors and researchers.

-
- The need to better utilize existing technologies and not always rush to adopt new technologies.
 - The importance of characterization as an element in the process of promoting sustainable use – for example in the identification of breeds with specific attributes such as disease resistance.
 - The need for information on breed characteristics to be effectively disseminated.
 - The problem of reconciling the short-term needs of farmers with the longer-term objectives associated with sustainable utilization.
- Responses and final comments of the authors included the following points:
- Better understanding of animal genetic resources is vital to achieving sustainable utilization.
 - Existing knowledge, including local knowledge, needs to be built upon and better integrated into management strategies.
 - Community-based structures are the building blocks of sustainability.
 - In addition to North–South collaboration, South–South cooperation in the utilization of animal genetic resources is needed.
 - Ensuring sufficient resources for the management of animal genetic resources requires better awareness of the importance of livestock production.
 - There are links between the utilization of crop and animal genetic resources, and research efforts in the two fields can be complementary.

Conservation of animal genetic resources: approaches and technologies for *in situ* and *ex situ* conservation

J.A. Woolliams¹, O. Matika¹ & J. Pattison²

¹Roslin Institute (Edinburgh), **Roslin**, Midlothian, EH25 9PS, United Kingdom

²Centre for African Studies, University of Edinburgh, School of Social and Political Studies, University of Edinburgh, 21 George Street, **Edinburgh** EH8 9LL, United Kingdom

Summary

Livestock production faces major challenges through the coincidence of major drivers of change, some with conflicting directions. These are:

1. An unprecedented global change in demands for traditional livestock products such as meat, milk and eggs.
2. Large changes in the demographic and regional distribution of these demands.
3. The need to reduce poverty in rural communities by providing sustainable livelihoods.
4. The possible emergence of new agricultural outputs such as bio-fuels making a significant impact upon traditional production systems.
5. A growing awareness of the need to reduce the environmental impact of livestock production.
6. The uncertainty in the scale and impact of climate change. This paper explores these challenges from a scientific perspective in the face of the large-scale and selective erosion of our animal genetic resources, and concludes that there is a stronger and more urgent need than ever before to secure the livestock genetic resources available to humankind through a comprehensive global conservation programme.

Résumé

La production animale se trouve face à des défis importants dus à la coïncidence de différents facteurs de changements, certains en évident conflit par rapport à leur orientation. C'est-à-dire:

1. Changement sans précédent de la demande au niveau mondial de produits traditionnels tels que la viande, le lait et les œufs.
2. Changements importants dans la distribution démographique et géographique de la demande.
3. Le besoin de réduire la pauvreté dans les communautés rurales en offrant un moyen d'existence durable.

4. L'émergence due à la possibilité de nouveaux produits de l'agriculture tels que le combustible biologique qui a un impact significatif sur les systèmes traditionnels de production.
5. Une majeure considération de la nécessité de réduire l'impact environnemental dû à la production animale.
6. L'incertitude sur le niveau et l'impact du changement climatique. Cet article étudie les défis du point de vue scientifique dans le cas d'une érosion sélective des ressources génétiques animales à large échelle. En conclusion, il existe plus que par le passé l'urgence d'assurer la disponibilité des ressources génétiques animales pour l'utilisation humaine à travers un programme mondial de conservation.

Resumen

La producción ganadera se enfrenta con importantes desafíos debido a la coincidencia de varios factores de cambio, algunos de los cuales en claro conflicto con respecto a su orientación. Estos son:

1. Un cambio sin precedentes en la demanda a nivel mundial de productos tradicionales tales como la carne, la leche y los huevos.
2. Importantes cambios en la distribución demográfica y geográfica de la demanda.
3. La necesidad de reducir la pobreza en las comunidades rurales ofreciendo una renta sostenible.
4. La emergencia debido a la posibilidad de nuevos productos de la agricultura tales como el combustible biológico que tienen un impacto significativo sobre los sistemas tradicionales de producción.
5. Una mayor conciencia de la necesidad de reducir el impacto ambiental debido a la producción ganadera.

6. La incertidumbre sobre el nivel y el impacto del cambio climático. Este artículo estudia estos desafíos desde un punto de vista científico en el caso de una amplia escala y erosión selectiva de los recursos zoogenéticos. En conclusión, existe más que nunca una fuerte y mayor urgencia de asegurar la disponibilidad de recursos zoogenéticos para uso humano a través un programa mundial de conservación.

Keywords: *Cryoconservation, Breed erosion, Climate change, Genomic revolution, Managing uncertainty, Inbreeding, Somatic cell nuclear transfer (SCNT).*

Darwin, Dylan and egg baskets: the scientific case for conservation

The first of this series of papers has described the trends that are operating on animal genetic resources for food and agriculture throughout the production systems of the world. A much simplified summary is that livestock are a focal point for many drivers of change related to their ability to lift people out of poverty and into sustainable livelihoods, to satisfy global demand for livestock products and promote international trade, and the need for livestock production to reduce its impact on the environment and its contribution to global warming. A broad conclusion is that there will be a need for sustainable intensification of livestock production. The other papers in this series have indicated the scientific background of how this might be better achieved both now and in the future, and the needs for scientific information to support decisions on animal genetic resources.

The current drivers of change have led to a large number of breeds slipping between the cracks as production environments change, and change rapidly. Production environments are now shaped in part, to a greater or lesser extent, by the economics of the current global market, both for inputs such as feed and water for animals, and outputs such as meat, milk and eggs. Broadly, breeds survive if they are fit for the market conditions that prevail, and decline towards extinction if they are not, a parallel of Darwin and natural selection. The decline in numbers further increases vulnerabilities to other catastrophic events, such as conflict, disease, flood or drought. box 1 examines the scale of erosion of the world's animal genetic resources using data from *The State*

of the World's Animal Genetic Resource for Food and Agriculture (FAO, 2007a) and concludes that as many as one in three breeds may be at risk of extinction, with a further one in ten already extinct.

Is this breed erosion a problem? Maybe not, if there is certainty and stability, but otherwise definitely yes. Unfortunately science tells us that, to quote Bob Dylan¹, “*the times they are a-changin’*” and that some of our past certainties may disappear. There is now an established scientific consensus that there will be a period of relatively rapid climatic warming over this century, and that human activity has contributed, and continues to contribute to this trend (IPPC, 2007).

As an example, box 2 shows the projected change in just one key agricultural parameter for one continent, the length of the growing season in Africa; other parameters such as the projected changes in the frequency and severity of droughts and floods are equally relevant. Box 2 illustrates an important additional point in that the degree of change and its agricultural and socio-economic consequences (see section 2.A.3 of *The State of the World* report for a brief overview of some of these) will depend on our future actions and their coordination on a global scale. These actions remain uncertain, but if they are limited or ineffective, more far-reaching consequences are expected. Further, as in all models, there are uncertainties resulting from limitations in our scientific knowledge and understanding: some scientists think the consensus position underestimates the extent of change, while others think the change is overestimated. Therefore, beyond the familiar uncertainties of market trends and the economic values of products, there is now an additional uncertainty of a magnitude and dimension that is beyond the experience of the modern world. In short, there is change rather than stability, with uncertainty writ large.

As a consequence of these developments, the chances are higher than ever before that what may fit the needs of today may not fit the needs of our children's children. Science shows how the genetic diversity that we have within any of our livestock species today can be regarded as being partitioned between breeds and within breeds. Estimates of the magnitude of the diversity between breeds are well

¹Bob Dylan, 1963. *The Times They Are A-Changin’*. Popular Song.

Box 1. A brief review of for mammalian and avian species based upon *The State of the World's Animal Genetic Resources for Food and Agriculture*

The table below summarizes the risk status of breeds in 2006, taking the data presented in tables 12 and 13 of Part 1 – Section B.5 of *The State of the World* report. At first sight it might be concluded that “only” one in five of all mammalian and avian breeds are “at risk” of extinction, although closer examination shows that only one in three can be viewed as “not at risk”. The discrepancy arises from the “unknown” category.

Risk Status	Mammalian		Avian		Total	
	Number	%	Number	%	Number	%
Critical	255	4.6	245	12.2	500	6.6
Critical-maintained	59	1.1	20	1.0	79	1.0
Endangered	406	7.3	287	14.3	693	9.2
Endangered-maintained	160	2.9	55	2.7	215	2.8
At risk	880	15.8	607	30.2	1 487	19.7
Not at risk	2 129	38.3	521	26.1	2 650	35.1
Unknown	1 907	34.3	825	41.3	2 732	36.1
Extinct	643	11.6	47	2.3	690	9.1
Total	5 559	100.0	2 000	100.0	7 559	100.0

It is possible to throw some light on the true state of the “unknown” breeds by analysis of the information on breeds that were “unknown” in 1999 but for which more precise information is now available. Examination of Tables 19, 21 and 22 of *The State of the World* report shows that a total of 238 breeds were classified as “unknown” in 1999 and classified as either “at risk”, “not at risk” or “extinct” in 2006. Of these 40 percent were “at risk”, 57 percent were “not at risk” and 3 percent were “extinct”. Using these figures as predictors of the true status of “unknown” breeds in 2006, the best estimates for all breeds in 2006 becomes 56 percent “not at risk”, 34 percent at risk” and 10 percent “extinct”, i.e. over one in three “at risk”, a further one in ten “extinct”, and just over one in two breeds “not at risk”. A further point to note is that among the breeds known to be at “at risk”, only one in five have some form of *in vivo* conservation measure in place. In conclusion, the position of global animal genetic resources is far from secure.


in excess of 50 percent of the total diversity for traits that are related to fitness for an environment (Cundiff *et al.*, 1986). This leads to the inescapable conclusion that the between-breed component of diversity is very important for addressing a broad range of environmental conditions. The concern is that the breeds thriving today are primarily those fitted to high inputs and high outputs. Given uncertainty over the production systems that livestock will face in the future, for example the possible diversion of crops to biofuels, the breeds thriving today may not meet all our needs for tomorrow. Experience shows that we cannot change

the genetic constitution of existing breeds rapidly enough to manage this uncertainty. Paradoxically, the ease of breed substitution which has placed so many breeds at risk is the primary reason why the *full* range of breeds we have today is so valuable for the future. There is a saying, “*don't put all your eggs in one basket*” and currently the world is moving towards a single basket of livestock.

The current state of insecurity of global animal genetic resources was discussed above. Global climate change might be anticipated to increase the insecurity of animal genetic resources, both directly through more extreme climatic events, even if the

Box 2. Scenarios illustrating the potential impact of global climate change on the length of the growing season in Africa, and the degree of uncertainty arising from differing assumptions

Brief descriptions of the scenarios are given in the notes below, but the two maps represent the extremes for this attribute taken from a range of scenarios considered by the authors. The colours, from deep red, light red, white, light green to green represent loss in excess of 20 percent, loss of 5-20 percent, change less than 5 percent, gain of 5-20 percent and gain in excess of 20 percent respectively.



Notes:

1. Regions gaining 5 percent or more in the growing season occupy considerably less than 1 percent of the coloured regions for either map; examples of such regions in both maps are a minority of the coloured region on the North African coast, and to the south of the Great Rift Valley in Ethiopia.
2. The maps are derived using the Hadley Centre Coupled Model version 3. The 2 scenarios shown are: on the left, A1F1, assuming very rapid global economic growth, global population peaking mid-century, rapid introduction of new and efficient technologies, with an emphasis on fossil fuel energy; on the right, B1, assuming rapid change globally to service and information economies, global population peaking mid-century, introduction of clean and efficient resource technologies, with global planning but no new climate initiatives.

The maps are reproduced from *Mapping climate vulnerability and poverty in Africa* by kind permission of P.K. Thornton.

animal genetic resources appear well-adapted today, and indirectly through competition for essential resources such as food and water leading to an increased risk of conflict.

There is a stronger case than ever before for action to secure animal genetic resources through conservation measures. This case is based on managing the uncertainties in future food security, and extends beyond our raised awareness of the need for managing genetic resources and ecosystems that flowed from the Convention on Biological Diversity. The scientific logic is to develop and implement a global conservation strategy to create a secure backup, a "second egg-basket". This is a conclusion reached by at a meeting of experts in Montpellier (Gibson *et al.*, 2006), and is one of the key action points identified by FAO in its *Global Plan of Action for Animal Genetic Resources* (FAO, 2007b). The underlying operational science will be returned to later in the paper.

Opening the conservation toolbox

Conservation can take different forms, depending on needs and resources. The major classification is whether or not the conservation is *in situ* or *ex situ*: the former describes a situation where conservation takes place in the environment in which the breed has been developed, and of necessity involves conserving live animals over generations. In contrast, *ex situ* conservation takes place outside the native environment. It may or may not involve live animals, as there is the possibility of storing gametes, sperm or oocytes, or cells with the potential to develop new animals, e.g. embryos, using the scientific advances of cryopreservation. There is a preference for *in situ* conservation, recognized by the Commission on Genetic Resources for Food and Agriculture (FAO, 1998a).

Why this preference for *in situ* conservation? The justification lies in the opportunity for the breed to continue to develop in its native environment, and in doing so the qualities that adapt it to the environment continue to be maintained through continued selection pressure. When the environment changes in one or more aspects, further selection builds upon an adapted foundation. Some adaptations, such as an ability to withstand drought or a resistance to a disease may be easily observed; others may be identified as part of the characterization process; others may be recognized unexpectedly and in crises. An

illustration of the potential importance of *in situ* conservation is the North Ronaldsay sheep, native to the United Kingdom, which was habitually kept in an environment where seaweed was important component of its diet. Upon removal from this environment a large proportion of sheep died from copper toxicity. Further investigation showed that the ability to extract copper from the seaweed, with high efficiency, was an important adaptation of the breed to their native diet. If there had been no recourse to an *in situ* population, the surviving *ex situ* population would have been strongly selected *against* the very adaptation which had made the breed potentially unique!

Given the potential benefits of *in situ* conservation, why there is a need to consider *ex situ* measures? The immediate answer is that resources and commitment of farmers may not be forthcoming in the face of the pressures that have led to the need for the breed to be conserved, as it is seen as failing to meet the current needs. Alternative *ex situ* options are therefore necessary. These may include the establishment of live populations of the breed in institutional or NGO environments that may differ from the native environment, or by adopting cryoconservation. The choice of conservation options is not a strict dichotomy, as combinations of *in situ* and *ex situ* may be used. In particular, the idea of *in situ* populations supported by cryoconservation has become the method of choice in many developed countries.

Cryoconservation has a significant profile in livestock conservation. The development, refinement and practice of the associated cryopreservation techniques has been driven by the interest of breeding organizations in many livestock species, because of the improved genetic progress that can be achieved by using these techniques within breeding programmes. Nevertheless, while cryoconservation is a powerful option for conserving animal genetic resources, there are significant limitations: first, there are major differences among the livestock species in terms of the ease and effectiveness of applying the techniques (discussed further in following sections of this paper); second, even in cattle where techniques are well developed success may be achieved only after a lot of time and resources, e.g. see Box 102, section 4.F.7 of *The State of the World* report and third, the cryopreservation of semen, oocytes and embryos requires the use of liquid nitrogen. Use of liquid nitrogen is not a universal option, as a significant number of countries have no, or only limited, experience of such procedures. Box 3 summarizes the information on this topic

presented in *The State of the World* report on this topic. It is clear that global conservation capabilities would be advanced if the capacity to use liquid nitrogen were to be made universal.

Conservation can be viewed as the creation of a gene bank containing live animals, or cryopreserved gametes and cells, or both. The gene banks secure the animal genetic resources, and in doing so provide new opportunities. One such opportunity is to enhance the exchange of animal genetic resources, and allow the benefits from animal genetic resources developed in one country to be shared elsewhere. Examples show that the importance of a breed can sometimes be more sustained in a country other than the one in which it is developed: for example the utilization of Sahiwal cattle (from South Asia) in Kenya.

development can improve matters further. The basic integration of scientific approaches to conservation is described in the “*Guidelines for management of small populations at risk*” developed by FAO (1998a), which cover all activities relating to conducting censuses and compiling inventories, through considering what conservation options may be appropriate for a single breed, how actions may be prioritized, through to the technical guidelines on setting up and managing gene banks of live animals and cryopreserved gametes. The techniques for cryopreserved gametes were reviewed and updated more recently by the European Regional Focal Point for animal genetic resources (ERFP, 2004). This paper will only introduce and discuss areas where the underlying science has developed or where new conservation needs have been identified.

Sharpening the tools: the contribution of science

This section will describe how science can help to make conservations more tractable and more effective, and how science currently under

How many minutes to midnight?

To be effective, conservation needs to be timely. Preserving the gametes of the last dodo would have had little impact in terms of preventing the loss of the species. The proactive identification of breeds at

Box 3. A brief review of worldwide practice of techniques relevant for cryoconservation

The following is based upon data contained in section 3.D.2 of *The State of the World* report¹ on use of artificial insemination (AI), which is a more widespread technology and is and more widely applicable across the range of livestock species than embryo transfer. Only 84 percent of the 148 countries providing data report the use of AI in routine practice, and those not using AI were primarily situated in SW Pacific, Africa and Asia regions. However, this fraction is an upper bound on the routine use of liquid nitrogen, since AI may be carried out with fresh semen, rather than frozen, without requiring cryopreservation. Furthermore, whilst the use of AI may indicate capacity for storage and use of cryopreserved semen, it need not imply the routine use of procedures for collection and cryopreservation of semen, both essential for cryoconservation, as in many cases it was reported that exotic semen was being used.

Section 3.D.2 also demonstrates that the practice of AI is primarily directed towards cattle: whilst only one of the 84 percent of countries reporting use of AI fails to mention cattle, only 34 percent and 21 percent of countries report the use of AI for pigs and sheep respectively, the two next most common species for AI use. As with cattle, these figures are upper bounds on the fraction of countries that routinely collect, store and use cryopreservation for these species.

risk for conservation actions is an important tool for monitoring animal genetic resources, yet it is a tool that remains blunt. One example is the identification of breeds that are at risk as a result of being confined to a small geographical area (a condition referred to as endemism), although they may be locally numerous. Such breeds may be at risk from catastrophic events. This was illustrated clearly in the United Kingdom during the foot and mouth disease epidemic of 2001. The regulations for controlling this disease within the European Union (EU) led the United Kingdom to cull livestock on a large scale, with the result that few individuals within the focus of the epidemic were left alive. Unfortunately, this focus closely coincided with the centre of population of the Herdwick sheep breed, which is numerous locally but restricted in its geographical spread. Recognition of the plight of this breed led to a number of emergency conservation actions during the epidemic.

Quantitative measures of the risk associated with endemism have not been formalized. Risk may vary across regions; for example the area affected by a catastrophic drought may be wider than the area affected by a catastrophic fire. Thus, an assessment of the risk associated with endemism requires careful analysis of the potential impact of catastrophic events in the region in question. Attempts have been made (Gandini *et al.*, 2004) to develop approaches to the calculation of risk status that are not merely functions of population numbers. Several such methods, of varying complexity, are in use, but these require further socio-economic and genetic inputs before they can be considered reliable. Limitations will remain, while it may be possible to obtain better data for quantifying some risk factors, such as the degree of cross-bred matings, other risks such as conflict may be harder to quantify objectively.

Turning safety nets into springboards

Conservation, particularly *in situ* conservation, has a dual purpose. It was introduced above as a "second egg-basket", a form of safety net. However, considerable socio-economic research has been carried out in an attempt to understand how this net can become a springboard for the return of a breed to the mainstream, in which no special measures beyond the market are required to maintain the population. In the FAO guidelines (1998a), the core approach to this transformation was establishing the true market value of a breed, emphasizing the need to consider lifetime

performance and lifetime contributions rather than simple measures of product yields under a regime conducive only to high outputs. This consideration and the options that exist for improving the recognition of full market value remain important. However, it is now widely accepted that a breed's value exceeds the expected market value of its products. Two further concepts can now be added to the valuation process to demonstrate this: first the contribution of a breed to managing climatic uncertainties and to recovery from environmental crises faced by farmers, and second the valuation of a wide range of potential non-market services. Box 4 illustrates why these concepts are important to maintaining breed populations and securing the livelihoods of farmers.

A more controversial area of economic science associated with conservation of live animals is the use of subsidy for maintaining breeds. An example of the complexity of this area is the mixed success of measures implemented by the EU, which has in the past supported such actions. While the subsidy halted the decline in census numbers of breeds covered by the scheme, there was a barrier to population growth caused by existence of a threshold population size (headage) below which a breed was considered eligible for subsidy: a trend existed for breeds to sit just below this threshold size for fear of losing subsidy. Therefore, subsidy is an effective safety net but an ineffective springboard! Consequently, subsequent EU support is more concerned with characterization and helping breeds to develop added values. This problem with headage barriers can also be faced by NGOs. One such example is the Rare Breeds Survival Trust in the United Kingdom, which has re-vamped its qualifying conditions to allow it to act more effectively as a springboard for moving breeds beyond "at risk" status.

The genomic revolution

The genomic revolution with its tools of complete genome sequence, dense high-throughput genotyping at increasingly affordable prices, and rapid detection of genetic polymorphism are primarily new tools of characterization - to go from sequence to consequence. These developments will lead to an advance of an order of magnitude beyond our current understanding, are addressed in the companion paper. However, in the context of this paper, DNA has "traditionally" been used as the source of DNA markers with which to measure a genetic distance between breeds, or to measure

Box 4. Beyond the expected product value

Managing uncertainty. This can be illustrated by the considerable variation that exists between breeds in terms of their ability to withstand drought, which is empirically seen to be much greater than variation within breeds. Farmers in many regions rely on their livestock as a means of maintaining livelihoods through droughts. A breed providing this service may perform a more vital role than a breed that provides better returns in the good times but fails in the bad times leaving the farmer without support. Therefore, the valuation of a breed's performance needs to take account of the foreseeable crises that affect the production environment in which it is kept, rather than the average conditions. This process of valuation does not need to involve a straight choice between one breed and another: farmers in many regions recognize the benefits of maintaining a mixed economy of breeds, maintaining highly productive breeds to capitalize on the good times, while maintaining the robust breeds as insurance for the bad times. This latter role maintains populations, while securing livelihoods in the fullest sense of the word "secure".

Non-market values. Many products and services generated by livestock breeds are not marketed; these often include: transportation and traction; manure as fertilizer or fuel; fibre and skin for clothing; household meat, milk and eggs. Breeds may differ in their ability to provide these services. In addition livestock provide financial and socio-cultural services.

Financial services (Dorward *et al.*, 2005) can depend on the animal having longevity in the environment in which it is maintained and retaining productivity in the harsher times of the production cycle. The ability to provide such services will clearly depend on the breed. Examples of financial services include:

- *buffering* (or consumption smoothing) whereby investments are made in livestock during periods when production or income exceeds consumption needs and then these investments are drawn upon later in the season when lower production and income are not sufficient to support consumption needs.
- *Saving*, whereby animals are kept explicitly to provide for some major expenditure (such as a major purchase or investment, or expenditure on school fees or an important social activity).
- *Insurance*, where animals are kept solely for the provision of insurance against unexpected events that either reduce income or require additional expenditure, such as accidents or illness.
- *Collateral* for borrowing.

Sociocultural services include important *social integration* functions in livestock keepers' society and culture. Traditional breeds may confer status on the individual owners, and may contribute to the sense of identity of whole communities through associations with traditional agricultural systems or landscapes, folklore, cuisine, ceremonies, and crafts. It should also be kept in mind that commercial breeders gain status when their animals are priced or exhibited.

genetic variation between and within breeds. These measures are then used to prioritize actions with a view to maximizing the diversity conserved (Eding & Bennewitz, 2007). There is an unresolved debate over the use of such methods, as there are sound arguments for basing actions on established and valued phenotypes rather than the measures based

on anonymous marker DNA. One reason for basing actions on phenotypes is that empirically there appears to be a poor correlation between quantitative measures of diversity based upon phenotypes and molecular measures of diversity (Reed & Frankham 2001). A future outcome of the genomic revolution may be to improve this

correlation through the richness of the information obtained with the new genomic tools. However if, as argued previously above, there is a need to set in place a comprehensive global conservation strategy, rather than one led by a process of choosing among breeds, then the issues surrounding prioritization among breeds may become more academic.

Dealing in diversity

One of the perennial concerns of managing populations *in vivo* in conservation schemes has been the fear of inbreeding and the loss of diversity. Inbreeding is an unavoidable and natural process present in all populations, and as Bryson (2005) points out, to avoid all inbreeding in humans back to the time of Julius Caesar would require more humans than have ever lived! There are considerable scientific arguments to show that problems associated with inbreeding are related to the rate at which it occurs, not the observed degree, with faster rates associated with higher risks of genetic problems. This was addressed in the FAO guidelines (1998a), but science has continued to advance in this area. New techniques have shown how this rate of inbreeding can be managed simultaneously with maximizing selection opportunities under a range of circumstances. Combining these twin objectives is important for the management of breeding within conservation schemes, as populations may need to have deleterious genes removed, which is a form of selection, or may be part of a selection programme to improve their economic viability. The same core technique can be modified to minimize the rate of

inbreeding given the resources available. Such techniques benefit from establishing the sires and dams of offspring each generation to build the pedigree. See box 5 for more details. In summary, these techniques move breeders from contemplating a win-lose "trade-off" between selection gain and inbreeding, to taking advantage of a win-win by obtaining the maximum gain whilst managing rates of inbreeding.

Not all the issues of inbreeding are concerned with live animals: in cryopreserved gene banks the diversity "put in" limits the diversity "taken out". The diversity put in depends on how donor animals are sampled- both how many and which ones. In the event of a crisis, expending time considering this may be a luxury. However, there are established techniques for identifying which individuals from a breed should contribute, and the size of their contribution, in order to maximize the genetic variation that can be mobilized from the cryopreserved bank, even where there are constraints on the numbers sampled. These are most easily applied if pedigrees are available, using the same core technology as for conservation schemes using live animals.

Managing expectations from cryoconservation

In a cryopreserved gene bank there is no interest on deposits - you only get out what you put in, at best! This observation is central to the design of cryopreserved gene banks. Such banks require funds, effort and commitment to collect samples and to maintain them ready for a time of need, and it is

Box 5. Managing rates of inbreeding in live animals

Breeding schemes may have conservation or selection objectives, but all schemes can be broadly classified into two groups: more sophisticated schemes with extensive pedigree recording and where genetic evaluations for selection are computed by combining information on a candidate and its relatives; and other schemes that are limited in their scope to accumulate full pedigrees on offspring, and/or rely on mass selection procedures. For the first group of schemes the sophistication of the scheme is sufficient to incorporate optimal contribution methods (Meuwissen, 2007) into selection procedures to manage rates of inbreeding. For the second group the rate of inbreeding can be managed with the use of a simple table, based on the ratio of number of breeding females to breeding males and the lifetime family size of a breeding female (Woolliams, 2007). The latter table is a more developed version of the T4.1 given in the FAO Guidelines for managing small populations at risk (FAO, 1998a).

vital that in the time of need the gene bank is fit for the purpose. The FAO guidelines (1998a) introduced clear and valuable objectives for setting up gene banks as templates for others to develop and customize to specific needs. There are now examples (Roughsedge *et al.*, 2006), such as the semen archive linked to the United Kingdom's National Scrapie Plan in which the sample numbers and sampling plan are linked to the objectives to be met in the future, *after* the semen is withdrawn from the bank. What was recognized by the FAO guidelines (1998a), and is now becoming more widely accepted, is that the amount of germplasm required for worthwhile objectives may be large and/or time consuming to acquire. It is essential that the managers of a cryopreserved gene bank recognize not only what the use of the stored material can achieve, but also what it *can't* achieve, as false expectations inevitably lead to poor strategic decisions.

Achieving more with less

It was already remarked in box 3 that most use of cryopreservation techniques for breeding surrounds cattle, with little use in some other species. So it is to be expected that the effectiveness of cryoconservation of gametes and their use post thaw varies widely between livestock species. This is illustrated by box 6, which is extracted from the FAO guidelines (1998a), which shows large differences between species in the time taken to collect sufficient semen to achieve the same package of measures defined by quantified outcomes from using the semen.

Furthermore, only for a minority of livestock species is it possible to routinely restore an animal with an intact genome of a breed produced entirely from cryopreserved material, i.e. an embryo, or cell, or gametes of both sexes. This is not yet possible in practice for any poultry species. The relevance of this is that for those species where it is not possible, re-establishment of a breed from cryopreserved material must involve another breed and repeated backcrossing. Important incremental advances continue to be made in the broad range of cryopreservation techniques, partly through the pull of mainstream animal breeding seeking new opportunities. Examples of notable improvement are the effective cryopreservation of oocytes in cattle, and the ability to collect and cryopreserve epididymal spermatozoa in several species. The latter adds a back-up tool of collecting male gametes from abattoirs, but such a course of action must not

risk breeding males or potential breeding males of a breed at risk.. However significant and important challenges remain and some are listed in box 7.

One important new opportunity in conserving breed diversity is the potential use of somatic cell nuclear transfer (SCNT), leading to cloning (Wilmot *et al.*, 1997). This is perhaps paradoxical, as cloning acts against diversity by creating individuals with identical genotypes! The explanation of this paradox is that the initial steps in the process involving the collection, preparation and storage of cells prior to nuclear transfer is a much more flexible technique, requiring fewer facilities, than the collection of gametes for cryopreservation (FAO, 1998b; Woolliams & Wilmot, 1999). FAO identified SCNT in 1997 as a viable option for emergency conservation actions where other more established techniques may be difficult to implement. Since then SCNT has been demonstrated in a wider range of livestock species, and its efficiency appears to be increasing in many parts of the world (Box 8). Given the developments in this field, the scope of application of SCNT and the recommended procedures for using SCNT in conservation actions should be reviewed and revised.











Ensuring best practice

The previous sections have demonstrated that science continues to make important and valuable advances in sharpening the tools conservation more effective in achieving a diverse set of objectives. The state of the art in this area was drawn together in 1998 by FAO to ensure best practice, and some aspects of cryopreservation were reviewed by ERF (2004). It would be timely to comprehensively refresh these guidelines.

Meeting the Challenge

Previously, it was argued that there is a need to establish a comprehensive conservation strategy for animal genetic resources in the face of the global trends and growing uncertainties described in the first paper in this series. Experience has shown that securing animal genetic resources is best carried out proactively, giving time for the development of effective *in situ* conservation schemes wherever possible. This will not be possible in all cases, and securing the *full* range of animal genetic resources, as argued previously, will require the resources to provide a cryoconserved backup of all breeds. As

Box 6. The time taken to acquire sufficient cryopreserved semen for achieving the FAO “default” package of objectives for ten livestock species (see Note 1 below). The numbers of samples required for the package are defined by requirements after use post-thawing, i.e. what is ultimately achieved from using the semen

		Days required to complete sample collection	
Mammals	Buffalo		60
	Cattle		11
	Goat		3
	Horse		208
	Pig		123
	Rabbit		10
	Sheep		10
Poultry	Chicken		153
	Duck		191
	Turkey		492

Notes

1. The “default” package is detailed in Section 5.2.6 of *Guidelines for management of small populations at risk* (FAO, 1998a) and includes semen for quantified sub-objectives involving re-establishment, supporting *in vivo* conservation, new breed development, and scientific research.
2. The numbers of samples required for the package are defined by requirements after use post-thawing, i.e. what is ultimately achieved from using the semen.
3. The times indicated are taken from *Guidelines for management of small populations at risk*, and are based upon leading technology current in 1998. While these times have been reduced for some species as a result of subsequent research, the large differences between species in required time will remain.

identified in box 3, such a strategy would require an extension of current capacities: cryopreservation techniques are not yet a global technology although routine in many countries, and species other than cattle would need to be addressed. There would be a need to refine the techniques for several species, with particular attention given to poultry. However, it is best to start now with current best practice rather than wait with animal genetic resources unsecured and at risk.

Coordination of gene banks will be needed either through multilateral or bilateral agreements. In this context, there is a need to resolve how cryoconserved material can be stored in duplicate (or more) locations, to reduce the risk of catastrophic

failure of one; how access and use can be made timely and traceable, with appropriate security to manage disease pathogens; and how replenishment of the gene bank can be achieved after access and use. These aspects are discussed in the FAO guidelines (1998a) and ERF guidelines (ERFP, 2004), but the principles contained therein need to be fleshed out. Of primary importance is the principle that such gene banks should encourage use – provided such use is equitable – as it is to the benefit of all.

Large-scale conservation cannot be achieved overnight for more than 7 000 breeds of domestic livestock! Operationally, in the face of the many drivers for change in what we require from animal

Box 7. Desirable advances in cryopreservation efficiency for the purpose of conservation

1. Reducing the scale of variation between species in the time taken to obtaining sufficient semen (or embryos) for delivering an identical quantified outcome post-thaw.
2. Developing a practical procedure to produce an intact genome of a poultry breed entirely from cryopreserved material.
3. Establishing reliable procedures in a range of species for obtaining thawed embryos for transfer that have little or no variation in the numbers of embryos per embryo (or oocyte) donor. Depending on the technique used for embryo or oocyte recovery this variation can be considerable and can create a serious lack of diversity in the resulting offspring. This is often ignored in simple formulae for number of embryos required, but the diversity “in” determines the diversity “out”!
4. Developing measures on semen pre-freezing to predict semen quality post-thaw. This would increase success rates per unit of stored semen, reduce numbers of doses stored and the reliability of outcomes post thaw. However the time taken to collect the semen might not be reduced.
5. Refining strategies for making best use of cryopreserved semen and embryos to re-establish extinct breeds (Boettecher *et al.*, 2005). More rapid re-establishment would encourage more use of gene bank material in such cases.

genetic resources a strategy is required to capture the diversity these breeds represent, and to ensure that few, if any, slip between the cracks. Some components of this strategy can be suggested. As breeds are more likely to get lost in more rapidly changing systems, an initial step would be for institutions funding development programmes to be proactive in requiring project proposals to identify conservation needs, and to supply costed and timebound plans for addressing these needs that would be available for review and eligible for funding. Such plans would be easier to draw up and organize if they were to be based upon “default” packages of quantified sub-objectives for the cryopreserved material, such as that suggested by FAO (1998a), or successor guidelines, which may then be customized to meet particular needs, if appropriate. A further important step is to identify an “emergency” package for geographically restricted breeds in the event of catastrophic events, such as drought and disease, and a fund for putting this into action when required. Such a package may require a range of options, including the collection of somatic cells, depending upon capacity in the affected area, the need and the time available. With these steps in operation, gaps in *ex situ* collections could be assessed to identify the need for further actions. None of these steps preclude the

development of regional or national initiatives based on their own priorities.

Conclusion

Livestock production faces major challenges through the coincidence of major drivers of change, some with conflicting directions. These are:

1. An unprecedented global change in demands for traditional livestock products such as meat, milk and eggs.
2. Large changes in the demographic and regional distribution of these demands.
3. The need to reduce poverty in rural communities by providing sustainable livelihoods.
4. The possible emergence of new agricultural outputs such as biofuels making a significant impact upon traditional production systems.
5. A growing awareness of the need to reduce the environmental impact of livestock production.
6. The uncertainty in the scale and impact of climate change. These challenges, with their inherent unpredictability, should be met by first securing the livestock genetic resources that are available to humankind.

Box 8. Somatic cell nuclear transfer and cloning

Somatic cell nuclear transfer (SCNT) was first demonstrated in sheep with the creation of Dolly by Wilmut and co-workers (Wilmut *et al.*, 1997) in 1997. In principle, this technique allows the creation of large numbers of animals with identical genotypes, by transferring a nucleus from a donor cell into an enucleated oocyte to create an embryo for transfer. Since Dolly the technique has been demonstrated in several mammalian livestock species: cattle (1998), goats (1999), pigs (2000), rabbits (2002) and horses (2003). The technique has also been demonstrated in rodents, dogs, cats and ferrets, leading to the hypothesis that SCNT may be feasible for all mammalian species. It has yet to be demonstrated in any avian species.

Although much of the public's attention has been drawn to its potential use for commercial cloning on demand, SCNT has properties that make it an attractive proposition for use in conservation schemes. An outline procedure for use in conservation would be to collect tissue samples, e.g. skin samples from live animals, prepare the cells for culture and store. When required for re-establishing a live animal, the cells would be thawed and used for nuclear transfer to create an embryo that could be then cultured *in vitro* and finally transferred to a recipient animal. Neither the donor of the enucleated oocyte nor the recipient need be the same breed as the nucleus donor.

The strengths of SCNT compared to gamete or embryo cryopreservation are primarily in the collection and storage of material:

- The cost of equipment and training required for collection and initial treatment of tissue samples is comparatively low.
- Samples that have been given an initial treatment can be transported back to a central laboratory for further processing and cryopreservation over a relatively long time period, unlike the near-immediate and on-site cryopreservation required for gametes and embryos.
- It may be possible to recover and re-process cell lines after accidental thawing, providing this is identified early enough, unlike thawed gametes and embryos.

The weaknesses of SCNT are primarily in the use of the cells post-thawing:

- low efficiency of providing viable embryos; and
- increased risks of disorders at birth, sometimes fatal, associated with sub-optimal embryo culture procedures.

As early as 1997, FAO had identified SCNT as a viable technique for emergency conservation actions. Since then, the technique has been shown to be feasible in several livestock species, as described above, and there is anecdotal evidence that the efficiency of producing viable embryos free of disorders can be considerably increased with experience. In conclusion, it would be timely to review the potential of this technique and to integrate it more firmly into conservation guidelines. It may be that SCNT can only be recommended as a desired option for a few livestock species in special circumstances; however it may be worth considering the cryopreservation of somatic cells even for poultry on the assumption that advances in technology may eventually make nuclear transfer viable in avian species. Groeneveld (2005) proposed to create national genebanks on the basis of somatic cells.

List of references

- Boettcher, P.J., Stella, A., Pizzi, F. & Gandini, G.** 2005. The combined use of embryos and semen for cryogenic conservation of mammalian livestock genetic resources. *Genetics Selection Evolution*, 37: 657–675.
- Bryson, B.** 2005. A short history of nearly everything. New York, USA, Broadway Books.
- Cundiff, L.V., Mac Neil, M.D., Gregory, K.E. & Koch, R.M.** 1986. Between and within-breed genetic analysis of calving traits, and survival to weaning in beef cattle. *Journal of Animal Science*, 63: 27–33
- Dorward, A., Kydd, J., Morrison, J. & Poulton, C.** 2005. Institutions, markets and economic co-ordination: Linking development policy to theory and praxis. *Development and Change*, 36: 1–25.
- Eding, H. & Bennewitz, J.** 2007. Measuring genetic diversity in farm animals. In K. Oldenbroek, (Ed.). *Utilisation and conservation of farm animal genetic resources*. Wageningen, the Netherlands, Wageningen Academic Publishers, pp. 232.
- ERFP.** 2004. Guidelines for the constitution of national cryopreservation programmes for farm animals, edited by S-J Hiemstra. Publication No. 1 of the European Regional Focal Point on Animal Genetic Resources. (also available at www.rfp-europe.org/files/file00244.pdf).
- FAO.** 1998a. Secondary Guidelines for Development of National Farm Animal Genetic Resources Management Plans. Management of Small Populations at Risk. Rome.
- FAO.** 1998b. New developments in biotechnology and their implications for the conservation of farm animal genetic resources: reversible DNA quiescence and somatic cell cloning. Report on a joint workshop of FAO and Istituto Sperimentale per la Zootechnia Monterotondo. Rome.
- FAO.** 2007a. The State of the World's Animal Genetic Resources for Food and Agriculture, edited by B. Rischkowsky & D. Pilling. Rome, (also available at <http://www.fao.org/docrep/010/a1250e/a1250e00.htm>), pp. 511.
- FAO.** 2007b. Global Plan of Action for Animal Genetic Resources. Rome.
- Gandini, G. & Oldenbroek, K.** 2007. Strategies for moving from conservation to utilization. In K. Oldenbroek, ed. *Utilisation and conservation of farm animal genetic resources*, pp. 29–54. Wageningen, the Netherlands, Wageningen Academic Publishers, pp. 232.
- Gandini, G.C., Ollivier, L., Danell, B., Distl, O., Georgoudis, A., Groeneveld, E., Martyniuk, E., van Arendonk, J.A.M. & Woolliams, J.A.** 2004. Criteria to assess the degree of endangerment of livestock breeds in Europe. *Livestock Production Science*, 91: 173–182.
- Gibson, J., Gamage, S., Hanotte, O., Iñiguez, L., Maillard, J.C., Rischkowsky, B., Semambo, D. & Toll, J.** 2006. Options and strategies for the conservation of farm animal genetic resources. Report of an International Workshop (7–10 November 2005, Montpellier, France). Rome, CGIAR System-wide Genetic Resources Programme (SGRP)/Bioversity International, pp. 53.
- Groeneveld, E.** 2005. A world wide emergency programme for the creation of national genebanks of endangered breeds in animal agriculture. *Animal Genetic Resources Information Bulletin* 36: 1–6.
- IPCC.** 2007: Summary for policymakers. In: S. Solomon, D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor & H.L. Miller, (Eds). *Climate change 2007: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK and New York, USA, Cambridge University Press.
- Meuwissen, T.H.E.** 2007. Operation of conservation schemes. In K. Oldenbroek, ed. *Utilisation and conservation of farm animal genetic resources*, pp. 167–193. Wageningen, the Netherlands, Wageningen Academic Publishers, pp. 232.
- Reed, D.H. & Frankham, R.** 2001. How closely correlated are molecular and quantitative measures of genetic variation? A meta-analysis. *Evolution*, 55: 1095–1103.
- Roughsedge, T., Villanueva, B. & Woolliams, J.A.** 2006. Determining the relationship between restorative potential and size of a gene bank to alleviate the risks inherent in a scrapie eradication breeding programme. *Livestock Science*, 100: 231–241.

.....

Thornton, P.K., Jones, P.G., Owiyo, T., Kruska, R.L., Herrero, M., Kristjanson, P., Notenbaert, A., Bekele, N. & Omolo, A., with contributions from Orindi, V., Ochieng, A., Otiende, B., Bhadwal, S., Anantram, K., Nair, S., Kumar, V. & Kelkar, U. 2006. Mapping climate vulnerability and poverty in Africa. Report to the Department for International Development. Nairobi, International Livestock Research Institute (ILRI). 200 pp. (also available at www.dfid.gov.uk/research/mapping-climate.pdf).

Wilmot, I., Schnieke, A.E., McWhir, J., Kind, A.J. & Campbell, K.H.S. 1997. Viable offspring derived from fetal and adult mammalian cells. *Nature*, 385: 810–813.

Woolliams, J.A. & Wilmot, I. 1999. New advances in cloning and their potential impact on genetic variation in livestock. *Animal Science*, 68: 245–256.

Woolliams, J.A. 2007. Genetic contributions and inbreeding. In K. Oldenbroek, ed. *Utilisation and conservation of farm animal genetic resources*, pp. 147–165. Wageningen, the Netherlands, Wageningen Academic Publishers, pp. 232.

Panellists' comments and discussion

Mr Arthur Mariante, EMBRAPA Genetic Resources and Biotechnology, Brazil

A very interesting paper, which brings back the old question: *in situ* or *ex situ*, with a different approach! John and his co-authors brought some new ingredients to this subject. Some of their core messages are:

- Diversity “put in” limits diversity “taken out”!
- How many and which ones to include?
- Gene Banks pay no interest! You get out only what you put in!
- What the use of the stored material can or cannot achieve?
- Use of SCNT for emergency conservation actions.

I would like to demonstrate some aspects of animal genetic resources in Brazil.

Most livestock are not indigenous to Brazil; animals were brought in by the settlers, have been

submitted to natural selection, and supported animal production in the country for centuries. At the beginning of the twentieth century, exotic breeds were imported and gradually replaced these adapted breeds.

To avoid the loss of this genetic material, in 1983 Embrapa decided to include conservation of animal genetic resources among its priorities. At that time, we decided to conserve those old breeds both ways, as suggested by John: *in situ* and *ex situ*. We agree with the authors that there is no dichotomy, and these two methods complement each other.

In situ conservation is carried out in nucleus herds (conservation nuclei), maintained in the habitats where the breeds have been naturally selected.

When there are human and physical resources in the nucleus, the collection and freezing of genetic material are carried out *on farm*. When it is not




Figure 1. Ex situ conservation - Semen and embryos are stored at the Animal Germplasm Bank located at our Experimental Farm located in Brasilia. 65 000 semen samples and 250 embryos are being stored at the Animal Gene Bank (AGB), located at the Experimental Farm. More than 8 000 DNA samples are being stored at the DNA Bank.

possible, some animals are temporarily transferred to the Experimental Farm for this collection.

We agree with the conclusions by the authors, when they list six major challenges that livestock production is facing nowadays. The mentioned unpredictability of these challenges should really be met by securing the animal genetic resources that are available to mankind.

This task should be shared by countries that have the facilities and human resources to do so, building capacities in regions where this is not yet being done.

The time may have arrived to establish regional gene banks, a huge project postponed by FAO in the early 1990s, due to different animal health legislation of countries within the same regions. The establishment of subregional gene banks could be the way to proceed in order to save endangered breeds of countries that are not yet prepared to do so. We are all responsible!

Ms Nitya S. Ghotge, ANTHRA, India

While on one hand the paper states that the position of global animal genetic resources is far from secure, it does not adequately address the relative merits and demerits of different approaches and technologies with reference to different nation states, which then brings one to the very crucial question of who will conserve the genetic material, where and how. The paper also does not touch on the very important aspect of the ownership of genes and genetic material.

Currently, the genetic diversity of domesticated livestock lies in the Southern, lesser-developed countries, often with farmers living in fragile and marginal livelihoods. Efforts to preserve this diversity must go in tandem with efforts to improve the livelihoods of these farmers, and this is where funds need to be channelled. The ownership of the genetic material must remain with the communities and not in the private hands of industries or institutes.

Our organization ANTHRA which is based in India works with small and marginal farmers – dalits, adivasis (indigenous communities), pastoralists and landless groups – especially with women from these marginalized communities. Our work focuses on production and farming systems, and within them the crops and fodder varieties, livestock and plant genetic resources, medicinal plants and health care traditions, land and water use, and the indigenous knowledge connected with these systems.

We support viable community-based livelihood-enriching interventions which use and strengthen peoples' knowledge systems in productive ways and make them less dependent on external forces. Towards this end we have been active in supporting local livestock production systems such as women and backyard poultry with a special focus on the Aseel, Nicobari and Kadaknath breeds; local goats – the Kanchu Meka breed; local cattle – the Dangi; and local pigs – the Nicobari for different adivasi (indigenous) communities, and the Deccani sheep for pastoral communities.

Mr Shakeel Bhatti, FAO International Treaty on Plant Genetic Resources for Food and Agriculture, Italy

Thank you, Mr Chairman.

As I am on this Panel, the only commentator from the plant genetic resource side and the only representative of an intergovernmental body, I would like to add some comments on the inter-relation between the important work that lies ahead for your Conference and the already existing work and intergovernmental processes in the field of plant genetic resources (PGR) – in particular, of course, the International Treaty on Plant Genetic Resources for Food and Agriculture (ITPGRFA).

Having heard the presentations by the authors, my basic observation is that plant and animal genetic resources are very distinct and cannot be forced into the same mould in legal and policy terms. And my basic argument would be that, in their distinctiveness, PGR and animal genetic resources (AnGR) policy *can* and *should* be complementary, mutually supportive and conceived and developed in a coordinated manner.

Conservation, which is addressed under Article 5 of the Treaty, is one of the basic objectives of the ITPGRFA. However, the ITPGRFA comes from the plant side and so what I am about to say has mostly elliptical value as a contribution to this debate.

Introduction to ITPGRFA

As the two areas are so different, let me, for those of you who are not familiar with the Treaty, recall some of the main features of the ITPGRFA. Historically, the work of the Global Plan of Action for Plant Genetic Resources was purely food crop-based. It was during the negotiations for the ITPGRFA that forages were brought in. The Treaty

establishes a multilateral system (MLS) for a fixed list of 64 crops and forages, established on the basis of the criteria of food security and interdependence. For these crops and forages the Treaty facilitates access and regulates benefit-sharing.

Farmers' Rights

As the paper mentions, there is a preference for *in situ* conservation recognized by the Commission on Genetic Resources for Food and Agriculture (CGRFA), the prime body for policy in agricultural genetic resources. As we heard this morning, that is where the rights of pastoralists and traditional livestock breeders, who are conserving AnGR diversity *in situ*, come in. You might be interested to know that in the context of the negotiations for the ITPGRFA, we had the same discussion on recognizing and incentivizing the enormous contribution of farmers to the *in situ* conservation of PGR. This discussion led to the adoption of Article 9 of the International Treaty, entitled "Farmers' Rights". So there has been a similar debate on the recognition of traditional communities in the conservation of PGR and the work on implementation of Farmers' Rights is still going on. There may be lessons to be learned there.

Linkages between PGR and AnGR

There may be a case to be made for working with ecosystemic approaches that integrate perspectives on PGR and AnGR to make overall production systems more effective. The linkages are, indeed, there in the production systems – AnGR production systems use crops and forages to produce. The coordination between PGR and AnGR policy may play a particularly important role in facilitating sustainable intensification in crop-based livestock production systems and for conservation in pastoralist production systems.

AnGR and PGR are very different: different biology, different production systems, different use and innovation patterns, etc. Thus, while recognizing that they are inter-related, the differences must be recognized. This is well reflected in the current policy and institutional framework of FAO, where – while they are both included in the Multi-year Programme of Work (MYPOW) – the process for plants is very different, being mostly contained in the framework of the ITPGRFA.

Lessons that can be learned

In light of rapid change and genetic erosion, there may be need for international regulation and cross-border controls to improve cooperation and development in the AnGR field. If you decide to go that way in this Conference, there are lessons which, I think, might be learned from the ITPGRFA and its negotiations. These lessons include the importance of multilateralism in designing appropriate policy and legal frameworks for genetic resources for food and agriculture. This is so important in agricultural genetic resources because of the millennia of open exchange of genetic resources in agriculture, both in both plant and animal kingdoms, which makes a bilateral approach very difficult to implement.

Another important consideration is the need for a Funding Strategy. The paper recommends that "an initial step" of a "conservation strategy to capture the diversity of breeds" could be for funding institutions to "require project proposals to identify conservation needs and to supply costed and timebound plans for such needs". I am pleased to inform you that an *Ad Hoc* Advisory Committee on the Funding Strategy of the Treaty has just identified some key priorities and eligibility for funding under the Funding Strategy of the Treaty. It has identified "on-farm conservation of PGRFA" in particular those listed in Annex I of the Treaty as one of the key priorities for funding of development projects. The Funding Strategy of the Treaty foresees all sorts of actors working together, including through other institutions.

Some concrete suggestions:

- The process following up from this Conference and the monitoring of the possible Global Plan of Action for Animal Genetic Resources can draw upon the Treaty process for support along the lines of the linkages outlined above. This would mean:
- Coordinating the processes of the Global Plan of Action for Animal Genetic Resources and the process of the ITPGRFA as far as their respective work on forages and pastures go.
- In a possible future revision of Annex I of the Treaty – which is done according to criteria of food security and interdependence – the needs of livestock production systems and their contribution to food security should be taken into account. This should take into account the importance of grasses and forages for livestock production systems and thereby for food security. This should apply especially to low- and medium-input livestock production systems.

- One target for the priorities under the Funding Strategy could be fodders and feeds – grasses (Africa) and legumes (South America).

Summary of plenary discussion

The meeting was then opened for general discussion and interventions from the floor. Key issues raised during this discussion included:

- The need to identify forces that drive breeds to extinction.
 - The need for guidelines to ensure that inappropriate restocking measures are avoided in the aftermath of catastrophes.
 - The need to consider policy and legal frameworks for conservation programmes.
 - The need to identify priorities for immediate action in the field of conservation.
- Responses and final comments of the authors included the following points:
- In general, action is most urgently required where the livestock sector is undergoing rapid changes.
 - *In situ* and *ex situ* conservation measures are complementary, but need to be coordinated to ensure that they achieve their objectives effectively.
 - Cooperation with conservation organizations interested in specific animal genetic resources is required.

Global Plan of Action for Animal Genetic Resources and the Interlaken Declaration

Commission on Genetic Resources for Food and Agriculture
FAO, Viale delle Terme di Caracalla, 00153 Rome, Italy


Published in 2008, pp. 46
ISBN 978-92-5-105848-0

The *Global Plan of Action for Animal Genetic Resources* (GPA) is the internationally agreed framework for the wise management of animal genetic resources for food and agriculture (AnGR), the culmination of an extended process involving the participation of 169 countries. It was adopted by 109 country delegations at the International Technical Conference on Animal Genetic Resources, held in Interlaken, Switzerland, from 3 to 7 September 2007. The countries also adopted the *Interlaken Declaration on Animal Genetic Resources*, by which they confirmed their common and individual responsibilities for the conservation, sustainable use and development of AnGR; for world food security; for improving human nutritional status; and for rural development. They committed themselves to facilitating access to these AnGR, and ensuring the fair and equitable sharing of the benefits from their use. The GPA is covered in three parts, I. The Rationale; II. The Strategic Priorities for Action; and III. Implementation and Financing.

Based on *The State of the World's Animal Genetic Resources* reviewed in AGRI 41, the first ever report of its kind, Part I of GPA gives reasons for proposing specific measures to reverse the ongoing trends of erosion and underutilization of AnGR through promoting the sustainable use and development of AnGR, conserving the important animal genetic resource diversity, promoting a fair and equitable sharing of the benefits arising from the use of AnGR, meeting the needs of pastoralists and farmers to have non-discriminatory access to genetic material, information, technologies, financial resources, research results, marketing systems, and natural resources; to assisting countries and institutions responsible for the management of AnGR, plus other means to enhance collaboration among countries.

Part II deals with the four strategic areas of:

1. Characterization, inventory and monitoring of trends and associated risks.
2. Sustainable use and development.
3. Conservation.
4. Policies, institutions and capacity-building, giving rationale for each strategy area and indicating actions required.



Part III discusses the important issues of funding the implementation of GPA, monitoring and reporting its progress and roles played in this regard at the national, regional and international levels. The GPA will set the agenda for AnGR management at national and international level for the next decade, therefore this publication is a required reading for planners and decision makers in the area of AnGR. It is published in five languages (Arabic, Chinese, English, French, Spanish) and being also translated to Russian. An electronic version is available at www.fao.org/ag/againfo/programmes/en/genetics/documents/Interlaken/GPA_en.pdf.

Animal electronic identification: experiences of Ministry for food, fishing and nutrition (in Spanish)

Identificación electrónica animal: Experiencias del Ministerio de Agricultura, Pesca y Alimentación
Ministerio de Agricultura, Pesca y Alimentación
MAPA, Alfonso XIII, 62 - 3º Pta. 28014 Madrid
Published in 2007, pp. 256
ISBN: 978-84-491-0791-7

In the last years, the cattle sector is experiencing outstanding changes in its management of production, originating, among other factors, by the technological and the scientific advances that are increasing the available technologies and that contribute to its development.

In this respect, one of the most considerable progresses has been the development of the radio frequency technology aimed to the identification of the animals. This techniques represents one of the priorities of the public authorities, since for its intrinsic safety and reliability, the electronic animal identification turns it into a key instrument to improve food safety.

Since the "Project IDEA" (1998-2001), Spain participated in the projects of the European Commission for the development of such technique. From the several conclusions originated form that project, many initiative were developed by the Spanish Breeding General Directorate (*Dirección General de Ganadería*)

The results of such initiatives are now collected in a book, published by the Ministry of Agriculture, that recalls this technology directly applied to the field. Initially, a short chapter describes the earlier techniques used in the sector followed by a clear description of the principles and utilities for a unique animal identification. The book shows, besides the bases of electronic animal identification, an exceptional summary of all these field experiences realized in Spain by the MAP, as well as the results of the parameters of retention and reading of each one of the tested devices. Clear photos and diagrams facilitate the reading and allow an easy understanding of even of the most complex points



Report of the FAO/WAAP Expert Meeting on Sustainable Utilization of Animal Genetic Resources


Edited by FAO/WAAP
FAO, Rome

Published in 2008, pp. 157

This is the report of the Expert Meeting that was jointly organized by the Food and Agriculture Organization (FAO) and World Association for Animal Production (WAAP) and held in Ferentillo, Italy, 2–4 July 2006. The two terms 'utilization' and 'sustainable' are keywords frequently used when discussing the development of animal genetic resources (AnGR). Although both have been formally defined by FAO, the CBD and others, clear understanding of the concepts underlying them, especially as regard AnGR development remains a challenge.

The main objectives of the Expert Meeting were to define the sustainable utilization of AnGR; identify guiding principles; and develop an outline for technical guidelines for sustainable utilization of AnGR. The report discusses the three guiding dimensions of the sustainable utilization of AnGR, i.e. economic, environmental and social.

The main body of the report occupies 27 pages covering five sections. Section 1 describes the background and objectives of the Expert Meeting. Section 2 sets out issues that were identified as important elements of the context for sustainable utilization, i.e. values of livestock genetic diversity, the changing animal production systems, and the current focus of livestock development efforts. Section 3 outlines some of the perspectives on sustainable utilization that emerged during the meeting and framed the discussion. Section 4 discusses guiding principles that need be considered if sustainable utilization of AnGR is to be achieved. Section 5 presents some more specific recommendations addressing the areas: information requirements for sustainable utilization; the development of AnGR and the production systems in which they are kept; market development; and the establishment of a favorable policy environment for sustainable utilization. Finally, some conclusions are offered, focusing particularly on what further work is required to clarify and develop the concept of sustainable utilization of AnGR.



The report annex includes 13 papers dealing with topics related to AnGR, ranging from touching base with FAO's work on the management of AnGR and production systems as they relate to AnGR to case studies and animal health and welfare issues. The report makes good reading for academicians and those concerned with the development of AnGR. It can be accessed at <http://dad.fao.org/cgi-bin/getblob.cgi?sid=-1,297>

EU beef farming systems and CAP regulations

P. Sarzeaud, A. Dimitriadou & M. Zjalic (Eds)

EAAP Technical Series No. 9

Wageningen Academic Publishers

Published in 2007, pp. 122

ISBN: 978-90-8686-058-6

The new Common Agricultural Policy, launched in 2003 and implemented since 2005, appears as a big change mainly due to the introduction of the Single Payment Scheme and the decoupling of subsidies from production. EU beef production is one of the sectors mostly affected by the existing CAP regulations. This book addresses what the impact could be on the future of European beef farming systems and in what ways could beef farmers adapt in order to ensure the sustainability of their farms.

Within the Cattle Network Working Group of EAAP, a group of European economists, The Beef Task Force, composed of experts and researchers from a number of EU countries, engaged in a common project with the object of monitoring and developing studies on policy impact and farm strategy analysis.

The BTF joined national studies into a common report, which is divided into four parts: the current picture of beef farming systems in Europe; CAP implementation at national level; first impact of reformed CAP on farming systems; future evolution of European beef production and land use.

This book highlights disparities in CAP implementation and presents the main developments regarding EU beef farming systems: size increase, farm/regional specialization, adaptation to market requirements and, eventually, social expectations. It is of interest to policy makers and all those who are involved in the cattle industry.



Management Guidelines Technical Annex
A. Iglesias, M. Moneo & A. Lopez Francos (Eds)
Options Méditerranéennes, Series B: Etudes et Recherches No. 58
CIHEAM Zaragoza / EC MEDAWater (Euro Mediterranean Regional
Programme for Local Water Management of the European Commission)
Published in 2007, pp. 496
ISSN: 1016-1228 ISBN: 2-85352-359-4

Natural rainfall variability is a recurrent characteristic of Mediterranean climate and intensive drought events have been more frequent in the last four decades with significant damages derived from water scarcity and low water quality at the local and national levels. In this context, Mediterranean countries have adapted to their climate and have extensive legislation, institutional capabilities, and technical resources to face drought. Nevertheless in most cases a proactive drought management based on anticipatory planning is not implemented. There is a need to formulate plans for drought management that shift from a reactive approach (crisis approach) to a proactive approach based on risk analysis, adapted to the current natural and social resources and taking into account the dynamics of the social and environmental pressures.

MEDROPLAN (www.iamz.cilieam.org/medroplan) is a Project funded by the European Commission within the framework of the EuroMediterranean Regional Programme for Local Water Management. The objectives of the MEDROPLAN Project are to provide Guidelines for Drought Preparedness Plans and the framework for the setting up of a Drought Preparedness Network for the Mediterranean countries.

The Technical Annex of the MEDROPLAN Drought Management Guidelines, that constitute the present document, compile the technical and scientific results of the MEDROPLAN Project. They are addressed to a specialized public, scientists and technicians interested in drought issues and in charge of dealing with drought problems in the water management and agricultural sectors. In the present publication, the reader can find in depth

information on the different components which constitute the drought planning methodology proposed by MEDROPLAN (the planning framework, the organizational, methodological, operational and public review components), and a compendium of examples of application in the six Partner Countries of the Project.



**Livestock biodiversity - Genetic resources
for the farming of the future**

S. Hall (Ed.)

Blackwell Science

Published in 2004, pp. 369


ISBN: 0-632-05499-9

Livestock biodiversity is integral to our culture, history, environment, economy and, most importantly, our future. Thousands of livestock breeds, from relatively small genetic pools, have evolved over time to suit particular environments and farming systems. This is both the result of natural processes and of human needs for specialized livestock - as our knowledge of genetics continues to increase we achieve a greater understanding of how this biodiversity evolved.

This book offers a detailed and comprehensive overview of livestock biodiversity. It explores the history behind it, shows how livestock biodiversity can be utilized as animal genetic resources through breed development and by crossbreeding, examines the state of world livestock biodiversity today, and emphasizes the importance of conserving and developing the biodiversity of livestock.

Special consideration is given to:

- How livestock biodiversity arose and is maintained in relation to human society.
- How it can be used in sustainable agricultural development.
- How it can be conserved for the benefit of present and future generations.
- Why conservation biologists, as well as agriculturists, should be involved in its protection.



Editorial policies and procedures

The mission of the Animal Genetic Resources Information Bulletin (AGRI) is the promotion of information on the better use of animal genetic resources of interest to food and agriculture production, under the Global Strategy for the Management of Farm Animal Genetic Resources. All aspects of the characterization, conservation and utilization of these resources are included, in accordance with the Convention on Biological Diversity.

AGRI will highlight information on the genetic, phenotypic and economic surveying and comparative description, use, development and maintenance of animal genetic resources; and on the development of operational strategies and procedures which enable their more cost-effective management. In doing this AGRI will give special attention to contributions dealing with breeds and procedures capable of contributing to the sustainable intensification of the world's medium to low input production environments (agro-ecosystems), which account for the substantial majority of the land area involved in livestock production; the total production of food and agriculture from livestock; and of our remaining farm animal genetic resources.

Views expressed in the paper published in AGRI represent the opinions of the author(s) and do not necessarily reflect those of the institutions which the authors are affiliated, FAO or the Editors.

The suitability of manuscripts for publication in AGRI is judged by the Editors and reviewers.

Electronic publication

AGRI is available in full electronically on the Internet, in addition to being published in hard copy, at: www.fao.org/dad-is

Types of articles

The following types of articles are published in AGRI.

Research articles

Findings of work on characterization, conservation and utilization of farm animal genetic resources (AnGR) in well described production environments, will be considered for publication in AGRI. Quality photographs of these genetic resources viewed in the primary production environment to which they are adapted, accompanying the manuscripts are encouraged.

Review articles

Unsolicited articles reviewing agro-ecosystems, country-level, regional or global developments on one or more aspects of the management of animal genetic resources, including state-of-the-art review articles on specific fields in AnGR, will be considered for publication in AGRI.

Position papers

Solicited papers on topical issues will also be published as deemed required.

Other published material

This includes book reviews, news and notes covering relevant meetings, training courses and major national, regional and international events and conclusions and recommendations associated with the outcomes of these major events. Readers are encouraged to send such items to the editors.

Guidelines for authors

Manuscript submission

Manuscripts prepared in English, French or Spanish with an English summary and another summary in either French or Spanish, should be submitted to AGRI Editor, AGAP, FAO, Viale delle Terme di Caracalla, 00153 Rome, Italy. Additionally the manuscript must be sent as a WinWord Electronic Mail attachment to agri-bulletin@fao.org.

Photographs, coloured or black and white, and figures must be always sent by mail.

Manuscripts should be typed double-spaced and with lines numbered in the left margin. All pages, including those of references, tables etc., must be consecutively numbered. The corresponding author is notified of the receipt of a manuscript.

For manuscripts that are accepted after revision, authors are encouraged to submit a last version (3½" disc format) in Word 6.0 for Windows of their revised manuscript along with the printed copy.

Preparation of the manuscript

The first page of the manuscript must include the running head (abbreviated title), title, names of authors, institutions, full addresses including postal codes and telephone number and other communication details (fax, e-mail, etc.) of the corresponding author. The running head not exceeding 45 characters plus spaces, should appear at the top of page 1 of the manuscript entirely in capital letters. The title of the manuscript is typed in upper and lower case letters. The title should be as brief as possible not exceeding 150 characters (including spaces) with species names when applicable. Authors, institutions and addresses are in upper and lower case italics. There is one blank line between the title and the authors. Addresses are typed as footnotes to the authors after leaving one blank line. Footnotes are designated numerically. Two lines are left below the footnotes.

Headings

Headings of sections, for example Summary, Introduction, etc., are left-justified. Leave two blank lines between addresses footnotes and Summary and between the heading Summary and its text. Summary should not exceed 200 words. It should be an objective summary briefly describing the procedures and findings and not simply stating that the study was carried on such and such and results are presented, etc. Leave one line between the summary text and Keywords which is written in italics as well as the keywords themselves. All headings of sections (14 regular) and sub-sections (12 regular) are typed bold and preceded and succeeded by one blank line and their text begins with no indentation. The heading of a sub-subsection

is written in italics, and ends with a dot after which the text follows on the same line. Keywords come immediately after the summaries. They should be no more than six, with no "and" or "&".

Tables and figures

Tables and figures must be enclosed with the paper and attached at the end of the text according their citation in the document. Photos will not be returned

Tables

Tables, including footnotes, should be preceded and succeeded by 2 blank lines. Table number and caption are written, above the table, in italics (12) followed by a dot, then one blank line. For each column or line title or sub-title, only the 1st letter of the 1st word is capitalized. Tables should be numbered consecutively in Arabic numerals. Tables and captions should be left justified as is the text. Use horizontal or vertical lines only when necessary. Do not use tabs or space-bar to create a table but only the appropriate commands.

Figures

Figures including titles and legends should be preceded and succeeded by two blank lines. Figure number and title are written, below the figure, in italics (12) and end with a dot. The term figures includes photos, line drawings, maps, diagrams etc.

All the submitted diagrams, must be accompanied with the original matrix of the data used to create them. It is strongly advised to submit diagrams in Word 6.0 or Excel 5.0. Figures should be numbered consecutively in Arabic numerals.

References

Every reference cited in the text should be included in the reference list and every reference in the reference list should have been mentioned in the text at least once. References should be ordered firstly alphabetically by the first author's surname and secondly by year.

-
- Example for reference in a periodical is:
Köhler-Rollefson, I. 1992. The camel breeds of India in social and historical perspective. *Animal Genetic Resources Information* 10, 53-64.
 - When there are more than one author:
Matos, C.A.P., D.L. Thomas, D. Gianola, R.J. Tempelman & L.D. Young. 1997. Genetic analysis of discrete reproductive traits in sheep using linear and nonnlinear models: 1. Estimation of genetic parameters 75, 76-87.
 - For a book or an ad hoc publication, e.g., reports, theses, etc.:
Cockrill, W.R. (Ed.). 1994. *The Husbandry and Health of the Domestic Buffalo*. FAO, Rome, Italy, pp. 993.
 - For an article in the proceedings of a meeting:
Hammond, K. 1996. FAO's programme for the management of farm animal genetic resources. In C. Devendra (Ed.), *Proceedings of IGA/FAO Round Table on the Global Management of Small Ruminant Genetic Resources*, Beijing, May 1996, FAO, Bangkok, Thailand, 4-13.
 - Where information included in the article has been obtained or derived from a World Wide Web site, then quote in the text, e.g. "derived from FAO. 1996" and in the References quote the URL standard form:
FAO. 1996. *Domestic Animal Diversity Information System*, <http://www.fao.org/dad-is/>, FAO, Rome, Italy.

For all future manuscript dispatch and correspondence regarding
AGRI, please use the following mailbox:

agri-bulletin@fao.org

Thanks for the collaboration

Normes et règles éditoriales

L'objectif du Bulletin d'information sur les ressources génétiques animales (AGRI) est la vulgarisation de l'information disponible sur la meilleure gestion des ressources génétiques animales d'intérêt pour la production alimentaire et agricole, d'après les recommandations de la Stratégie mondiale pour la gestion des ressources génétiques des animaux domestiques. Tous les aspects relatifs à la caractérisation, la conservation et l'utilisation de ces ressources seront pris en considération, suivant les normes de la Convention pour la Biodiversité.

AGRI désire diffuser de l'information sur la génétique, les enquêtes phénotypiques et économiques et les descriptions comparatives, l'utilisation et la conservation des ressources génétiques animales, ainsi que toute information sur le développement de stratégies opérationnelles et de normes qui puissent permettre une meilleure gestion de la relation coût/efficacité. C'est pour cela que AGRI prendra spécialement en considération toutes les contributions référées aux races et aux normes capables de permettre une intensification durable des milieux (agroécosystèmes) à revenus moyens et bas dans le monde; qui comprennent la majeure partie des terres consacrées à l'élevage, à la production totale des aliments et l'agriculture provenant de l'élevage; et tout ce qui reste comme ressources génétiques des animaux domestiques.

Les opinions exprimées dans les articles publiés dans AGRI appartiennent seulement aux auteurs et donc ne représentent pas nécessairement l'opinion des instituts pour lesquels ils travaillent, la FAO ou les éditeurs.

L'opportunité ou non de publier un article dans AGRI sera jugée par les éditeurs et les réviseurs.

Publication électronique

En plus de sa version imprimée, la version totale de AGRI se trouve disponible sur Internet, sur le site:

<http://www.fao.org/dad-is/>

Types d'articles

Les articles suivants pourront être publiés sur AGRI.

Articles de recherche

Seront prises en considération pour leur publication sur AGRI les études sur la caractérisation, la conservation et l'utilisation des ressources génétiques des animaux domestiques (AnGR) accompagnées d'une bonne description du milieu. On encourage les auteurs à envoyer des photographies de bonne qualité qui montrent les races en question dans leur milieu naturel de production.

Révisions

Occasionnellement, des articles contenant une révision des agroécosystèmes, au niveau national, régional ou mondial, avec un ou plusieurs aspects se rapportant à la gestion des ressources génétiques animales, y compris les mises à jour des différentes zones de AnGR, seront pris en considération.

Articles spécifiques

Ponctuellement, des articles sur des thèmes spécifiques pourront être demandés pour la publication d'éditions spéciales.

Autre matériel pour publication

Ceci comprend la révision de livres, nouvelles et notes de réunions importantes, cours de formation et principaux événements nationaux, régionaux et internationaux; ainsi que les conclusions et recommandation par rapport aux objectifs des ces principaux événements. Les auteurs sont priés d'envoyer ce genre de matériel aux éditeurs.

Guide pour les auteurs

Présentation du manuscrit

Les articles se présenteront en anglais, français ou espagnol, avec un résumé en anglais et sa traduction en français ou en espagnol; ils seront envoyés à l'éditeur de AGRI, AGAP, FAO, Viale

delle Terme di Caracalla, 00153 Rome, Italie. En outre, l'article devra être envoyé par courrier électronique comme document attaché en version WinWord à *agri-bulletin@fao.org*. Les photographies, en couleur ou en blanc et noir, seront toujours envoyées par courrier normal.

Les manuscrits se présenteront à double interligne et avec le numéro correspondant à chaque ligne sur la marge gauche. Toutes les pages seront numérotées, y compris celles avec les références bibliographiques, les tableaux, etc. L'auteur recevra une lettre lui donnant bonne réception de son document.

Lorsqu'un article, après sa révision, sera accepté, on demandera à l'auteur d'envoyer la version finale révisée sur disquette (format 31/2") en Word 6.0 x Windows, ainsi qu'une copie sur papier.

Préparation du manuscrit

Sur la première page du manuscrit on indiquera le titre de l'article en abrégé, le titre et noms des auteurs, des institutions, les adresses complètes (y compris code postal et numéro de téléphone); ainsi que tout autre moyen de contact tel que télécopie, courriel, etc. avec l'auteur principal. Le titre abrégé ne devra pas dépasser 45 caractères, plus les espaces nécessaires, et s'écrira sur la partie supérieure de la page 1 du manuscrit en majuscules. Le titre en entier du manuscrit sera écrit en majuscules et minuscules; il devra être aussi bref que possible, sans dépasser 150 caractères (y compris les espaces nécessaires), et avec l'indication des noms des espèces. Les noms des auteurs, des institutions et les adresses seront en italique et en lettres majuscules et minuscules. On laissera un espace en blanc entre le titre et les noms des auteurs. Les adresses seront indiquées comme de bas à pied de page pour chacun des auteurs après avoir laissé un espace en blanc après les noms. Chaque note de bas de page sera numérotée. On laissera deux espaces en blanc après les adresses.

Titres

Les titres de chaque chapitre, par exemple Résumé, Introduction, etc. seront alignés à gauche. Laisser deux espaces en blanc entre les notes de bas de page avec les adresses et le Résumé, et entre le titre Résumé et le texte qui suit. Le résumé ne devra pas dépasser les 200 mots. Il s'agira d'un résumé objectif faisant une brève description des processus

utilisés et des résultats obtenus, et non pas une simple présentation du travail réalisé avec une description générale des résultats. Laisser un espace en blanc entre la fin du texte du résumé et les mots clés, qui seront écrits en italique ainsi que le titre Mots clés. Les mots clés seront au maximum six et il ne devra pas y avoir de et ou &. Tous les titres principaux de chapitre (14 regular) et sous-chapitre (12 regular) seront en gras avec un espace en blanc avant et après. Le texte commencera sans retrait. Un titre à l'intérieur d'un sous-chapitre s'écrira en italique, suivi d'un point, avec le texte à continuation.

Tableaux et figures

Les tableaux et les figures iront à la fin du texte en suivant l'ordre d'apparition dans le texte. Les photographies ne seront pas dévolues aux auteurs.

Tableaux

Les tableaux, y compris les notes de bas de page, devront avoir un espace en blanc avant et après. Le numéro du tableau et le titre s'écriront sur la partie supérieure en italique (12) avec un point à la fin et un espace en blanc en dessous. Sur chaque colonne, titre d'en-tête ou sous-titre, seulement la première lettre du premier mot sera en majuscule. Les tableaux et leur titre seront alignés à gauche, ainsi que le texte. Les lignes verticales et horizontales seront utilisées seulement si nécessaire. Ne pas utiliser les "tabs" ou la barre d'espacement pour créer un tableau.

Figures

Les figures, y compris les titres et les légendes, seront précédés et suivis de deux espaces en blanc. Le numéro de la figure et le titre s'écriront sur la partie supérieure en italique (12) avec un point à la fin. Sous la rubrique figure on trouvera les photographies, les graphiques, les cartes, les diagrammes, etc. Dans le cas des diagrammes, la matrice originale avec les données utilisées pour son élaboration devra être envoyée. On recommande l'utilisation de Word 6.0 ou Excel 5.0 pour la présentation des diagrammes.

Références

Toute référence présente dans le texte devra apparaître sur la liste des références, et chaque référence de la liste aura été citée au moins une fois dans le texte. Les références iront en ordre alphabétique du nom de l'auteur, suivi de l'année.

- Exemple dans le cas d'une référence sur une revue:
Köhler-Rollefson, I. 1992. The camel breeds of India in social and historical perspective. *Animal Genetic Resources Information* 10, 53-64.
- Lorsqu'il s'agit de plus d'un auteur:
Matos, C.A.P., D.L. Thomas, D. Gianola, R.J. Tempelman & L.D. Young. 1997. Genetic analysis of discrete reproductive traits in sheep using linear and nonlinear models: 1. *Estimation of genetic parameters* 75, 76-87.
- Dans le cas d'un livre ou d'une publication ad hoc, par exemple un rapport, une thèse, etc.:
Cockrill, W.R. (Ed.). 1994. *The Husbandry and Health of the Domestic Buffalo*. FAO, Rome, Italy, pp. 993.
- S'il s'agit d'un acte d'une réunion:
Hammond, K. 1996. FAO's programme for the management of farm animal genetic resources. In C. Devendra (Ed.), *Proceedings of IGA/FAO Round Table on the Global Management of Small Ruminant Genetic Resources*, Beijing, May 1996, FAO, Bangkok, Thailand, 4-13.
- Lorsque l'information contenue dans l'article ait été obtenue ou dérive d'un site World Wide Web, il faudra mettre le texte entre guillemets; par exemple "tiré de la FAO. 1996" et indiquer dans les Références la forme standard URL:
FAO. 1996. *Domestic Animal Diversity Information System*, <http://www.fao.org/dad-is/>, FAO, Rome, Italy.

Pour tout envoi de manuscrits ou correspondance au sujet d'AGRI, vous êtes prié d'utiliser l'adresse suivante:

agri-bulletin@fao.org

Merci pour votre collaboration

Reglas y normas editoriales

El objetivo del Boletín de Información sobre Recursos Genéticos Animales (AGRI) es la divulgación de la información sobre una mejor gestión de los recursos genéticos animales de interés para la producción alimentaria y agrícola, siguiendo la Estrategia Mundial para la Gestión de los Recursos Genéticos de los Animales Domésticos. Todos los aspectos referidos a la caracterización, la conservación y el uso de estos recursos serán tomados en consideración, de acuerdo con el Convenio sobre la diversidad biológica.

AGRI publicará información sobre genética, encuestas fenotípicas y económicas y descripciones comparativas, uso, desarrollo y conservación de los recursos genéticos animales, así como sobre el desarrollo de estrategias operacionales y normas que permitan una gestión más eficaz de la relación costo/eficacia. Por ello, AGRI prestará especial atención a las contribuciones referidas a razas y normas capaces de contribuir a la intensificación sostenible de los medios (agroecosistemas) con ingresos medios y bajos en el mundo, que comprenden casi la mayor parte de las tierras dedicadas a la producción ganadera; la producción total de alimentos y agricultura provenientes de la ganadería; y el resto de los recursos genéticos de animales domésticos.

Los puntos de vista expresados en los artículos publicados en AGRI son solamente las opiniones de los autores y, por tanto, no reflejan necesariamente la opinión de las instituciones para las cuales trabajan dichos autores, de la FAO o de los editores.

La oportunidad o no de publicar un artículo en AGRI será juzgada por los editores y revisores.

Publicación electrónica

Además de su publicación impresa, la versión íntegra de AGRI se encuentra disponible electrónicamente en Internet, en el sitio:

www.fao.org/dad-is/

Tipos de artículos

Serán publicados en AGRI los siguientes tipos de artículos:

Artículos sobre investigación

Se tomarán en consideración para su publicación en AGRI los estudios sobre la caracterización, conservación y uso de los recursos genéticos de los animales domésticos (AnGR) con una buena descripción del entorno. Se agradecerá el envío de fotografías de calidad que presenten a las razas en cuestión en su ambiente natural de producción.

Artículos de revisión

Se podrán tomar en consideración ocasionalmente aquellos artículos que presenten una revisión de los agroecosistemas, a nivel nacional, regional o mundial, con el desarrollo de uno o más aspectos referidos a la gestión de los recursos genéticos animales, incluidas las revisiones sobre el estado actual de las distintas áreas de AnGR.

Artículos específicos

Se solicitarán puntualmente artículos sobre temas específicos para ediciones especiales.

Otro material para publicación

Incluye la revisión de libros, noticias y notas referidas a reuniones importantes, cursos de formación y principales eventos nacionales, regionales e internacionales, así como conclusiones y recomendaciones relacionadas con los objetivos de estos principales eventos. Se invita a los lectores a enviar este tipo de material a los editores.

Guía para los autores

Presentación del manuscrito

Los artículos se presentarán en inglés, francés o español, junto con un resumen en inglés y su traducción en francés o español, y se enviarán al editor de AGRI, AGAP, FAO, Viale delle Terme di Caracalla, 00153 Roma, Italia. El artículo deberá ser enviado en versión WinWord en fichero adjunto por

correo electrónico a *agri-bulletin@fao.org*. Las fotografías, color o en blanco y negro, se enviarán siempre por correo normal.

Los manuscritos se presentarán con doble espacio y con el número correspondiente a cada línea en el margen izquierdo. Todas las páginas serán numeradas, incluidas las de las referencias bibliográficas, cuadros, etc. El autor recibirá una notificación sobre la recepción de su documento.

En el caso de aceptación de un artículo después de su revisión, se solicitará al autor una versión final de su artículo revisado en disquete (formato 31/2") en Word 6.0 x Windows, así como una copia impresa del mismo.

Preparación del manuscrito

En la primera página del manuscrito se indicará el título abreviado del artículo, títulos y nombres de los autores, instituciones, direcciones completas (incluido código postal y número de teléfono); así como otros medios de contacto tales como fax, correo electrónico, etc. del autor principal. El título abreviado no deberá sobrepasar los 45 caracteres más los espacios correspondientes, y aparecerá en la parte superior de la página 1 del manuscrito en mayúsculas. El título entero del manuscrito se escribirá en mayúsculas y minúsculas. Dicho título debe ser lo más breve posible y no sobrepasar los 150 caracteres (incluidos los espacios necesarios), con los nombres de las especies, si necesario. Los nombres de los autores, instituciones y direcciones se escribirán en cursiva y en letras mayúsculas y minúsculas. Se dejará una línea en blanco entre el título y los nombres de los autores. Las direcciones se escribirán como notas de pie de página de cada autor después de dejar una línea en blanco entre los nombres y éstas. Cada nota de pie de página con la dirección será indicada numéricamente. Se dejarán dos líneas en blanco después de las direcciones.

Títulos

Los títulos de cada sección, por ejemplo Resumen, Introducción, etc., serán alineados a la izquierda. Dejar dos líneas en blanco entre las notas de pie de página con las direcciones y el Resumen y entre el título Resumen y el texto que sigue. El resumen no deberá exceder de 200 palabras. Deberá ser un resumen objetivo que describa brevemente los procesos y logros obtenidos, y no una presentación de cómo se ha llevado a cabo el estudio y una descripción genérica de los resultados. Dejar una

línea en blanco entre el final del texto del resumen y las palabras clave, que se escribirán en cursiva así como el título Palabras clave. No deberán ser más de seis y no deberán contener "y" o "&". Todos los títulos principales de capítulo (14 regular) y subcapítulo (12 regular) serán en negrita e irán precedidos y seguidos de una línea en blanco. El texto correspondiente empezará sin sangrado. Un título dentro de un subcapítulo se escribirá en cursiva e irá seguido de un punto con a continuación el texto correspondiente.

Cuadros y figuras

Los cuadros y las figuras se incluirán al final del texto siguiendo el orden de cita dentro del mismo. Las fotografías no serán devueltas a sus autores.

Cuadros

Los cuadros, incluidas las notas de pie de página, deberán ir precedidos y seguidos por dos líneas en blanco. El número del cuadro y su título se escribirán en la parte superior en cursiva (12) con un punto al final y seguido de una línea en blanco. En cada columna o título de encabezamiento o subtítulo, sólo la primera letra de la primera palabra irá en mayúscula. Los cuadros irán numerados de forma consecutiva con números árabes. Los cuadros y sus títulos se alinearán a la izquierda, así como el texto. Se utilizarán líneas horizontales o verticales sólo cuando sea necesario. No utilizar tabuladores o la barra espaciadora para crear un cuadro.

Figuras

Las figuras, incluidos los títulos y leyendas, irán precedidas y seguidas de dos líneas en blanco. El número de la figura y el título se escribirán en la parte superior en cursiva (12) con un punto al final. La palabra figura incluye las fotografías, los gráficos, los mapas, los diagramas, etc. En el caso del diagrama se enviará la matriz original con los datos utilizados para crearlo. Se recomienda encarecidamente la utilización de Word 6.0 o Excel 5.0 para la presentación de los diagramas.

Referencias

Toda referencia presente en el texto deberá aparecer en la lista de referencias y, de la misma manera, cada referencia de la lista deberá haber sido citada por lo menos una vez en el texto. Las referencias deben ir en orden alfabético del apellido del autor, seguido por el año.

- Ejemplo en el caso de una referencia de una revista:
Köhler-Rollefson, I. 1992. The camel breeds of India in social and historical perspective. *Animal Genetic Resources Information* 10, 53-64.
- Cuando se trate de más de un autor:
Matos, C.A.P., D.L. Thomas, D. Gianola, R.J. Tempelman & L.D. Young. 1997. Genetic analysis of discrete reproductive traits in sheep using linear and nonnlinear models: 1. Estimation of genetic parameters 75, 76-87.
- En el caso de un libro o de una publicación ad hoc, por ejemplo informes, tesis, etc.:
Cockrill, W.R. (Ed.). 1994. *The Husbandry and Health of the Domestic Buffalo*. FAO, Rome, Italy, pp. 993.
- Cuando se trate de un artículo dentro de las actas de una reunión:
Hammond, K. 1996. FAO's programme for the management of farm animal genetic resources. In C. Devendra (Ed.), *Proceedings of IGA/FAO Round Table on the Global Management of Small Ruminant Genetic Resources*, Beijing, May 1996, FAO, Bangkok, Thailand, 4-13.
- Cuando la información contenida en el artículo haya sido obtenida o derive de un sitio World Wide Web, poner el texto entre comillas; por ejemplo "sacado de la FAO. 1996" e indicar en las Referencias la forma estándar URL:
FAO. 1996. Domestic Animal Diversity Information System, <http://www.fao.org/dad-is/>, FAO, Rome, Italy.

Se ruega enviar los manuscritos o la correspondencia relativa a AGRI a la dirección siguiente:

agri-bulletin@fao.org

Gracias por su colaboración



ISBN 978-92-5-005952-5 ISSN 1014-2339



9 789250 059525

TC/D/I0072Tri/1/03.08/2000