

# COMMISSION DU CODEX ALIMENTARIUS



Organisation des Nations  
Unies pour l'alimentation  
et l'agriculture



Organisation  
mondiale de la Santé

F

Viale delle Terme di Caracalla, 00153 Rome, Italie - Tél: (+39) 06 57051 - Fax: (+39) 06 5705 4593 - E-mail: [codex@fao.org](mailto:codex@fao.org) - [www.codexalimentarius.net](http://www.codexalimentarius.net)

CX 4/40.2

CL 2010/55 - PR  
décembre 2010

DESTINATAIRE: Intermédiaire du Codex  
Organisations internationales intéressées

EXPEDITEUR: Secrétariat,  
Commission du Codex Alimentarius, Programme mixte FAO/OMS sur les normes alimentaires  
FAO, Viale delle Terme di Caracalla, 00153 Rome, Italie

OBJET : DEMANDE DE COMMENTAIRES SUR LES RECOMMANDATIONS DE LA RÉUNION MIXTE  
FAO/OMS SUR LES RÉSIDUS DE PESTICIDES (JMPR)<sup>1</sup> ET LES LMR POUR LES PESTICIDES À  
L'ÉTAPE 3 DE LA PROCÉDURE

DATE LIMITE : 1<sup>er</sup> mars 2011

COMMENTAIRES : À : Ms Lifang DUAN  
Division Résidus  
Institut pour le Contrôle des produits  
agrochimiques  
Ministère de l' Agriculture (ICAMA)  
No. 18, Maizidian Street, Chaoyang District  
Beijing 100125, R.P. de Chine  
Fax: +86 10 5919 4252  
E-mail: [ccpr@agri.gov.cn](mailto:ccpr@agri.gov.cn) (de préférence)

Copie à : Secrétariat  
Commission Codex Alimentarius  
Programme mixte FAO/OMS sur les normes  
alimentaires  
FAO  
Viale delle Terme di Caracalla  
00153 Rome, Italie  
Fax: +39 06 5705 4593  
E-mail: [codex@fao.org](mailto:codex@fao.org) (de préférence)

## CONTEXTE

### A. LMR A L'ÉTAPE 3 DE LA PROCÉDURE

1. La réunion mixte annuelle de la FAO/OMS sur les résidus de pesticides (JMPR) s'est tenue à Rome, Italie, du 21 au 30 septembre 2010. Les résumés suivants des résultats de la réunion annuelle de la JMPR sont fournis et rendus accessibles à une date précoce aux parties intéressées.
2. La réunion a évalué 23 pesticides, dont 8 étaient de nouveaux composés et 5 étaient des réévaluation dans le cadre du programme de révision périodique du Comité Codex sur les résidus de pesticides (CCPR). La réunion a établi des doses journalières admissibles (DJA) ainsi que des doses de référence aiguës (DrfA).
3. La réunion a estimé des niveaux maxima de résidus, qu'elle a recommandés pour être utilisés comme limites maximales de résidus (LMR) par le CCPR. Elle a aussi évalué des médianes de résidus en essais contrôlés (MREC) et les plus hauts niveaux de résidus (HR) comme base pour l'estimation des doses journalières de résidus des pesticides révisés. L'application des niveaux HR est expliquée dans le rapport de la réunion de 1999 (section 2.4). Les attributions et estimations sont reprises au tableau.
4. Les pesticides pour lesquels les doses journalières estimées peuvent, sur base des informations disponibles, dépasser la DJA sont indiquées avec des notes de bas de page en détail dans le rapport de la réunion 1999 (section 2.2.). Les notes de bas de page s'appliquent aussi aux produits spécifiques lorsque les informations disponibles indiquent que la DrfA d'un pesticide risque d'être dépassée lorsque le produit est consommé. Il faut noter que toutes ces distinctions s'appliquent uniquement aux nouveaux composés et aux composés réévalués dans le cadre du programme de révision périodique du CCPR.
5. Dans le tableau sont indiqués les numéros de référence Codex des composés et les numéros de la classification Codex (NCC) des produits, pour faciliter la référence aux LMR Codex pour les résidus de pesticides (*Codex Alimentarius*, Vol. 2B) et d'autres documents et documents de travail de la Commission du Codex Alimentarius. À la fois les composés et les produits sont listés par ordre alphabétique.
6. Outre les abréviations reprises ci-dessus, les qualifications suivantes sont utilisées dans le tableau.

<sup>1</sup> Les recommandations de la JMPR pour les limites maximales de résidus correspondent à l'étape 3 de la procédure du Codex.

* (après le nom du pesticide)	Nouveau composé
** (après le nom du pesticide)	Composé révisé dans le cadre du programme de révision périodique du CCPR
* (après la LMR recommandée)	A la limite de quantification ou à la limite approximative de quantification
HR-P	Plus haute valeur dans un produit transformé, en mg/kg en multipliant le HR dans le produit frais par le facteur de transformation
Po	La recommandations intègre le traitement après récolte du produit
PoP (après recommandation pour les aliments transformés (classes D et E de la classification Codex)	La recommandation intègre le traitement après récolte dans le produit alimentaire primaire.
MREC-P	Une MREC pour un produit transformé calculée en appliquant à la MREC calculée pour le produit frais agricole, le facteur de concentration ou de réduction pour la transformation.
W (au lieu de la LMR recommandée)	La recommandation précédente est retirée, ou retrait de la LMR recommandée ou existante du Codex ou un avant-projet de LMR est recommandé.

7. L'annexe est aussi disponible sur les sites Internet ci-dessous :

lien internet FAO : <http://www.fao.org/agriculture/crops/core-themes/theme/pests/pm/jmpr/en/>

lien internet OMS : <http://www.who.int/ipcs/food/jmpr/summaries/en/index.html>

8. En cas de problème lors du téléchargement des documents repris ci-dessus, veuillez contacter les secrétariats de la JMPR ou de la FAO aux adresses suivantes afin d'obtenir une copie par courriel :

Mme Yong Zhen YANG  
Secrétaire FAO JMPR  
Division de la production et protection des Plantes  
FAO des Nations Unies  
Viale delle Terme di Caracalla  
00153 Rome, Italie  
Tél: +39 06 57054246  
Fax: +39 06 570 53224  
Courriel: [YongZhen.Yang@fao.org](mailto:YongZhen.Yang@fao.org)

Dr Philippe Verger (Secrétariat OMS JMPR)  
Secrétaire OMS JMPR  
GEMS/Food Programme  
Département de la sécurité alimentaire et des zoonoses  
(FOS)  
Organisation mondiale de la santé  
1211 Genève 27, Suisse  
Tél : +41 22 791 3053  
Fax: +41 22 791 4807  
Courriel: [vergerp@who.int](mailto:vergerp@who.int)

#### DEMANDE DE COMMENTAIRES

9. Les gouvernements membres et organisations internationales intéressés désireux de soumettre des commentaires sur les avant-projets de LMR nouvellement proposés correspondant à l'étape 3 de la procédure Codex comme proposés par la JMPR 2010 ainsi que sur d'autres recommandations pertinentes pour le travail de la 43<sup>ème</sup> session du Comité Codex sur les Résidus de Pesticides (voir tableau ci-dessous) sont priés de le faire par écrit, conformément aux procédures pour l'élaboration des Normes Codex et textes connexes (*Manuel de procédure du Codex Alimentarius*), de préférence par courriel, aux adresses indiquées en page de couverture et ce avant le 1<sup>er</sup> mars 2011.

**B. LMR A L'ÉTAPE 6 DE LA PROCÉDURE**

10. La 33<sup>ème</sup> session de la Commission a adopté à l'étape 5 les avant-projets de LMR tels que proposés à l'Annexe IV de ALINORM 0/33/24 et les a avancés à l'étape 6, (voir ALINORM 10/33/REP, par. 69 et Annexe IV) qui ont été mis au point par la 42<sup>ème</sup> session du CCPR, tout en prenant note des réserves émises par la Communauté européenne et la Norvège sur les LMR pour fluopicolide (235) et haloxyfop (194)

11. Par ailleurs, la 42<sup>ème</sup> session du CCPR a renvoyé à l'étape 6 quelques projets de LMR tels que présentés à l'Annexe VII de ALINORM 10/33/24 pour de nouveaux commentaires et examen lors de sa prochaine session (voir ALINORM 10/33/24, par. 36, 50 et 54).

12. Ces documents ont été précédemment diffusés aux intermédiaires du Codex et sont disponibles sur les sites Internet suivants : <http://www.codexalimentarius.net> sous Rapports, réunions et événements.

13. Les gouvernements et organisations internationales intéressés désireux de soumettre des commentaires sur les projets de LMR à l'étape 6 de la procédure Codex sont priés de la faire par écrit conformément aux procédures pour l'élaboration de normes Codex et textes connexes (*Manuel de procédure du Codex Alimentarius*), de préférence par courriel, aux adresses indiquées en page de couverture et ce avant le 1<sup>er</sup> mars 2011.

**ANNEX I**  
**Established ADI and ARfD values and recommended MRL, STMR and HR values**

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
Bifenazate (219) ADI: 0–0.01 mg/kg bw  ARfD: Unnecessary	VD 0071	Beans (dry)	0.3		0.01	
	FB 0264	Blackberries	7		2.25	
	FB 0266	Dewberries (including Boysenberry and Loganberry)	7		2.25	
	FB 0272	Raspberries, Red, Black	7		2.25	
	VP 0060	Legume vegetables	7		1.5	
<p>Definition of the residue (for compliance with the MRL for plant and animal commodities and for estimation of dietary intake for plant and animal commodities): Sum of bifenazate and bifenazatediazene (diazencarboxylic acid, 2-(4-methoxy-[1,1'-biphenyl-3-yl] 1-methylethyl ester), expressed as bifenazate.</p> <p>The residue is fat-soluble.</p>						
Bifenthrin (178)** ADI: 0–0.01 mg/kg bw ARfD: 0.01 mg/kg bw	FI 0327	Banana	0.1		0.01	0.01
	GC 0640	Barley	W	0.05 *		
	AS 0640	Barley straw and fodder, dry	W	0.5		
	FB 0264	Blackberries	1		0.29	0.51
	VB 0040	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	0.3		0.115	0.19
	MF 0812	Cattle fat	W	0.5		
	MO 1280	Cattle kidney	W a	0.05*		
	MO 1281	Cattle liver	W a	0.05*		
	MM 0812	Cattle meat	W a	0.5 (fat)		
	ML 0812	Cattle milk	W a	0.05*		
	PE 0840	Chicken eggs	W	0.01*		
	PF 0840	Chicken fat	W	0.05*		
	PM 0840	Chicken meat	W	0.05* (fat)		
	PO 0840	Chicken, Edible offal of	W	0.05*		
	FC 0001	Citrus fruits	0.05		0.05	0.05
	SO 0691	Cotton seed	0.5		0.05	
	AB 1203	Cotton seed meal			0.003	
	OR 0691	Cotton seed oil, edible			0.005	
	FB 0266	Dewberries (including Boysenberry and Loganberry)	1		0.29	0.51
	MO 0105	Edible offal (Mammalian)	0.2		0.07	0.165
	VO 0440	Egg plant	0.3		0.05	0.1
	FC 0203	Grapefruit	W b	0.05*		
	DH 1100	Hops, dry Beer	20	10	1.9 0.011	
	FC 0204	Lemon	W b	0.05*		
	GC 0645	Maize	0.05*	0.05*	0	
	AS 0645	Maize fodder	15	0.2	2.2 dw	5.5 dw
	OC 0645	Maize oil, crude			0	
OR 0645	Maize oil, edible			0		

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	CF 1255	Maize flour			0	
		Maize grits			0	
		Maize starch			0	
	FI 0345	Mango	0.5 c		0.01	0.01
	MM 0095	Meat (from mammals other than marine mammals)	3 (fat)		0.59 fat 0.07 muscle	1.9 fat 0.104 muscle
	FM 0183	Milk fats	3		0.49	
	ML 0106	Milks	0.2		0.053	
	VL 0485	Mustard greens	4		1.16	2.1
	VO 0442	Okra	0.2		0.07	0.11
	FC 0208	Orange, sweet	W b	0.05*		
	FI 0350	Papaya	0.4 c		0.01	0.01
	AL 0072	Pea hay or Pea fodder (dry)	0.7		0.093 dw	0.39 dw
	FP 0230	Pear	W	0.5		
	VO 0051	Peppers	0.5		0.14	0.31
	HS 0444	Peppers, Chili, dried	5		1.4	
	VR 0589	Potato	W d	0.05*		
	VD 0070	Pulses	0.3		0.05	
	VL 0494	Radish leaves (including Radish tops)	4		1.75	2.3
	SO 0495	Rape seed	0.05		0.05	
	OR 0495	Rape seed oil, edible	0.1		0.08	
		Rape seed meal			0.027	
	FB 0272	Raspberries, Red, Black	1		0.29	0.51
	VR 0075	Root and tuber vegetables	0.05		0.05	0.05
	AB 1265	Soya bean meal			0.01	
	OR 0541	Soya bean oil, refined			0.05	
	FB 0275	Strawberry e	3	1	0.46	2.3
	DT 1114	Tea, Green, Black (black, fermented and dried)	30		5.2	
	VO 0448	Tomato	0.3		0.06	0.15
	VW 0448	Tomato paste			0.04	
		Tomato puree			0.04	
	TN 0085	Tree nuts	0.05		0.05	0.05
	GC 0654	Wheat	0.5 Po	0.5 Po	0.25	0.4
	CM 0654	Wheat bran, unprocessed	2 PoP	2 PoP	0.79 PoP	1.26 PoP
	CF 1211	Wheat flour	W f	0.2 PoP		
	CF 1210	Wheat germ	1 Po		0.45 PoP	0.72 PoP
	AS 0654	Wheat straw and fodder, dry	W	0.5		
	CF 1212	Wheat wholemeal	W f	0.5 PoP		

Definition of the residue (for compliance with the MRL for plant and animal commodities and for estimation of dietary intake for plant and animal commodities): bifenthrin (sum of isomers).

The residue is fat-soluble.

a The recommendations for cattle kidney and cattle liver are withdrawn, to be replaced by a recommendation for mammalian edible offal. Recommendations for cattle fat, meat and cattle milk are withdrawn and replaced by recommendations for mammalian meat and milks.

b The recommendations for grapefruit, lemon and orange, sweet are withdrawn to be replaced by recommendation for citrus fruits.

c The recommendations for mango, okra and papaya are based on reported use conditions provided appropriate protection of the crop, but were not supported by official information on uses.

d The recommendation for potato is withdrawn to be replaced by recommendation for root and tuber vegetables.

e For strawberry, the ARfD is exceeded. No alternative GAP is available.

f The recommendations for maximum residue levels for wheat flour and whole meal are withdrawn, because they are covered by the recommendation for wheat.

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
Boscalid (221) ADI: 0–0.04 mg/kg bw ARfD: Unnecessary	FC 0001	Citrus fruits	2		0.05	
	AB 0001	Citrus pulp, dry	6		1.5	
	DH 1100	Hops, dry	60		21.5	
	VL 0053	Leafy vegetables	40	30	3.65	
		Orange juice			0.0108	
	VS 0078	Stalk and stem vegetables	30		8.55	
		Citrus oil	50		27.7	
Definition of the residue (for compliance with the MRL for plant and animal commodities and for estimation of dietary intake for plant commodities): boscalid.						
Definition of the residue (for estimation of dietary intake for animal commodities): sum of boscalid, 2-chloro-N-(4'-chloro-5-hydroxybiphenyl-2-yl)nicotinamide including its conjugate, expressed as boscalid.						
The residue is fat soluble.						
Cadusafos (174) **	FI 0327	Banana	0.01	0.01	0.005	0.005
	VR 0589	Potato	W	0.02		
ADI: 0–0.0005 mg/kg bw ARfD: 0.001 mg/kg bw						
Definition of the residue (for compliance with the MRL for plant and animal commodities and for estimation of dietary intake for plant and animal commodities): Cadusafos						
The residue is not fat-soluble.						
Chlorantraniliprole (230) ADI: 0–2 mg/kg bw ARfD: Unnecessary	AL 1020	Alfalfa fodder	50		17.3	
	FB 0018	Berries and other small fruits	1		0.336	
		Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	2		0.385	
	FC 0001	Citrus fruits	0.5		0.07	
	MO 0105	Edible offal (Mammalian)	0.2	0.01*	0.03 kidney 0.047 liver	
		Eggs	0.1	0.01*	0.023	
	PE 0112	Eggs	0.1	0.01*	0.023	
	FB 0269	Grapes	W	1		
	AS 0645	Maize fodder	25		3.1	
	MM 0095	Meat (from mammals other than marine mammals)	0.2 (fat)	0.01 * fat	0.049 fat 0.009 muscle	
		Milk fats	0.2	0.1	0.048	
	ML 0106	Milks	0.05	0.01*	0.006	
	HH 0738	Mints	15		4.6	
	PO 0111	Poultry, Edible offal of	0.01*		0.0016	
	PM 0110	Poultry meat	* (fat)		0.0008 fat 0.00007 muscle	
		Sugar cane	0.5		0.145	
	VO 0447	Sweet corn (corn-on-the-cob)	0.01*		0.01	
TN 0085	Tree nuts	0.02		0.01		

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
Definition of the residue (for compliance with the MRL and for estimation of dietary intake) for plant and animal commodities: chlorantraniliprole						
The residue is fat-soluble						
Chlorothalonil (081)** ADI: 0–0.02 mg/kg bw ARfD: 0.6 mg/kg bw	FI 0327 GC 0640 AS 0640	Banana Barley Barley straw and fodder, dry	W W W	0.01* c 0.1 20		
4-Hydroxy-2,5,6-trichloroisophthalonitrile a ADI: 0–0.008 mg/kg bw	VD 0071	Beans (dry)	W	0.2		
	FB 0018	Berries and other small fruit (except grapes)			SDS-3701: 0.01	SDS-3701: 0.06
ARfD: 0.03 mg/kg bw	VB 0040	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas			SDS-3701: 0.01	SDS-3701: 0.02
	VB 0400	Broccoli	W	5		
3-amido-2,4,5-trichlorobenzoic acid b	VB 0402	Brussels sprouts	6	5	Chlorothalonil: 1.5	Chlorothalonil: 2.8
	VA 0035	Bulb vegetables			SDS-3701: 0.01	SDS-3701: 0.04
	VB 0041	Cabbages, Head	W	1		
	VR 0577	Carrot	W	1		
	VB 0404	Cauliflower	W	1		
	VX 0624	Celery	20	10	Chlorothalonil: 2.65	Chlorothalonil: 7.5
	HH 0624	Celery leaves	W	3		
	GC 0080	Cereal grains			SDS-3701: 0.02	
	FS 0013	Cherries	W	0.5		
	VP 0526	Common bean (pods an/or immature seeds)	W	5		
	FB 0265	Cranberry	W	5		
	VC 0424	Cucumber	3	5	Chlorothalonil: 0.41	Chlorothalonil: 1.3
	FB 0021	Currants, Black, Red, White	20	5	Chlorothalonil: 20 d	Chlorothalonil: 20 d
	DF 0269	Dried grapes (= currants, Raisins and Sultanas)			Chlorothalonil: 0.248	Chlorothalonil: 0.416
					SDS-3701: 0.0079	SDS-3701: 0.19
	MO 0105	Edible offal (Mammalian)	0.2		SDS-3701: 0.16	SDS-3701: 0.18
	PE 0112	Eggs		0.05	SDS-3701: 0.031	SDS-3701: 0.04
	VB 0042	Flowerhead brassicas (includes Broccoli, Broccoli, Chinese and Cauliflower)	5		Chlorothalonil: 5 c	Chlorothalonil: 5 c
	VC 0045	Fruiting vegetables, Cucurbits			SDS-3701: 0.015	SDS-3701: 0.06
	VO 0050	Fruiting vegetables, other than Cucurbits			SDS-3701: 0.015	SDS-3701: 0.06

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	VC 0425	Gherkin	3		Chlorothalonil: 0.41	Chlorothalonil: 1.3
	FB 0268	Gooseberry	20		Chlorothalonil: 20 d	Chlorothalonil: 20 d
	FB 0269	Grapes	3	0.5	Chlorothalonil: 0.955 SDS-3701: 0.01	Chlorothalonil: 1.6 SDS-3701: 0.15
	JF 0269	Grape juice			Chlorothalonil: 0.134 SDS-3701: 0.0027	
	AB 0269	Grape pomace, dry			Chlorothalonil: 0.745 SDS-3701: 0.031	
		Grape, pomace wet			Chlorothalonil: 1.24 SDS-3701: 0.012	
	HH 0092	Herbs			SDS-3701: 0.02	SDS-3701: 0.19
	VL 0053	Leafy vegetables			SDS-3701: 0.02	SDS-3701: 0.19
	VA 0384	Leek	40		Chlorothalonil: 17.5	Chlorothalonil: 22
	AL 0157	Legume animal feeds			SDS-3701: 0.03	SDS-3701: 0.03
	VP 0060	Legume vegetables	W	5	SDS-3701: 0.01	SDS-3701: 0.02
	MF 0100	Mammalian fats (except milk fats)	0.07		SDS-3701: 0.025	SDS-3701: 0.05
	MM 0095	Meat (from mammals other than marine mammals)	0.02		SDS-3701: 0.01	SDS-3701: 0.012
	VC 0046	Melons, except Watermelon	2	2	Chlorothalonil: 0.04	Chlorothalonil: 0.21
	ML 0106	Milks	0.07		SDS-3701: 0.05	
	SO 0088	Oilseed			SDS-3701: 0.02	
	VA 0385	Onion, Bulb	W	0.5		
	VA 0386	Onion, Chinese	10		Chlorothalonil: 0.835	Chlorothalonil: 7.5
	VA 0387	Onion, Welsh	10		Chlorothalonil: 0.835	Chlorothalonil: 7.5
	FI 0350	Papaya	20		Chlorothalonil: 2.3 SDS-3701: 0.01	Chlorothalonil: 6.4 SDS-3701: 0.01
	FS 0247	Peach	W	0.2		

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	SO 0697	Peanut	0.1	0.05	Chlorothalonil: 0.01	
	HS 0444	Peppers Chili, dried	W	70		
	VO 0445	Pepper, sweet (including Pimento or pimiento)	W	7		
	VR 0589	Potato	W	0.2		
	PF 0111	Poultry fats	0.01		SDS-3701: 0.01	SDS-3701: 0.01
	PM 0110	Poultry meat	0.01		SDS-3701: 0.01	SDS-3701: 0.01
	PO 0113	Poultry skin	0.01		SDS-3701: 0.01	SDS-3701: 0.01
	PO 0111	Poultry, edible offal of	0.07		SDS-3701: 0.039	SDS-3701: 0.05
	VD 0070	Pulses	1		Chlorothalonil: 0.19 SDS-3701: 0.02	
	VR 0075	Root and tuber vegetables	0.3		Chlorothalonil: 0.3 d SDS-3701: 0.02	Chlorothalonil: 0.3 d SDS-3701: 0.03
		Root and tuber vegetables, tops and leaves			SDS-3701: 0.02	SDS-3701: 0.04
	VA 0389	Spring onion	10		Chlorothalonil: 0.835	Chlorothalonil: 7.5
	VC 0431	Squash, Summer	3	5	Chlorothalonil: 0.41	Chlorothalonil: 1.3
	VS 0078	Stalk and stem vegetables			SDS-3701: 0.01	SDS-3701: 0.02
	AS 0081	Straw and fodder (dry) of cereal grains			SDS-3701: 0.03	SDS-3701: 0.08
	FB 0275	Strawberry	5		Chlorothalonil: 2.05	Chlorothalonil: 3
	VO 0447	Sweet Corn (corn-on-the-cob)	W	0.01*		
	VO 0448	Tomato	W	10		
	GC 0654	Wheat	W	0.1		
	AS 0654	Wheat, straw and fodder, dry	W	20		
		Wine			Chlorothalonil: 0.0096 SDS-3701: 0.019	
	VC 0433	Winter squash	W	5		

Definition of the residue (for compliance with MRL) for plant commodities: chlorothalonil

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
Definitions of the residue (for estimation of dietary intake) for plant commodities: - chlorothalonil - SDS-3701 (2,5,6-trichloro-4-hydroxyisophthalonitrile) all considered separately						
Definition of the residue (for compliance with MRL and for estimation of dietary intake) for animal commodities: SDS-3701 (2,5,6-trichloro-4-hydroxyisophthalonitrile)						
The residue is not fat-soluble.						
a Company Code SDS-3701						
b 3-carbamyl-2,4,5-trichlorobenzoic acid (R611965) - ADI and ARfD considered unnecessary as covered by the parent compound						
c Based on bagged bananas						
d Based on the maximum residue level						
Clothianidin (238)* ADI: 0–0.1 mg/kg bw ARfD: 0.6 mg/kg bw	FC 0001	Citrus fruits	0.07 (T)		0.02	0.02
	FP 0009	Pome fruits	0.4 (C,t)		0.10	0.20
	FS 0012	Stone fruits	0.2 (cT)		0.04	0.12
	DF 0014	Prunes	0.2 (cT)		0.07	0.21
	FB 0018	Berries and other small fruits (except grapes)	0.07 (c,T)		0.01	0.05
	FB 0269	Grapes	0.7 (C,t)		0.12	0.41
	DF 0269	Dried grapes (= currants, Raisins and Sultanas)	1 (C,t)		0.31	1.066
	JF 0269	Grape juice	0.2 (C,t)		0.18	-
	FI 0327	Banana	0.02 (C,t)		0.02	0.02
	FI 0350	Papaya	0.01* (T)		0	0
	FI 0353	Pineapple	0.01* (T)		0	0
	VB 0040	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	0.2 (T)		0.015	0.04
	VC 0045	Fruiting vegetables, Cucurbits	0.02* (T)		0.02	0.02
	VO 0050	Fruiting vegetables, other than cucurbits (except sweet corn)	0.05 (T)		0.02	0.03
	VO 0447	Sweet corn (corn-on-the-cob)	0.01* (C,T)		0.01	0.01
	HS 0444	Pepper Chili, dried	0.5 (T)		0.2	0.3
	VL 0053	Leafy vegetables	2 (T)		0.52	0.80
	VP 0060	Legume vegetables	0.01* (T)		0.01	0.01
	VD 0070	Pulses	0.02 (T)		0.02	-
VR 0075	Root and tuber vegetables	0.2 (C,T)		0.02	0.15	
VS 0078	Stalk and stem vegetables (except artichoke and celery)	0.04 (C)		0.01	0.025	
VS 0620	Artichoke, Globe	0.05 (T)		0.024	0.029	
VS 0624	Celery	0.04 (T)		0.01	0.02	
GC 0640	Barley	0.04 (cT)		0.01	-	
GC 0645	Maize	0.02 (cT)		0.02	-	
GC 0656	Popcorn	0.01* (c, T)		0.01	-	
GC 0649	Rice	0.5 (C)		0.145	-	
GC 0651	Sorghum	0.01* (C)		0.01	-	

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	GC 0654	Wheat	0.02*(c,T)		0.02	-
	GS 0659	Sugar cane	0.4 (C)		0.03	0.14
	TN 0672	Pecan	0.01*(T)		0.01	0.01
	SO 0088	Oilseed	0.02*(c,T)		0.02	-
	SB 0715	Cacao beans	0.02*(T)		0.02	-
	SB 0716	Coffee beans	0.05 (T)		0.015	-
	AL 0072	Pea hay or Pea fodder (dry)	0.2, dw (T)		0.05 dw	0.10 dw
	AS 0640	Barley straw and fodder, dry	0.2, dw (T,c)		0.05 dw	0.14 dw
	AS 0645	Maize fodder	0.01 * dw (T)		0.01 dw	0.01 dw
	AS 0651	Sorghum straw and fodder, dry	0.01* dw (C)		0.01 dw	0.01 dw
	AS 0654	Wheat straw and fodder, dry	0.2 dw (T,c)		0.05 dw	0.14 dw
	DT 1114	Tea, Green, Black (black, fermented and dried)	0.7 (T)		0.12	-
	MM 0095	Meat (from mammals other than marine mammals)	0.02* (C, t)		0.02	0.02
	MF 0100	Mammalian fats (except milk fats)	0.02* (C, t)		0.02	0.02
	MO 0105	Edible offal (Mammalian)(except liver)	0.02* (C, t)		0.02	0.02
	MO 0099	Liver of cattle, goats, pigs and sheep	0.2 (c, T)			
	ML 0106	Milks	0.02		0.002	-
	PM 0110	Poultry meat	0.01* (C, t)		0.01	0.01
	PF 0111	Poultry fats	0.01* (C, t)		0.01	0.01
	PO 0111	Poultry, edible offal of	0.1 (T, c)		0.018	0.05
	PE 0112	Eggs	0.01* (C, t)		0.01	0.01
Definition of the residue for compliance with the MRL and for estimation of dietary intake for plant commodities: sum of clothianidin and its Z-isomers.						
Definition of the residue for compliance with the MRL and for estimation of dietary intake for animal commodities: sum of clothianidin and its Z-isomers.						
The residue is not fat-soluble.						
Cyproconazole (239)* ADI: 0–0.02 mg/kg bw ARfD: 0.06 mg/kg bw	VD 0071	Beans (dry)	0.02*		0.02	0.02
	GC 0080	Cereal grains (except maize, rice and sorghum)	0.08		0.02	0.07
	MO 0105	Edible offal (Mammalian)	0.5		0.14	0.46
	PE 0112	Eggs	0.01*		0.01	0.01
	GC 0645	Maize	0.01*		0.01	0.01
	AS 0645	Maize fodder	2		0.28	1.5
	MM 0095	Meat (from mammals other than marine mammals)	0.02 (fat)		0.003 muscle 0.003 fat	0.003 muscle 0.02 fat
	ML 0106	Milks	0.01		0.009	
	VD 0072	Peas (dry)	0.02*		0.02	0.02

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	VP 0064	Peas, shelled (succulent seeds)	0.01		0.01	0.01
	PO 0111	Poultry, edible offal of	0.01*		0	0.01
	PM 0110	Poultry meat	0.01*		0.01 muscle 0.01 fat	0.01 muscle 0.01 fat
	SO 0495	Rape seed	0.4		0.065	0.23
	OR 0495	Rape seed oil, edible			0.0052	
	VD 0541	Soya bean (dry)	0.07		0.02	0.05
	AL 0541	Soya bean fodder	3		0.66	1.9
	OR 0541	Soya bean oil, refined	0.1		0.036	
	AB 1265	Soya bean meal			0.013	
	AS 0081	Straw and fodder (dry) of cereal grains (except maize, rice and sorghum)	5		0.785	3.6
	VR 0596	Sugar beet	0.05		0.02	0.04
Definition of the residue for compliance with the MRL and for estimation of dietary intake for plant commodities: Cyproconazole.						
Definition of the residue for compliance with the MRL for animal commodities: Cyproconazole						
Definition of the residue for estimation of dietary intake for animal commodities except milk: Cyproconazole.						
Definition of the residue for estimation of dietary intake of milk: sum of cyproconazole and metabolites M21 ((5-(4-chlorophenyl)-5-hydroxy-4-methyl-6-[1,2,4]triazol-1-yl-hex-2-enoic acid) and M36 ( $\delta$ -(4-chlorophenyl)- $\beta$ , $\delta$ -dihydroxy- $\gamma$ -methyl-1H-1,2,4-triazole-1-hexenoic acid) expressed as cyproconazole.						
The residue is fat-soluble.						
Dicamba (240)* ADI: 0–0.3 mg/kg bw	VS 0621 GC 0640	Asparagus Barley	5 7		0.87 1.7 1.6 a	3.3
ARfD: 0.5 mg/kg bw	AS 0640	Barley straw and fodder, dry	50		3.65 a	30 a
	SO 0691	Cotton seed	0.04 *		0.04	
	OR 0691	Cottonseed oil, edible			0.008	
	AS 0162	Hay or fodder (dry) of grasses	30		6.3 a	19 a
	MO 0105	Edible offal (Mammalian)	0.7		0.160 kidney 0.028 liver	0.331 kidney 0.082 liver
	GC 0645	Maize	0.01 *		0.02 0.01 a	
	AS 0645	Maize fodder	0.6		0.06 a	0.33 a
	OC 0645	Maize oil, crude			0.00058	
	MF 0100	Mammalian fats (except milk fats)	0.07		0.023	0.036
	MM0095	Meat (from mammals other than marine mammals)	0.03		0.01	0.02
	ML 0106	Milks	0.2		0.021	
	PF 0111	Poultry fats	0.04		0.01	0.01
	PM 0110	Poultry meat	0.02		0.01	0.012
	PO 0111	Poultry, edible offal of	0.07		0.01 Liver	0.044 Liver
	PE 0112	Eggs	0.01 *		0.01	0.01

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	GC 0651	Sorghum	4		2.0 1.0 a	
	AS 0651	Sorghum straw and fodder, dry	8		1.3 a	5.4 a
	GS 0659	Sugar cane	1		0.095	1.1
	DM 0659	Sugar cane molasses			3.4 4.0 a	
	VO 1275	White sugar			0.05	
	GC 0654	Sweet corn (kernels)	0.02		0.04	0.04
	GC 0654	Wheat	2		0.26 0.22 a	
	CF 0654	Wheat bran, processed			0.26	
	CF 1211	Wheat flour			0.02	
	AS 0654	Wheat straw and fodder, dry	50		3.8 a	30 a
<p>Definition of the residue for compliance with the MRL for plant commodities: dicamba            Definition of the residue for estimation of dietary intake for plant commodities: sum of dicamba and 5-OH dicamba expressed as dicamba            Definition of the residue for compliance with the MRL and for estimation of dietary intake for animal commodities: sum of dicamba and 3,6-dichlorosalicylic acid (DCSA) expressed as dicamba</p> <p>The residue is not fat-soluble            a highest residue and median residue for the estimation of animal dietary burden expressed on a dry weight basis (residues of dicamba only)</p>						
Difenoconazole (224) ADI: 0–0.01 mg/kg bw ARfD: 0.3 mg/kg bw	AM 0660	Almond hulls			1.24	3.22
	VP 0060	Legume vegetables	0.7		0.07	0.5
	MO 0105	Edible offal (Mammalian)	0.2		0.041	0.12
	VR 0604	Ginseng	0.5		0.02	0.36
	MM 0095	Meat (from mammals other than marine mammals)	0.05 (fat) a		0.01 muscle 0.012 fat	0.021 muscle 0.031 fat
	ML 0106	Milks	0.005* a		0.001	
	FI 0350	Papaya	0.3b		0.065	0.13
	FI 0351	Passion fruit	0.05		0.01	0.04
	TN 0085	Tree nuts	0.03		0.01	0.02
<p>Definition of the residue (for compliance with the MRL and for estimation of dietary intake) for plant commodities: difenoconazole.            Definition of the residue (for compliance with the MRL and for estimation of dietary intake) for animal commodities: sum of difenoconazole and 1-[2-chloro-4-(4-chloro-phenoxy)-phenyl]-2-(1,2,4-triazol)-1-yl-ethano), expressed as difenoconazole.</p> <p>The residue is fat-soluble</p> <p>a The maximum residue limit recommended by the 2007 JMPR remained the same.            b The recommendation is based on reported use conditions to provide appropriate protection of the crop, but it is not supported by official information on use</p>						
<p>Dithianon (180) **            ADI: 0–0.01 mg/kg bw            ARfD: 0.1 mg/kg bw</p>						
Endosulfan (032) ADI: 0–0.006 mg/kg bw ARfD: 0.08 mg/kg bw	DT 1114	Tea, Green, Black (black, fermented and dried)	10	W	4.1	

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
Definition of the residue (for compliance with the MRL and for estimation of the dietary intake) for plant commodities: sum of alpha endosulfan, beta endosulfan and endosulfan sulfate.						
The residue is fat-soluble.						
Etoazole (241)* ADI: 0–0.05 mg/kg bw ARfD: Unnecessary	AM 0660 FC 0001 JF 0001 VC 0424 FB 0269 DF 0269 JF 0269 MO 0105 DH 1100 MM 0095 ML 0106 HH 0738 DT 1114 TN 0085	Almond hulls Citrus fruits Citrus juice Cucumber Grapes Dried grapes (= currants, Raisins and Sultanas) Grape juice Edible offal (mammalian) Hops, dry Meat (from mammals other than marine mammals) Milks Mints Mint oil Tea, Green, Black (black, fermented and dried) Tree nuts	3 0.1 0.02 0.5 0.044 0.068 0.01* 15 0.01* 0.01* 15 15 0.01*		0.23 0.01 0.005 0.01 0.04 0.044 0.068 0 4.2 0 0 4.9 7.8 4.75 0	
Definition of the residue (for compliance with the MRL and for estimation of the dietary intake) for plant and animal commodities: etoazole						
The residue is fat-soluble						
Fenpyroximate (193) ADI: 0–0.01 mg/kg bw ARfD: 0.02 mg/kg bw	FP 0226 FC 0001 VC 0424 DF 0269 FB 0269 VO 0050 VC 0046 FC 0004 HS 0444 FP 0009 TN 0085	Apple Citrus fruits Cucumber Dried grapes (= Currants, Raisins and Sultanas) Grapes Fruiting vegetables, other than Cucurbits (except sweet corn and mushrooms) Melons, except Watermelon Oranges, Sweet, Sour (including Orange-like hybrids): several cultivars Peppers Chili, dried Pome fruits Tree nuts	W a 0.5 0.03 0.3 0.1 0.2 0.05 W a 1 0.3 0.05 *	0.3 1 0.2	0.034 0.01 0.06 0.02 0.06 0.05 0.05 0.09 0.05 *	0.067 0.02 0.14 0.05 0.14 0.05 0.9 0.16 0.05 *
Definition of the residue (for compliance with the MRL and for estimation of dietary intake) and for plant and animal commodities: fenpyroximate						
a Replaced by commodity group maximum residue level recommendation						

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
Flubendiamide (242)* ADI: 0–0.02 mg/kg bw  ARfD: 0.2 mg/kg bw	AM 0660	Almond hulls	10		2.45	
	VB 0040	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	4		0.365	2.7
	VS 0624	Celery	5		1.7	2.6
	SO 0691	Cotton seed	1.5		0.15	
	VC 0045	Fruiting vegetables, Cucurbits	0.2		0.045	0.09
	MO 0105	Edible offal (Mammalian)	1		0.32	0.57
	FB 0269	Grapes	2		0.42	0.81
	GC 0645	Maize	0.02		0.01	
	CF 1255	Maize flour			0.021	
	VP 0060	Legume vegetables	2		0.43	0.90
	VL 0482	Lettuce, Head	5		0.875	2.2
	VL 0483	Lettuce, leaf	7		1.7	4.0
	MM 0095	Meat (from mammals other than marine mammals) (fat)	2 (fat)		0.06 muscle 0.62 fat	0.13 muscle 1.2 fat
	ML 0106	Milks	0.1		0.066	
	FM 0183	Milk fats	5		1.6	4.0
	AL 0072	Pea hay or Pea fodder (dry)	40		13.5	26
	VO 0051	Peppers	0.7		0.09	0.37
	HS 0444	Peppers Chili, dried	7		0.9	
	FP 0009	Pome fruits	0.8		0.25	0.59
	VD 0070	Pulses	1		0.18	
	AL 0541	Soya bean fodder	60		27.5	41
	FS 0012	Stone fruits	2		0.585	1.0
	VO 0447	Sweet corn (corn-on-the-cob)	0.02		0.01	0.01
	DT 1114	Tea, Green, Black (black, fermented and dried)	50		23	29
	VO 0448	Tomato	2		0.35	0.63
	TN 0085	Tree Nuts	0.1		0.015	0.05
	Definition of the residue (for compliance with the MRL and for estimation of dietary intake) for animal and plant commodities: flubendiamide					
The residue is fat-soluble						
Fludioxonil (211) ADI: 0–0.4 mg/kg bw ARfD: Unnecessary	FC 0001	Citrus fruits	10 Po	7 Po	0.41	
	FI 0355	Pomegranate	2 Po		1.0	
	VR 0508	Sweet potato	10 Po		3.5	
	VR 0600	Yams	10 Po		3.5	
Definition of the residue (for compliance with the MRL and for estimation of dietary intake) for plant commodities: fludioxonil.						
Definition of the residue (for compliance with the MRL and for estimation of dietary intake) for animal commodities: fludioxonil and metabolites determined as 2,2-difluoro-1,3-benzodioxole-4-carboxylic acid and calculated as fludioxonil.						
The residue is fat-soluble.						
Fluopyram (243)* ADI: 0–0.01 mg/kg bw	VC 0424	Cucumber	0.5		0.19	0.11
	FB 0269	Grapes	2		1	0.58

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg	
			New	Previous			
ARfD: 0.5 mg/kg bw	DF 0269	Dried grapes (= currants, Raisins and Sultanas)	5		2.9	1.68	
	MO 0105	Edible offal (mammalian)	0.7		0.574 liver 0.059 kidney	0.472 liver 0.051 kidney	
	MM 0095	Meat (from mammals other than marine mammals)	0.1		0.054 muscle 0.076 fat	0.043 muscle 0.061 fat	
	ML 0106	Milks	0.07			0.039	
	AB 0269	Grape pomace, dry Wine				12.4 0.1	
	JF 0269	Grape juice				0.012	
Definition of the residue (for compliance with the MRL and for estimation of dietary intake) for plant commodities: fluopyram							
Definition of the residue (for compliance with the MRL) for animal commodities: Sum of fluopyram and 2-(trifluoromethyl)benzamide, expressed as fluopyram.							
Definition of the residue (for estimation of dietary intake) for animal commodities: Sum of fluopyram, 2-(trifluoromethyl)benzamide and the combined residues of the E-olefine and Z-olefine isomers of fluopyram, all expressed as fluopyram.							
Although fluopyram (parent compound) is fat-soluble, the 2-(trifluoromethyl)benzamide metabolite (the major component of the residue) is not fat soluble.							
Meptyldinocap (244)* ADI: 0–0.02mg/kg bw ARfD: Unnecessary	VC 0431	Squash, Summer	0.07 a		0.02		
	VC 0424	Cucumber	0.07 a		0.02		
	VC 0046	Melons, except Watermelon	0.5 a		0.005		
	FB 0269	Grapes	0.2 a		0.025		
	JF 0269	Grape juice Wine			0.002 0.00072		
	FB 0275	Strawberry Strawberry jam Strawberry preserve	0.3 b		0.085 0.024 0.024		
	Definition of the residue (for compliance with the MRL and for estimation of dietary intake) for plant commodities: the sum of meptyldinocap, and the corresponding phenol 2, 4-DNOP, expressed as parent meptyldinocap.						
	a The maximum residue level accommodates the residues derived from the use of dinocap on fruiting vegetables, cucumbers. The Meeting recommended to re-evaluate the current CXL of 0.05*.						
b The current dinocap Codex MRL of 0.5 mg/kg covers the use of meptyldinocap.							
Novaluron (217) ADI: 0–0.01 mg/kg bw ARfD: Unnecessary	VD 0071	Beans (dry)	0.1		0.05		
	FB 0020	Blueberries	7		2.1		
	VB 0040	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassica	0.7		0.105		
	VP 0526	Common bean (pods and/or immature seeds)	0.7		0.165		
	MO 0105	Edible offal (Mammalian)	0.7	0.7	0.13		
	PE 0112	Eggs	0.1	0.01*	0.029		
	VC 0045	Fruiting vegetables, Cucurbits	0.2		0.05		

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	VO 0050	Fruiting vegetables, other than Cucurbits (except sweet corn)	0.7		0.1	
	MM 0095	Meat (from mammals other than marine mammals)	10 (fat)	10 (fat)	0.08 muscle 1.7 fat	
	ML 0106	Milks	0.4	0.4	0.13	
	FM 0183	Milk fats	7	7	2.6 cream	
	VL 0485	Mustard greens	25		3.6	
	PM 0110	Poultry meat	0.5 (fat)	0.01* (fat)	0.005 muscle 0.13 fat	
	PO 0111	Poultry, edible offal of	0.1		0.015	
	DF 0014	Prunes	3		1.27	
	FS 0012	Stone fruits	7		2.2	
	FB 0275	Strawberry	0.5		0.15	
	GS 0659	Sugar cane	0.5		0.08	
	VL 0464	Chard	15		4.0	
	VO 0448	Tomato	W a	0.02 *		
		Tomato puree			0.073	
	VW 0448	Tomato paste			0.11	
Definition of the residue (for compliance with the MRL and for estimation of dietary intake) for plant and animal commodities: Novaluron						
The residue is fat-soluble.						
a Replaced by commodity group MRL.						
Tebuconazole (189)**						
ADI: 0–0.03 mg/kg bw						
ARfD: 0.3 mg/kg bw						
Definition of the residue (for compliance with the MRL and for estimation of dietary intake) for plant and animal commodities: tebuconazole						
Thiamethoxam (245)*	VS 0620	Artichoke, Globe	0.5		0.23	0.24
ADI: 0–0.08 mg/kg bw	FI 0327	Banana	0.02*		0.02	0.02
ARfD: 1 mg/kg bw	GC 0640	Barley	0.4		0.12	
	AS 0640	Barley straw and fodder, dry	2		0.39	1.7
	FB 0018	Berries and other small fruits	0.5		0.055	0.26
	VB 0040	Brassica (cole or cabbage) vegetables, Head cabbages, Flowerhead brassicas	5		0.53	1.1
	SB 0715	Cacao beans	0.02*		0.02	
	VS 0624	Celery	1		0.21	0.43
	FC 0001	Citrus fruits	0.5		0.028	0.104
	SB 0716	Coffee beans	0.2		0.035	
	MO 0105	Edible offal (Mammalian)	0.01*		0.01	0.01
	PE 0112	Eggs	0.01*		0.01	0.01
	VC 0045	Fruiting vegetables, Cucurbits	0.5		0.105	0.29

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
	VO 0050	Fruiting vegetables, other than Cucurbits (except sweet corn)	0.7		0.08	0.47
	VL 0053	Leafy vegetables	3		0.54	1.9
	VP 0060	Legume vegetables	0.01*		0.01	0.01
	GC 0645	Maize	0.05		0.02	
	AS 0645	Maize fodder	0.05		0.01	0.04
	MM 0095	Meat (from mammals other than marine mammals)	0.02		0.01 muscle 0.01 fat	0.01 muscle 0.01 fat
	ML 0106	Milks	0.05		0.006	
	SO 0088	Oilseed	0.02*		0.02	
	FI 0350	Papaya	0.01*		0	0
	AL 0072	Pea hay or Pea fodder (dry)	0.3		0.05	0.24
	TN 0672	Pecan	0.01*		0.01	0.01
	HS 0444	Peppers Chili, dried	7		0.8	4.7
	FI 0353	Pineapple	0.01*		0	0
	FP 0009	Pome fruits	0.3		0.07	0.15
	GC 0656	Popcorn	0.01*		0.01	
	PM 0110	Poultry meat	0.01*		0.01	0.01
	PO 0111	Poultry, Edible offal of	0.01*		0.016	0.042
	VD 0070	Pulses	0.04		0.02	
	VR 0075	Root and tuber vegetables	0.3		0.01	0.20
	FS 0012	Stone fruits	1		0.195	0.60
	VO 0447	Sweet corn (corn-on-the-cob)	0.01*		0.01	0.01
	DT 1114	Tea, Green, Black (black, fermented and dried)	20		4.1	
	GC 0654	Wheat	0.05		0.02	
	AS 0654	Wheat straw and fodder, dry	2		0.39	1.7
		Apple juice			0.065	
		Barley flour			0.010	
		Barley, pearled			0.030	
		Coffee, roasted			0.0049	
		Cotton seed oil, Refined			0.0004	
		Orange juice			0.007	
		Prunes, dried			0.16	0.50
		Semolina			0.014	
	JF 0048	Tomato juice			0.054	
	VW 0448	Tomato paste			0.24	
		Tomato pulp			0.08	
		Wheat bran			0.020	
		Wheat bread			0.014	
		Wheat flour			0.014	
		Wine			0.055	

Definition of the residue (for compliance with the MRL) for plant and animal commodities: thiamethoxam.

Definition of the residue (for the estimation of dietary intake) for plant and animal commodities (except poultry): thiamethoxam and CGA 322704 (CGA 322704 to be included with clothianidin and considered separately from thiamethoxam)

Definition of the residue (for the estimation of dietary intake) for poultry: sum of thiamethoxam, CGA 322704 and MU3 and CGA322704 (CGA 322704 to be included with clothianidin and considered separately from thiamethoxam)

See also clothianidin

Pesticide (Codex reference number)	CCN	Commodity	Recommended MRL mg/kg		STMR or STMR-P mg/kg	HR or HR-P mg/kg
			New	Previous		
The residue is not fat-soluble.						
Triazophos (143)	CM 0649	Rice, husked	2		0.421	1.19
ADI: 0–0.001 mg/kg bw	VP 0541	Soya bean (immature seeds)	0.5		0.07	0.15
ARfD: 0.001 mg/kg bw						
Definition of the residue for compliance with the MRL for all commodities and for estimation of dietary intake for plant and animal commodities: triazophos						
The residue is not fat-soluble.						
Recommended MRLs, STMRs and HR values for Spices						
Codex Number	Commodity	Pesticide	Recommended MRL mg/kg		Median residue mg/kg	HR mg/kg
			New	Previous		
028B	Fruit or berry	Carbaryl	0.8		0.1	0.78
		Carbendazim	0.1		0.1	0.1
		Cypermethrin	0.5	0.2	0.05	0.43
		Aldicarb	0.07		0.07	0.07
		Bifenthrin	0.03		0.03	0.03
		Carbosulfan	0.07		0.07	0.07
		Cyfluthrin	0.03		0.03	0.03
		Cyhalothrin	0.03		0.03	0.03
		Deltamethrin	0.03		0.03	0.03
		Fenvalerate	0.03		0.03	0.03
		Methidathion	0.02		0.02	0.02
		Methiocarb	0.07		0.07	0.07
		Methomyl	0.07		0.07	0.07
		Omethoate	0.02		0.02	0.02
		Oxamyl	0.07		0.07	0.07
		Profenofos	0.07		0.07	0.07
		Triazophos	0.07		0.07	0.07
028D	Root or rhizome	Deltamethrin	0.5		0.05	0.33
		Aldicarb	0.02		0.02	0.02
		Bifenthrin	0.05		0.05	0.05
		Captan	0.05		0.05	0.05
		Carbaryl	0.1		0.1	0.1
		Carbendazim	0.1		0.1	0.1
		Carbosulfan	0.1		0.1	0.1
		Cyfluthrin	0.05		0.05	0.05
		Cyhalothrin	0.05		0.05	0.05
		Fenvalerate	0.05		0.05	0.05
		Methidathion	0.05		0.05	0.05
		Methiocarb	0.1		0.1	0.1
		Omethoate	0.05		0.05	0.05
		Oxamyl	0.05		0.05	0.05
		Profenofos	0.05		0.05	0.05
		Triazophos	0.1		0.1	0.1