



**JOINT FAO/WHO FOOD STANDARDS PROGRAMME**  
**CODEX COMMITTEE ON RESIDUES OF VETERINARY DRUGS IN FOODS**  
**25<sup>th</sup> Session**  
**(Virtual)**  
**12-16 and 20 July 2021**

**SUPPORT DOCUMENT TO THE DISCUSSION ON THE MRLs AND RMRs FOR RESIDUES OF VETERINARY DRUGS**

(Prepared by the Codex Secretariat)

## **INTRODUCTION**

This working document is in support to the discussion on the residues of veterinary drugs in foods of the 25<sup>th</sup> Session of the Codex Committee on Residues of Veterinary Drugs in Foods (CCRVDF25). The document includes:

- Part 1 - Codex Maximum Residue Limits (MRLs) and Risk Management Recommendations (RMRs) for Veterinary Drugs as adopted by the Codex Alimentarius Commission as its 41<sup>st</sup> Session (July 2018); and
- Part 2 - Draft and proposed draft MRLs and RMRs arising from the 88<sup>th</sup> Meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA88).

**ACRONYMS**

ADI	Acceptable Daily Intake
ARfD	Acute Reference Dose
BMDL10	Benchmark Dose for a 10% Response
CCRVDF	Codex Committee on Residues of Veterinary Drugs in Foods
EDI	Estimated Dietary Exposure
GEADE	Global Estimate of Acute Dietary Exposure
GECDE	Global Estimate of Chronic Dietary Exposure
JECFA	Joint FAO/WHO Expert Committee on Food Additives
JMPR	Joint FAO/WHO Meeting on Pesticide Residues
LOAEL	Lowest-Observed-Adverse-Effect Level
mADI	Microbiological ADI
MRL	Maximum Residue Limit
NOAEL	No-Observed-Adverse-Effect Level
TMDI	Theoretical Maximum Daily Intake
TRR	Total Radioactive Residue

**Part 1****A) MAXIMUM RESIDUE LIMITS FOR VETERINARY DRUGS IN FOODS**

<b>ABAMECTIN</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		45 (1995); 47 (1996)		
<b>ADI:</b>		0-2 µg/kg body weight (1997). Established for the sum of abamectin and (Z)-8,9 isomer by JMPR (1997)		
<b>Residue Definition:</b>		Avermectin B1a		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Liver	100	26 <sup>th</sup> (2003)	
Cattle	Kidney	50	26 <sup>th</sup> (2003)	
Cattle	Fat	100	26 <sup>th</sup> (2003)	

<b>ALBENDAZOLE</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		34 (1989)		
<b>ADI:</b>		0-50 µg/kg body weight (JECFA34, 1989)		
<b>Residue Definition:</b>		Except milk, 2-aminosulfone metabolite; Milk, not yet identified.		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Not specified	Muscle	100	20 <sup>th</sup> (1993)	
Not specified	Liver	5000	20 <sup>th</sup> (1993)	
Not specified	Kidney	5000	20 <sup>th</sup> (1993)	
Not specified	Fat	100	20 <sup>th</sup> (1993)	
Not specified	Milk (µg/l)	100	20 <sup>th</sup> (1993)	

<b>AMOXICILLIN (antimicrobial agent)</b>				
<b>JECFA Evaluation:</b>		75 (2011); 85 (2017)		
<b>ADI:</b>		0-0.07 µg/kg body weight on the basis of microbiological effects (JECFA75, 2011)		
<b>mADI:</b>		0–0.002 mg/kg body weight (bw) based on the effects of amoxicillin on the intestinal microbiota (JECFA85, 2017)		
<b>ARfD:</b>		0.005 mg/kg bw based on microbiological effects on the intestinal microbiota (JECFA85, 2017)		
<b>EDI:</b>		JECFA75 (2001) did not calculate an EDI for amoxicillin owing to the small number of quantifiable residue data points. Using the model diet of 300 g muscle, 100 g live, 50 g kidney, 50 g fat and 1.5 liter of milk with the MRLs recommended, the TMDI is 31 µg/person, which represents 74% of the upper bound of the ADI.		
<b>GECDE:</b>		0.14 µg/kg bw per day (for the general population), which represents 7% of the upper bound of the mADI (JECFA85, 2017).		
<b>GEADE:</b>		1.4 µg/kg bw (for the general population), which represents 28% of the microbiological ARfD. (JECFA85, 2017). 1.6 µg/kg bw (for children), which represents 31% of the <b>microbiological ARfD</b> (JECFA85, 2017)		
<b>Residue Definition:</b>		Amoxicillin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	50	35 <sup>th</sup> (2012)	
Cattle	Liver	50	35 <sup>th</sup> (2012)	
Cattle	Kidney	50	35 <sup>th</sup> (2012)	
Cattle	Fat	50	35 <sup>th</sup> (2012)	
Cattle	Milk	4	35 <sup>th</sup> (2012)	
Sheep	Muscle	50	35 <sup>th</sup> (2012)	
Sheep	Liver	50	35 <sup>th</sup> (2012)	
Sheep	Kidney	50	35 <sup>th</sup> (2012)	
Sheep	Fat	50	35 <sup>th</sup> (2012)	
Sheep	Milk	4	35 <sup>th</sup> (2012)	
Pigs	Muscle	50	35 <sup>th</sup> (2012)	
Pigs	Liver	50	35 <sup>th</sup> (2012)	
Pigs	Kidney	50	35 <sup>th</sup> (2012)	
Pigs	Fat/Skin	50	35 <sup>th</sup> (2012)	
Finfish <sup>a</sup>	Fillet <sup>b</sup>	50	41 <sup>st</sup> (2018)	
Finfish <sup>a</sup>	Muscle	50	41 <sup>st</sup> (2018)	

<sup>a</sup> The term “finfish” includes all fish species.

<sup>b</sup> Muscle plus skin in natural proportion.

<b>AMPICILLIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		85 (2017)		
<b>mADI:</b>		0–0.003 mg/kg bw based on a NOAEL equivalent to 0.025 mg/kg bw per day for increase in population(s) of ampicillin-resistant bacteria in the gastrointestinal tract in humans, and using a safety factor of 10 (for the variability in the composition of the intestinal microbiota within and between individuals) (JECFA85, 2017).		
<b>ARfD:</b>		0.012 mg/kg bw based on the microbiological end-point (JECFA85, 2017)		
<b>GECDE:</b>		0.29 µg/kg bw per day (for the general population), which represents 10% of the upper bound of the ADI (JECFA85, 2017)		
<b>GEADE:</b>		1.9 µg/kg bw per day (for the general population), which represents 16% of the ARfD (JECFA85, 2017) 1.7 µg/kg bw per day (for children), which represents 14% of the ARfD (JECFA85, 2017)		
<b>Residue Definition:</b>		Ampicillin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Finfish <sup>a</sup>	Fillet <sup>b</sup>	50	41 <sup>st</sup> (2018)	JECFA85 recommended an MRL of 50 µg/kg for ampicillin in finfish muscle and in finfish muscle plus skin in natural proportion, the same as that recommended for amoxicillin, because the modes of action, the physicochemical properties and the toxicological and pharmacokinetic profiles of amoxicillin and ampicillin are very similar.
Finfish <sup>a</sup>	Muscle	50	41 <sup>st</sup> (2018)	

<sup>a</sup> The term “finfish” includes all fish species.

<sup>b</sup> Muscle plus skin in natural proportion.

<b>AVILAMYCIN (antimicrobial agent)</b>				
<b>JECFA Evaluation:</b>		70 (2008)		
<b>ADI:</b>		0-2 mg/kg body weight on the basis of a NOAEL of 150 mg avilamycin activity/kg body weight per day and a safety factor of 100 and rounding to one significant figure (JECFA70, 2008)		
<b>Residue Definition:</b>		Dichloroisoeverninic acid (DIA)		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Pigs	Muscle	200	32 <sup>nd</sup> (2009)	
Pigs	Liver	300	32 <sup>nd</sup> (2009)	
Pigs	Kidney	200	32 <sup>nd</sup> (2009)	
Pigs	Fat/Skin	200	32 <sup>nd</sup> (2009)	
Chicken	Muscle	200	32 <sup>nd</sup> (2009)	
Chicken	Liver	300	32 <sup>nd</sup> (2009)	
Chicken	Kidney	200	32 <sup>nd</sup> (2009)	
Chicken	Fat/Skin	200	32 <sup>nd</sup> (2009)	
Turkey	Muscle	200	32 <sup>nd</sup> (2009)	
Turkey	Liver	300	32 <sup>nd</sup> (2009)	
Turkey	Kidney	200	32 <sup>nd</sup> (2009)	
Turkey	Fat/Skin	200	32 <sup>nd</sup> (2009)	
Rabbits	Muscle	200	32 <sup>nd</sup> (2009)	
Rabbits	Liver	300	32 <sup>nd</sup> (2009)	
Rabbits	Kidney	200	32 <sup>nd</sup> (2009)	
Rabbits	Fat/Skin	200	32 <sup>nd</sup> (2009)	

<b>AZAPERONE (tranquilizing agent)</b>				
<b>JECFA Evaluation:</b>		38 (1991); 43 (1994); 50 (1998); 52 (1999)		
<b>ADI:</b>		0-6 µg/kg body weight (JECFA50, 1998)		
<b>Residue Definition:</b>		Sum of azaperone and azaperol		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Pig	Muscle	60	23 <sup>rd</sup> (1999)	
Pig	Liver	100	23 <sup>rd</sup> (1999)	
Pig	Kidney	100	23 <sup>rd</sup> (1999)	
Pig	Fat	60	23 <sup>rd</sup> (1999)	

<b>BENZYLPENICILLIN/PROCAINE BENZYLPENICILLIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		36 (1990); 50 (1998)		
<b>ADI:</b>		30 µg-penicillin/person/day (JECFA50, 1998). Residues of benzylpenicillin and procaine benzylpenicillin should be kept below this level.		
<b>Residue Definition:</b>		Benzylpenicillin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	50	23 <sup>rd</sup> (1999)	
Cattle	Liver	50	23 <sup>rd</sup> (1999)	
Cattle	Kidney	50	23 <sup>rd</sup> (1999)	
Cattle	Milk (µg/l)	4	23 <sup>rd</sup> (1999)	
Chicken	Muscle	50	23 <sup>rd</sup> (1999)	Applies to procaine benzylpenicillin only
Chicken	Liver	50	23 <sup>rd</sup> (1999)	Applies to procaine benzylpenicillin only
Chicken	Kidney	50	23 <sup>rd</sup> (1999)	Applies to procaine benzylpenicillin only
Pig	Muscle	50	23 <sup>rd</sup> (1999)	
Pig	Liver	50	23 <sup>rd</sup> (1999)	
Pig	Kidney	50	23 <sup>rd</sup> (1999)	

<b>CARAZOLOL</b> (beta-adreniceptor-blocking agent)				
<b>JECFA Evaluation:</b>		38 (1991); 43 (1994); 52 (1999)		
<b>ADI:</b>		0-0.1 µg/kg body weight (JECFA43, 1994). ADI based on the acute pharmacological effects of carazolol.		
<b>Residue Definition:</b>		Carazolol		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Pig	Muscle	5	26 <sup>th</sup> (2003)	The concentration at the injection site two hours after treatment may result in an intake that exceeds the ARfD and therefore, an appropriate withdrawal period should be applied.
Pig	Liver	25	26 <sup>th</sup> (2003)	
Pig	Kidney	25	26 <sup>th</sup> (2003)	
Pig	Fat/Skin	5	26 <sup>th</sup> (2003)	The concentration at the injection site two hours after treatment may result in an intake that exceeds the ARfD and therefore, an appropriate withdrawal period should be applied.

<b>CEFTIOFUR (antimicrobial agent)</b>				
<b>JECFA Evaluation:</b>		45 (1995); 48 (1997)		
<b>ADI:</b>		0-50 µg/kg body weight (JECFA45, 1995)		
<b>Residue Definition:</b>		Desfuroylceftiofur		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	1000	23 <sup>rd</sup> (1999)	
Cattle	Liver	2000	23 <sup>rd</sup> (1999)	
Cattle	Kidney	6000	23 <sup>rd</sup> (1999)	
Cattle	Fat	2000	23 <sup>rd</sup> (1999)	
Cattle	Milk (µg/l)	100	23 <sup>rd</sup> (1999)	
Pig	Muscle	1000	23 <sup>rd</sup> (1999)	
Pig	Liver	2000	23 <sup>rd</sup> (1999)	
Pig	Kidney	6000	23 <sup>rd</sup> (1999)	
Pig	Fat	2000	23 <sup>rd</sup> (1999)	

<b>CHLORTETRACYCLINE/OXYTETRACYCLINE/TETRACYCLINE (antimicrobial agent)</b>				
<b>JECFA Evaluation:</b>		45 (1995); 47 (1996); 50 (1998); 58 (2002)		
<b>ADI:</b>		0-30 µg/kg body weight (JECFA50, 1998). Group ADI for chlortetracycline, oxytetracycline and tetracycline.		
<b>Residue Definition:</b>		Parent drugs, singly or in combination		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	200	26 <sup>th</sup> (2003)	
Cattle	Liver	600	26 <sup>th</sup> (2003)	
Cattle	Kidney	1200	26 <sup>th</sup> (2003)	
Cattle	Milk (µg/l)	100	26 <sup>th</sup> (2003)	
Fish	Muscle	200	26 <sup>th</sup> (2003)	Applies only to oxytetracycline
Giant prawn ( <i>Paeneus monodon</i> )	Muscle	200	26 <sup>th</sup> (2003)	Applies only to oxytetracycline
Pig	Muscle	200	26 <sup>th</sup> (2003)	
Pig	Liver	600	26 <sup>th</sup> (2003)	
Pig	Kidney	1200	26 <sup>th</sup> (2003)	
Poultry	Muscle	200	26 <sup>th</sup> (2003)	
Poultry	Liver	600	26 <sup>th</sup> (2003)	
Poultry	Kidney	1200	26 <sup>th</sup> (2003)	
Poultry	Eggs	400	26 <sup>th</sup> (2003)	
Sheep	Muscle	200	26 <sup>th</sup> (2003)	
Sheep	Liver	600	26 <sup>th</sup> (2003)	
Sheep	Kidney	1200	26 <sup>th</sup> (2003)	
Sheep	Milk (µg/l)	100	26 <sup>th</sup> (2003)	



<b>CLENBUTEROL</b> (adrenoceptor agonist)				
<b>JECFA Evaluation:</b>		47 (1996)		
<b>ADI:</b>		0-0.004 µg/kg body weight (JECFA47, 1996)		
<b>Residue Definition:</b>		Clenbuterol		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	0.2	26 <sup>th</sup> (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Cattle	Liver	0.6	26 <sup>th</sup> (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Cattle	Kidney	0.6	26 <sup>th</sup> (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Cattle	Fat	0.2	26 <sup>th</sup> (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Cattle	Milk (µg/l)	0.05	26 <sup>th</sup> (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Horse	Muscle	0.2	26 <sup>th</sup> (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Horse	Liver	0.6	26 <sup>th</sup> (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Horse	Kidney	0.6	26 <sup>th</sup> (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.
Horse	Fat	0.2	26 <sup>th</sup> (2003)	Due to the potential abuse of this drug, the MRLs are recommended only when associated with a nationally approved therapeutic use, such as tocolysis or as an adjunct therapy in respiratory diseases.

<b>CLOSANTEL</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		36 (1990); 40 (1992)		
<b>ADI:</b>		0-30 µg/kg body weight (JECFA40, 1992)		
<b>Residue Definition:</b>		Closantel		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	1000	20 <sup>th</sup> (1993)	
Cattle	Liver	1000	20 <sup>th</sup> (1993)	
Cattle	Kidney	3000	20 <sup>th</sup> (1993)	
Cattle	Fat	3000	20 <sup>th</sup> (1993)	
Sheep	Muscle	1500	20 <sup>th</sup> (1993)	
Sheep	Liver	1500	20 <sup>th</sup> (1993)	
Sheep	Kidney	5000	20 <sup>th</sup> (1993)	
Sheep	Fat	2000	20 <sup>th</sup> (1993)	

<b>COLISTIN (antimicrobial agent)</b>				
<b>JECFA Evaluation:</b>		66 (2006)		
<b>ADI:</b>		0-7 µg/kg body weight (JECFA66, 2006)		
<b>Residue Definition:</b>		Sum of colistin A and colistin B		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	150	31 <sup>st</sup> (2008)	
Cattle	Liver	150	31 <sup>st</sup> (2008)	
Cattle	Kidney	200	31 <sup>st</sup> (2008)	
Cattle	Fat	150	31 <sup>st</sup> (2008)	
Cattle	Milk	50	31 <sup>st</sup> (2008)	
Sheep	Muscle	150	31 <sup>st</sup> (2008)	
Sheep	Liver	150	31 <sup>st</sup> (2008)	
Sheep	Kidney	200	31 <sup>st</sup> (2008)	
Sheep	Fat	150	31 <sup>st</sup> (2008)	
Sheep	Milk	50	31 <sup>st</sup> (2008)	
Goat	Muscle	150	31 <sup>st</sup> (2008)	
Goat	Liver	150	31 <sup>st</sup> (2008)	
Goat	Kidney	200	31 <sup>st</sup> (2008)	
Goat	Fat	150	31 <sup>st</sup> (2008)	
Pig	Muscle	150	31 <sup>st</sup> (2008)	
Pig	Liver	150	31 <sup>st</sup> (2008)	
Pig	Kidney	200	31 <sup>st</sup> (2008)	
Pig	Fat	150	31 <sup>st</sup> (2008)	The MRL includes skin + fat
Chicken	Muscle	150	31 <sup>st</sup> (2008)	
Chicken	Liver	150	31 <sup>st</sup> (2008)	
Chicken	Kidney	200	31 <sup>st</sup> (2008)	
Chicken	Fat	150	31 <sup>st</sup> (2008)	The MRL includes skin + fat
Chicken	Eggs	300	31 <sup>st</sup> (2008)	
Turkey	Muscle	150	31 <sup>st</sup> (2008)	
Turkey	Liver	150	31 <sup>st</sup> (2008)	
Turkey	Kidney	200	31 <sup>st</sup> (2008)	
Turkey	Fat	150	31 <sup>st</sup> (2008)	The MRL includes skin + fat
Rabbit	Muscle	150	31 <sup>st</sup> (2008)	
Rabbit	Liver	150	31 <sup>st</sup> (2008)	
Rabbit	Kidney	200	31 <sup>st</sup> (2008)	
Rabbit	Fat	150	31 <sup>st</sup> (2008)	

<b>CYFLUTHRIN</b> (insecticide)				
<b>JECFA Evaluation:</b>		48 (1997)		
<b>ADI:</b>		0-20 µg/kg body weight (JECFA48, 1997)		
<b>Residue Definition:</b>		Cyfluthrin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	20	26 <sup>th</sup> (2003)	
Cattle	Liver	20	26 <sup>th</sup> (2003)	
Cattle	Kidney	20	26 <sup>th</sup> (2003)	
Cattle	Fat	200	26 <sup>th</sup> (2003)	
Cattle	Milk (µg/l)	40	26 <sup>th</sup> (2003)	

<b>CYHALOTHRIN</b> (insecticide)				
<b>JECFA Evaluation:</b>		54 (2000); 58 (2002); 62 (2004)		
<b>ADI:</b>		0-5 µg/kg body weight (JECFA62, 2004)		
<b>Residue Definition:</b>		Cyhalothrin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	20	28 <sup>th</sup> (2005)	
Cattle	Liver	20	28 <sup>th</sup> (2005)	
Cattle	Kidney	20	28 <sup>th</sup> (2005)	
Cattle	Fat	400	28 <sup>th</sup> (2005)	
Cattle	Milk	30	28 <sup>th</sup> (2005)	
Pig	Muscle	20	28 <sup>th</sup> (2005)	
Pig	Liver	20	28 <sup>th</sup> (2005)	
Pig	Kidney	20	28 <sup>th</sup> (2005)	
Pig	Fat	400	28 <sup>th</sup> (2005)	
Sheep	Muscle	20	28 <sup>th</sup> (2005)	
Sheep	Liver	50	28 <sup>th</sup> (2005)	
Sheep	Kidney	20	28 <sup>th</sup> (2005)	
Sheep	Fat	400	28 <sup>th</sup> (2005)	

<b>CYPERMETHRIN AND ALPHA-CYPERMETHRIN (insecticide)</b>				
<b>JECFA Evaluation:</b>		62 (2004)		
<b>ADI:</b>		JECFA established a common ADI of 0-20 µg/kg bw for both cypermethrin and alpha-cypermethrin (JECFA62, 2004)		
<b>Residue Definition:</b>		Total of cypermethrin residues (resulting from the use of cypermethrin or alpha-cypermethrin as veterinary drugs)		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Note</b>
Cattle	Muscle	50	29 <sup>th</sup> (2006)	
Cattle	Liver	50	29 <sup>th</sup> (2006)	
Cattle	Kidney	50	29 <sup>th</sup> (2006)	
Cattle	Fat	1000	29 <sup>th</sup> (2006)	
Cattle	Milk	100	29 <sup>th</sup> (2006)	
Sheep	Muscle	50	29 <sup>th</sup> (2006)	
Sheep	Liver	50	29 <sup>th</sup> (2006)	
Sheep	Kidney	50	29 <sup>th</sup> (2006)	
Sheep	Fat	1000	29 <sup>th</sup> (2006)	

<b>DANOFLOXACIN (antimicrobial agent)</b>				
<b>JECFA Evaluation:</b>		48 (1997)		
<b>ADI:</b>		0-20 µg/kg body weight (JECFA48, 1997)		
<b>Residue Definition:</b>		Danofloxacin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	200	24 <sup>th</sup> (2001)	
Cattle	Liver	400	24 <sup>th</sup> (2001)	
Cattle	Kidney	400	24 <sup>th</sup> (2001)	
Cattle	Fat	100	24 <sup>th</sup> (2001)	
Chicken	Muscle	200	24 <sup>th</sup> (2001)	
Chicken	Liver	400	24 <sup>th</sup> (2001)	
Chicken	Kidney	400	24 <sup>th</sup> (2001)	
Chicken	Fat	100	24 <sup>th</sup> (2001)	Fat/skin in normal proportion
Pig	Muscle	100	24 <sup>th</sup> (2001)	
Pig	Liver	50	24 <sup>th</sup> (2001)	
Pig	Kidney	200	24 <sup>th</sup> (2001)	
Pig	Fat	100	24 <sup>th</sup> (2001)	

<b>DELTA METHRIN</b> (insecticide)				
<b>JECFA Evaluation:</b>		52 (1999); 60 (2003)		
<b>ADI:</b>		0-10 µg/kg body weight (1982). Established by JMPR (1982)		
<b>Residue Definition:</b>		Deltamethrin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	30	26 <sup>th</sup> (2003)	
Cattle	Liver	50	26 <sup>th</sup> (2003)	
Cattle	Kidney	50	26 <sup>th</sup> (2003)	
Cattle	Fat	500	26 <sup>th</sup> (2003)	
Cattle	Milk	30	26 <sup>th</sup> (2003)	
Chicken	Muscle	30	26 <sup>th</sup> (2003)	
Chicken	Liver	50	26 <sup>th</sup> (2003)	
Chicken	Kidney	50	26 <sup>th</sup> (2003)	
Chicken	Fat	500	26 <sup>th</sup> (2003)	
Chicken	Eggs	30	26 <sup>th</sup> (2003)	
Salmon	Muscle	30	26 <sup>th</sup> (2003)	
Sheep	Muscle	30	26 <sup>th</sup> (2003)	
Sheep	Liver	50	26 <sup>th</sup> (2003)	
Sheep	Kidney	50	26 <sup>th</sup> (2003)	
Sheep	Fat	500	26 <sup>th</sup> (2003)	

<b>DERQUANTEL</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		75 (2011); 78 (2013)		
<b>ADI:</b>		0-0.3 µg/kg body weight on the basis of a LOAEL of 0.1 mg/kg body weight per day for acute clinical observations in dogs, consistent with antagonistic activity on the nicotinic acetylcholine receptors. A safety factor of 300 was applied to the LOAEL (JECFA75, 2011).		
<b>EDI:</b>		There were insufficient data to calculate an EDI, and the TMDI approach was used. Using the model diet and the MT:TR approach, these MRLs result in an estimated dietary exposure of 6.8 µg/person, which represents approximately 38% of the upper bound of the ADI (JECFA78, 2013).		
<b>Residue Definition:</b>		Derquantel		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Sheep	Muscle	0.3	38 <sup>th</sup> (2015)	
Sheep	Liver	0.8	38 <sup>th</sup> (2015)	
Sheep	Kidney	0.4	38 <sup>th</sup> (2015)	
Sheep	Fat	7.0	38 <sup>th</sup> (2015)	

<b>DEXAMETHASONE</b> (glucocorticosteroid)				
<b>JECFA Evaluation:</b>		70 (2008)		
<b>ADI:</b>		0-0.015 µg/kg body weight (JECFA42, 1995)		
<b>Residue Definition:</b>		Dexamethasone		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	1.0	32 <sup>nd</sup> (2009)	
Cattle	Liver	2.0	32 <sup>nd</sup> (2009)	
Cattle	Kidney	1.0	32 <sup>nd</sup> (2009)	
Cattle	Milk (µg/l)	0.3	32 <sup>nd</sup> (2009)	
Pig	Muscle	1.0	32 <sup>nd</sup> (2009)	
Pig	Liver	2.0	32 <sup>nd</sup> (2009)	
Pig	Kidney	1.0	32 <sup>nd</sup> (2009)	
Horses	Muscle	1.0	32 <sup>nd</sup> (2009)	
Horses	Liver	2.0	32 <sup>nd</sup> (2009)	
Horses	Kidney	1.0	32 <sup>nd</sup> (2009)	

<b>DICLAZURIL</b> (antiprotozoal agent)				
<b>JECFA Evaluation:</b>		45 (1995); 50 (1998)		
<b>ADI:</b>		0-30 µg/kg body weight (JECFA50, 1998)		
<b>Residue Definition:</b>		Diclazuril		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Poultry	Muscle	500	23 <sup>rd</sup> (1999)	
Poultry	Liver	3000	23 <sup>rd</sup> (1999)	
Poultry	Kidney	2000	23 <sup>rd</sup> (1999)	
Poultry	Fat/Skin	1000	23 <sup>rd</sup> (1999)	
Rabbit	Muscle	500	23 <sup>rd</sup> (1999)	
Rabbit	Liver	3000	23 <sup>rd</sup> (1999)	
Rabbit	Kidney	2000	23 <sup>rd</sup> (1999)	
Rabbit	Fat	1000	23 <sup>rd</sup> (1999)	
Sheep	Muscle	500	23 <sup>rd</sup> (1999)	
Sheep	Liver	3000	23 <sup>rd</sup> (1999)	
Sheep	Kidney	2000	23 <sup>rd</sup> (1999)	
Sheep	Fat	1000	23 <sup>rd</sup> (1999)	

<b>DICYCLANIL</b> (insecticide)				
<b>JECFA Evaluation:</b>		54 (2000); 60 (2003)		
<b>ADI:</b>		0-7 µg/kg body weight (JECFA54, 2000)		
<b>Residue Definition:</b>		Dicyclanil		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Sheep	Muscle	150	28 <sup>th</sup> (2005)	
Sheep	Liver	125	28 <sup>th</sup> (2005)	
Sheep	Kidney	125	28 <sup>th</sup> (2005)	
Sheep	Fat	200	28 <sup>th</sup> (2005)	

<b>DIHYDROSTREPTOMYCIN/STREPTOMYCIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		43 (1994); 48 (1997); 52 (1999); 58 (2002)		
<b>ADI:</b>		0-50 µg/kg body weight (JECFA48, 1997). Group ADI for combined residues of dihydrostreptomycin and streptomycin.		
<b>Residue Definition:</b>		Sum of dihydrostreptomycin and streptomycin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	600	24 <sup>th</sup> (2001)	
Cattle	Liver	600	24 <sup>th</sup> (2001)	
Cattle	Kidney	1000	24 <sup>th</sup> (2001)	
Cattle	Fat	600	24 <sup>th</sup> (2001)	
Cattle	Milk	200	26 <sup>th</sup> (2003)	
Chicken	Muscle	600	24 <sup>th</sup> (2001)	
Chicken	Liver	600	24 <sup>th</sup> (2001)	
Chicken	Kidney	1000	24 <sup>th</sup> (2001)	
Chicken	Fat	600	24 <sup>th</sup> (2001)	
Pig	Muscle	600	24 <sup>th</sup> (2001)	
Pig	Liver	600	24 <sup>th</sup> (2001)	
Pig	Kidney	1000	24 <sup>th</sup> (2001)	
Pig	Fat	600	24 <sup>th</sup> (2001)	
Sheep	Muscle	600	24 <sup>th</sup> (2001)	
Sheep	Liver	600	24 <sup>th</sup> (2001)	
Sheep	Kidney	1000	24 <sup>th</sup> (2001)	
Sheep	Fat	600	24 <sup>th</sup> (2001)	
Sheep	Milk	200	26 <sup>th</sup> (2003)	



<b>DIMINAZENE</b> (trypanocide)				
<b>JECFA Evaluation:</b>		34 (1989); 42 (1994)		
<b>ADI:</b>		0-100 µg/kg body weight (JECFA42, 1994)		
<b>Residue Definition:</b>		Diminazene		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	500	22 <sup>nd</sup> (1997)	
Cattle	Liver	12000	22 <sup>nd</sup> (1997)	
Cattle	Kidney	6000	22 <sup>nd</sup> (1997)	
Cattle	Milk (µg/l)	150	22 <sup>nd</sup> (1997)	Limit of quantitation of the analytical method

<b>DORAMECTIN</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		45 (1995); 52 (1999); 58 (2002); 62 (2004)		
<b>ADI:</b>		0-1 µg/kg body weight (JECFA58, 2002)		
<b>Residue Definition:</b>		Doramectin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	10	22 <sup>nd</sup> (1997)	High concentration of residues at the injection site over a 35 day period after subcutaneous or intramuscular administration of the drug at the recommended dose
Cattle	Liver	100	22 <sup>nd</sup> (1997)	
Cattle	Kidney	30	22 <sup>nd</sup> (1997)	
Cattle	Fat	150	22 <sup>nd</sup> (1997)	High concentration of residues at the injection site over a 35 day period after subcutaneous or intramuscular administration of the drug at the recommended dose
Cattle	Milk	15	29 <sup>th</sup> (2006)	Depending on the route and/or time of administration the use of doramectin in dairy cows may result in extended withdrawal periods in milk. This may be addressed in national/regional regulatory programmes.
Pig	Muscle	5	24 <sup>th</sup> (2001)	
Pig	Liver	100	24 <sup>th</sup> (2001)	
Pig	Kidney	30	24 <sup>th</sup> (2001)	
Pig	Fat	150	24 <sup>th</sup> (2001)	

<b>EMAMECTIN BENZOATE</b> (antiparasitic agent)				
<b>JECFA Evaluation:</b>		78 (2013)		
<b>ADI:</b>		ADI of 0–0.5 µg/kg body weight established by JMPR (2011), based on an overall NOAEL of 0.25 mg/kg body weight per day for neurotoxicity from 14- and 53-week studies in dogs, supported by an overall NOAEL of 0.25 mg/kg body weight per day from 1- and 2-year studies in rats. An uncertainty factor of 500 was applied to the NOAEL, which includes an additional uncertainty factor of 5 to account for the steep dose–response curve and irreversible histopathological effects in neural tissues at the LOAEL in dogs, as used by JMPR and confirmed by the current Committee (JECFA78, 2013).		
<b>EDI:</b>		11 µg/person per day, which represents approximately 37% of the upper bound of the ADI (JECFA78, 2013).		
<b>Residue Definition:</b>		Emamectin B1a.		
Species	Tissue	MRL (µg/kg)	CAC	Notes
Salmon	Muscle	100	38 <sup>th</sup> (2015)	
Salmon	Fillet	100	38 <sup>th</sup> (2015)	Muscle plus skin in natural proportion
Trout	Muscle	100	38 <sup>th</sup> (2015)	
Trout	Fillet	100	38 <sup>th</sup> (2015)	Muscle plus skin in natural proportion

<b>EPRINOMECTIN</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		50 (1998)		
<b>ADI:</b>		0-10 µg/kg body weight (JECFA50, 1998).		
<b>Residue Definition:</b>		Eprinomectin B1a.		
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	100	26 <sup>th</sup> (2003)	
Cattle	Liver	2000	26 <sup>th</sup> (2003)	
Cattle	Kidney	300	26 <sup>th</sup> (2003)	
Cattle	Fat	250	26 <sup>th</sup> (2003)	
Cattle	Milk (µg/l)	20	26 <sup>th</sup> (2003)	

<b>ERYTHROMYCIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		66 (2006)		
<b>ADI:</b>		0-0.7 µg/kg body weight (JECFA66, 2006)		
<b>Residue Definition:</b>		Erythromycin A		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Chicken	Muscle	100	31 <sup>st</sup> (2008)	
Chicken	Liver	100	31 <sup>st</sup> (2008)	
Chicken	Kidney	100	31 <sup>st</sup> (2008)	
Chicken	Fat	100	31 <sup>st</sup> (2008)	The MRL includes skin + fat
Chicken	Eggs	50	31 <sup>st</sup> (2008)	
Turkey	Muscle	100	31 <sup>st</sup> (2008)	
Turkey	Liver	100	31 <sup>st</sup> (2008)	
Turkey	Kidney	100	31 <sup>st</sup> (2008)	
Turkey	Fat	100	31 <sup>st</sup> (2008)	The MRL includes skin + fat

<b>ESTRADIOL-17BETA</b> (production aid)				
<b>JECFA Evaluation:</b>		25 (1981); 32 (1987); 52 (1999)		
<b>ADI:</b>		Unnecessary (JECFA32, 1987); 0-0.05 µg/kg body weight (JECFA52, 1999).		
<b>Residue Definition:</b>		Estradiol-17beta		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	Unnecessary	21 <sup>st</sup> (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health
Cattle	Liver	Unnecessary	21 <sup>st</sup> (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health
Cattle	Kidney	Unnecessary	21 <sup>st</sup> (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health
Cattle	Fat	unnecessary	21 <sup>st</sup> (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health

<b>FEBANTEL/FENBENDAZOLE/OXFENDAZOLE</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		38 (1991); 45 (1995); 50 (1998)		
<b>ADI:</b>		0-7 µg/kg body weight (JECFA50, 1998). Group ADI.		
<b>Residue Definition:</b>		Sum of fenbendazole, oxfendazole and oxfendazole sulphone, expressed as oxfendazole sulphone equivalents.		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	100	23 <sup>rd</sup> (1999)	
Cattle	Liver	500	23 <sup>rd</sup> (1999)	
Cattle	Kidney	100	23 <sup>rd</sup> (1999)	
Cattle	Fat	100	23 <sup>rd</sup> (1999)	
Cattle	Milk (µg/l)	100	23 <sup>rd</sup> (1999)	
Goat	Muscle	100	23 <sup>rd</sup> (1999)	
Goat	Liver	500	23 <sup>rd</sup> (1999)	
Goat	Kidney	100	23 <sup>rd</sup> (1999)	
Goat	Fat	100	23 <sup>rd</sup> (1999)	
Horse	Muscle	100	23 <sup>rd</sup> (1999)	
Horse	Liver	500	23 <sup>rd</sup> (1999)	
Horse	Kidney	100	23 <sup>rd</sup> (1999)	
Horse	Fat	100	23 <sup>rd</sup> (1999)	
Pig	Muscle	100	23 <sup>rd</sup> (1999)	
Pig	Liver	500	23 <sup>rd</sup> (1999)	
Pig	Kidney	100	23 <sup>rd</sup> (1999)	
Pig	Fat	100	23 <sup>rd</sup> (1999)	
Sheep	Muscle	100	23 <sup>rd</sup> (1999)	
Sheep	Liver	500	23 <sup>rd</sup> (1999)	
Sheep	Kidney	100	23 <sup>rd</sup> (1999)	
Sheep	Fat	100	23 <sup>rd</sup> (1999)	
Sheep	Milk (µg/l)	100	23 <sup>rd</sup> (1999)	

<b>FLUAZURON</b> (insecticide)				
<b>JECFA Evaluation:</b>		48 (1997)		
<b>ADI:</b>		0-40 µg/kg body weight (JECFA48, 1997)		
<b>Residue Definition:</b>		Fluazuron.		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	200	23 <sup>rd</sup> (1999)	
Cattle	Liver	500	23 <sup>rd</sup> (1999)	
Cattle	Kidney	500	23 <sup>rd</sup> (1999)	
Cattle	Fat	7000	23 <sup>rd</sup> (1999)	

<b>FLUBENDAZOLE</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		40 (1992)		
<b>ADI:</b>		0-12 µg/kg body weight (JECFA40, 1992)		
<b>Residue Definition:</b>		Flubendazole		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Pig	Muscle	10	21 <sup>st</sup> (1995)	
Pig	Liver	10	21 <sup>st</sup> (1995)	
Poultry	Muscle	200	21 <sup>st</sup> (1995)	
Poultry	Liver	500	21 <sup>st</sup> (1995)	
Poultry	Eggs	400	21 <sup>st</sup> (1995)	

<b>FLUMEQUINE</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		42 (1994); 48 (1997); 54 (2000); 60 (2002); 62 (2004); 66 (2006)		
<b>ADI:</b>		0-30 µg/kg body weight (JECFA62, 2004)		
<b>Residue Definition:</b>		Flumequine		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	500	28 <sup>th</sup> (2005)	
Cattle	Liver	500	28 <sup>th</sup> (2005)	
Cattle	Kidney	3000	28 <sup>th</sup> (2005)	
Cattle	Fat	1000	28 <sup>th</sup> (2005)	
Chicken	Muscle	500	28 <sup>th</sup> (2005)	
Chicken	Liver	500	28 <sup>th</sup> (2005)	
Chicken	Kidney	3000	28 <sup>th</sup> (2005)	
Chicken	Fat	1000	28 <sup>th</sup> (2005)	
Pig	Muscle	500	28 <sup>th</sup> (2005)	
Pig	Liver	500	28 <sup>th</sup> (2005)	
Pig	Kidney	3000	28 <sup>th</sup> (2005)	
Pig	Fat	1000	28 <sup>th</sup> (2005)	
Sheep	Muscle	500	28 <sup>th</sup> (2005)	
Sheep	Liver	500	28 <sup>th</sup> (2005)	
Sheep	Kidney	3000	28 <sup>th</sup> (2005)	
Sheep	Fat	1000	28 <sup>th</sup> (2005)	
Trout	Muscle	500	28 <sup>th</sup> (2005)	Muscle including normal proportion of skin

<b>GENTAMICIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		43 (1994); 48 (1997); 50 (1998)		
<b>ADI:</b>		0-20 µg/kg body weight (JECFA50, 1998)		
<b>Residue Definition:</b>		Gentamicin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	100	24 <sup>th</sup> (2001)	
Cattle	Liver	2000	24 <sup>th</sup> (2001)	
Cattle	Kidney	5000	24 <sup>th</sup> (2001)	
Cattle	Fat	100	24 <sup>th</sup> (2001)	
Cattle	Milk (µg/l)	200	24 <sup>th</sup> (2001)	
Pig	Muscle	100	24 <sup>th</sup> (2001)	
Pig	Liver	2000	24 <sup>th</sup> (2001)	
Pig	Kidney	5000	24 <sup>th</sup> (2001)	
Pig	Fat	100	24 <sup>th</sup> (2001)	

<b>IMIDOCARB</b> (antiprotozoal agent)				
<b>JECFA Evaluation:</b>		50 (1998); 60 (2003)		
<b>ADI:</b>		0-10 µg/kg body weight (JECFA50, 1998)		
<b>Residue Definition:</b>		Imidocarb		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	300	28 <sup>th</sup> (2005)	
Cattle	Liver	1500	28 <sup>th</sup> (2005)	
Cattle	Kidney	2000	28 <sup>th</sup> (2005)	
Cattle	Fat	50	28 <sup>th</sup> (2005)	
Cattle	Milk	50	28 <sup>th</sup> (2005)	

<b>ISOMETAMIDIUM</b> (trypanocide)				
<b>JECFA Evaluation:</b>		34 (1989); 40 (1992)		
<b>ADI:</b>		0-100 µg/kg body weight (JECFA40, 1992)		
<b>Residue Definition:</b>		Isometamidium		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	100	21 <sup>st</sup> (1995)	
Cattle	Liver	500	21 <sup>st</sup> (1995)	
Cattle	Kidney	1000	21 <sup>st</sup> (1995)	
Cattle	Fat	100	21 <sup>st</sup> (1995)	
Cattle	Milk (µg/l)	100	21 <sup>st</sup> (1995)	

<b>IVERMECTIN</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		36 (1990); 40 (1992); 54 (2000); 58 (2002); 81 (2015)		
<b>ADI:</b>		0-10 µg/kg body weight on the basis of a NOAEL of 0.5 mg/kg body weight per day for neurological effects (mydriasis) and retardation of weight gain in a 14-week dog study, with application of an uncertainty factor of 50 (5 for interspecies differences based on pharmacokinetic studies in dogs and humans and 10 for intraspecies differences). The previous ADI of 0-1 µg/kg body weight was withdrawn. (JECFA81, 2015).		
<b>ARfD:</b>		0.2 mg/kg body weight, based on a NOAEL of 1.5 mg/kg body weight, the highest dose tested in a safety, tolerability and pharmacokinetics study in healthy human subjects, with application of an uncertainty factor of 10 for intraspecies variability (JECFA81, 2015).		
<b>GECDE:</b>		<p>The EDI is 38 µg/person per day, based on a 60 kg individual, which represents 6% of the upper bound of the ADI.</p> <p>The GECDE for the general population is 0.9 µg/kg body weight per day, which represents 9% of the upper bound of the ADI.</p> <p>The GECDE for children is 1.5 µg/kg body weight per day, which represents 15% of the upper bound of the ADI.</p> <p>The GECDE for infants is 1.3 µg/kg body weight per day, which represents 13% of the upper bound of the ADI. (JECFA81, 2015)</p>		
<b>GEADE:</b>		A combined analysis of all studies submitted showed that after 14 days, the maximum values of residues found at injection sites led to a GEADE of 52 µg/kg bw for the general population and 87 µg/kg bw for children, corresponding, respectively, to 27% and 43% of the ARfD. (JECFA81, 2015)		
<b>Residue Definition:</b>		Ivermectin B <sub>1a</sub>		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	30	40 <sup>th</sup> (2017)	
Cattle	Liver	800	40 <sup>th</sup> (2017)	
Cattle	Kidney	100	40 <sup>th</sup> (2017)	
Cattle	Fat	400	40 <sup>th</sup> (2017)	
Cattle	Milk	10	26 <sup>th</sup> (2003)	
Pig	Liver	15	20 <sup>th</sup> (1993)	
Pig	Fat	20	20 <sup>th</sup> (1993)	
Sheep	Liver	15	20 <sup>th</sup> (1993)	
Sheep	Fat	20	20 <sup>th</sup> (1993)	

<b>LASALOCID SODIUM</b> (antiparasitic agent)				
<b>JECFA Evaluation:</b>		78 (2013)		
<b>ADI:</b>		0-5 µg/kg body weight on the basis of a NOAEL of 0.5 mg/kg body weight per day from a developmental toxicity study in rabbits and a multigeneration reproductive toxicity study in rats, with application of an uncertainty factor of 100 for interspecies and intraspecies variability (JECFA78, 2013).		
<b>EDI:</b>		80 µg/person per day was calculated, which represents approximately 27% of the upper bound of the ADI (JECFA78, 2013).		
<b>Residue Definition:</b>		Lasalocid A		
<b>Note:</b>		JECFA78 extended the MRLs in chicken to turkey and quail and extrapolated the MRLs in chicken to pheasant. No information was available for duck, including on approved uses. As the compound is not registered for use in laying hens, according to the sponsor, it is not appropriate to recommend MRLs for egg.		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Chicken	Muscle	400	40 <sup>th</sup> (2017)	
Chicken	Liver	1200	40 <sup>th</sup> (2017)	
Chicken	Kidney	600	40 <sup>th</sup> (2017)	
Chicken	Skin + Fat	600	40 <sup>th</sup> (2017)	
Turkey	Muscle	400	40 <sup>th</sup> (2017)	
Turkey	Liver	1200	40 <sup>th</sup> (2017)	
Turkey	Kidney	600	40 <sup>th</sup> (2017)	
Turkey	Skin + Fat	600	40 <sup>th</sup> (2017)	
Quail	Muscle	400	40 <sup>th</sup> (2017)	
Quail	Liver	1200	40 <sup>th</sup> (2017)	
Quail	Kidney	600	40 <sup>th</sup> (2017)	
Quail	Skin + Fat	600	40 <sup>th</sup> (2017)	
Pheasant	Muscle	400	40 <sup>th</sup> (2017)	
Pheasant	Liver	1200	40 <sup>th</sup> (2017)	
Pheasant	Kidney	600	40 <sup>th</sup> (2017)	
Pheasant	Skin + Fat	600	40 <sup>th</sup> (2017)	



<b>LEVAMISOLE</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		36 (1990); 42 (1994)		
<b>ADI:</b>		0-6 µg/kg body weight (JECFA42, 1994)		
<b>Residue Definition:</b>		Levamisole		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	10	22 <sup>nd</sup> (1997)	
Cattle	Liver	100	22 <sup>nd</sup> (1997)	
Cattle	Kidney	10	22 <sup>nd</sup> (1997)	
Cattle	Fat	10	22 <sup>nd</sup> (1997)	
Pig	Muscle	10	22 <sup>nd</sup> (1997)	
Pig	Liver	100	22 <sup>nd</sup> (1997)	
Pig	Kidney	10	22 <sup>nd</sup> (1997)	
Pig	Fat	10	22 <sup>nd</sup> (1997)	
Poultry	Muscle	10	22 <sup>nd</sup> (1997)	
Poultry	Liver	100	22 <sup>nd</sup> (1997)	
Poultry	Kidney	10	22 <sup>nd</sup> (1997)	
Poultry	Fat	10	22 <sup>nd</sup> (1997)	
Sheep	Muscle	10	22 <sup>nd</sup> (1997)	
Sheep	Liver	100	22 <sup>nd</sup> (1997)	
Sheep	Kidney	10	22 <sup>nd</sup> (1997)	
Sheep	Fat	10	22 <sup>nd</sup> (1997)	

<b>LINCOMYCIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		54 (2000); 58 (2002); 62 (2004)		
<b>ADI:</b>		0-30 µg/kg body weight (JECFA54, 2000)		
<b>Residue Definition:</b>		Lincomycin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Milk	150	26 <sup>th</sup> (2003)	
Chicken	Muscle	200	26 <sup>th</sup> (2003)	
Chicken	Liver	500	26 <sup>th</sup> (2003)	
Chicken	Kidney	500	26 <sup>th</sup> (2003)	
Chicken	Fat	100	26 <sup>th</sup> (2003)	Additional MRL for skin with adhering fat of 300 µg/kg
Pig	Muscle	200	26 <sup>th</sup> (2003)	
Pig	Liver	500	26 <sup>th</sup> (2003)	
Pig	Kidney	1500	26 <sup>th</sup> (2003)	
Pig	Fat	100	26 <sup>th</sup> (2003)	Additional MRL for skin with adhering fat of 300 µg/kg

<b>LUNEFURON</b> (insecticide)				
<b>JECFA Evaluation:</b>		85 (2017)		
<b>ADI:</b>		0–0.02 mg/kg bw based on the NOAEL of 1.93 mg/kg bw per day for tonic-clonic seizures and findings in lungs, gastrointestinal tract, liver and urinary tract in a 2-year dietary study in rats, and using a safety factor of 100 (10 for interspecies variability and 10 for intraspecies variability) (JECFA85, 2017).		
<b>ARfD:</b>		Unnecessary, in view of lufenuron low acute oral toxicity and the absence of developmental toxicity and other toxicological effects likely to be elicited by a single dose (JECFA85, 2017)		
<b>GECDE:</b>		1.1 µg/kg bw per day (for the general population), which represents 5.5% of the upper bound of the ADI. As lufenuron is also used as pesticide the overall dietary exposure was estimated. The assumptions and detailed results will be displayed in the JECFA85 report. Results below are only for use as veterinary drug (JECFA85, 2017).		
<b>Residue Definition:</b>		Lunefuron		
Species	Tissue	MRL (µg/kg)	CAC	Notes
Salmon	Fillet <sup>a</sup>	1350	41 <sup>st</sup> (2018)	
Trout	Fillet <sup>a</sup>	1350	41 <sup>st</sup> (2018)	

<sup>a</sup> Muscle plus skin in natural proportion.

<b>MELENGESTROL ACETATE</b> (production aid)				
<b>JECFA Evaluation:</b>		54 (2000); 58 (2002); 62 (2004); 66 (2006) 70 (2008)		
<b>ADI:</b>		0-0.03 µg/kg body weight (JECFA54, 2000)		
<b>Residue Definition:</b>		Melengestrol acetate		
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	1	32 <sup>nd</sup> (2009)	
Cattle	Liver	10	32 <sup>nd</sup> (2009)	
Cattle	Kidney	2	32 <sup>nd</sup> (2009)	
Cattle	Fat	18	32 <sup>nd</sup> (2009)	

<b>MONENSIN (antimicrobial agent)</b>				
<b>JECFA Evaluation:</b>		70 (2008); 75 (2011)		
<b>ADI:</b>		0–10 µg/kg body weight on the basis of a NOAEL of 1.14 mg/kg body weight per day and a safety factor of 100 and rounding to one significant figure (JECFA70, 2008).		
<b>Estimated Dietary Exposure:</b>		Using the revised MRL, the TMDI from JECFA70 was recalculated, resulting in a value of 481 µg/person, which represents 80% of the upper bound of the ADI (JECFA75, 2011).		
<b>Residue Definition:</b>		Monensin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	10	32 <sup>nd</sup> (2009)	
Cattle	Liver	100	35 <sup>th</sup> (2012)	
Cattle	Kidney	10	32 <sup>nd</sup> (2009)	
Cattle	Fat	100	32 <sup>nd</sup> (2009)	
Cattle	Milk	2	32 <sup>nd</sup> (2009)	
Sheep	Muscle	10	32 <sup>nd</sup> (2009)	
Sheep	Liver	20	32 <sup>nd</sup> (2009)	
Sheep	Kidney	10	32 <sup>nd</sup> (2009)	
Sheep	Fat	100	32 <sup>nd</sup> (2009)	
Goats	Muscle	10	32 <sup>nd</sup> (2009)	
Goats	Liver	20	32 <sup>nd</sup> (2009)	
Goats	Kidney	10	32 <sup>nd</sup> (2009)	
Goats	Fat	100	32 <sup>nd</sup> (2009)	
Chicken	Muscle	10	32 <sup>nd</sup> (2009)	
Chicken	Liver	10	32 <sup>nd</sup> (2009)	
Chicken	Kidney	10	32 <sup>nd</sup> (2009)	
Chicken	Fat	100	32 <sup>nd</sup> (2009)	
Turkey	Muscle	10	32 <sup>nd</sup> (2009)	
Turkey	Liver	10	32 <sup>nd</sup> (2009)	
Turkey	Kidney	10	32 <sup>nd</sup> (2009)	
Turkey	Fat	100	32 <sup>nd</sup> (2009)	
Quail	Muscle	10	32 <sup>nd</sup> (2009)	
Quail	Liver	10	32 <sup>nd</sup> (2009)	
Quail	Kidney	10	32 <sup>nd</sup> (2009)	
Quail	Fat	100	32 <sup>nd</sup> (2009)	

<b>MONEPANTEL</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		75 (2011); 78 (2013); 85 <sup>th</sup> (2017)		
<b>ADI:</b>		0–0.02 mg/kg bw based on the NOAEL of 1.93 mg/kg bw per day for tonic-clonic seizures and findings in lungs, gastrointestinal tract, liver and urinary tract in a 2-year dietary study in rats, and using a safety factor of 100 (10 for interspecies variability and 10 for intraspecies variability) (JECFA85, 2017).		
<b>ARfD:</b>		Unnecessary		
<b>Estimated Dietary Exposure:</b>		Using the model diet and marker residue to total residue ratios of 1.00 for muscle and 0.66 for fat, liver and kidney, and applying a correction factor of 0.94 to account for the mass difference between monepantel sulfone (the marker residue) and monepantel, the EDI is 446 µg/person per day, which represents approximately 37% of the upper bound of the ADI (JECFA78, 2013).		
<b>GECDE:</b>		13.7 µg per kg bw per day (for the general population), which represents 68% of the upper bound of the ADI (JECFA85, 2017). 5.0 µg per kg bw per day (for children), which represents 22% of the upper bound of the ADI (JECFA85, 2017). 4.4 µg per kg bw per day (for infants), which represents 25% of the upper bound of the ADI (JECFA85, 2017).		
<b>Residue Definition:</b>		Monepantel sulfone, expressed as monepantel.		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	300	41 <sup>st</sup> (2018)	
Cattle	Liver	2000	41 <sup>st</sup> (2018)	
Cattle	Kidney	1000	41 <sup>st</sup> (2018)	
Cattle	Fat	7000	41 <sup>st</sup> (2018)	
Sheep	Muscle	500	38 <sup>th</sup> (2015)	
Sheep	Liver	7000	38 <sup>th</sup> (2015)	
Sheep	Kidney	1700	38 <sup>th</sup> (2015)	
Sheep	Fat	13000	38 <sup>th</sup> (2015)	

<b>MOXIDECTIN</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		45 (1995); 47 (1996); 48 (1998); 50 (1998)		
<b>ADI:</b>		0-2 µg/kg body weight (JECFA45, 1995)		
<b>Residue Definition:</b>		Moxidectin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	20	22 <sup>nd</sup> (1997)	Very high concentration and great variation in the level of residues at the injection site in cattle over a 49 day period after dosing.
Cattle	Liver	100	22 <sup>nd</sup> (1997)	
Cattle	Kidney	50	22 <sup>nd</sup> (1997)	
Cattle	Fat	500	22 <sup>nd</sup> (1997)	
Deer	Muscle	20	23 <sup>rd</sup> (1999)	
Deer	Liver	100	23 <sup>rd</sup> (1999)	
Deer	Kidney	50	23 <sup>rd</sup> (1999)	
Deer	Fat	500	23 <sup>rd</sup> (1999)	
Sheep	Muscle	50	22 <sup>nd</sup> (1997)	
Sheep	Liver	100	22 <sup>nd</sup> (1997)	
Sheep	Kidney	50	22 <sup>nd</sup> (1997)	
Sheep	Fat	500	22 <sup>nd</sup> (1997)	

<b>NARASIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		70 (2008); 75 (2011)		
<b>ADI:</b>		0-5 µg/kg body weight on the basis of a NOAEL of 0.5 mg/kg body weight per day and a safety factor of 100 (JECFA70, 2008)		
<b>Residue Definition:</b>		Narasin A		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	15	35 <sup>th</sup> (2012)	
Cattle	Liver	50	35 <sup>th</sup> (2012)	
Cattle	Kidney	15	35 <sup>th</sup> (2012)	
Cattle	Fat	50	35 <sup>th</sup> (2012)	
Chicken	Muscle	15	32 <sup>nd</sup> (2009)	
Chicken	Liver	50	32 <sup>nd</sup> (2009)	
Chicken	Kidney	15	32 <sup>nd</sup> (2009)	
Chicken	Fat	50	32 <sup>nd</sup> (2009)	
Pig	Muscle	15	34 <sup>th</sup> (2011)	
Pig	Liver	50	34 <sup>th</sup> (2011)	
Pig	Kidney	15	34 <sup>th</sup> (2011)	
Pig	Fat	50	34 <sup>th</sup> (2011)	

<b>NEOMYCIN (antimicrobial agent)</b>				
<b>JECFA Evaluation:</b>		43 (1994); 47 (1996); 52 (1999); 58 (2002); 60 (2003)		
<b>ADI:</b>		0-60 µg/kg body weight (JECFA47, 1996)		
<b>Residue Definition:</b>		Neomycin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	500	23 <sup>rd</sup> (1999)	
Cattle	Liver	500	28 <sup>th</sup> (2005)	
Cattle	Kidney	10000	28 <sup>th</sup> (2005)	
Cattle	Fat	500	23 <sup>rd</sup> (1999)	
Cattle	Milk	1500	28 <sup>th</sup> (2005)	
Chicken	Muscle	500	23 <sup>rd</sup> (1999)	
Chicken	Liver	500	23 <sup>rd</sup> (1999)	
Chicken	Kidney	10000	23 <sup>rd</sup> (1999)	
Chicken	Fat	500	23 <sup>rd</sup> (1999)	
Chicken	Eggs	500	23 <sup>rd</sup> (1999)	
Duck	Muscle	500	23 <sup>rd</sup> (1999)	
Duck	Liver	500	23 <sup>rd</sup> (1999)	
Duck	Kidney	10000	23 <sup>rd</sup> (1999)	
Duck	Fat	500	23 <sup>rd</sup> (1999)	
Goat	Muscle	500	23 <sup>rd</sup> (1999)	
Goat	Liver	500	23 <sup>rd</sup> (1999)	
Goat	Kidney	10000	23 <sup>rd</sup> (1999)	
Goat	Fat	500	23 <sup>rd</sup> (1999)	
Pig	Muscle	500	23 <sup>rd</sup> (1999)	
Pig	Liver	500	23 <sup>rd</sup> (1999)	
Pig	Kidney	10000	23 <sup>rd</sup> (1999)	
Pig	Fat	500	23 <sup>rd</sup> (1999)	
Sheep	Muscle	500	23 <sup>rd</sup> (1999)	
Sheep	Liver	500	23 <sup>rd</sup> (1999)	
Sheep	Kidney	10000	23 <sup>rd</sup> (1999)	
Sheep	Fat	500	23 <sup>rd</sup> (1999)	
Turkey	Muscle	500	23 <sup>rd</sup> (1999)	
Turkey	Liver	500	23 <sup>rd</sup> (1999)	
Turkey	Kidney	10000	23 <sup>rd</sup> (1999)	
Turkey	Fat	500	23 <sup>rd</sup> (1999)	

<b>NICARBAZIN</b> (antiprotozoal agent)				
<b>JECFA Evaluation:</b>		50 (1998)		
<b>ADI:</b>		0-400 µg/kg body weight (JECFA50, 1998)		
<b>Residue Definition:</b>		N,N'-bis(4-nitrophenyl)urea		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Chicken	Muscle	200	23 <sup>rd</sup> (1999)	Broilers
Chicken	Liver	200	23 <sup>rd</sup> (1999)	Broilers
Chicken	Kidney	200	23 <sup>rd</sup> (1999)	Broilers
Chicken	Fat/Skin	200	23 <sup>rd</sup> (1999)	Broilers

<b>PHOXIM</b> (insecticide)				
<b>JECFA Evaluation:</b>		52 (1999); 62 (2004)		
<b>ADI:</b>		0-4 µg/kg body weight (JECFA52, 1999)		
<b>Residue Definition:</b>		Phoxim		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Goat	Muscle	50	26 <sup>th</sup> (2003)	
Goat	Liver	50	26 <sup>th</sup> (2003)	
Goat	Kidney	50	26 <sup>th</sup> (2003)	
Goat	Fat	400	26 <sup>th</sup> (2003)	
Pig	Muscle	50	26 <sup>th</sup> (2003)	
Pig	Liver	50	26 <sup>th</sup> (2003)	
Pig	Kidney	50	26 <sup>th</sup> (2003)	
Pig	Fat	400	26 <sup>th</sup> (2003)	
Sheep	Muscle	50	26 <sup>th</sup> (2003)	
Sheep	Liver	50	26 <sup>th</sup> (2003)	
Sheep	Kidney	50	26 <sup>th</sup> (2003)	
Sheep	Fat	400	26 <sup>th</sup> (2003)	

<b>PIRLIMYCIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		62 (2004)		
<b>ADI:</b>		0-8 µg/kg bw (JECFA62, 2004)		
<b>Residue Definition:</b>		Pirlimycin		
<b>Species</b>	<b>Tissue</b>	<b>MRLs (µg/kg)</b>	<b>CAC</b>	<b>Note</b>
Cattle	Muscle	100	29 <sup>th</sup> (2006)	
Cattle	Liver	1000	29 <sup>th</sup> (2006)	
Cattle	Kidney	400	29 <sup>th</sup> (2006)	
Cattle	Fat	100	29 <sup>th</sup> (2006)	
Cattle	Milk	100	29 <sup>th</sup> (2006)	JECFA evaluated the effect of pirlimycin residues on starter cultures and for this reason recommended an MRL of 100 µg/kg of milk. Codex Members may therefore adapt national/regional MRLs in order to address this technological aspect for trade of fresh liquid milk intended for processing using starter culture.

<b>PORCINE SOMATOTROPIN</b> (production aid)				
<b>JECFA Evaluation:</b>		52 (1999)		
<b>ADI:</b>		Not Specified (JECFA52, 1999)		
<b>Residue Definition:</b>		Not applicable		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Pig	Muscle	not specified	26 <sup>th</sup> (2003)	
Pig	Liver	not specified	26 <sup>th</sup> (2003)	
Pig	Kidney	not specified	26 <sup>th</sup> (2003)	
Pig	Fat	not specified	26 <sup>th</sup> (2003)	



<b>PROGESTERONE</b> (production aid)				
<b>JECFA Evaluation:</b>		25 (1981); 32 (1987); 52 (1999)		
<b>ADI:</b>		0-30 µg/kg body weight (JECFA52, 1999)		
<b>Residue Definition:</b>		Progesterone		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	Unnecessary	21 <sup>st</sup> (2005)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Liver	Unnecessary	21 <sup>st</sup> (2005)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Kidney	Unnecessary	21 <sup>st</sup> (2005)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Fat	Unnecessary	21 <sup>st</sup> (2005)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.

<b>RACTOPAMINE</b> (production aid)				
<b>JECFA Evaluation:</b>		40 (1992); 62 (2004); 66 (2006)		
<b>ADI:</b>		0-1 µg/kg body weight (JECFA66, 2006)		
<b>Residue Definition:</b>		Ractopamine		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	10	35 <sup>th</sup> (2012)	
Cattle	Liver	40	35 <sup>th</sup> (2012)	
Cattle	Kidney	90	35 <sup>th</sup> (2012)	
Cattle	Fat	10	35 <sup>th</sup> (2012)	
Pig	Muscle	10	35 <sup>th</sup> (2012)	
Pig	Liver	40	35 <sup>th</sup> (2012)	
Pig	Kidney	90	35 <sup>th</sup> (2012)	
Pig	Fat	10	35 <sup>th</sup> (2012)	The MRL includes skin + fat

<b>SARAFLOXACIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		50 (1998)		
<b>ADI:</b>		0-0.3 µg/kg body weight (JECFA50, 1998)		
<b>Residue Definition:</b>		Sarafloxacin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Chicken	Muscle	10	24 <sup>th</sup> (2001)	
Chicken	Liver	80	24 <sup>th</sup> (2001)	
Chicken	Kidney	80	24 <sup>th</sup> (2001)	
Chicken	Fat	20	24 <sup>th</sup> (2001)	
Turkey	Muscle	10	24 <sup>th</sup> (2001)	
Turkey	Liver	80	24 <sup>th</sup> (2001)	
Turkey	Kidney	80	24 <sup>th</sup> (2001)	
Turkey	Fat	20	24 <sup>th</sup> (2001)	

<b>SPECTINOMYCIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		42 (1994); 50 (1998)		
<b>ADI:</b>		0-40 µg/kg body weight (JECFA42, 1994)		
<b>Residue Definition:</b>		Spectinomycin		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	500	23 <sup>rd</sup> (1999)	
Cattle	Liver	2000	23 <sup>rd</sup> (1999)	
Cattle	Kidney	5000	23 <sup>rd</sup> (1999)	
Cattle	Fat	2000	23 <sup>rd</sup> (1999)	
Cattle	Milk (µg/l)	200	23 <sup>rd</sup> (1999)	
Chicken	Muscle	500	23 <sup>rd</sup> (1999)	
Chicken	Liver	2000	23 <sup>rd</sup> (1999)	
Chicken	Kidney	5000	23 <sup>rd</sup> (1999)	
Chicken	Fat	2000	23 <sup>rd</sup> (1999)	
Chicken	Eggs	2000	23 <sup>rd</sup> (1999)	
Pig	Muscle	500	23 <sup>rd</sup> (1999)	
Pig	Liver	2000	23 <sup>rd</sup> (1999)	
Pig	Kidney	5000	23 <sup>rd</sup> (1999)	
Pig	Fat	2000	23 <sup>rd</sup> (1999)	
Sheep	Muscle	500	23 <sup>rd</sup> (1999)	
Sheep	Liver	2000	23 <sup>rd</sup> (1999)	
Sheep	Kidney	5000	23 <sup>rd</sup> (1999)	
Sheep	Fat	2000	23 <sup>rd</sup> (1999)	

<b>SPIRAMYCIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		38 (1991); 43 (1994); 47 (1996); 48 (1997)		
<b>ADI:</b>		0-50 µg/kg body weight (JECFA43, 1994)		
<b>Residue Definition:</b>		Cattle and chickens, sum of spiramycin and neospiramycin; Pigs, spiramycin equivalents (antimicrobially active residues).		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	200	22 <sup>nd</sup> (1997)	
Cattle	Liver	600	22 <sup>nd</sup> (1997)	
Cattle	Kidney	300	22 <sup>nd</sup> (1997)	
Cattle	Fat	300	22 <sup>nd</sup> (1997)	
Cattle	Milk (µg/l)	200	22 <sup>nd</sup> (1997)	
Chicken	Muscle	200	22 <sup>nd</sup> (1997)	
Chicken	Liver	600	22 <sup>nd</sup> (1997)	
Chicken	Kidney	800	22 <sup>nd</sup> (1997)	
Chicken	Fat	300	22 <sup>nd</sup> (1997)	
Pig	Muscle	200	22 <sup>nd</sup> (1997)	
Pig	Liver	600	22 <sup>nd</sup> (1997)	
Pig	Kidney	300	22 <sup>nd</sup> (1997)	
Pig	Fat	300	22 <sup>nd</sup> (1997)	

<b>SULFADIMIDINE</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		34 (1989); 38 (1991); 42 (1994)		
<b>ADI:</b>		0-50 µg/kg body weight (JECFA42, 1994)		
<b>Residue Definition:</b>		Sulfadimidine		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Milk (µg/l)	25	21 <sup>st</sup> (1995)	
Not specified	Muscle	100	21 <sup>st</sup> (1995)	
Not specified	Liver	100	21 <sup>st</sup> (1995)	
Not specified	Kidney	100	21 <sup>st</sup> (1995)	
Not specified	Fat	100	21 <sup>st</sup> (1995)	

<b>TEFLUBENZURON</b> (insecticide)				
<b>JECFA Evaluation:</b>		81 (2015)		
<b>ADI:</b>		0-5 µg/kg body weight on the basis of a lower 95% confidence limit on the BMDL10 of 0.54 mg/kg body weight per day for hepatocellular hypertrophy in male mice observed in a carcinogenicity study, with application of an uncertainty factor of 100 to account for interspecies and intraspecies variability (JECFA81, 2015).		
<b>GECDE:</b>		The EDI is 42.9 µg/person per day, on the basis of a 60 kg individual, which represents approximately 14% of the upper bound of the ADI. The GECDE for the general population is 1.6 µg/kg body weight per day, which represents 31% of the upper bound of the ADI. The GECDE for children is 2.1 µg/kg body weight per day, which represents 43% of the upper bound of the ADI. The GECDE for infants is 0.9 µg/kg body weight per day, which represents 18% of the upper bound of the ADI. (JECFA81, 2015)		
<b>Residue Definition:</b>		Teflubenzuron		
Species	Tissue	MRL (µg/kg)	CAC	Notes
Salmon	Muscle	400	40 <sup>th</sup> (2017)	
Salmon	Fillet	400	40 <sup>th</sup> (2017)	Muscle plus skin in natural proportion

<b>TESTOSTERONE</b> (production aid)				
<b>JECFA Evaluation:</b>		25 (1981); 32 (1987); 52 (1999)		
<b>ADI:</b>		0-2 µg/kg body weight (JECFA52, 1999)		
<b>Residue Definition:</b>		Testosterone		
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	Unnecessary	21 <sup>st</sup> (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Liver	Unnecessary	21 <sup>st</sup> (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Kidney	Unnecessary	21 <sup>st</sup> (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.
Cattle	Fat	Unnecessary	21 <sup>st</sup> (1995)	Residues resulting from the use of this substances as a growth promoter in accordance with good animal husbandry practice are unlikely to pose a hazard to human health.

<b>THIABENDAZOLE</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		40 (1992); 48 (1997); 58 (2002)		
<b>ADI:</b>		0-100 µg/kg body weight (JECFA40, 1992)		
<b>Residue Definition:</b>		Sum of thiabendazole and 5-hydroxythiabendazole		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Cattle	Liver	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Cattle	Kidney	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Cattle	Fat	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Cattle	Milk (µg/l)	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Goat	Muscle	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Goat	Liver	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Goat	Kidney	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Goat	Fat	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Goat	Milk (µg/l)	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Pig	Muscle	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Pig	Liver	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Pig	Kidney	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Pig	Fat	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Sheep	Muscle	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.

Species	Tissue	MRL (µg/kg)	CAC	Notes
Sheep	Liver	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Sheep	Kidney	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.
Sheep	Fat	100	21 <sup>st</sup> (1995)	The MRL also covers residues derived from feed containing the residues resulted from agricultural use.

TILMICOSIN (antimicrobial agent)				
JECFA Evaluation:		47 (1996); 54 (2000); 70 (2008)		
ADI:		0-40 µg/kg body weight (JECFA47, 1996)		
Residue Definition:		Tilmicosin		
Species	Tissue	MRL (µg/kg)	CAC	Notes
Cattle	Muscle	100	23 <sup>rd</sup> (1999)	
Cattle	Liver	1000	23 <sup>rd</sup> (1999)	
Cattle	Kidney	300	23 <sup>rd</sup> (1999)	
Cattle	Fat	100	23 <sup>rd</sup> (1999)	
Chicken	Muscle	150	34 <sup>th</sup> (2011)	
Chicken	Liver	2400	34 <sup>th</sup> (2011)	
Chicken	Kidney	600	34 <sup>th</sup> (2011)	
Chicken	Skin/Fat	250	34 <sup>th</sup> (2011)	
Pig	Muscle	100	23 <sup>rd</sup> (1999)	
Pig	Liver	1500	23 <sup>rd</sup> (1999)	
Pig	Kidney	1000	23 <sup>rd</sup> (1999)	
Pig	Fat	100	23 <sup>rd</sup> (1999)	
Sheep	Muscle	100	23 <sup>rd</sup> (1999)	
Sheep	Liver	1000	23 <sup>rd</sup> (1999)	
Sheep	Kidney	300	23 <sup>rd</sup> (1999)	
Sheep	Fat	100	23 <sup>rd</sup> (1999)	
Turkey	Muscle	100	34 <sup>th</sup> (2011)	
Turkey	Kidney	1200	34 <sup>th</sup> (2011)	
Turkey	Liver	1400	34 <sup>th</sup> (2011)	
Turkey	Skin/Fat	250	34 <sup>th</sup> (2011)	

<b>TRENBOLONE ACETATE</b> (growth promoter)				
<b>JECFA Evaluation:</b>		26 (1982); 27 (1983); 32 (1987); 34 (1989)		
<b>ADI:</b>		0-0.02 µg/kg body weight (JECFA34, 1989)		
<b>Residue Definition:</b>		Cattle muscle, beta-Trenbolone; Cattle liver, alpha-Trenbolone.		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	2	21 <sup>st</sup> (1995)	
Cattle	Liver	10	21 <sup>st</sup> (1995)	

<b>TRICHLORFON (Metrifonate)</b> (insecticide)				
<b>JECFA Evaluation:</b>		54 (2000); 60 (2003); 66 (2006)		
<b>ADI:</b>		0-2 µg/kg bw (JECFA60, 2003)		
<b>Residue Definition:</b>		JECFA confirmed the MRL for cows's milk and the guidance levels for muscle, liver, kidney and fat of cattle recommended at the 54 <sup>th</sup> meeting (WHO TRS 900, 2001)		
<b>Species</b>	<b>Tissue</b>	<b>MRLs (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Milk	50	29 <sup>th</sup> (2006)	

<b>TRICLABENDAZOLE</b> (anthelmintic agent)				
<b>JECFA Evaluation:</b>		40 (1992); 66 (2006); 70 (2008)		
<b>ADI:</b>		0-3 µg/kg body weight (JECFA40, 1993).		
<b>Residue Definition:</b>		Ketotriclabnedazole		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	250	32 <sup>nd</sup> (2009)	
Cattle	Liver	850	32 <sup>nd</sup> (2009)	
Cattle	Kidney	400	32 <sup>nd</sup> (2009)	
Cattle	Fat	100	32 <sup>nd</sup> (2009)	
Sheep	Muscle	200	32 <sup>nd</sup> (2009)	
Sheep	Liver	300	32 <sup>nd</sup> (2009)	
Sheep	Kidney	200	32 <sup>nd</sup> (2009)	
Sheep	Fat	100	32 <sup>nd</sup> (2009)	

<b>TYLOSIN</b> (antimicrobial agent)				
<b>JECFA Evaluation:</b>		70 (2008)		
<b>ADI:</b>		0-30 µg/kg body weight based on a microbiological end-point derived from in vitro MIC susceptibility testing and faecal binding data (MICcalc = 1.698) (JECFA70, 2008).		
<b>Residue Definition:</b>		Tylosin A		
<b>Species</b>	<b>Tissue</b>	<b>MRLs (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	100	32 <sup>nd</sup> (2009)	
Cattle	Liver	100	32 <sup>nd</sup> (2009)	
Cattle	Kidney	100	32 <sup>nd</sup> (2009)	
Cattle	Fat	100	32 <sup>nd</sup> (2009)	
Cattle	Milk	100	32 <sup>nd</sup> (2009)	
Pig	Muscle	100	32 <sup>nd</sup> (2009)	
Pig	Liver	100	32 <sup>nd</sup> (2009)	
Pig	Kidney	100	32 <sup>nd</sup> (2009)	
Pig	Fat	100	32 <sup>nd</sup> (2009)	
Chicken	Muscle	100	32 <sup>nd</sup> (2009)	
Chicken	Liver	100	32 <sup>nd</sup> (2009)	
Chicken	Kidney	100	32 <sup>nd</sup> (2009)	
Chicken	Fat/Skin	100	32 <sup>nd</sup> (2009)	
Chicken	Eggs	300	32 <sup>nd</sup> (2009)	

<b>ZERANOL</b> (growth promoter)				
<b>JECFA Evaluation:</b>		26 (1982); 27 (1983); 32 (1987)		
<b>ADI:</b>		0-0.5 µg/kg body weight (JECFA32, 1987)		
<b>Residue Definition:</b>		Zeranol		
<b>Species</b>	<b>Tissue</b>	<b>MRL (µg/kg)</b>	<b>CAC</b>	<b>Notes</b>
Cattle	Muscle	2	21 <sup>st</sup> (1995)	
Cattle	Liver	10	21 <sup>st</sup> (1995)	



**B) RISK MANAGEMENT RECOMMENDATIONS (RMRs) FOR RESIDUES OF VETERINARY DRUGS****CARBADOX** (growth promoter)

---

**JECFA evaluation:** 36<sup>th</sup> (1990) and 60<sup>th</sup> (2003)**CAC37** (2014)**Recommended risk management measures**

In view of the JECFA conclusions on the available scientific information, there is no safe level of residues of carbadox or its metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of carbadox in food. This can be accomplished by not using carbadox in food producing animals.

**CHLORAMPHENICOL** (antimicrobial agent)

---

**JECFA evaluation:** 12<sup>th</sup> (1968), 32<sup>nd</sup> (1987), 42<sup>nd</sup> (1994) and 62<sup>nd</sup> (2004)**CAC37** (2014)**Recommended risk management measures**

In view of the JECFA conclusions on the available scientific information, there is no safe level of residues of chloramphenicol or its metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of chloramphenicol in food. This can be accomplished by not using chloramphenicol in food producing animals.

**CHLORPROMAZINE** (tranquilliser agent)

---

**JECFA evaluation:** 38<sup>th</sup> (1991)**CAC37** (2014)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of chlorpromazine or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of chlorpromazine in food. This can be accomplished by not using chlorpromazine in food producing animals.

**DIMETRIDAZOLE** (antiprotozoal agent)

---

**JECFA evaluation:** 34<sup>th</sup> (1989)**CAC38** (2015)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of dimetridazole or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of dimetridazole in food. This can be accomplished by not using dimetridazole in food producing animals.

**FURAZOLIDONE** (antimicrobial agent)

---

**JECFA evaluation:** 40<sup>th</sup> (1992)**CAC37** (2014)**Recommended risk management measures**

In view of the JECFA conclusions on the available scientific information, there is no safe level of residues of furazolidone or its metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of furazolidone in food. This can be accomplished by not using furazolidone in food producing animals.

**GENTIAN VIOLET** (antibacterial, antifungal and anthelmintic agent)

---

**JECFA evaluation:** 78<sup>th</sup> (2013)**CAC41** (2018)**Recommended risk management measures**

In view of the JECFA conclusions on the available scientific information, there is no safe level of residues of gentian violet or its metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of gentian violet in food. This can be accomplished by not using gentian violet in food producing animals.

---

**IPRONIDAZOLE** (antiprotozoal agent)

---

**JECFA evaluation:** 34<sup>th</sup> (1989)**CAC38** (2015)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of ipronidazole or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of ipronidazole in food. This can be accomplished by not using ipronidazole in food producing animals.

---

**MALACHITE GREEN** (antifungal and antiprotozoal agent)

---

**JECFA evaluation:** 70<sup>th</sup> (2008)**CAC37** (2014)**Recommended risk management measures**

In view of the JECFA conclusions on the available scientific information, there is no safe level of residues of malachite green or its metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of malachite green in food. This can be accomplished by not using malachite green in food producing animals.

---

**METRONIDAZOLE** (antiprotozoal agent)

---

**JECFA evaluation:** 34<sup>th</sup> (1989)**CAC38** (2015)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of metronidazole or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of metronidazole in food. This can be accomplished by not using metronidazole in food producing animals.

---

**NITROFURAL** (antimicrobial agent)

---

**JECFA evaluation:** 40<sup>th</sup> (1992)**CAC37** (2014)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of nitrofurazone or its metabolites<sup>1</sup> in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of nitrofurazone in food. This can be accomplished by not using nitrofurazone in food producing animals.

<sup>1</sup> Semicarbazide is not a unique indicator of nitrofurazone use and low levels can be associated with other legitimate sources.

---

**OLAQUINDOX** (antibacterial agent)

---

**JECFA evaluation:** 36<sup>th</sup> (1990) and 42<sup>nd</sup> (1994)**CAC37** (2014)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of olaquinoxid or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of olaquinoxid in food. This can be accomplished by not using olaquinoxid in food producing animals.

---

**RONIDAZOLE** (antiprotozoal agent)

---

**JECFA evaluation:** 34<sup>th</sup> (1989) and 42<sup>nd</sup> (1994)**CAC38** (2015)**Recommended risk management measures**

In view of the JECFA conclusions, although insufficient data were available or there was a lack of data to establish a safe level of residues of ronidazole or its metabolites in food representing an acceptable risk to consumers, significant health concerns were identified. For this reason, competent authorities should prevent residues of ronidazole in food. This can be accomplished by not using ronidazole in food producing animals.

---

**STILBENES** (growth promoter)

---

**JECFA evaluation:** 5<sup>th</sup> (1960)

**IARC evaluation:** monograph 100A (2012)

**CAC37** (2014)

**Recommended risk management measures**

In view of the available scientific information, there is no safe level of residues of stilbenes or their metabolites in food that represents an acceptable risk to consumers. For this reason, competent authorities should prevent residues of stilbenes in food. This can be accomplished by not using stilbenes in food producing animals.

**Part 2****A) PROPOSED DRAFT MAXIMUM RESIDUE LIMITS FOR VETERINARY DRUGS IN FOODS CURRENTLY UNDER CONSIDERATION BY CCRVDF**

<b>DIFLUBENZURON (insecticide)</b>				
<b>ADI</b>		0–0.02 mg/kg body weight (bw) – based on a NOAEL of 2 mg/kg bw per day for increased methaemoglobin and sulfhaemoglobin levels in a 2-year study of toxicity and carcinogenicity in rats; and increased methaemoglobin and sulfhaemoglobin levels, platelet counts and hepatic pigmentation in a 1-year study of toxicity in dogs – applying a safety factor of 100 (10 for interspecies variability and 10 for intraspecies variability) (JECFA88, 2019).		
<b>ARfD:</b>		Unnecessary (JECFA85 reiterated the conclusion of JECFA81 that it was not necessary to establish an ARfD, in view of the low acute oral toxicity and the absence of developmental toxicity, and any other toxicological effects likely to be elicited by a single dose (JECFA88, 2019).		
<b>GECDE:</b>		0.84 µg/kg bw per day (for the general population), which represents 4% of the upper bound of the ADI (JECFA88, 2019). 2.85 µg/kg bw per day (for children), which represents 14% of the upper bound of the ADI. (JECFA88, 2019).		
<b>GEADE:</b>		Not estimated because JECFA88 concluded that it was not necessary to establish an ARfD.		
<b>Residue Definition:</b>		Diflubenzuron and the ratio of the MR to the TRR of 0.9 established at its 81st meeting (JECFA88, 2019).		
<b>Species</b>	<b>Tissue</b>	<b>MRLs (µg/kg) recommended by JECFA85</b>	<b>Step</b>	<b>JECFA</b>
Salmon	Muscle plus skin in natural proportions	10	3	88

<b>FLUMETHRIN (insecticide)</b>				
<b>ADI:</b>		0–0.004 mg/kg bw based on the NOAEL of 0.37 mg/kg bw per day for skin lesions in parental animals and reduced survival and body-weight gain in pups in a two-generation toxicity study in rats, and using a safety factor of 100 (10 for interspecies variability and 10 for intraspecies variability).		
<b>ARfD:</b>		0.005 mg/kg bw based on the NOAEL of 0.5 mg/kg bw for salivation in dams in a developmental toxicity study in rats, and using a safety factor of 100 (10 for interspecies variability and 10 for intraspecies variability).		
<b>GECDE:</b>		0.008 µg/kg bw per day (for the general population), which represents 0.2% of the upper bound of the ADI. 0.006 µg/kg bw per day (for children), which represents 0.2% of the upper bound of the ADI. <u>Note:</u> As flumethrin is also used as pesticide the overall dietary exposure was estimated. The assumptions and detailed results will be displayed in the JECFA85 report. Results below are only for use as veterinary drug.		
<b>GEADE:</b>		0.1 µg/kg bw per day (for the general population), which represents 2.2% of the ARfD. 0.1 µg/kg bw per day (for children), which represents 2.2% of the ARfD.		
<b>Residue Definition:</b>		Flumethrin (trans-Z1 and trans Z2 diastereomers at a ratio of approximately 60:40)		
<b>Species</b>	<b>Tissue</b>	<b>MRLs (µg/kg) recommended by JECFA85</b>	<b>Step</b>	<b>JECFA</b>
	Honey	6	5	85

Note: JECFA85 set an MRL for honey of 6 µg/kg, which is twice the limit of quantification (LOQ; 3 µg/kg) of the most reliable analytical method (liquid chromatography coupled with tandem mass spectrometry; LC–MS/MS) used in the residues studies

<b>HALQUINOL</b> (broad-spectrum antimicrobial)				
<b>ADI:</b>		0–0.2 mg/kg bw, based on histopathological changes in the kidney, accompanied by increases in absolute and relative renal weight in a 1-year chronic toxicity study in rats, applying a safety factor of 100 (10 for interspecies variability and 10 for intraspecies variability) (JECFA88, 2019).		
<b>ARfD:</b>		0.3 mg/kg bw, based on a NOAEL of 30 mg/kg bw for clinical signs in dams observed in a developmental toxicity study in mice, with application of a safety factor of 100 (10 for interspecies variability and 10 for intraspecies variability) (JECFA88, 2019).		
<b>GECDE:</b>		5.9 µg/kg bw per day (for general population), which represents 3% of the upper bound of the ADI (JECFA88, 2019). 6.9 µg/kg bw per day (for children), which represents 3.4% of the upper bound of the ADI (JECFA88, 2019).		
<b>GEADE:</b>		2–224 µg/kg bw per day, which represents 0.5–75% of the ARfD (comparable for children and adults) (JECFA88, 2019).		
<b>Residue Definition:</b>		The sum of 5-chloroquinolin-8-ol (5-CL), 5,7-dichloroquinolin-8-ol 5,7-DCL (5,7-DCL) and their glucuronide metabolites: 5-CLG (expressed as 5-CL equivalents) and 5,7-DCLG (expressed as 5,7-DCL equivalents) (JECFA88, 2019).		
<b>Species</b>	<b>Tissue</b>	<b>MRLs (µg/kg) recommended by JECFA85</b>	<b>Step</b>	<b>JECFA</b>
Swine	Muscle	40	3	88
Swine	Liver	500	3	88
Swine	Kidney	900	3	88
Swine	Skin plus fat	350	3	88

<b>IVERMECTIN</b> (broad-spectrum antiparasitic agent)				
<b>ADI:</b>		0–10 µg/kg bw. (JECFA81, 2015)		
<b>ARfD:</b>		0.2 mg/kg bw. (JECFA81, 2015)		
<b>GECDE:</b>		0.41 µg/kg bw per day (for general population), which represents 4% of the upper bound of the ADI. (JECFA88, 2019). 0.59 µg/kg bw per day (for children), which represents 5.9% of the upper bound of the ADI. (JECFA88, 2019).		
<b>GEADE:</b>		87 µg/kg bw per day (for general population), which represents 43% of the ARfD, from consumption of cattle muscle, and of 1.1 µg/kg bw, which represents 0.6% of the ARfD (JECFA88, 2019). 82 µg/kg bw per day (for children), which represents 41% of the ARfD, from consumption of cattle muscle and of 1.0 µg/kg bw, which represents 0.5% of the ARfD, from consumption of sheep muscle (JECFA88, 2019).		
<b>Residue Definition:</b>		Ivermectin B <sub>1a</sub> (H <sub>2</sub> B <sub>1a</sub> , or 22,23-dihydroavermectin B <sub>1a</sub> ) in sheep and goat (JECFA88, 2019).		
<b>Species</b>	<b>Tissue</b>	<b>MRLs (µg/kg) recommended by JECFA85</b>	<b>Step</b>	<b>JECFA</b>
Pig	Muscle	10	3	88
Pig	Liver	15	3	88
Pig	Kidney	15	3	88
Pig	Fat	20	3	88
Sheep	Muscle	10	3	88
Sheep	Liver	15	3	88
Sheep	Kidney	15	3	88
Sheep	Fat	20	3	88
Goat	Muscle	10	3	88
Goat	Liver	15	3	88
Goat	Kidney	15	3	88
Goat	Fat	20	3	88

<b>ZILPATEROL HYDROCHLORIDE</b> ( $\beta$ 2-adrenoceptor agonist)				
<b>ADI:</b>		0-0.04 $\mu\text{g}/\text{kg}$ body weight established at JECFA78 (WHO TRS No. 988, 2014) and reaffirmed at JECFA81 (JECFA81, 2015).		
<b>ARfD:</b>		0.04 $\mu\text{g}/\text{kg}$ body weight based on a lowest-observed-adverse-effect level (LOAEL) of 0.76 $\mu\text{g}/\text{kg}$ body weight for acute pharmacological effects observed in a single-dose human study, with application of an uncertainty factor of 20, comprising a default uncertainty factor of 10 for human individual variability and an additional uncertainty factor of 2 to account for use of a LOAEL for a slight effect instead of a NOAEL (JECFA81, 2015).		
<b>GEADE:</b>		1.9 $\mu\text{g}/\text{day}$ for the general population, which represents approximately 80% of the ARfD. The GEADE is 0.57 $\mu\text{g}/\text{day}$ for children, which represents approximately 94% of the ARfD. (JECFA81, 2015).		
<b>Residue Definition:</b>		Zilpaterol (free base) in muscle, liver and kidney.		
<b>Species</b>	<b>Tissue</b>	<b>MRLs (<math>\mu\text{g}/\text{kg}</math>) recommended by JECFA81</b>	<b>Step</b>	<b>JECFA</b>
Cattle	Kidney	3.3	4	81
	Liver	3.5	4	81
	Muscle	0.5	4	81



**C) MRLS HELD AT STEP 8 BY THE CODEX ALIMENTARIUS COMMISSION**

BOVINE SOMATOTROPINS (production aid)						
<b>JECFA Evaluation:</b>		40 (1992); 50 (1998)				
<b>ADI:</b>		Not specified (1992). The ADI applies to somagrebove, sometribove, somavubove, somidobove.				
<b>Residue Definition:</b>		Not applicable				
Species	Tissue	MRL ( $\mu\text{g}/\text{kg}$ )		Step	JECFA	CCRVDF
Cattle	Muscle	Not specified	1/	8	40, 50	7IV, 8II
Cattle	Liver	Not specified	1/	8	40	7IV, 8II
Cattle	Kidney	Not specified	1/	8	40	7IV, 8II
Cattle	Fat	Not specified	1/	8	40	7IV, 8II
Cattle	Milk	Not specified	1/	8	40	7IV, 8II

ADI "not specified" means that available data on the toxicity and intake of the veterinary drug indicate a large margin of safety for consumption of residues in food when the drug is used according to good practice in the use of veterinary drugs. For that reason, and for the reasons stated in the individual evaluation, JECFA concluded that use of the veterinary drugs does not represent a hazard to human and that there is no need to specify a numerical ADI.

1/ MRL "not specified" means that available data on the identity and concentration of residues of the veterinary drug in animal tissues indicate a wide margin of safety for consumption of residues in food when the drug is used according to good practice in the use of veterinary drugs. For that reason, and for the reasons stated in the individual evaluation, JECFA concluded that the presence of drug residues in the named animal product does not present a health concern and that there is no need to specify a numerical MRL.