

CODEX ALIMENTARIUS COMMISSION



Food and Agriculture
Organization of the
United Nations



World Health
Organization

E

Viale delle Terme di Caracalla, 00153 Rome, Italy - Tel: (+39) 06 57051 - E-mail: codex@fao.org - www.codexalimentarius.org

Agenda Item 10

CX/CF 20/14/10
February 2020

JOINT FAO/WHO FOOD STANDARDS PROGRAMME CODEX COMMITTEE ON CONTAMINANTS IN FOODS

14th Session

Utrecht, The Netherlands, 20 – 24 April 2020

PROPOSED DRAFT MAXIMUM LEVELS FOR TOTAL AFLATOXINS IN CERTAIN CEREALS AND CEREAL-BASED PRODUCTS INCLUDING FOODS FOR INFANTS AND YOUNG CHILDREN (At Step 4)

(Prepared by the Electronic Working Group chaired by Brazil and co-chaired by India)

Codex members and observers wishing to submit comments at Step 3 on this document should do so as instructed in CL 2020/23/OCS-CF available on the Codex webpage/Circular Letters:
<http://www.fao.org/fao-who-codexalimentarius/resources/circular-letters/en/>.

BACKGROUND

1. The Codex Committee on Contaminants in Foods (CCCF) has been discussing the establishment of maximum levels (MLs) for total aflatoxins (AFs, namely the sum of aflatoxins B₁, B₂, G₁ and G₂) in cereals and cereal-based foods since 2013. At the 13th Session of CCCF (CCCF13, 2019) a discussion paper was presented to the Committee with data available in the GEMS/Food database on the occurrence of AFs in cereal and cereal-based products, including cereal-based food for infants and young children, and focusing on maize, rice, sorghum, wheat and flours of these cereals.
2. The discussion paper showed¹ that there is a large dataset available on the occurrence of AFs in cereals and cereal-based products in the GEMS/Food database (more than 17000 samples), submitted mainly by the European Union (EU), Singapore and Canada. The discussion paper also demonstrated that the establishment of any MLs for AFs in maize grain, flour, meal, semolina and flakes derived from maize, husked and polished rice, wheat grain, flour, meal, semolina and flakes derived from wheat could greatly reduce total AFs exposure worldwide, as already stated by the Joint FAO/WHO Committee on Food Additives (JECFA) (TRS 1002-JECFA 83/11).
3. While there was general support for the establishment of maximum levels (MLs), observations were made that the work should be based on more geographically representative data. It was noted that occurrence data in cereals used for the analysis and the subsequent proposal for new work, relied heavily on data from only a few countries and regions. Although calls for data on the occurrence of AFs in cereals and cereal-based products have been made since 2014, the Committee pointed out that the available data were not sufficiently representative of cereal-based foods from all GEMS/Food cluster diets.
4. CCCF13 therefore agreed to establish an Electronic Working Group (EWG) chaired by Brazil and co-chaired by India to present at its next session proposals for MLs for total AFs in maize grain destined for further processing, flour, meal, semolina and flakes derived from maize, husked and polished rice (excluding parboiled rice), cereal-based food for infants and young children and sorghum. The Committee further agreed to include sorghum in the list noting that it was a staple food in many parts of the world and that once the work on the MLs for the food categories mentioned above were completed, the proposal of MLs for other cereals and cereal-based products should be considered. There was also agreement that a call for data should be issued on whole wheat flour and parboiled rice to better assess whether these food categories should be added later.²

¹ CX/CF 19/13/15

² REP19/CF, paras. 146-155, Appendix IX

5. The 42nd Session of the Codex Alimentarius Commission (CAC42, 2019) approved³ the new work on the establishment of MLs for aflatoxins in certain cereals and cereal-based products including foods for infants and young children (i.e. maize grain destined for further processing, flour, meal, semolina and flakes derived from maize, husked and polished rice, sorghum grain destined for further processing and cereal-based food for infants and young children). The proposals for MLs will consider both the impact on the AF exposure and the sample rejection rate.

KEY POINTS DISCUSSED IN THE ELETRONIC WORKING GROUP

6. In developing this discussion paper, the following points were raised by the EWG:

- Some countries questioned the geographic representation of the samples

Data available at the GEMS/Food database came mostly from the United States of America (USA) and European Union, even though calls for data on the occurrence of AFs in cereals and cereals-based products have been made since 2014. Analysis of data grouped by continent, countries and years of sampling showed that the mean level of AFs (lower bound) did not significantly vary between them as to impact on the MLs proposed for each food category. Moreover, the Committee must consider the toxicological relevance of AFs and how the establishment of MLs for these food categories could greatly reduce human exposure to these mycotoxins.

- Some countries questioned the rationale used to propose MLs for each food category

The rationale used to propose the different MLs was based on the profile of contamination of each food category. After creating histograms and determining the P95 for the AFs occurrence in samples submitted to the GEMS/Food database, MLs were proposed considering a maximum rejection rate of 5%. A preliminary exposure assessment was carried out to illustrate the intake reduction of each ML proposed and to support the risk management decisions. After that, a ML was recommended based on the combination of intake reduction and a minimum sample rejection.

- Some countries questioned about the presence of outliers on the dataset

Considering that the Committee have not yet harmonized a procedure to deal with outliers in datasets of heterogeneous distributed contaminants and considering the possibility of samples being contaminated with high levels of AFs it was decided not to remove the possible outliers from this document. Furthermore, the presence of the possible outliers in the dataset did not impact the proposal of MLs since they did not stretch the 95 percentiles.

RECOMMENDATIONS

7. CCCF is invited to consider the proposed MLs for the selected food categories as shown in Appendix as well as the issues raised under other matters taking into account the information provided in paragraph 6 and Appendix II, and comments submitted in reply to CL 2020/23-CF.

³ REP19/CF, Appendix V

APPENDIX I
(For comments)

PROPOSED DRAFT MAXIMUM LEVELS FOR TOTAL AFLATOXINS IN CERTAIN CEREALS AND CEREAL-BASED PRODUCTS INCLUDING FOODS FOR INFANTS AND YOUNG CHILDREN

Food category	Proposal 1		Proposal 2	
	ML	Sample rejection (%)	ML	Sample rejection (%)
Maize grain, destined for further processing ^a	20 µg/kg	4.5	15 µg/kg	5.4
Flour, meal, semolina and flakes derived from maize	15 µg/kg	1.1	10 µg/kg	1.5
Husked rice	20 µg/kg	2.2	15 µg/kg	2.7
Polished rice	8 µg/kg	0.5	4 µg/kg	1.2
Sorghum grain, destined for further processing ^a	10 µg/kg	2.0	8 µg/kg	2.7
Cereal-based Food for infants and young children ^b	2 µg/kg	0.4	1 µg/kg	0.7

^aDestined for further processing” means intended to undergo an additional processing/treatment that has proven to reduce level of AFs before being used as an ingredient in foodstuffs, otherwise processed or offered for human consumption; ^bAll cereal-based foods intended for infants (up to 12 months) and young children (12 to 36 months).

OTHER MATTERS

Codex members and observers are also invited to provide comments or information on the following:

- a) Whether sampling plans and performance criteria for the analysis of total aflatoxins for the food categories listed above, considering each of the MLs should be developed;

In the affirmative, please consider the following questions:

- b) Whether performance criteria for AFs should consider 70% of total aflatoxins would be AFB1 and the remaining 30% would be distributed equally between AFB2, AFG1 and AFG2;
- c) Provide information on analytical methods and sampling plans for the analysis of AFs in cereals and cereal-based products in order to inform the discussion on the associated sampling plans and performance criteria.

APPENDIX II

(For information)

INTRODUCTION

1. Aflatoxins (AFs) are considered the most important naturally occurring group of mycotoxins in the world's food supply. AFs (B₁, B₂, G₁ and G₂) were classified as human liver carcinogens by an evaluation conducted by the JECFA, with AFB₁ being considered the most potent one (FAO/WHO, 1998; FAO/WHO, 2017). No tolerable daily intake was proposed since they are genotoxic carcinogens. JECFA noted, at its last toxicological evaluation on aflatoxins (FAO/WHO, 2017), that rice, wheat and sorghum needed to be addressed in future risk management activities for aflatoxins, considering their contribution to aflatoxin exposure in some parts of the world where these cereals are consumed as staple foods in the diet.
2. Since the complete elimination of aflatoxins from the food supply is not feasible, measures should be taken to control and manage worldwide contamination. At CCCF13 (2019), it was noted that the Code of Practice for the Prevention and Reduction of Mycotoxin Contamination in Cereals (CXC 55 -2004) was adopted in 2003 and revised in 2017 and the logical next step for the CCCF was to establish MLs for aflatoxins in some cereal and cereal based products. Maximum Levels (MLs) for total aflatoxins have been established by the Codex Alimentarius Commission for almonds, Brazil nuts, hazelnuts, peanuts intended for further processing, pistachios and dried figs (CXS 193-1995). The focus of this document is to review occurrence data submitted to the GEMS/Food database and to propose additional MLs for total aflatoxins in cereals and cereal-based products, including food for infants and young children.

DATA ANALYSIS

3. Data on aflatoxins levels in maize grain for further processing, flour, meal, semolina and flakes derived from maize, husked and polished rice, sorghum grain and cereal-based food for infants and small children were obtained from the GEMS/Food database. Data for samples analysed between 2007 and 2019 were extracted from the database for analysis. Worldwide occurrence of aflatoxins in cereals and products thereof was evaluated using data extracted from the GEMS/Food database as of November 2019.
4. First, data were individually analysed and grouped into categories according to their listed "food name, food code and local food name". Final food categories were created considering the data available in the GEMS/Food database and the CCCF grouping recommendations. The following data were removed from the dataset:
 - a. Data that did not meet the basic criteria - For example, samples classified as maize grain but described in the local food name as canned maize (i.e. sweet corn consumed as a vegetable rather than as a cereal grain);
 - b. Aggregated samples (i.e. samples reported as summary statistics rather than individually);
 - c. Samples that were cooked before analysis - since Codex MLs are proposed for raw foods, the form in which they are internationally traded;
 - d. Samples that did not report LOQ or LOD values and that did not have quantifiable results;
 - e. Samples that did not report the exactly quantifiable result when the value was higher than the LOQ – For example, samples that reported results as less than a numerical value, but the value was higher than the LOQ reported (Results \leq 20 $\mu\text{g}/\text{kg}$; LOQ=5);
 - f. Samples that were analysed using methods that had higher LOQs than the highest hypothetical ML considered for each food category in this document;
 - g. Outliers were not removed since aflatoxins distribution are not homogeneous and therefore it is not unlikely that samples with high AFs concentration could be found in the market. Besides that, the few high values maintained in the dataset did not impacted the proposal of MLs since they did not stretch the 95 percentiles. The outliers data treatment for mycotoxins should be further discussed taking into account mycotoxins heterogeneous distribution in food samples.
5. For aflatoxins, some samples included information on individual aflatoxin (AFB₁, AFB₂, AFG₁, AFG₂), the sum of AFB₁ plus AFB₂ and total aflatoxins, which generated up to 6 entries per sample. In such cases, data were gathered according to the "serial number" provided. Samples that reported results only for AFB₂, AFG₁ or AFG₂ were excluded when it was not possible to sum individual concentrations to yield a total aflatoxin concentration using the "serial number". Considering this information, it was not possible to keep a record of the samples excluded from the dataset, since just one sample could lead to the insertion of up to six lines in the dataset.

6. Only samples intended for human consumption were maintained in the dataset, i.e. animal feed samples were not included in the analysis. Lower bound AFs concentrations were estimated considering samples below the reported LOQ as zero, since the positive detection rate for almost all food categories were less than 20%.

PROPOSED DRAFT OF MAXIMUM LEVELS FOR TOTAL AFLATOXINS IN CERTAIN CEREALS AND CEREAL-BASED PRODUCTS, INCLUDING FOOD FOR INFANTS AND YOUNG CHILDREN

7. In order to propose ML for total aflatoxins, data for each food category were organized in three different tables, containing information on the worldwide AFs occurrence, the seasonality during the period analysed and the effects of the implementation of different hypothetical MLs on AFs intake and sample rejection. Different MLs were proposed according to the contaminant distribution profile of each food group.
8. Since the risk assessment for AFs was conducted by JECFA in 2017 (JECFA49), dietary exposure to aflatoxins was estimated in this document only to support the risk management decisions. Dietary exposure to aflatoxins through the consumption of maize grain for further processing, flour, meal, semolina and flakes derived from maize, husked and polished rice, and sorghum grain for further processing was estimated using the GEMS/Food occurrence data and mean consumption data obtained from the 17 GEMS/Food Cluster Diets. Consumption data were chosen in order to best represent the food categories evaluated. Annex I of Appendix I shows countries that belong to each GEMS/Food Cluster and consumption data for each cluster diets can be found in Annex II. Dietary exposure to AFs through the consumption of food for infants and young children was not evaluated since there were no consumption data available for the GEMS/Food Cluster Diets for such foods.
9. Tables 1, 2 and 3 show data on the occurrence and concentration of AFs in maize grain destined for further processing. A total of 1,189,587 samples were analysed and 10% were positive for one or more AFs. The mean of positive samples was 60.7 µg/kg, mean and the 95th percentile (P95) of the lower bound were, respectively, 6.1 µg/kg and 18 µg/kg. Most samples analysed came from the USA (99.6%). The highest lower-bound mean concentrations were found in samples submitted by Finland (400 µg/kg), the USA (6.1 µg/kg), Saudi Arabia (4.4 µg/kg), Philippines (3.8 µg/kg) and Indonesia (3.3 µg/kg). 2012, 2013 and 2011 showed the highest incidence levels of AFs, with respectively, 27.5%, 14.6% and 13.4% of samples containing detectable concentrations of one or more AFs. Table 3 shows that mean of the lower bound ranged from 1.0 µg/kg in samples submitted by Asian countries to 6.1 µg/kg on samples from American countries.

Table 1. GEMS/Food data on the occurrence and concentrations of AFs in maize grain destined for further processing.

Country	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
Belgium	1/19 (5.3)	2.0 (2.0)	0.1	-
Brazil	0/36 (0)	<LOQ	<LOQ	-
Bulgaria	0/3 (0)	<LOQ	<LOQ	-
Canada	29/64 (45.3)	0.1 (0.1-90)	2.9	7.9
Cyprus	1/9 (11.1)	0.8 (0.8)	0.1	-
European Union	1,070/4,045 (26.5)	7.5 (0.02-226)	2.0	6.7
Finland	2/2 (100)	400.4 (0.8-800)	400.4	-
France	0/11 (0)	<LOQ	<LOQ	-
Germany	0/7 (0)	<LOQ	<LOQ	-
Hungary	1/12 (8.3)	4.4 (4.4)	0.4	-
Indonesia	14/20 (70.0)	4.7 (0.3-16.2)	3.3	16.2
Ireland	1/4 (25)	1.0 (1.0)	0.3	-
Italy	2/8 (25)	6.6 (6.6)	1.7	-
Philippines	3/7 (42.9)	8.8 (2.0-14.8)	3.8	-
Poland	0/10 (0)	<LOQ	<LOQ	-
Romania	64/148 (43.2)	3.9 (0.1-41.3)	1.7	4.8
Saudi Arabia	4/37 (10.8)	3.8 (0.1-9.9)	4.4	-
Singapore	0/27 (0)	<LOQ	<LOQ	-
Slovakia	0/3 (0)	<LOQ	<LOQ	-
Slovenia	0/25 (0)	<LOQ	<LOQ	-
Spain	0/5 (0)	<LOQ	<LOQ	-
Thailand	0/20 (0)	<LOQ	<LOQ	-
USA	181,161/1,185,065 (10.0)	61.2 (0.02-9,928)	6.1	18.0
Total	119,352/1,189,587 (10.0)	60.7 (0.02-9,928)	6.1	18.0

^aSamples analysed with methods with LOQ higher than 20 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 2. GEMS/Food data on the occurrence and concentrations of AFs in maize grain destined for further processing organized by year of sampling.

Year	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
2007	13/20 (65.0)	3.4 (0.07-16.2)	3.3	16.2
2008	0/6 (0)	<LOQ	<LOQ	-
2009	10/60 (16.7)	11.5 (0.4-56.2)	1.9	7.1
2010	2,542/37,624 (6.8)	15.3 (2.0-29.8)	4.1	7.0
2011	21,481/160,769 (13.4)	5.6 (0.2-186.2)	10.4	62.0
2012	44,480/161,623 (27.5)	7.4 (0.1-800)	23.0	96.0
2013	22,129/15,244 (14.6)	10.7 (0.1-319.6)	5.6	20.0
2014	5,642/102,865 (5.5)	2.1 (0.2-14.8)	0.9	5.3
2015	3,929/102,824 (3.8)	14.8 (0.001-226)	1.8	0.0
2016	4,690/120,291 (3.9)	10.3 (0.02-113.3)	1.5	0.0
2017	5,408/121,017 (4.5)	4.1 (0.3-47.9)	1.9	0.0
2018	5,943/144,886 (4.1)	2.7 (1.6-4.5)	0.8	0.0
2019	3,085/86,319 (3.6)	19.1 (3.9-34.9)	0.6	0.0
NS	0/39 (0)	<LOQ	<LOQ	-
Total	119,352/1,189,587 (10.0)	60.7 (0.02-9,928)	6.1	18.0

NS: year of sampling was not specified; ^aSamples analysed with methods with LOQ higher than 20 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 3. GEMS/Food data on the occurrence and concentrations of AFs in maize grain destined for further processing organized by continent.

Continent	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
America	118,190/1,185,165 (10.0)	61.2 (0.02-9928)	6.1	18
Asia	21/111 (18.9)	5.1 (0.05-16.2)	1.0	5.4
Europe	1,142/4,311 (26.5)	8.0 (0.02-800)	2.1	6.6
Total	119,352/1,189,587 (10.0)	60.7 (0.02-9928)	6.1	18.0

^aSamples analysed with methods with LOQ higher than 20 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

10. Table 4 shows the impact of the implementation of MLs on exposure and on rejection rates for AFs in maize grain destined for further processing. The intake reduction was estimated for the Cluster Diet with the highest consumption of the food category being examined (worst case scenario -G06) and the sample rejection rate was calculated using all samples in the data set. Four different hypothetical MLs were considered, based on the AFs contamination profile of maize grain data submitted to the GEMS/Food database. Among the four values considered, the establishment of an ML of 20 µg/kg seems to be the most adequate value, both for intake reduction (90.5%; G06) as well as the sample rejection rate (4.5%).

Table 4. Effect of hypothetical MLs on aflatoxins intake through the consumption of maize grain for Cluster G06 (highest consumption pattern).

ML (µg/kg)	Mean AF (µg/kg)	Intake (ng/kg bw per day) ^a	Intake reduction (%)	Sample rejection (%) ^b
No limits	6.1	1.25	-	-
20	0.6	0.1	90.5	4.5
15	0.4	0.09	93.2	5.4
10	0.3	0.05	95.7	6.6
8	0.2	0.04	97.0	7.4

^aConsumption data used: maize, raw; G06=12.33 g/person (mean consumption). ^bPercentage of samples above proposed MLs for AFs considering samples from all Clusters Diets for this food category.

11. Considering the adoption of a ML of 20 µg/kg for maize grain, the rejection rate would not exceed 5% for any of the countries that submitted samples to the GEMS/Food at this time and would be the following for all samples collected in these years: 2011 (8.2%) and 2012 (17.3%).
12. Tables 5, 6 and 7 show data on the occurrence and concentration of AFs in flour, meal, semolina and flakes derived from maize. A total of 3,265 samples were submitted to the GEMS/Food database and 13% were positive for one or more AFs. Mean of positive samples was 13.6 µg/kg, mean and the P95 of the lower bound were respectively 1.8 µg/kg and 1.7 µg/kg. Most samples analysed came from the European Union (55%) and the USA (30%). The highest mean level of the lower bound was found in samples submitted by Singapore (13.9 µg/kg) and Philippines (4.9 µg/kg). The years of 2008 and 2013 showed the highest incidence levels of AFs, with, respectively, 100% (2 of 2 samples) and 28.3%, of samples being positive.

Table 5. GEMS/Food data on the occurrence and concentrations of AFs in flour, meal, semolina and flakes derived from maize.

Country	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
Argentina	1/81 (1.2)	0.1 (0.1)	0.002	-
Brazil	0/30 (0)	<LOQ	<LOQ	-
Canada	32/209 (15.3)	6.1 (0.3-18.7)	0.9	8.8
European Union	175/1,799 (9.7)	5.8 (0.01-790)	0.6	0.6
Philippines	1/1 (100)	4.9 (4.9)	4.9	-
Singapore	86/165 (52.1)	26.7 (0.05-476)	13.9	25.7
USA	131/980 (13.4)	17.4 (0.4-277.9)	2.3	5.6
Total	426/3,265 (13.0)	13.6 (0.01-790)	1.8	1.7

^aSamples analysed with methods with LOQ higher than 15 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 6. GEMS/Food data on the occurrence and concentrations of AFs in flour, meal, semolina and flakes derived from maize, organized by year of sampling.

Year	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
2008	2/2 (100)	0.4 (0.4)	0.4	-
2009	20/136 (14.7)	4.7 (0.2-19.8)	0.7	5.2
2010	8/120 (6.7)	1.2 (0.2-4.4)	0.1	-
2011	20/141 (14.2)	2.1 (0.2-5.0)	0.3	2.8
2012	56/529 (10.6)	1.5 (0.03-10.1)	0.2	0.6
2013	52/184 (28.3)	0.9 (0.1-4.9)	0.3	1.1
2014	43/248 (17.3)	26.6 (0.07-476)	4.6	1.2
2015	15/224 (6.7)	18.1 (0.02-221)	1.2	0.0
2016	96/546 (17.6)	29.9 (0.01-790)	5.3	3.1
2017	48/566 (8.5)	16.5 (0.06-394)	1.4	0.9
2018	30/254 (11.8)	7.7 (0.84-52.9)	0.9	3.0
2019	7/155 (4.5)	2.3 (0.1-6.6)	0.1	-
NS	29/160 (18.1)	6.3 (0.1-18.7)	1.1	9.7
Total	426/3,265 (13.0)	13.6 (0.01-790)	1.8	1.7

NS: year of sampling was not specified; ^aSamples analysed with methods with LOQ higher than 15 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 7. GEMS/Food data on the occurrence and concentrations of AFs in flour, meal, semolina and flakes derived from maize, organized by continent.

Continent	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
America	164/1,300 (12.6)	15.1 (0.1-277.9)	1.9	4.5
Asia	87/166 (52.4)	26.4 (0.1-476)	13.8	24.9
Europe	175/1,799 (9.7)	5.8 (0.01-790)	0.6	0.6
Total	426/3,265 (13.0)	13.6 (0.01-790)	1.8	1.7

^aSamples analysed with methods with LOQ higher than 15 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

13. Table 8 shows the impact of hypothetical MLs for AFs in flour, meal, semolina and flakes derived from maize. Among the five values tested, the data available suggest the establishment of a ML of 10 µg/kg, considering both the intake reduction (90%; G13) as well as the sample rejection rate (1.5%). Considering the adoption of a ML of 10 µg/kg flour, meal, semolina and flakes derived from maize, the rejection rate would exceed 5% only for samples submitted by Singapore (6.1%). The ML of 20 µg/kg was not considered viable since previous discussion papers on aflatoxins in cereals have already showed the effects of processing on the reduction of total AFs content.

Table 8. Effect of hypothetical MLs on aflatoxins intake through the consumption of flour, meal, semolina and flakes derived from maize for cluster G13 (highest consumption pattern).

ML (µg/kg)	Mean AF (µg/kg)	Intake (ng/kg bw per day) ^a	Intake reduction (%)	Sample rejection (%) ^b
No limits	1.8	2.8	-	-
20	0.3	0.4	84.4	1.0
15	0.25	0.4	85.9	1.1
10	0.2	0.3	88.5	1.5
8	0.18	0.3	89.6	1.7
4	0.09	0.1	94.8	3.3

^aConsumption data used: maize, flour (white flour and wholemeal); G13= 94.34 g/person (mean consumption). ^bPercentage of samples above proposed MLs for AFs considering samples from all Clusters Diets for this food category.

14. Tables 9, 10 and 11 show data on the occurrence and concentration of AFs in husked rice. 22.5% of the 692 samples submitted to the GEMS/Food database were positive for at least one aflatoxin. Mean of positive samples was 9.4 µg/kg, mean and the P95 of the lower bound were 2.1 µg/kg and 8.0 µg/kg. USA, European Union and Thailand contributed with the largest dataset of husked rice, representing 42%, 28% and 13% of the samples, respectively. The highest mean level of the lower bound was found in samples submitted by Finland (66.8 µg/kg), Thailand (3.4 µg/kg), and USA (2.9 µg/kg). The highest incidence levels of AFs were found in the years of 2017 (43%), 2008 (42%), 2009 (36%) and 2010 (31%).

Table 9. GEMS/Food data on the occurrence and concentrations of AFs in husked rice.

Country	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
Austria	1/2 (50)	0.2 (0.2)	0.1	-
Brazil	2/19 (10.5)	0.3 (0.3)	0.03	-
Canada	16/43 (37.2)	0.8 (0.01-7.1)	0.3	1.4
European Union	63/195 (32.3)	1.8 (0.1-10.3)	0.6	4.2
Finland	3/3 (100)	66.8 (0.2-200)	66.8	-
France	1/2 (50)	4.2 (4.2)	2.1	-
Lithuania	0/3 (0)	<LOQ	<LOQ	-
Romania	0/1 (0)	<LOQ	<LOQ	-
Singapore	2/35 (5.7)	0.1 (0.1-0.18)	0.01	-
Slovakia	1/6 (16.7)	0.4 (0.4)	0.06	-
Spain	0/2 (0)	<LOQ	<LOQ	-
Sweden	0/1 (0)	<LOQ	<LOQ	-
Thailand	20/90 (22.2)	15.5 (0.3-104)	3.4	13.6
USA	47/290 (16.2)	17.8 (0.6-132)	2.9	11.1
Total	156/692 (22.5)	9.4 (0.01-200)	2.1	8.0

^aSamples analysed with methods with LOQ higher than 15 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 10. GEMS/Food data on the occurrence and concentrations of AFs in husked rice organized by year of sampling.

Year	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
2008	10/24(41.7)	1.1 (0.01-7.1)	0.5	1.8
2009	14/39 (35.9)	0.3 (0.004-1.4)	0.09	0.3
2010	15/49 (30.6)	0.3 (0.2-0.4)	0.08	0.3
2011	0/2 (0)	<LOQ	<LOQ	-
2012	6/27 (22.2)	36.6 (3.6-200)	8.1	-
2013	16/60 (26.7)	4.9 (0.7-10.3)	1.3	9.5
2014	0/37 (0)	<LOQ	<LOQ	-
2015	4/44 (9.1)	22.3 (1.3-82.1)	2.0	-
2016	5/62 (8.1)	3.4 (0.2-6.8)	0.3	-
2017	26/61 (42.6)	0.7 (0.1-4.9)	0.3	0.5
2018	17/64 (26.6)	16.2 (0.3-104)	4.3	26.0
2019	7/75 (9.3)	7.1 (0.3-34.5)	0.7	-
NS	37/148 (25)	19.4 (2.0-132)	4.9	17.0
Total	156/692 (22.5)	9.4 (0.01-200)	2.1	8.0

NS: year of sampling was not specified; ^aSamples analysed with methods with LOQ higher than 15 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 11. GEMS/Food data on the occurrence and concentrations of AFs in husked rice, organized by continent.

Continent	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
America	65/352 (18.5)	13.1 (0.01-132)	2.4	9.0
Asia	22/125 (17.6)	14.1 (0.1-104)	2.5	3.1
Europe	69/215 (32.1)	4.6 (0.1-200)	1.5	4.2
Total	156/692 (22.5)	9.4 (0.01-200)	2.1	8.0

^aSamples analysed with methods with LOQ higher than 15 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

15. Table 12 shows the impact of hypothetical MLs for husked rice. The establishment of a ML of 15 µg/kg seems the most adequate value, considering a reduction of 74% in AFs intake for cluster G03, the cluster with the highest reported consumption of rice, and a sample rejection rate of 2.7%.

Table 12. Effect of hypothetical MLs on aflatoxins intake through the consumption of husked rice for cluster G03 (highest consumption pattern).

ML (µg/kg)	Mean AF (µg/kg)	Intake (ng/kg bw per day) ^a	Intake reduction (%)	Sample rejection (%) ^b
No limits	2.14	1.11	-	-
20	0.65	0.34	69.7	2.2
15	0.55	0.29	74.2	2.7
12	0.53	0.28	75.2	2.9
10	0.47	0.24	78.1	3.5
8	0.34	0.17	84.2	4.9

^aConsumption data used: rice, husked, dry (incl. paddy rice); G03=31.05 g/person (mean consumption).

^bPercentage of samples above proposed MLs for AFs considering samples from all Clusters Diets for this food category.

16. If the committee agrees on the adoption of a ML of 15 µg/kg for husked rice, samples collected in 2018 and with no information on sampling date would exceed 5% of the rejection rate, representing, respectively, 11% and 6.1% of the samples available on the dataset.
17. Data on the occurrence and concentration of AFs in polished rice are shown in Tables 13, 14 and 15. A total of 7261 samples were submitted to the GEMS/Food database, being 20% positive for one or more AFs. Mean of positive samples was 3.0 µg/kg, mean and the P95 of the lower bound were, respectively, 0.6 µg/kg and 1.1 µg/kg. Most samples analysed came from European Union (73%), the USA (8.8%) and Thailand (8.3%). The highest mean level of the lower bound was found in samples submitted by Finland (109.6 µg/kg), followed by Czech Republic and Luxembourg (0.8 µg/kg). The highest incidence of AFs was found in 2008 (56%) and 2009 (56%), followed by the years of 2013 (34%), 2010 (28%) and 2011 (28%).

Table 13. GEMS/Food data on the occurrence and concentrations of AFs in polished rice.

Country	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
Brazil	1/71 (1.4)	4.9 (4.9)	0.07	-
Bulgaria	0/10 (0)	<LOQ	<LOQ	-
Canada	46/80 (57.5)	0.4 (0.002-2.9)	0.2	1.6
Czech Republic	2/3 (66.7)	1.2 (1.2)	0.8	-
European Union	1,249/5,271 (23.7)	1.2 (0.01-251)	0.3	1.2
Finland	22/22 (100)	109.6 (0.2-800)	109.6	770.0
Hungary	0/10 (0)	<LOQ	<LOQ	-
Ireland	0/1 (0)	<LOQ	<LOQ	-
Luxembourg	2/2 (100)	0.8 (0.09-1.5)	0.8	-
Romania	2/5 (40)	1.2 (0.08-2.3)	0.5	-
Saudi Arabia	39/401 (9.7)	2.9 (0.01-27.1)	0.3	0.7
Singapore	3/53 (5.7)	0.1 (0.06-0.16)	0.01	-
Slovakia	0/84 (0)	<LOQ	<LOQ	-
Spain	0/1 (0)	<LOQ	<LOQ	-
Thailand	82/602 (13.6)	1.5 (0.3-28.9)	0.2	0.6
USA	28/645 (4.3)	8.7 (0.6-88)	0.4	0.0
Total	1,476/7,261 (20.3)	3.0 (0.002-800)	0.6	1.1

^aSamples analysed with methods with LOQ higher than 12 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 14. GEMS/Food data on the occurrence and concentrations of AFs in polished rice organized by year of sampling.

Year	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
2008	24/43 (55.8)	0.4 (0.01-2.9)	0.2	1.5
2009	210/377 (55.7)	0.9 (0.002-13.0)	0.5	2.7
2010	164/582 (28.2)	1.1 (0.02-13.6)	0.3	1.4
2011	173/623 (27.8)	1.4 (0.01-17.0)	0.4	1.6
2012	87/689 (12.6)	28.6 (0.03-800)	3.6	0.9
2013	220/650 (33.8)	0.7 (0.01-7.0)	0.2	0.8
2014	178/991 (18.0)	0.9 (0.01-9.0)	0.2	0.8
2015	100/616 (16.2)	3.8 (0.01-251)	0.6	0.9
2016	125/857 (14.6)	1.4 (0.01-27.1)	0.2	0.8
2017	105/624 (16.8)	1.0 (0.01-6.2)	0.2	1.1
2018	64/463 (13.8)	1.9 (0.3-28.9)	0.3	0.9
2019	1/46 (2.2)	0.50 (0.5)	0.01	0.0
NS	25/700 (3.6)	9.25 (0.06-88.0)	0.3	0.0
Total	1,476/7,261 (20.3)	3.0 (0.002-800)	0.6	1.1

NS: year of sampling was not specified; ^aSamples analysed with methods with LOQ higher than 12 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 15. GEMS/Food data on the occurrence and concentrations of AFs in polished rice, organized by continent.

Continent	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
America	75/796 (9.4)	3.52 (0.002-88)	0.33	0.2
Asia	124/1,056 (11.7)	1.93 (0.01-29)	0.23	0.6
Europe	1,277/5,409 (23.6)	3.1 (0.01-800)	0.72	1.2
Total	1,476/7,261 (20.3)	3.0 (0.002-800)	0.6	1.1

^aSamples analysed with methods with LOQ higher than 12 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

18. The impact of hypothetical MLs for AFs in polished rice is shown in Table 16. Considering the data available, the implementation of a ML of 8 µg/kg seems suitable since it will reduce AFs intake in 70% (G09) and would generate a rejection rate of only 0.5%. If the Committee agrees with the ML suggested (8 µg/kg), the rejection rate would exceed 5% only for samples submitted by Finland (27%; 6 samples ≥ 200 µg/kg).

Table 16. Effect of hypothetical MLs on aflatoxins intake through the consumption of polished rice for cluster G09 (highest consumption pattern).

ML (µg/kg)	Mean AF (µg/kg)	Intake (ng/kg bw per day) ^a	Intake reduction (%)	Sample rejection (%) ^b
No limits	0.63	2.99	-	-
12	0.201	0.96	68.0	0.28
10	0.196	0.93	68.8	0.32
8	0.187	0.89	70.2	0.4
4	0.14	0.68	77.4	1.2

^aConsumption data used: rice, polished, dry; G09= 262.1 g/person (mean consumption). ^bPercentage of samples above proposed MLs for AFs considering samples from all Clusters Diets for this food category.

19. Tables 17, 18 and 19 show data on the occurrence and concentration of AFs in sorghum grain destined for further processing. 6% of the 13,168 samples submitted to the GEMS/Food database were positive for at least one aflatoxin. Mean of positive samples was 12.6 µg/kg, and the P95 of the lower bound were 0.7 µg/kg and 6.0 µg/kg. Almost all data of sorghum grain were submitted by the USA (99% of the samples). The highest mean level of the lower bound was found in samples submitted by Indonesia (9.9 µg/kg). The highest incidence levels of AFs were found in the years of 2010 (90%) and 2009 (33%).

Table 17. GEMS/Food data on the occurrence and concentrations of AFs in sorghum grain destined for further processing.

Country	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
Indonesia	17/17 (100)	9.9 (2.3-13.9)	9.9	13.8
Japan	1/9 (11.1)	0.4 (0.4)	0.04	-
Republic of Korea	5/93 (5.4)	4.4(0.3-10.8)	0.2	-
USA	749/13,049 (5.7)	12.7 (5.0-204)	0.7	5.0
Total	772/13,168 (5.9)	12.6 (0.3-204)	0.7	6.0

^aSamples analysed with methods with LOQ higher than 20 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 18. GEMS/Food data on the occurrence and concentrations of AFs in sorghum grain destined for further processing organized by year of sampling.

Year	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
2008	0/1 (0)	<LOQ	<LOQ	-
2009	1/3 (33.3)	0.4 (0.4)	0.1	-
2010	18/20 (90.0)	9.4 (0.3-13.9)	8.5	13.8
2011	0/12 (0)	<LOQ	<LOQ	-
2012	4/84 (4.8)	5.5 (0.6-10.8)	0.3	-
NS	749/13,048 (5.7)	12.7 (5.0-204)	0.7	5.0
Total	772/13,168 (5.9)	12.6 (0.3-204)	0.7	6.0

NS: year of sampling was not specified; ^aSamples analysed with methods with LOQ higher than 20 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 19. GEMS/Food data on the occurrence and concentrations of AFs in sorghum grain destined for further processing, organized by continent.

Continent	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
America	749/13,049 (5.7)	12.7 (5.0-204)	0.7	5.0
Asia	23/119 (19.3)	8.3 (0.3-13.9)	1.6	13.6
Total	772/13,168 (5.9)	12.6 (0.3-204)	0.7	6.0

^aSamples analysed with methods with LOQ higher than 20 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

20. Table 20 shows the impact of hypothetical MLs for sorghum grain destined for further processing. The establishment of a ML of 8 µg/kg seems to be reasonable, considering a reduction of 73% in AFs intake for cluster G12 and a sample rejection rate of 2.7%.

Table 20. Effect of hypothetical MLs on aflatoxins intake through the consumption sorghum grain destined for further processing for cluster G12 (highest consumption pattern).

ML (µg/kg)	Mean AF (µg/kg)	Intake (ng/kg bw per day) ^a	Intake reduction (%)	Sample rejection (%) ^b
No limits	0.7	0.09	-	-
20	0.5	0.06	32.9	0.4
15	0.4	0.05	45.6	1.0
10	0.3	0.03	63.7	2.0
8	0.2	0.02	72.6	2.7

^aConsumption data used: sorghum, raw (incl flour. incl beer); G12= 7.12 g/person (mean consumption).

^bPercentage of samples above proposed MLs for AFs considering samples from all Clusters Diets for this food category

21. If the committee agrees on the adoption of a ML of 8 µg/kg for sorghum grain destined for further processing, samples submitted by Indonesia and samples collected on 2010 would exceed 5% of the rejection rate, representing, respectively, 70% and 60% of the samples available on the dataset of the category analysed.
22. Data on the occurrence and concentration of AFs in food for infants and young children are shown in Tables 21, 22 and 23. A total of 4,532 samples were submitted to the GEMS/Food database, being 5% positive for one or more AFs. Mean of positive samples was 2.8 µg/kg, mean and the P95 of the lower bound were, respectively, 0.13 µg/kg and 0.0 µg/kg. Most samples analysed were submitted by the European Union (76%), Singapore (7%), USA (5%) and Poland (5%). The highest mean level of the lower bound was found in samples submitted by Finland (38.5 µg/kg). The highest incidence of AFs was found in 2008 (29%), followed by the years of 2009 (14%) and 2013 (11%).

Table 21. GEMS/Food data on the occurrence and concentrations of AFs in cereal-based food for infants and young children.

Country	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
Argentina	0/4	<LOQ	<LOQ	-
Brazil	0/38 (0)	<LOQ	<LOQ	-
Bulgaria	0/2 (0)	<LOQ	<LOQ	-
Canada	0/50 (0)	<LOQ	<LOQ	-
Cyprus	0/1 (0)	<LOQ	<LOQ	-
Czech Republic	0/13 (0)	<LOQ	<LOQ	-
European Union	151/3,461 (4.4)	0.2 (0.006-2.1)	0.01	0.0
Finland	13/13 (100)	38.5 (0.3-50)	38.5	50.0
France	0/7 (0)	<LOQ	<LOQ	-
Germany	0/40 (0)	<LOQ	<LOQ	-
Hong Kong	6/20 (30)	0.2 (0.01-1.0)	0.05	-
Hungary	0/30 (0)	<LOQ	<LOQ	-
Ireland	0/14 (0)	<LOQ	<LOQ	-
Italy	0/1 (0)	<LOQ	<LOQ	-
Lithuania	1/1 (100)	2.1 (2.1)	0.05	-
Luxembourg	1/2 (50)	0.01 (0.01)	0.003	-
Malta	1/12 (8.3)	0.07 (0.07)	0.01	-
Poland	1/226 (0.4)	0.02 (0.02)	0.0001	-
Portugal	0/2 (0)	<LOQ	<LOQ	-
Republic of Korea	0/21 (0)	<LOQ	<LOQ	-
Romania	0/1 (0)	<LOQ	<LOQ	-
Saudi Arabia	0/14 (0)	<LOQ	<LOQ	-
Singapore	18/306 (5.9)	0.2 (0.05-0.7)	0.01	0.1
Slovenia	0/27 (0)	<LOQ	<LOQ	-
Spain	0/13 (0)	<LOQ	<LOQ	-
USA	18/231 (7.8)	3.0 (1.0-7.4)	0.2	0.5
Total	210/4,550 (4.6)	2.8 (0.006-50)	0.1	0.0

^aSamples analysed with methods with LOQ higher than 8µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 22. GEMS/Food data on the occurrence and concentrations of AFs in cereal-based food for infants and young children organized by year of sampling.

Year	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
2008	2/7 (28.6)	2.1 (2.1)	0.6	-
2009	25/181 (13.8)	0.2 (0.05-0.3)	0.03	0.4
2010	29/527 (5.5)	0.2 (0.05-0.7)	0.01	0.1
2011	6/319 (1.9)	0.07 (0.05-0.2)	0.001	-
2012	15/834 (1.8)	33.3 (0.02-50)	0.6	0.0
2013	26/250 (10.4)	0.1 (0.006-0.2)	0.01	0.1
2014	49/562 (8.7)	0.2 (0.01-1.5)	0.02	0.1
2015	9/796 (1.1)	0.05 (0.01-0.1)	0.001	-
2016	28/320 (8.8)	2.2 (0.02-7.4)	0.13	0.1
2017	13/364 (3.6)	0.04 (0.01-0.1)	0.001	0.0
2018	0/27 (0)	<LOQ	<LOQ	-
2019	0/2 (0)	<LOQ	<LOQ	-
NS	8/361 (2.2)	0.2 (0.2-0.3)	0.005	-
Total	210/4,550 (4.6)	2.8 (0.006-50)	0.1	0.0

NS: year of sampling was not specified; ^aSamples analysed with methods with LOQ higher than 8µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

Table 23. GEMS/Food data on the occurrence and concentrations of AFs in cereal-based food for infants and young children, organized by continent.

Continent	Number and proportion of positive samples ^a (%)	Mean of positive samples (range) - µg/kg	Lower bound ^b (µg/kg)	
			Mean	P95 ^c
America	18/323 (5.6)	3.0 (1.1-7.4)	0.1	0.0
Asia	24/361 (6.6)	0.2 (0.01-1.0)	0.01	0.05
Europe	168/3,866 (4.3)	3.2 (0.01-50)	0.1	0.0
Total	210/4,550 (4.6)	2.8 (0.01-50.0)	0.1	0.0

^aSamples analysed with methods with LOQ higher than 8 µg/kg were removed; ^bLB: mean of all samples (samples below LOQ were considered as zero); ^cP95 was only estimated when the number of positive samples were ≥10.

23. The impact of hypothetical MLs for AFs in food for infants and young children is shown in Table 24. Dietary exposure to AFs through the consumption of food for infants and young children was not estimated since this food category is intended for consumption by a specific population group and worldwide consumption data for this group is not available. However, infants and young children are of great concern regarding contaminants exposure and, therefore, the effect of establishment of a ML on sample rejection was evaluated for this food category.
24. Considering the data available and the susceptibility of infants and young children, the implementation of a ML of 1 µg/kg seems suitable since would result in a rejection rate of only 0.7% of samples available at the international trade level. If the Committee agrees with the ML suggested (1 µg/kg), the rejection rate would exceed 5% only for samples submitted by Finland (87%; 7 samples = 50 µg/kg) and Lithuania (100%; 1 sample = 2.1 µg/kg) and would also exceed 5% in samples collected in 2008 (29%; 2 samples = 2.1 µg/kg).

Table 24. Effect of the implementation of different MLs for aflatoxins in cereal-based food for infants and young children (only cereal based foods).

ML ($\mu\text{g}/\text{kg}$)	Mean AF ($\mu\text{g}/\text{kg}$)	Sample rejection (%)
No limits	0.13	-
8	0.02	0.2
6	0.014	0.2
4	0.011	0.3
2	0.008	0.4
1	0.005	0.7

25. Considering all data available at the GEMS/Food database and the scenarios tested above, the following MLs are being suggested for total AFs. The proposed MLs for each food category were based both on the intake reduction and sample rejection (less than 5%). Those MLs are a reasonable choice for the food categories selected, since they greatly contributed to AFs intake reduction and did not result in a large withdrawal of samples from international trade.

Table 25. MLs proposed for total aflatoxins in cereals and cereal-based products.

Food category	Proposal 1		Proposal 2	
	ML	Sample rejection (%)	ML	Sample rejection (%)
Maize grain, destined for further processing ^a	20 $\mu\text{g}/\text{kg}$	4.5	15 $\mu\text{g}/\text{kg}$	5.4
Flour, meal, semolina and flakes derived from maize	15 $\mu\text{g}/\text{kg}$	1.1	10 $\mu\text{g}/\text{kg}$	1.5
Husked rice	20 $\mu\text{g}/\text{kg}$	2.2	15 $\mu\text{g}/\text{kg}$	2.7
Polished rice	8 $\mu\text{g}/\text{kg}$	0.5	4 $\mu\text{g}/\text{kg}$	1.2
Sorghum grain, destined for further processing ^a	10 $\mu\text{g}/\text{kg}$	2.0	8 $\mu\text{g}/\text{kg}$	2.7
Cereal-based Food for infants and young children ^b	2 $\mu\text{g}/\text{kg}$	0.4	1 $\mu\text{g}/\text{kg}$	0.7

^a "Destined for further processing" means intended to undergo an additional processing/treatment that has proven to reduce level of AFs before being used as an ingredient in foodstuffs, otherwise processed or offered for human consumption; ^bAll cereal foods intended for infants (up to 12 months) and young children (12 to 36 months).

26. The fact that the MLs suggested above were proposed based on data available at the GEMS/Food database, submitted mainly by European Union and the USA is a drawback, since it may not be representative of AFs occurrence in cereal-based staple foods across all the GEMS/Food Cluster Diets. However, considering that calls for data on AFs in cereals and cereals-based products have been issued repeatedly since 2014, and a more representative dataset did not become available, it is reasonable that MLs for these food groups should be established based on the present dataset despite its shortcomings, considering the toxicological relevance of the implementation of these maximum levels in order to reduce AFs exposure worldwide.
27. Table 26 shows the profile of aflatoxins content in food categories evaluated in this paper. Data available showed that AFB1 is the most prevalent mycotoxin, representing up to 90% of total aflatoxins found in samples analysed.

Table 26. Profile of aflatoxins content in food categories evaluated in this paper.

Food category	% AFB1/AFs^a
Maize grain, destined for further processing ^a	95
Flour, meal, semolina and flakes derived from maize	90
Husked rice	78
Polished rice	92
Sorghum grain	95
Cereal-based Food for infants and young children ^b	92

^a typical proportion of aflatoxin B1 (AFB1) occurrence in naturally contaminated samples according to the data submitted to the GEMS/Food database. AFs = AFB1+AFB2+AFG1+AFG2

Annex I of Appendix II: GEMS/Food 17 Cluster**Table 1.** Countries included in each GEMS/Food Cluster Diets.

Cluster	Countries
G01	Afghanistan, Algeria, Azerbaijan, Iraq, Jordan, Libya, Mauritania, Mongolia, Morocco, Occupied Palestinian Territory, Pakistan, Syrian Arab Republic, Tunisia, Turkmenistan, Uzbekistan, Yemen
G02	Albania, Bosnia and Herzegovina, Georgia, Kazakhstan, Kyrgyzstan, Montenegro, Republic of Moldova, Ukraine
G03	Angola, Benin, Burundi, Cameroon, Congo, Côte d'Ivoire, Democratic Republic of the Congo, Ghana, Guinea, Liberia, Madagascar, Mozambique, Paraguay, Togo, Zambia
G04	Antigua and Barbuda, Bahamas, Barbados, Brunei Darussalam, French Polynesia, Grenada, Israel, Jamaica, Kuwait, Netherlands Antilles, Saint Kitts and Nevis, Saint Lucia, Saint Vincent and the Grenadines, Saudi Arabia, United Arab Emirates
G05	Argentina, Bolivia, Brazil, Cape Verde, Chile, Colombia, Costa Rica, Djibouti, Dominican Republic, Ecuador, El Salvador, Guatemala, Guyana, Honduras, India, Malaysia, Maldives, Mauritius, Mexico, New Caledonia, Nicaragua, North Macedonia, Panama, Peru, Seychelles, South Africa, Suriname, Tajikistan, Trinidad and Tobago, Venezuela
G06	Armenia, Cuba, Egypt, Greece, Iran, Lebanon, Turkey
G07	Australia, Bermuda, Finland, France, Iceland, Luxembourg, Norway, Switzerland, United Kingdom, Uruguay
G08	Austria, Germany, Poland, Spain
G09	Bangladesh, Cambodia, China, Democratic People's Republic of Korea, Guinea Bissau, Indonesia, Lao People's Democratic Republic, Myanmar, Nepal, Philippines, Sierra Leone, Thailand, Timor Leste, Viet Nam
G10	Belarus, Bulgaria, Canada, Croatia, Cyprus, Estonia, Italy, Japan, Latvia, Malta, New Zealand, Republic of Korea, Russian Federation, United States of America
G11	Belgium, Netherlands
G12	Belize, Dominica
G13	Botswana, Burkina Faso, Central African Republic, Chad, Eswatini, Ethiopia, Gambia, Haiti, Kenya, Malawi, Mali, Namibia, Niger, Nigeria, Senegal, Somalia, Sudan, United Republic of Tanzania, Zimbabwe
G14	Comoros, Fiji Islands, Kiribati, Papua New Guinea, Solomon Islands, Sri Lanka, Vanuatu
G15	Czechia, Denmark, Hungary, Ireland, Lithuania, Portugal, Romania, Serbia, Slovakia, Slovenia, Sweden
G16	Gabon, Rwanda, Uganda
G17	Samoa, São Tome and Principe

Annex II of Appendix II: GEMS/Food Consumption Data**Table 1a.** Consumption data obtained from the GEMS/Food Cluster Diets - G01 to G08 (g/person/day).

Food category	G01	G02	G03	G04	G05	G06	G07	G08
Maize raw	0.6	NC	0.6	NC	1.2	12.3	NC	NC
Maize flour	22.7	35.6	87.3	34.9	46.7	49.1	14.3	12.9
Rice husked	1.2	1.3	31.1	4.8	0.3	2.2	2.4	1.6
Rice polished	34.2	10.4	41.7	82.4	150.2	70.5	13.4	10.8
Sorghum raw	0.0	0.01	0.0	0.01	0.0	0.0	0.0	0.0

NC= no consumption data available.

Table 1b. Consumption data obtained from the GEMS/Food Cluster Diets - G09 to G17 (g/person/day).

Food category	G09	G10	G11	G12	G13	G14	G15	G16	G17
Maize raw	1.4	NC	NC	NC	NC	0.01	0.03	NC	NC
Maize flour	19.7	12.5	4.2	52.3	94.3	8.1	28.0	56.0	28.1
Rice husked	0.4	1.1	0.0	5.0	13.5	3.5	2.0	0.01	8.8
Rice polished	266.1	57.2	12.8	62.8	30.2	218.3	12.8	15.2	51.3
Sorghum raw	0.01	1.2	0.0	7.1	0.0	0.0	0.0	0.0	0.0

NC= no consumption data available.

REFERENCES

Codex Alimentarius Commission (CAC), 1995. Codex general standard for contaminants and toxins in food and feed – Codex Standard 193-1995. Available at: <http://tinyurl.com/mpkehpr>.

Codex Alimentarius Commission, 2019. DISCUSSION PAPER ON THE ESTABLISHMENT OF MAXIMUM LEVELS FOR TOTAL AFLATOXINS IN CEREALS (WHEAT, MAIZE, SORGHUM AND RICE), FLOUR AND CEREAL-BASED FOODS FOR INFANTS AND YOUNG CHILDREN – CX/CF 19/13/15. Available at: encurtador.com.br/cnPSU

Codex Alimentarius Commission, 2019. REPORT OF THE 13rd SESSION OF THE CODEX COMMITTEE ON CONTAMINANTS IN FOODS, REP19/CF. Available at: encurtador.com.br/hsTVW

Codex Alimentarius Commission, 2019. Forty-second Session, REP19/CAC. Available at: encurtador.com.br/gHTZ8

FAO/WHO, 1998. Joint FAO/WHO Expert Committee on Food Additives - Evaluation of certain food additives and contaminants: forty-ninth report of the Joint FAO/WHO Expert Committee on Food Additives. vol. 40. WHO Food Additives Series, p. 73.

FAO/WHO, 2017. Joint FAO/WHO Expert Committee on Food Additives (JECFA) - Evaluation of certain food contaminants: eighty-third report of the Joint FAO/WHO Expert Committee on Food Additives. vol. 1002. WHO technical report series, Rome, Italy, p. 182

APPENDIX III
LIST OF PARTICIPANTS

CHAIR**Brazil**

Lígia Lindner Schreiner
Health Regulation Expert
Brazilian Health Regulatory Agency

Larissa Bertollo Gomes Pôrto
Health Regulation Expert
Brazilian Health Regulatory Agency

Co-chair:**India**

Dr. S. Vasanthi, Scientist E
National Institute of Nutrition ICMR

Mr Perumal Karthikeyan
Assistant Director
Food Safety and Standards Authority of India
E-mail: baranip@yahoo.com

Argentina

Argentina's Codex Contact Point
PUNTO FOCAL CODEX
Codex Secretariat
Agroindustry Secretariat

Brazil

Carolina Araujo Vieira
Health Regulation Expert
Brazilian Health Regulatory Agency

Ms Patricia Diniz Andrade
Professor
Brasília Federal Institute of Education, Science
and Technology - IFB
Brasília
Brazil

Canada

Ian Richard
Scientific Evaluator, Food Contaminants Section
Bureau of Chemical Safety, Health Canada

China

Yi Shao

Yongning Wu

Di Wu
FAO/WHO
Yangzte Delta Region Institute of TsingHua Univ.

Congo

Rolande Ingrid Rachel FOUEMINA
ACONOQ
Codex Secretariat
Agence Congolaise de Normalisation et de la
Qualité

Costa Rica

Amanda Lasso Cruz
Ministerio de Economía Industria y Comercio

Cuba

Roberto Dair Garcia de la Rosa
Public Health Ministry

Ecuador

Ana Gabriela Escobar
AGROCALIDAD

European Union

Ms Veerle Vanheusden
Administrator
DG SANTE
European Commission
Brussels

Belgium

India

Dr. K. K. Sharma
Network Coordinator All India Network Project on
Pesticide Residues
Indian Agricultural Research Institute New Delhi –

Dr. Rajesh R
Assistant Director (Tech)
Export Inspection Agency-Kolkata

Iran

Mansooreh Mazaheri
ISIRI-Standard Research Institute

Japan

Mr. Tsuyoshi ARAI
Deputy Director
Food Safety Standards and Evaluation Division,
Pharmaceutical Safety and Environmental Health
Bureau Ministry of Health, Labour and Welfare of
Japan

Kazakhstan

Zhanar Tolysbayeva
The Ministry of Healthcare

Mexico

Tania Daniela Fosado Soriano
Punto de Contacto CODEX México
Secretaría de Economía
Mexico, Mexico

Nigeria

IBITAYO Femi James

North Macedonia

Maja Lukareva
FAO/WHO
Food and Veterinary Agency

Republic of Korea

Seong Yeji
MFDS

Republic of Korea Codex Secretariat
Ministry of Agriculture, Food and Rural Affairs

Lee Geun Pil
Ministry of Agriculture, Food and Rural affairs

Saudi Arabia

Mohammed Alhuthiel
Saudi Food and Drug Authority

Lam Almaiman
Saudi Food and Drug Authority

Thailand

Standards officer, Office of Standard
Development,
National Bureau of Agricultural Commodity and
Food Standards,
Bangkok Thailand

Chutiwan Jatupornpong

Korwadee Phonkliang
Codex Secretariat
Ministry of Agriculture and Cooperatives

Turkey

Sinan ARSLAN
Republic of Turkey Ministry of Food, Agriculture

United Kingdom

Mark Willis
Food Standards Agency

United States of America

Henry Kim
U.S. Food and Drug Administration
Center for Food Safety and Applied Nutrition

Lauren Posnick Robin
U.S. Delegate to CCCF
U.S. Food and Drug Administration
Center for Food Safety and Applied Nutrition

International Confectionery Association (ICA)

Eleonora Alquati
Belgium
E-mail: eleonora.alquati@caobisco.eu

**International Council of Grocery
Manufacturers Associations (ICGMA)**

Nichole Mitchell
Analyst, Ingredient Safety

Nancy Wilkins

United States

**International Special Dietary Foods Industries
(ISDI)**

Jean Christophe Kremer

Belgium

Institute of Food Technologists (IFT)

Rosetta Newsome

United States