

CODEX ALIMENTARIUS COMMISSION



Food and Agriculture
Organization of the
United Nations



World Health
Organization

Viale delle Terme di Caracalla, 00153 Rome, Italy - Tel: (+39) 06 57051 - E-mail: codex@fao.org - www.codexalimentarius.org

CL 2025/35-PR

July 2025

TO: Codex Contact Points
Contact Points of international organizations having observer status with Codex

FROM: Secretariat, Codex Alimentarius Commission,
Joint FAO/WHO Food Standards Programme

SUBJECT: **Request for comments at Step 3 on the recommendations of the Joint FAO/WHO Meeting on Pesticide Residues (JMPR) (2024)¹**

DEADLINE: **25 August 2025**

Background

1. The Joint FAO/WHO Meeting on Pesticide Residues (JMPR) was held from 17 to 26 September 2024.
2. During the meeting, the FAO Panel of Experts was responsible for reviewing residue and analytical aspects of the pesticides under consideration, including data on their metabolism, fate in the environment and use patterns, and for estimating the maximum levels of residues that might occur as a result of use of the pesticides according to good agricultural practice (GAP). Maximum residue levels and supervised trials median residue (STMR) values were estimated for commodities of animal origin. The WHO Core Assessment Group was responsible for reviewing toxicological and related data to establish acceptable daily intakes (ADIs) and acute reference doses (ARfDs), where necessary.
3. The Meeting evaluated 37 pesticides, including six new compounds and six compounds that were re-evaluated within the periodic review programme of the Codex Committee on Pesticide Residues (CCPR), for toxicity or residues, or both.
4. The Meeting established ADIs and ARfDs, estimated maximum residue levels and recommended them for use by CCPR, and estimated STMR and highest residue (HR) levels as a basis for estimating dietary intake.
5. The Meeting also estimated the dietary exposures (both short-term and long-term) of the pesticides reviewed and, on this basis, performed a dietary risk assessment concerning the relevant ADI and, where necessary, ARfD. Cases in which ADIs or ARfDs may be exceeded were clearly indicated to facilitate the decision-making process by CCPR.
6. Pesticides for which the estimated dietary exposure might, based on the available information, exceed their ADIs are marked with footnotes, which are also applied to specific commodities when the available information indicates that the ARfD of a pesticide might be exceeded when the commodity is consumed. The allocations and estimates are shown in the tables in the Annex.
7. The tables include the Codex reference numbers of the compounds and the Codex classification numbers (CCNs) of the commodities to facilitate reference to the Codex MRLs and other Codex documents. Compounds are listed in alphabetical order.
8. Apart from the abbreviations indicated above, the following qualifications are used in the tables.

¹ The recommendations of the JMPR for pesticide maximum residue limits correspond to Step 3 of the Codex Procedure.

(*) (following the recommended maximum residue level)	At or about the limit of quantification
(**)	Compound evaluated under the periodic review programme.
(***)	New compound
ADI	Acceptable daily intake
ARfD	Acute reference dose
CCN	Codex classification number
Dw	The value is reported in the dry weight of the feed commodity.
HR	Highest residue
HR-P	Highest residue in a processed commodity, in mg/kg, calculated by multiplying the HR in the raw commodity by the processing factor.
Po	The recommendation accommodates post-harvest treatment of the commodity.
PoP (following recommendation for processed foods (classes D and E in the Codex classification))	The recommendation accommodates post-harvest treatment of the primary food commodity.
RAC	Raw agricultural commodity
STMR	Supervised trial(s) median residue
STMR-P	An STMR for a processed commodity calculated by applying the concentration or reduction factor for the process to the STMR calculated for the raw agricultural commodity.
W (in place of a recommended MRL)	The previous recommendation is withdrawn, or withdrawal of the recommended MRL or existing Codex or draft MRL is recommended.

9. The report of the 2024 JMPR Meeting (including the complete Annex I) is available in English only at:

FAO: <https://openknowledge.fao.org/handle/20.500.14283/CD5918EN>
<https://doi.org/10.4060/cd5918en>

Should there be problems in downloading the above documents, please contact the FAO and/or WHO JMPR Secretariats at the following addresses to get a copy as an email attachment:

FAO JMPR Secretariat
E-mail: Pesticide-Management@fao.org

WHO JMPR Secretariat
E-mail: JMPR@WHO.INT

REQUEST FOR COMMENTS

10. Codex members and observer international organizations having granted observer status in Codex wishing to submit comments on the proposed MRLs that correspond to Step 3 of the Codex Procedure as proposed by the 2024 JMPR Meeting and also on other recommendations which are relevant to the work of CCPR56 (see tables in the Annex), as well as concern forms, should do so in writing, in conformity with the Procedures for the Elaboration of Codex Standards and Related Texts (*Codex Alimentarius Procedural Manual*) by the deadline indicated on cover page.

11. Concern forms should be sent separately to the Codex Secretariat (codex@fao.org) with a copy to the CCPR Secretariat (ccpr@agri.gov.cn) in a Word file to facilitate their compilation.
12. Circular letters are available on the Codex website² (Circular Letters) and on the CCPR website³.
13. Codex members and observers are invited to provide comments on the MRLs as shown in the Annex to this CL, which is uploaded to the Codex Online Commenting System (OCS): <https://ocs.codexalimentarius.org/>, as per the guidance below, while taking into account the data and information provided in the report of the JMPR (2023).

GUIDANCE ON THE PROVISION OF COMMENTS

14. Comments should be submitted through the Codex Contact Points of Codex members and observers using the OCS.
15. Contact Points of Codex members and observers may login to the OCS and access the document open for comments by selecting “Enter” in the “My reviews” page, available after login to the system.
16. Other OCS resources, including [Frequently Asked Questions \(FAQ\)](#), as well as the user manual and short guide, can be found at the following link: <http://www.fao.org/fao-who-codexalimentarius/resources/circular-letters/en/>.
17. For questions on the OCS, please contact Codex-OCS@fao.org.

² <http://www.fao.org/fao-who-codexalimentarius/circular-letters/en>

³ <https://www.fao.org/fao-who-codexalimentarius/committees/committee/related-circular-letters/en/?committee=CCPR>

ANNEX
ACCEPTABLE DAILY INTAKES, ACUTE REFERENCE DOSES, RECOMMENDED MAXIMUM RESIDUE LEVELS,
SUPERVISED TRIALS MEDIAN RESIDUE VALUES, AND OTHER VALUES RECORDED BY THE 2024 JMPR MEETING
AVAILABLE IN ENGLISH ONLY

ACETAMIPRID (246)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Acetamiprid (246) ADI: 0–0.07 mg/kg bw ARfD: 0.1 mg/kg bw (2011)	VD 0536	Mung beans (dry)	0.4	–	0.032	–
	VD 2065	Subgroup of dry beans, except soya beans and mung beans	0.2	–	0.025	–
	VD 2066	Subgroup of dry peas	0.24	–	0.025	–
<p>Definition of the residue for compliance with the MRL and for dietary risk assessment for plant commodities: Acetamiprid. Definition of the residue for compliance with the MRL and dietary risk assessment for animal commodities: Sum of acetamiprid and desmethyl-acetamiprid, expressed as acetamiprid. The residue is not fat-soluble.</p>						

ACIBENZOLAR-S-METHYL (228)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Acibenzolar-S-methyl (228) ADI: 0–0.08 mg/kg bw ARfD: 0.5 mg/kg bw (2016)	FP 0226	Apple	0.02	0.3	0.01	0.01
	DF 0226	Apple, dried	0.06	–	0.031	0.031
	VS 0623	Cardoon	0.2	–	0.073	0.10
	VS 0624	Celery	0.2	–	0.073	0.10
	VS 0625	Celtuce	0.2	–	0.073	0.10
	VS 0380	Fennel, bulb	0.2	–	0.073	0.10
	PO 0111	Group of avian, edible offal of	0.02*	–	0	0
	PO 0111	Poultry, edible offal of	W	0.02*	–	–
	PF 0111	Group of avian fats	0.02*	–	0	0
	PF 0111	Poultry fats	W	0.02*	–	–
	PM 0110	Group of avian muscle	0.02*	–	0	0
	PM 0110	Poultry meat	W	0.02*	–	–
	MO 0105	Group of edible offal (mammalian)	0.02*	–	0	0
	MO 0105	Edible offal (mammalian)	W	0.02*	–	–
	PE 0112	Group of eggs	0.02*	–	0	0
	PE 0112	Eggs	W	0.02*	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.02*	–	0	0
	MF 0100	Mammalian fats (except milk fats)	W	0.02*	–	–
	ML 0106	Group of milks	0.01*	–	0	0
	ML 0106	Milks	W	0.01*	–	–
MM 0095	Group of muscle (from mammals other than marine mammals)	0.02*	–	0	0	
MM 0095	Meat (from mammals other than marine mammals)	W	0.02*	–	–	
VS 0627	Rhubarb	0.2	–	0.073	0.10	

Definition of the residue for compliance with the MRL for plant and animal commodities: Sum of acibenzolar-S-methyl and 1,2,3benzothiadiazole-7-carboxylic acid (acibenzolar acid) (free and conjugates), expressed as acibenzolar-S-methyl.

Definition of the residue for estimation of dietary intake for plant commodities: Sum of acibenzolar-S-methyl and 1,2,3benzothiadiazole-7-carboxylic acid (acibenzolar acid), (free and conjugated) and 1,2,3-benzothiadiazole-4-hydroxy-7-carboxylic acid (4-OH acibenzolar acid) (free and conjugated), expressed as acibenzolar-S-methyl.

Definition of the residue for estimation of dietary intake for animal commodities: Sum of acibenzolar-S-methyl and 1,2,3benzothiadiazole-7-carboxylic acid (acibenzolar acid) (free and conjugates), expressed as acibenzolar-S-methyl.

The residue is not fat-soluble.

ACYNONAPYR (333)***

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Acynonapyr (333)*** ADI: 0–0.1 mg/kg ARfD: Unnecessary (2024)			JMPR (2024) did not have enough time to conclude the residue definitions, so it decided to postpone the evaluation to the 2025 JMPR.			
Definitions of residue: Not concluded.						

AZOXYSTROBIN (229)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Azoxystrobin (229) ADI: 0–0.03 mg/kg bw ARfD: Unnecessary (2024)	FI 0326	Avocado	1.5	–	0.06 (pulp)	–
	VC 0045	Fruiting vegetables, cucurbits	W	1	–	–
	VC 0045	Fruiting vegetables, cucurbits, except melons and watermelons	1	–	0.17 (whole fruit) 0.2 (pulp)	–
	PO 0111	Group of avian, edible offal of	0.01	–	0	–
	PO 0111	Poultry liver	W	0.01	–	–
	PF 0111	Group of avian fats	0.01	–	0	–
	PF 0111	Poultry fat	W	0.01	–	–
	PM 0110	Group of avian muscle	0.01	–	0	–
	PM 0110	Poultry meat	W	–	–	–
	FC 0001	Group of citrus fruit	–	–	0.49 (pulp)	–
	MO 0105	Group of edible offal (mammalian)	0.07	–	0.02	–
	MO 0105	Edible offal (mammalian)	W	0.07	–	–
	PE 0112	Group of eggs	0.01	–	0	–
	PE 0112	Eggs	W	0.01	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.05	–	0.01	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.01	–	0.01	–
	MM 0095	Meat (from mammals other than marine mammals)	W	0.05	–	–
	MU 1100	Hops, dried	40	30	8.4	–
	VC 0046	Melon	5	–	0.02 (pulp)	–
	FI 0353	Pineapple	2	–	0.01 (pulp)	–
JF 0341	Pineapple juice	–	–	0.026	–	
VC 0432	Watermelon	5	–	0.02	–	

Definition of residues for compliance with the MRL for plant and animal commodities: Azoxystrobin.
Definition of residues for dietary intake assessment for plant and animal commodities: Azoxystrobin.
The residue is fat-soluble.

BUPROFEZIN (173)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Buprofezin (173) ADI: 0–0.009 mg/kg bw ARfD: 0.5 mg/kg bw (2008)	PO 0111	Group of avian, edible offal	0.05*	–	0.05 (liver) 0.05 (kidney)	0.05 (liver) 0.05 (kidney)
	PO 0111	Poultry, edible offal of	W	0.01*	–	–
	PF 0111	Group of avian fats	0.05*	–	0.05	0.05
	PF 0111	Poultry fats	W	0.01*	0.05	0.05
	PM 0110	Group of avian muscle	0.05*	–	0.05	0.05
	PM 0110	Poultry meat	W	0.01*	–	–
	MO 0105	Group of edible offal (mammalian)	0.05*	–	0.05 (liver) 0.05 (kidney)	0.05 (liver) 0.05 (kidney)
	MO 0105	Edible offal (mammalian)	W	0.01*	–	–
	PE 0112	Group of eggs	0.05*	–	0.05	0.05
	PE 0112	Eggs	W	0.01*	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.05*	–	0.05	0.05
	MF 0100	Mammalian fats (except milk fats)	W	0.01*	–	–
	ML 0106	Group of milks	0.01*	–	0.01	–
	ML 0106	Milks	W	0.01*	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.05*	–	0.05	0.05
MM 0095	Meat (from mammals other than marine mammals)	W	0.01*	–	–	

Definition of the residue for compliance with the MRL for plant and animal commodities: Buprofezin.
Definition of the residue for dietary intake assessment for plant and animal commodities: Buprofezin.
The residue is not fat-soluble.

CARFENTRAZONE-ETHYL (338)***

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Carfentrazone-ethyl (338)*** ADI: 0–0.03 mg/kg bw ARfD: 2 mg/kg bw (2024)			As the meeting could not conclude on residue definition for plant and animal commodities, maximum residue levels in plant and animal commodities could not be estimated.			
Definition of residues for compliance with MRL and dietary risk assessment for plant and animal commodities: Not concluded.						

CHLORPYRIFOS (17)**

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Chlorpyrifos (17)** Insufficient toxicological data to establish ADI or ARfD.			As the meeting could not conclude on residue definition for plant and animal commodities, maximum residue levels in plant and animal commodities could not be estimated.			
Definition of the residue for compliance with the MRL and dietary risk assessment for plant and animal commodities: Not concluded.						

CHLORMEQUAT (015)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Chlormequat (015) ADI: 0–0.05 mg/kg bw ARfD: 0.05 mg/kg bw (2017)	GC 0640	Barley	5	2	0.59	–
	GCT 7074	Barley, beer	–	–	0.12	–
	CF 0640	Barley bran, processed	–	–	0.55	–
	CF 3511	Barley, flour	–	–	0.11	–
	GCT 7013	Barley, malt	–	–	0.12	–
	CM 0640	Barley, pearled	–	–	0.32	–
	PO 0111	Group of avian, edible offal of	0.2	–	0.043	0.085
	PO 0111	Poultry, edible offal of	W	0.2	–	–
	PF 0111	Group of avian fats	0.04*	–	0.04	0.04
	PF 0111	Poultry fats	W	0.04*	–	–
	PM 0111	Group of avian muscle	0.04*	–	0.04	0.04
	PM 0111	Poultry meat	W	0.04*	–	–
	MO 0105	Group of edible offal (mammalian)	0.5	–	0.036 (liver) 0.20 (kidney)	0.11 (liver) 0.40 (kidney)
	MO 0105	Edible offal (mammalian)	W	0.5	–	–
	PE 0269	Group of eggs	0.2	–	0.049	0.094
	PE 0269	Eggs	W	0.2	–	–
	MF 0100	Group of mammalian fats (except milk fat)	0.1	–	0.04	0.043
	MF 0100	Mammalian fats (except milk fat)	W	0.1	–	–
	ML 0095	Group of milks	0.2	–	0.069	–
	ML 0095	Milks	W	0.2	–	–
MM 0095	Group of muscle (from mammals other than marine mammals)	0.2	–	0.04	0.085	
MM 0095	Meat (from mammals other than marine mammals)	W	0.2	–	–	

Definition of the residue for compliance with the MRL for plant and animal commodities: Chlormequat cation.
Definition of the residue for dietary intake assessment for plant and animal commodities: Chlormequat cation.
The residue is not fat-soluble.

CYCLOBUTRIFLURAM (339)***

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Cyclobutrifluram (339)*** ADI: 0–0.2 mg/kg bw ARfD: 0.5 (women of childbearing-age) (2024)	FI 0327	Banana	0.01*	–	0	–

Definition of the residue for compliance with the MRL for plant commodities: Cyclobutrifluram.

Definition of the residue for compliance with the MRL for animal commodities: Sum of residues of cyclobutrifluram and 2(trifluoromethyl)pyridine-3-carboxamide (SYN510275), expressed as cyclobutrifluram.

Definition of the residue for dietary risk assessment for plant commodities: Sum of residues of cyclobutrifluram (free and conjugated), N-[(1S,2R)-2-(2,4-dichlorophenyl)-2-hydroxy-cyclobutyl]-2-(trifluoromethyl)pyridine-3-carboxamide (SYN549104) (free and conjugated), 2-(trifluoromethyl)pyridine-3-carboxylic acid (SYN510260) (free and conjugated) and 2-(trifluoromethyl)pyridine-3-carboxamide (SYN510275) (free and conjugated), expressed as cyclobutrifluram.

Definition of the residue for dietary risk assessment for animal commodities: Sum of residues of cyclobutrifluram, N-[(1S,2R)-2-(2,4-dichlorophenyl)-2-hydroxy-cyclobutyl]-2-(trifluoromethyl)pyridine-3-carboxamide (SYN549104) (free and conjugated) and 2(trifluoromethyl)pyridine-3-carboxamide (SYN510275) (free and conjugated), expressed as cyclobutrifluram.
The residue is not fat-soluble.

CYPROCONAZOLE (239)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Cyproconazole (239) ADI: 0–0.02 mg/kg bw ARfD: 0.06 mg/kg bw (2010)	VD 2065	Dry beans, subgroup of (except soya bean)	0.02	–	0.01	–
	VD 0071	Beans (dry)	W	0.02	–	–
	VD 2066	Dry peas, subgroup of	0.02	–	0.01	–
	VD 0072	Peas (dry)	W	0.02	–	–
	PO 0111	Group of avian, edible offal of	0.01*	–	0.01	0.01
	PO 0111	Poultry, edible offal of	W	0.01*	–	–
	PF 0111	Group of avian fats	0.01*	–	0.01	0.01
	PM 0110	Group of avian muscle	0.01*	–	0.01	0.01
	PM 0110	Poultry meat	W	0.01*	–	–
	MO 0105	Group of edible offal (mammalian)	0.5	–	0.14	0.46
	MO 0105	Edible offal (mammalian)	W	0.5	–	–
	PE 0112	Group of eggs	0.01*	–	0.01	0.01
	PE 0112	Eggs	W	0.01*	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.02	–	0.003	0.02
	ML 0106	Group of milks	0.01	–	0.009	–
	ML 0106	Milks	W	0.01	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.01	–	0.003	0.003
MM 0095	Meat (from mammals other than marine mammals)	W	0.01	–	–	
AL 3301	Products of legume feeds with low water (<20%) content (hay), except soya bean and lentil	0.3	–	–	–	

Definition of the residue for compliance with the MRL for plant and animal commodities: Cyproconazole.

Definition of the residue for dietary intake for plant commodities: Cyproconazole.

Definition of the residue for dietary intake for animal commodities: Free and conjugated cyproconazole.

The residue is fat-soluble.

ETHOXYQUIN (035)**

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Ethoxyquin (035)** Insufficient toxicological data were provided to establish ADI and ARfD.	FP 0230	Pear	W	3 (Po)	–	–

Definition of the residue for compliance with the MRL for plant commodities: Ethoxyquin.

Definition of the residue for dietary risk assessment for plant commodities: Not concluded.

ETO FENPROX (184)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Etofenprox (184) ADI: 0–0.03 mg/kg bw ARfD: 1 mg/kg bw (2011)	PO 0111	Group of avian, edible offal of	0.02	–	0.013	0.013
	PO 0111	Poultry, edible offal of	W	0.01*	–	–
	PF 0111	Group of avian fats	0.5	–	0.4	0.4
	PM 0110	Group of avian muscle	0.01*	–	0.003	0.003
	PM 0110	Poultry meat	W	0.01*	–	–
	MO 0105	Group of edible offal (mammalian)	0.1	–	0.072	0.093
	MO 0105	Edible offal (mammalian)	W	0.05*	–	–
	PE 0112	Group of eggs	0.1	–	0.07	0.07
	PE 0112	Eggs	W	0.01*	–	–
	MF 0100	Group of mammalian fats (except milk fats)	3	–	1.5	2.4
	ML 0106	Group of milks	0.1	–	0.096	–
	ML 0106	Milks	W	0.02	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.07	–	0.055	0.062
	MM 0095	Meat (from mammals other than marine mammals)	W	0.5 (fat)	–	–
	GC 0649	Rice	9	0.01*	3.1	–
CM 0649	Rice, husked	0.3	–	0.09	–	
CM 1205	Rice, polished	0.04	–	0.01	–	
<p>Definition of the residue for compliance with the MRL for plant and animal commodities: Etofenprox. Definition of the residue for estimation of dietary intake for plant and animal commodities: Etofenprox. The residue is fat-soluble.</p>						

FENPROPIDIN (340)***

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Fenpropidin (340)*** ADI: 0–0.02 mg/kg bw ARfD: 0.03 mg/kg bw (women of child-bearing age) (2024)	FI 0327	Banana	9	–	0.097 (pulp)	1.2 (pulp)
	GC 0640	Barley	0.15	–	0.055	–
	–	Barley, beer	–	–	0.011	–
	–	Barley, malt	–	–	0.05	–
	PO 0111	Group of avian, edible offal of	0.08	–	0.039	0.074
	PF 0111	Group of avian fats	0.02	–	0.019	0.021
	PM 0110	Group of avian muscle	0.02	–	0.019	0.021
	MO 0105	Group of edible offal (mammalian)	0.4	–	0.3 (liver) 0.0056 (kidney)	0.96 (liver) 0.24 (kidney)
	PE 0112	Group of eggs	0.02	–	0.019	0.021
	MF 0100	Group of mammalian fats (except milk fats)	0.02	–	0.022	0.046
	ML 0106	Group of milks	0.02*	–	0.017	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.02	–	0.036	0.044
	AM 1051	Fodder beet, roots	0.03	–	–	–
	AM 3573	Fodder beet, leaves or tops	30 (dw)	–	–	–
	AS 0081	Straw and hay of cereal grains	4 (dw)	–	–	–
	VR 0596	Sugar beet	0.03	–	–	–
	AM 3599	Sugar beet, dried pulp	0.2	–	–	–
	AV 0596	Sugar beet, leaves or tops (dry)	30 (dw)	–	–	–
	DM 0596	Sugar beet, molasses	–	–	0.0025	–
	DM 3523	Sugar beet, raw/refined sugar	–	–	0.0025	–
	GC 0653	Triticale	0.05	–	0.01	–
	GC 0654	Wheat	0.05	–	0.01	–
	CF 0654	Wheat bran, processed	0.3	–	0.043	–
CF 1211	Wheat, flour	–	–	0.002	–	
–	Wheat, whole-grain bread	–	–	0.0098	–	
–	Wheat, whole-meal flour	–	–	0.0011	–	

Definition of the residue for compliance with the MRL and for dietary risk assessment for plant commodities: Fenpropidin.

Definition of the residue for compliance with the MRL for animal commodities: Sum of fenpropidin and CGA 289267 {2-methyl-2-[4-(2-methyl-3-piperidin-1-yl-propyl)-phenyl]propionic acid}, expressed as fenpropidin.

Definition of the residue for dietary risk assessment for animal commodities: Sum of fenpropidin, CGA 289267 {2-methyl-2-[4-(2-methyl-3-piperidin-1-yl-propyl)-phenyl]propionic acid}, and SYN 515213 {3-hydroxy-2-methyl-2-[4-(2-methyl-3-piperidin-1-yl-propyl)phenyl]-propionic acid} (free and conjugated), expressed as fenpropidin.

The residue is not fat-soluble.

FENPYROXIMATE (193)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Fenpyroximate (193) ADI: 0–0.005 mg/kg bw ARfD: 0.005 mg/kg bw (2021)	FP 0226	Apple	0.1	0.2	0.033	0.072
	DF 0226	Apple, dried	0.5	1	0.15	0.32
	JF 0226	Apple, juice	–	–	0.0053	–
	–	Apple, pasteurized sauce	–	–	0.0059	–
	AB 0226	Apple pomace, dried	1 (dw)	–	2.3	–
	VO 2700	Cherry tomato	0.2	–	0.06	0.12
	VC 0424	Cucumber	0.04	–	0.02	0.03
	MO 0105	Group of edible offal (mammalian)	0.8 ^a	–	0.40	0.77
	MO 0105	Edible offal (mammalian)	W	0.8	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.2	–	0.063	0.13
	MF 0100	Mammalian fats	W	0.2	–	–
	ML 0106	Group of milks	0.01	–	0.005	–
	ML 0106	Milks	W	0.01	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.05	–	0.015	0.041
	MM 0095	Mammalian meats	W	0.2 (fat)	–	–
	AB 0003	Mandarins, dried pulp	0.8 (dw)	–	0.25	–
	OR 0003	Mandarins, oil, edible	25	150	7.8	–
	VC 2039	Subgroup of cucumbers and summer squashes	W	0.3	–	–
	FC 0003	Subgroup of mandarins (including mandarin-like hybrids)	0.15	1	0.05 (RAC) 0.02 (pulp)	0.10 (RAC) 0.03 (pulp)
	–	Subgroup of mandarins, juice	–	–	0.0050	–
	–	Subgroup of mandarins, marmalade	–	–	0.0025	–
	FC 0004	Subgroup of oranges, sweet, sour (including orange-like hybrids)	0.15	0.7	0.055 (RAC) 0.02 (pulp)	0.07 (RAC) 0.02 (pulp)
	–	Subgroup of oranges, marmalade	–	–	0.0027	–
	VO 2045	Subgroup of tomatoes	W	0.3	–	–
	VO 0448	Tomato	0.2	–	0.06	0.12
	AB 0004	Oranges, dried pulp	0.8 (dw)	4 (dw)	0.28	–
	JF 0004	Orange, juice	–	–	0.0054	–
	OR 0004	Orange oil, edible	25	100	8.6	–
	HS 3382	Orange, peel (fresh)	0.5	–	0.155	0.22
	HS 3383	Satsuma mandarin, peel (fresh)	0.6	–	0.185	0.30
	–	Tomato canned	–	–	0.024	0.047
	JF 0448	Tomato, juice	–	–	0.038	–
DM 0448	Tomato puree	–	–	0.043	–	

Definition of the residue for compliance with MRLs for plant commodities: Fenpyroximate.

Definition of the residue for dietary risk assessment for plant commodities: Sum of fenpyroximate and tert-butyl (Z)- α -(1,3-dimethyl-5-phenoxy-pyrazol-4-yl)methylene-amino-oxy)-p-toluate (its Z-isomer M-1), expressed as fenpyroximate.

Definition of the residue for compliance with the MRL for animal commodities: Sum of fenpyroximate and (E)-4-[(1,3-dimethyl-5-phenoxy-pyrazol-4-yl)methyleneaminooxymethyl]benzoic acid (M-3), expressed as fenpyroximate.

Definition of the residue for dietary risk assessment for animal commodities: Sum of fenpyroximate, 2-hydroxymethyl-2-propyl (E)-4-[(1,3-dimethyl-5-phenoxy-pyrazol-4-yl)-methyleneaminooxymethyl]benzoate (Fen-OH), 2-hydroxy-2-methylpropyl (E)- α -(1,3-dimethyl-5-phenoxy-pyrazol-4-yl)methyleneamino-oxy)-p-toluate (R-UL-1) and (E)-4-[(1,3-dimethyl-5-phenoxy-pyrazol-4-yl)methyleneaminooxymethyl]benzoic acid (M-3), expressed as fenpyroximate.

The residue is fat-soluble.

FIPRONIL (202)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Fipronil (202) ADI: 0–0.0002 mg/kg bw ARfD: 0.03 mg/kg bw (2021)	FI 0327	Banana	0.004*	0.004*	0	0
	GC 2087	Barley, similar grains, and pseudocereals with husks, subgroup of	0.002*	0.004*	0.002	–
	AS 0640	Barley straw and fodder, dry	0.05 (dw)	0.07 (dw)	–	–
	HH 0722	Basil	W	0.8	–	–
	SO 0691	Cotton seed	0.003	0.01	0.002	–
	OC 0691	Cotton seed oil, crude	–	–	0.0008	–
	OR 0691	Cotton seed oil, refined	–	–	0.0006	–
	VD 2065	Dry beans, subgroup of (except soya beans)	0.008	0.01	0.001	–
	VD 2066	Dry peas, subgroup of	0.008	–	0.001	–
	PO 0111	Group of avian, edible offal of	0.02	–	0.0303	0.0380
	PO 0111	Poultry, edible offal of	W	0.03	–	–
	PF 0111	Group of avian fats	0.05	–	0.0382	0.0922
	PF 0111	Poultry fats	W	0.07	–	–
	PM 0110	Group of avian muscle	0.02	–	0.0107	0.0288
	PM 0110	Poultry meat	W	0.007	–	–
	MO 0105	Group of edible offal (mammalian)	0.05	–	0.0141 (liver) 0.0047 (kidney)	0.0590 (liver) 0.0248 (kidney)
	MO 0105	Edible offal (mammalian)	W	0.1	–	–
	PE 0112	Group of eggs	0.04	0.04	0.0322	0.0566
	MF 0100	Group of mammalian fats (except milk fats)	0.15	–	0.0680	0.1912
	MF 0100	Mammalian fats (except milk fats)	W	0.4	–	–
	ML 0106	Group of milks	0.02	–	0.0044	–
	ML 0106	Milks	W	0.03	–	–
	FM 0106	Group of milk fat	0.3	–	0.0892	–
	FM 0183	Milk fat	W	0.3	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.015	–	0.0043	0.0221
	MM 0095	Meat (from mammals other than marine mammals)	W	0.03	–	–
	VL 0053	Leafy vegetables, group of	0.01	0.01	0	0.02919
VP 0060	Legume vegetables, group of	W	0.01	–	–	

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
	GC 2091	Maize cereals, subgroup of	0.01	0.01	0.004	–
	AS 0647	Oat straw and fodder, dry	0.05 (dw)	0.07 (dw)	–	–
	VA 0385	Onion, bulb	0.03	0.03	0.02	0.033
	VR 0589	Potato	0.05	0.05	0.00495	0.0296
	–	Potato, cooked peeled	–	–	0.00121	0.00725
	–	Potato peeled	–	–	0.00158	0.00947
	–	Potato flakes	–	–	0.00223	–
	–	Potato, French fries	–	–	0.00183	0.01095
	–	Potato, microwave unpeeled	–	–	0.00334	0.01998
	–	Potato washed	–	–	0.00245	0.01465
	GC 0649	Rice	0.002	0.4	0.002	–
	CM 0649	Rice, husked	0.001*	0.4	0.002	–
	CM 1205	Rice, polished	0.001*	0.15	0.002	–
	CM 1206	Rice bran, unprocessed	W	2	–	–
	AS 0649	Rice straw and fodder, dry	W	0.6 (dw)	–	–
	VR 0075	Root and tuber vegetables, group of (except potato and sugar beet)	0.002	0.002	0	0.00212
	AS 0650	Rye straw and fodder dry	0.05 (dw)	0.05 (dw)	–	–
	VD 0541	Soya bean (dry)	0.002	0.01	0.002	–
	AB 0541	Soya bean hulls	0.015	0.06	–	–
	OC 0541	Soya bean oil, crude	0.01	0.05	0.0088	–
	DM 3524	Sugar (from sugarcane)	–	–	0.00007	–
	VR 0596	Sugar beet	0.004	0.01	0.004	–
	GS 0659	Sugar cane	0.01	0.01	0.003	0.0073
	–	Sugar cane juice	–	–	0.0018	–
	DM 0659	Sugar cane molasses	–	–	0.00007	–
	SO 2091	Sunflower seeds, subgroup of	0.01	0.004*	0.008	–
	VO 2045	Tomato, subgroup of	0.004*	0.01*	0.008	0.008
	AS 0653	Triticale straw and fodder, dry	0.05 (dw)	0.05 (dw)	–	–
	GC 2086	Wheat, similar grains, and pseudocereals with husks, subgroup of	0.03	0.004*	0.008	–
	AS 0654	Wheat straw and fodder, dry	0.05 (dw)	0.05 (dw)	–	–

Definition of the residue for compliance with the MRL for plant and animal commodities: Sum of fipronil and 5-amino-3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-4-trifluoromethylsulfonylpyrazole (MB46136) expressed in terms of fipronil.

Definition of the residue for dietary risk assessment for plant and animal commodities: Sum of fipronil and 5-amino-3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-4-trifluoromethylsulfonylpyrazole (MB46136), 5-amino-3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-4-trifluoromethylthiopyrazole (MB45950) and 5-amino-3-cyano-1-(2,6-dichloro-4-trifluoromethylphenyl)-4-trifluoromethylpyrazole (MB46513) expressed in terms of fipronil.

The residue is fat-soluble.

FLORPYRAUXIFEN-BENZYL (341)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Florpyrauxifen-benzyl (341) ADI: Unnecessary ARfD: Unnecessary (2024)	PE 0112	Eggs	0.03*	–	–	–
	PO 0111	Group of avian, edible offal of	0.03*	–	–	–
	PF 0111	Group of avian fats	0.03*	–	–	–
	PM 0110	Group of avian muscle	0.03*	–	–	–
	MO 0105	Group of edible offal (mammalian)	0.09	–	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.03*	–	–	–
	ML 0106	Group of milks	0.03*	–	–	–
	MM 0095	Group of muscle (from mammals, other than marine mammals)	0.03*	–	–	–
	AS 0162	Hay and/or straw of grasses	5 (dw)	–	–	–
	GC 0645	Maize	0.01*	–	–	–
	AS 3557	Maize, hay and/or straw	0.01* (dw)	–	–	–
	GC 0649	Rice	0.3	–	–	–
	AS 0649	Rice, hay and/or straw	2 (dw)	–	–	–
	CM 0649	Rice, husked	0.01*	–	–	–
<p>Definition of the residue for compliance with the MRL for plant commodities: Florpyrauxifen-benzyl.</p> <p>Definition of the residue for compliance with the MRL for animal commodities: Sum of florpyrauxifen-benzyl, X11966341 and X11438848, expressed as florpyrauxifen-benzyl.</p> <p>Definition of the residue for dietary risk assessment for plant and animal commodities: Not necessary.</p> <p>The residue is fat-soluble.</p>						

FLUAZINAM (306)***

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Fluazinam (306)*** ADI: 0–0.01 mg/kg bw ARfD: 0.07 mg/kg bw (women of child-bearing age) (2024)	Not considered for residues by the present meeting.					

FLUBENDIAMIDE (242)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Flubendiamide (242) ADI: 0–0.02 mg/kg bw ARfD: 0.2 mg/kg bw (2010)	MO 0105	Group of edible offal (mammalian)	1	–	0.31 (liver) 0.32 (kidney)	0.56 (liver) 0.57 (kidney)
	MO 0105	Edible offal (mammalian)	W	1	–	–
	MF 0100	Group of mammalian fats (except milk fats)	2	–	0.62	1.2
	FM 0106	Group of milk fats	5	–	1.6	4.0
	FM 0106	Milk fats	W	5	–	–
	ML 0106	Group of milks	0.1	–	0.066	–
	ML 0106	Milks	W	0.1	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.2	–	0.06	0.13
	MM 0095	Meat (from mammals other than marine mammals)	W	2 (fat)	–	–
	GC 0649	Rice	4	–	1.15	–
	CM 0649	Rice, husked	0.1	–	0.03	–
	CM 1205	Rice, polished	0.01*	–	0.01	–
<p>Definition of the residue for compliance with the MRL and dietary risk assessment for plant commodities: Flubendiamide. Definition of the residue for compliance with the MRL for animal commodities: Flubendiamide. Definition of the residue for dietary risk assessment for animal commodities: Flubendiamide and flubendiamide-iodophthalimide. The residue is fat-soluble.</p>						

FLUOXAPIPROLIN (342)***

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Fluoxapiprolin (342)*** ADI: Unnecessary ARfD: Unnecessary	VO 2700	Cherry tomato	0.1	–	–	–
	FB 0269	Grapes	0.15	–	–	–
	DF 0269	Grapes, dried	0.5	–	–	–
Fluoxapiprolin-pyrazolealanine-oxopropanoic acid (BCS-DE72761) ADI: 0–0.5 mg/kg bw ARfD: Unnecessary	PO 0111	Group of avian edible offal	0.01*	–	–	–
	PF 0111	Group of avian fats	0.01*	–	–	–
	PM 0110	Group of avian muscle	0.01*	–	–	–
Fluoxapiprolin-BDMpyrazole (BCS-BP32808) ADI: 0–1.5 µg/kg bw ARfD: 0.02 mg/kg bw	MO 0105	Group of edible offal (mammalian)	0.01*	–	–	–
	PE 0112	Group of eggs	0.01*	–	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.01*	–	–	–
Fluoxapiprolin-pyrazolecarboxylic acid (BCS-CZ38260) ADI: 0–1.5 µg/kg bw ARfD: 0.02 mg/kg bw (2024)	ML 0106	Group of milks	0.01*	–	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.01*	–	–	–
	VA 0385	Onion, bulb	0.03	–	–	–
	VR 0589	Potato	0.01*	–	–	–
	VO 0448	Tomato	0.07	–	–	–
–	Tomato, dried fruit	0.32	–	–	–	
<p>Definition of the residue for compliance with the MRL for plant and animal commodities: Fluoxapiprolin. Definition of the residue for dietary risk assessment for plant and animal commodities: Not necessary. The residue is fat-soluble.</p>						

FLUPYRADIFURONE (285)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Flupyradifurone (285) ADI: 0–0.08 mg/kg bw ARfD: 0.2 mg/kg bw (2015)	PO 0111	Group of avian, edible offal of	1	–	0.39	0.88
	PO 0111	Poultry, edible offal of	W	1	–	–
	PF 0111	Group of avian fats	0.3	–	0.11	0.24
	PF 0111	Poultry fat	W	0.3	–	–
	PM 0110	Group of avian muscle	0.8	–	0.27	0.64
	PM 0110	Poultry meat	W	0.8	–	–
	MO 0105	Group of edible offal, mammalian	4	–	0.81 (liver) 0.87 (kidney)	2.75 (liver) 3.4 (kidney)
	MO 0105	Edible offal, mammalian	W	4	–	–
	PE 0112	Group of eggs	0.7	–	0.15	0.42
	PE 0112	Eggs	W	0.7	–	–
	MF 0100	Group of mammalian fats (except milk fats)	1	–	0.15	0.86
	MF 0100	Mammalian fats (except milk fats)	W	1	–	–
	ML 0106	Group of milks	0.7	–	0.11	0.48
	ML 0106	Milks	W	0.7	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	1.5	–	0.30	1.27
	MM 0095	Meat (from mammals other than marine mammals)	W	1.5	–	–
	SO 0305	Olives for oil production	5	–	0.495	–
	SO 0495	Rape seeds	0.4	–	0.36	–
FT 0305	Table olives	5	–	0.495	3.3	
–	Olive oil, crude	–	–	0.09415	–	
–	Olive oil, refined	–	–	0.05435	–	
–	Olive oil, solvent extracted refined	–	–	0.05435	–	

Definition of the residue for compliance with the MRL in plant commodities: Flupyradifurone.

Definition for estimation of dietary intake for plant commodities: Sum of flupyradifurone, difluoroacetic acid and 6-chloronicotinic acid, expressed as parent equivalents.

Definition for compliance with the MRL and estimation of dietary intake for animal commodities: Sum of flupyradifurone and difluoroacetic acid, expressed as parent equivalents.

The residue is not fat-soluble.

FOLPET (041)**

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Folpet (041)** ADI: 0–0.1 mg/kg bw (also applies to phthalamic acid expressed as folpet) ARfD: Unnecessary Phthalic acid ADI: 0–0.1 mg/kg bw (2024)	FP 0226	Apple	W	10	–	–
	FI 0327	Banana	2	–	0.060 0.054 (phthalic acid)	–
	GC 0640	Barley	1.5	–	1.5 0.79 (phthalic acid)	–
	AS 0640	Barley, hay and/or straw	40 (dw)	–	–	–
	FB 0020	Blueberries	–	–	1.7 (phthalic acid) 8.0 (phthalamic acid from phosmet use)	–
	FB 0265	Cranberries	–	–	7.4 (phthalic acid) 1.8 (phthalamic acid from phosmet use)	–
	VC 0424	Cucumber	W	1	–	–
	FB 0269	Grapes	W	10	–	–
	DF 0269	Grapes, dried (i.e., currants, raisins, and sultanas)	W	40	–	–
	JF 0269	Grape juice	–	–	3.5 7.7 (phthalic acid)	–
	AB 0269	Grape pomace, dried	20 (dw)	–	–	–
	PO 0111	Group of avian, edible offal of	0.01*	–	0.16 0.053 (phthalic acid)	–
	PF 0111	Group of avian, fats	0.01*	–	0.040 0 (phthalic acid)	–
	PM 0110	Group of avian muscle	0.01*	–	0.040 0 (phthalic acid)	–
	MO 0105	Group of edible offal (mammalian)	0.01*	–	0.058 (liver) 0.33 (kidney) 0.051 (phthalic acid)	–

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
	PE 0112	Group of eggs	0.01*	–	0.10 0.050 (phthalic acid)	–
	MF 0100	Group of mammalian fats (except milk fats)	0.01*	–	0.029 0.064 (phthalic acid)	–
	ML 0106	Group of milks	0.01*	–	0.024 0 (phthalic acid)	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.01*	–	0.029 0 (phthalic acid)	–
	VL 0482	Lettuce, head	W	50	–	–
	VC 0046	Melons, except watermelon	W	3	–	–
	VA 0385	Onion, bulb	W	1	–	–
	VR 0589	Potato	W	0.1	–	–
	VR 0589	Potato	–	–	0.23 (phthalic acid) 0.78 (phthalamic acid from phosmet use)	–
	FB 0275	Strawberry	W	5	–	–
	VO 0448	Tomato	W	3	–	–
	GC 0654	Wheat	0.04	–	0.38 0.32 (phthalic acid)	–
	AS 0654	Wheat, hay and/or straw	40 (dw)	–	–	–
	–	Wine	–	–	0.73 5.0 (phthalic acid)	–
	FB 1236	Wine grapes	15	–	8.7 4.9 (phthalic acid)	–

Definition of the residue for compliance with the MRL for plant and animal commodities: Folpet.

Definition of the residue for folpet dietary risk assessment for plant and animal commodities: Sum of folpet and phthalamic acid, expressed as folpet.

Definition of the residue for phthalic acid dietary risk assessment for plant and animal commodities: Phthalic acid.

The residue is not fat-soluble.

FOSETYL ALUMINIUM (302)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Fosetyl aluminum (302) ADI: 0–1 mg/kg bw ARfD: Unnecessary (2017)	OR 0001	Citrus oil, edible	–	–	0.68	–
	PO 0111	Group of avian, edible offal of	0.05*	–	0	–
	PO 0111	Poultry edible offal	W	0.05*	–	–
	PF 0111	Group of avian fats	0.05*	–	0	–
	PF 0111	Poultry fat	W	0.05*	–	–
	PM 0110	Group of avian muscle	0.05*	–	0	–
	PM 0110	Poultry meat	W	0.05*	–	–
	MO 0105	Group of edible offal (mammalian)	0.5	–	0.21 (liver) 0.33 (kidney)	–
	MM 0105	Edible offal (mammalian)	W	0.5	–	–
	PE 0112	Group of eggs	0.05	–	0.02	–
	PE 0112	Eggs	W	0.05*	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.3	–	0.13	–
	MF 0100	Mammalian fats (except milk fats)	W	0.2	–	–
	ML 0106	Group of milks	0.1	–	0.051	–
	MM 0106	Milks	W	0.1	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.15	–	0.062	–
	MM 0095	Meat (from mammals other than marine mammals)	W	0.15	–	–
	–	Marmalade	–	–	5.9	–
	AB 0004	Oranges, dried pulp	150	–	41.2	–
	JF 0004	Orange, juice	–	–	9.8	–
GC 0649	Rice	40	–	12.55	–	
CM 1205	Rice, polished	40	–	8.55	–	
–	Sterilized canned fruit	–	–	6.1	–	
FC 0004	Subgroup of oranges, sweet, sour	50	20	18 (pulp)	–	

Definition of residue for compliance with MRLs and dietary risk assessment for plant commodities: Sum of fosetyl, phosphonic acid and their salts, expressed as phosphonic acid.

Definition of residue for compliance with MRLs and dietary risk assessment for animal commodities: Phosphonic acid.
The residue is not fat-soluble.

HEXYTHIAZOX (176)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Hexythiazox (176) ADI: 0–0.03 mg/kg bw ARfD: Unnecessary (2008)	FB 2005	Caneberries, subgroup of	4	–	1.0	–
	PO 0111	Group of avian, edible offal of	0.05	–	0.01	–
	PO 0111	Poultry, edible offal of	W	0.05	–	–
	PF 0111	Group of avian fat	0.05	–	0.002	–
	PM 0110	Group of avian muscle	0.05*	–	0	–
	PM 0110	Poultry meat	W	0.05* (F)	–	–
	MO 0105	Group of edible offal, mammalian	0.05	–	0.01	–
	MO 0105	Edible offal, mammalian	W	0.05	–	–
	PE 0112	Group of eggs	0.05	–	0.002	–
	PE 0112	Eggs	W	0.05	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.05	–	0.01	–
	MF 0100	Mammalian fats (except milk fats)	W	0.05	–	–
	FM 0106	Group of milk fats	0.05	–	0.01	–
	FM 0183	Milk fats	W	0.05	–	–
	ML 0106	Group of milks	0.05	–	0.01	–
	ML 0106	Milks	W	0.05	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.05*	–	0	–
	MM 0095	Meat (from mammals other than marine mammals)	W	0.05	–	–
–	Hops, beer	–	–	0.21	–	
DH 1100	Hops, dried	20	3	6.0	–	
–	Hops, spent	–	–	0.75	–	

Definition of the residue for compliance with the MRL in commodities: Hexythiazox.

Definition for estimation of dietary intake for plant commodities: Sum of hexythiazox and all metabolites containing the trans-5-(4-chlorophenyl)-4-methyl-2-oxothiazolidine-moiety (PT-1-3), expressed as hexythiazox.

Definition for compliance with the MRL and for estimation of dietary intake for animal commodities: Sum of hexythiazox and all metabolites containing the trans-5-(4-chlorophenyl)-4-methyl-2-oxothiazolidine-moiety (PT-1-3), expressed as hexythiazox. The residue is fat-soluble.

LAMBDA-CYHALOTHRIN (146)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Lambda-cyhalothrin (146) ADI: 0–0.02 mg/kg bw ARfD: 0.02 mg/kg bw (2007)	Residue evaluation was not performed at the present meeting.					
Definition of the residue for compliance with the MRL for plant and animal commodities: Cyhalothrin, sum of isomers. Definition of the residue for estimation of the dietary exposure for plant and animal commodities: Cyhalothrin, sum of isomers. The residue is fat-soluble.						

MALEIC HYDRAZIDE (102)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Maleic hydrazide (102) Insufficient toxicologic data were provided to establish ADI and ARfD.	VA 0381	Garlic	W	15	–	–
	VA 0385	Onion, bulb	W	15	–	–
	VR 0589	Potato	W	50	–	–
	VA 0388	Shallot	W	15	–	–
Definition of the residue for compliance with the MRL for plant and animal commodities: Maleic hydrazide (free). Definition of the residue for dietary intake assessment for plant and animal commodities: Not concluded. The residue is not fat-soluble.						

METHOPRENE (147)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Methoprene (147) ADI: 0–0.09 mg/kg bw ARfD: Unnecessary	PO 0111	Group of avian, edible offal of	0.02	–	0.007	–
	PO 0111	Poultry, edible offal of	W	0.02	–	–
S-methoprene ADI: 0–0.05 mg/kg bw ARfD: Unnecessary (2001)	PF 0111	Group of avian fats	0.02	–	0.007	–
	PM 0110	Group of avian muscle	0.02	–	0.007	–
	PM 0110	Poultry meat	W	0.02	–	–
	MO 0105	Group of edible offal (mammalian)	0.02	–	0.014	–
	MO 0105	Edible offal (mammalian)	W	0.02	–	–
	PE 0112	Group of eggs	0.02	–	0.006	–
	PE 0112	Eggs	W	0.02	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.2	–	0.092	–
	ML 0106	Group of milks	0.1	–	0.044	–
	ML 0106	Milks	W	0.1 (fat)	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.02	–	0.007	–
MM 0095	Meat (from mammals other than marine mammals)	W	0.2 (fat)	–	–	
TN 0085	Tree nuts	3 (Po)	–	2.1	–	
Definition of residue for compliance with MRL for methoprene and S-methoprene for plant and animal: Methoprene. Definition of residue for dietary intake assessment for methoprene and S-methoprene for plant and animal: Methoprene. The residue is fat-soluble.						

NOVALURON (217)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Novaluron (217) ADI: 0–0.01 mg/kg bw ARfD: Unnecessary (2005)	AM 0660	Almond hulls	15	–	4.46	–
	PO 0111	Group of avian, edible offal of	0.1	–	0.0943	–
	PO 0111	Poultry, edible offal of	W	0.1	–	–
	PF 0111	Group of avian fats	1.5	–	0.919	–
	PM 0110	Group of avian muscle	0.04	–	0.0297	–
	PM 0110	Poultry muscle	0.04	–	0.0297	–
	PM 0110	Poultry meat	W	0.5 (fat)	–	–
	MO 0105	Group of edible offal (mammalian)	0.2	–	0.15	–
	MO 0105	Edible offal (mammalian)	W	0.7	–	–
	PE 0112	Group of eggs	0.3	–	0.1845	–
	PE 0112	Eggs	W	0.1	–	–
	MF 0100	Group of mammalian fats (except milk fats)	3	–	0.599	–
	MF 0106	Group of milk fats	3	–	2.337	–
	FM 0183	Milk fats	W	7	–	–
	ML 0106	Group of milks	0.2	–	0.1176	–
	ML 0106	Milks	W	0.4	–	–
	TN 0085	Group of tree nuts	0.08	–	0.015	–
	MM 0095	Muscle (from mammals other than marine mammals)	0.2	–	0.096	–
	MM 0095	Meat (from mammals other than marine mammals) [in the fat]	W	10 (fat)	–	–
	GC 0649	Rice	5	–	0.6	–
CM 1205	Rice, polished	0.15	–	0.015	–	
<p>Definition of residues for compliance with MRL in plant and animal commodities: Novaluron. Definition of residues for compliance with MRL and for the estimation of dietary exposure in plant and animal commodities: Novaluron. The residue is fat-soluble.</p>						

PERMETHRIN (120)**

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Permethrin (120)** Insufficient data were provided to establish ADI and ARfD			Not considered for residues by the present meeting.			

PHOSPHONIC ACID (301)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Phosphonic acid (301) ADI: 0–1 mg/kg bw ARfD: Unnecessary (2017)			See fosetyl aluminium.			
<p>Definition of residue for compliance with MRLs and dietary risk assessment for plant commodities: Sum of fosetyl, phosphonic acid and their salts, expressed as phosphonic acid.</p> <p>Definition of residue for compliance with MRLs and dietary risk assessment for animal commodities: Phosphonic acid. The residue is not fat-soluble.</p>						

PHOSMET (103)**

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Phosmet (103)** ADI: 0–0.006 mg/kg bw ARfD: 0.03 mg/kg bw Phthalamic acid ADI: 0–0.1 mg/kg bw Phthalic acid (and its anhydride) ADI: 0–0.1 mg/kg bw (2024)	FS 0240	Apricots	W	10	–	–
	FB 0020	Blueberries	20a	10	4.5 7.4 (phthalic acid) 4.1 (phthalamic acid as folpet)	17
	SO 0691	Cottonseed	W	0.05	–	–
	FB 0265	Cranberries	3	3	1.15 1.7 (phthalic acid) 0.94 (phthalamic acid as folpet)	2.2
	FB 0269	Grapes	W	10	–	–
	FC 0001	Group of citrus fruit	W	3	–	–
	FP 0009	Group of pome fruits	W	10	–	–
	TN 0085	Group of tree nuts	W	0.2	–	–
	MM 0095	Meat (from mammals other than marine mammals)	W	1 (fat)	–	–
	ML 0106	Milks	W	0.02	–	–
	FS 0245	Nectarine	W	10	–	–
	FS 0247	Peach	W	10	–	–
	VR 0589	Potato	0.05*	0.05*	0 0.44 (phthalic acid) 0.78 (phthalamic acid as folpet)	0

Definition of the residue for compliance with the MRL for plant and animal commodities: Phosmet.

Definition of the residue for phosmet long-term dietary risk assessment for plant commodities: Sum of phosmet plus 6-times 2(dimethoxyphosphoryl-sulfanylmethyl)isoindole-1,3-dione (phosmet-oxon), expressed as phosmet.

Definition of the residue for phosmet acute dietary risk assessment for plant commodities: Sum of phosmet plus 25-times 2(dimethoxyphosphoryl-sulfanylmethyl)isoindole-1,3-dione (phosmet-oxon), expressed as phosmet.

Definition of the residue for phthalic acid dietary risk assessment for plant and animal commodities: Phthalic acid.

Definition of the residue for phthalamic acid dietary risk assessment for plant and animal commodities: Phthalamic acid, expressed as folpet.

The residue is not fat-soluble.

PROCHLORAZ (142)**

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Prochloraz (142)** ADI: 0–0.02 mg/kg bw ARfD: 0.2 mg/kg bw (2023)	FI 0030	Assorted tropical and subtropical fruits – inedible peel	W	7	–	–
	FI 0326	Avocado	5 (Po)	–	1.6	2.9
	GC 0640	Barley	0.6	–	0.0245	–
	–	Barley beer	–	–	0.00588	–
	–	Barley malt	–	–	0.02107	–
	–	Barley, pearly	–	–	0.0049	–
	–	Barley, pot	–	–	0.00588	–
	GC 0080	Cereal grains	W	2	–	–
	FC 0001	Citrus fruits	W	10	–	–
	PO 0111	Group of avian, edible offal of	0.08	–	0.1	0.13
	PO 0111	Poultry, edible offal of	W	0.2	–	–
	PF 0111	Group of avian fat	0.01	–	0.01	0.013
	PM 0110	Group of avian muscle	0.01*	–	0.014	0.017
	PM 0110	Poultry meat	W	0.05	–	–
	MO 0105	Group of edible offal, mammalian	0.4	–	0.22 (liver) 0.072 (kidney)	0.27 (liver) 0.091 (kidney)
	MO 0105	Edible offal, mammalian	W	10	–	–
	PE 0112	Group of eggs	0.2	–	0.13	0.18
	PE 0112	Eggs	W	0.1	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.02	–	0.015	0.019
	ML 0106	Group of milks	0.02	–	0.0057	–
	ML 0106	Milks	W	0.05	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.01	–	0.0041	0.005
	MM 0095	Meat (from mammals other than marine mammals)	W	0.5	–	–
	SO 0693	Linseed	W	0.05	–	–
	VO 0450	Mushroom	W	3	–	–
	GC 0647	Oats	0.5	–	0.024	–
	HS 0790	Pepper, black, white	W	10	–	–
	SO 0495	Rape seed	W	0.7	–	–
	GC 0650	Rye	0.15	–	0.012	–
	AS 3560	Rye, hay and/or straw	15 (dw)	–	–	–
AS 0081	Straw and hay of cereal grains except pseudocereals	40 (dw)	40 (dw)	–	–	
OR 0702	Sunflower seed oil, edible	W	1	–	–	
SO 0702	Sunflower seeds	W	0.5	–	–	
GC 0653	Triticale	0.15	–	–	–	
GC 0654	Wheat	0.4	–	0.035	–	

Definition of the residue for compliance of MRL for plant and animal commodities: Sum of prochloraz, N-propyl-N-[2-(2,4,6trichlorophenoxy)ethyl]urea (BTS 44595) and N-formyl-N-propyl-N-[2-(2,4,6-trichlorophenoxy)ethyl]urea (BTS 44596), expressed as prochloraz.

Definition of the residue for dietary intake assessment for plant commodities: Sum of prochloraz and N-formyl-N-propyl-N-[2-(2,4,6trichlorophenoxy)ethyl]urea (BTS 44596), expressed as prochloraz.

Definition of the residue for dietary intake assessment for animal commodities: Sum of prochloraz, N-formyl-N-propyl-N-[2-(2,4,6trichlorophenoxy)ethyl]urea (BTS 44596) and 2,4,6-trichlorophenoxyacetic acid (BTS 9608) (free and conjugated), all expressed as prochloraz.

The residue is fat-soluble.

PROPICONAZOLE (160)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Propiconazole (160) ADI: 0–0.07 mg/kg bw ARfD: 0.3 mg/kg bw (2004)	PO 0111	Group of avian, edible offal	0.01*	–	–	–
	PO 0111	Poultry, edible offal of	W	0.01*	–	–
	PF 0111	Group of avian fats	0.01*	–	–	–
	PF 0111	Poultry fats	W	0.01*	–	–
	PM 0110	Group of avian muscle	0.01*	–	–	–
	PM 0110	Poultry meat	W	0.01* (fat)	–	–
	MO 0105	Group of edible offal (mammalian)	0.2	–	–	–
	MO 0105	Edible offal (mammalian)	W	0.2	–	–
	PE 0112	Group of eggs	0.01*	–	–	–
	PE 0112	Eggs	W	0.01*	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.05	–	–	–
	MF 0100	Mammalian fats (except pt milk fats)	W	0.05	–	–
	ML 0106	Group of milks	0.01*	–	–	–
	ML 0106	Milks	W	0.01*	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.01*	–	–	–
	MM 0095	Meat (from mammals other than marine mammals)	W	0.01* (fat)	–	–
CM 1205	Rice, polished	3	10	1.95	–	

Definition of the residue for compliance with the MRL for plant and animal commodities: Propiconazole.

Definition of the residue for dietary risk assessment for plant and animal commodities: Propiconazole plus all metabolites convertible to 2,4-dichlorobenzoic acid (2,4-DCBA), expressed as propiconazole.

The residue is fat-soluble.

PYDIFLUMETOFEN (309)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Pydiflumetofen (309) ADI: 0–0.1 mg/kg bw ARfD: 0.3 mg/kg bw (2018)	VL 0482	Lettuce, head	20	–	5.25	10
	VL 0483	Lettuce, leaf	30	–	5.3	18
	SB 0716	Coffee bean	0.2	–	0.01	0.01
	–	Coffee, instant	–	–	0.0077	–
	AM 3587	Cotton gin trash	7	–	1.7	3.4
	SO 0691	Cottonseed	0.6	–	0.063	–
	OR 0691	Cotton seed oil, edible	–	–	0.0019	–
	PO 0111	Group of avian, edible offal of	0.01*	–	0.02	0.02
	PO 0111	Poultry, edible offal of	W	0.01*	–	–
	PF 0111	Group of avian fats	0.01*	–	0.02	0.02
	PF 0111	Poultry fats	W	0.01*	–	–
	PM 0110	Group of avian muscle	0.01*	–	0.02	0.02
	PM 0110	Poultry meat	W	0.01*	–	–
	MO 0105	Group of edible offal (mammalian)	0.1	–	0.09 (liver) 0.09 (kidney)	0.44 (liver) 0.30 (kidney)
	MO 0105	Edible offal (mammalian)	W	0.1	–	–
	PE 0112	Group of eggs	0.02	–	0.02	0.023
	PE 0112	Eggs	W	0.02	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.1	–	0.02	0.07
	MF 0100	Mammalian fats (except milk fats)	W	0.1	–	–
	ML 0106	Group of milks	0.01*	–	0.02	–
	ML 0106	Milks	W	0.01*	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.01*	–	0.02	0.02
	MM 0095	Meat (from mammals other than marine mammals)	W	0.01*	–	–
FI 0345	Mango	0.08	–	0.01	0.01	
FI 2540	Pitaya	0.9	–	0.01	0.03	
SM 0716	Roasted coffee	–	–	0.0077	–	
FB 2005	Subgroup of cane berries	4	–	0.815	2.6	

Definition of the residue for compliance with the MRL for plant and animal commodities: Pydiflumetofen.

Definition of the residue for dietary risk assessment for plant commodities: Pydiflumetofen.

Definition of the residue for dietary risk assessment for animal commodities other than mammalian liver and kidney: Sum of pydiflumetofen and 2,4,6-TCP (2,4,6-trichlorophenol) and its conjugates, expressed as pydiflumetofen.

Definition of the residue for dietary risk assessment for mammalian liver and kidney: Sum of pydiflumetofen, 2,4,6-trichlorophenol (2,4,6-TCP) and its conjugates, and SYN547897 and its conjugates, expressed as pydiflumetofen.

The residue is fat-soluble.

SPINOSAD (203)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Spinosad (203) ADI: 0–0.02 mg/kg bw ARfD: Unnecessary (2001)	MF 0812	Cattle fat	3	–	0.565	–
	MM 0812	Cattle meat	W	3	–	–
	MM 0812	Cattle muscle	0.3	–	0.078	–
	PO 0111	Group of avian, edible offal of	–	–	0.01	–
	PF 0111	Group of avian fats	0.2	–	0.05	–
	PM 0110	Group of avian muscle	0.01	–	0.01	–
	PM 0110	Poultry meat	W	0.2	–	–
	MO 0105	Group of edible offal (mammalian) [except cattle]	0.5	–	0.064 (liver) 0.032 (kidney)	–
	MO 0105	Edible offal (mammalian) [except cattle]	W	0.5	–	–
	PE 0112	Group of eggs	0.01	–	0.01	–
	PE 0112	Eggs	W	0.01	–	–
	MF 0100	Group of mammalian fats (except milk fats) [except cattle]	2	–	0.32	–
	MF 0100	Mammalian fat	W	–	0.32	–
	MM 0095	Group of muscle (from mammals other than marine mammals) [except cattle]	0.07	–	0.01	–
	MM 0095	Meat (from mammals other than marine mammals) [except cattle]	W	2	0.01 (muscle) 0.32 (fat)	–
	FI 0345	Mango	0.01*	–	0	–
DT 1114	Tea, green or black, fermented and dried, (including concentrates)	10	–	0.325	–	
–	Tea infusion	–	–	0.0004	–	

Definition of the residue for compliance with the MRL in plant and animal commodities: Sum of spinosyn A and spinosyn D.

Definition of the residue for dietary intake assessment in plant and animal commodities: Sum of spinosyn A and spinosyn D.

The residue is fat-soluble for residues in meat but not fat-soluble in milk.

TEBUCONAZOLE (189)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Tebuconazole (189) ADI: 0–0.03 mg/kg bw ARfD: 0.3 mg/kg bw (2010)	HS 0780	Cumin seed	0.9	–	0.22	–
Definition of the residue for compliance with the MRL in plant and animal commodities: Tebuconazole. Definition of the residue for dietary intake assessment in plant and animal commodities: Tebuconazole. The residue is not fat-soluble.						

TEBUFENOZIDE (196)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Tebufenozide (196) ADI: 0–0.02 mg/kg bw (1996)	ML 0812	Cattle milk	W	0.05	–	–
	PO 0111	Group of avian edible offal	0.02*	–	0.02	0.02
	PF 0111	Group of avian fat	0.02*	–	0.02	0.02
	PM 0111	Group of avian muscle	0.02*	–	0.02	0.02
	PM 0111	Poultry meat	W	0.02*	–	–
	MO 0105	Group of edible offal (mammalian)	0.06	–	0.028 (liver) 0.02 (kidney)	0.053 (liver) 0.024 (kidney)
	MO 0105	Edible offal (mammalian)	W	0.02*	–	–
	PE 0112	Group of eggs	0.02*	–	0.02	0.02
	MF 0100	Group of mammalian fats (except milk fats)	0.2	–	0.054	0.17
	ML 0106	Group of milks	0.02	–	0.014	–
	ML 0106	Milks	W	0.01*	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.03	–	0.02	0.029
	MM 0095	Meat (from mammals other than marine mammals)	W	0.05 (fat)	–	–
	GC 0649	Rice	15	–	4.7	–
	CM 0649	Rice, husked	0.6	0.1	0.175	–
CM 1205	Rice, polished	0.3	–	0.045	–	
Definition of the residue for compliance with the MRL in plant and animal commodities: Tebufenozide. Definition of the residue for dietary intake assessment in plant and animal commodities: Tebufenozide. The residue is fat-soluble.						

TETRANILIPROLE (324)

Compound	CCN	Commodity	Recommended maximum residue level (mg/kg)		STMR or STMR-P (mg/kg)	HR or HR-P (mg/kg)
			New	Previous		
Tetraniliprole (324) ADI: 0–0.2 mg/kg bw ARfD: Unnecessary (2021)	AS 0640	Barley, hay and/or straw	0.08 (dw)	–	–	–
	PO 0111	Group of avian, edible offal of	0.01*	–	0	–
	PO 0111	Poultry, edible offal	W	0.01*	–	–
	PF 0111	Group of avian fats	0.01*	–	0	–
	PF 0111	Poultry, fats	W	0.01*	–	–
	PM 0110	Group of avian muscle	0.01*	–	0	–
	PM 0110	Poultry, meat	W	0.01*	–	–
	MO 0105	Group of edible offal (mammalian)	1.0	–	0.43	–
	MO 0105	Edible offal (mammalian)	W	1.0	–	–
	PE 0112	Group of eggs	0.01*	–	0	–
	PE 0112	Eggs	W	0.01*	–	–
	MF 0100	Group of mammalian fats (except milk fats)	0.15	–	0.26	–
	MF 0100	Mammalian fats (except milk fats)	W	0.15	–	–
	ML 0106	Group of milks	0.15	–	0.12	–
	ML 0106	Milks	W	0.15	–	–
	MM 0095	Group of muscle (from mammals other than marine mammals)	0.07	–	0.047	–
	MM 0095	Meat (from mammals other than marine mammals)	W	0.1	–	–
	GC 2088	Rice cereals, subgroup of	0.5	0.02	0.011	–
	AS 0649	Rice, hay and/or straw	9 (dw)	20 (dw)	–	–
	GM 0649	Rice, husked	0.03	0.01*	0.01	–
GC 2087	Subgroup of barley, similar grains, and pseudocereals with husks	0.01*	–	0.01	–	
GC 2086	Subgroup of wheat, similar grains, and pseudocereals without husks	0.01*	–	0.01	–	
AS 0654	Wheat, hay and/or straw	0.08 (dw)	–	–	–	

Definition of the residue for compliance with the MRL for plant and animal commodities: Tetraniliprole.

Definition of the residue for dietary risk assessment for plant commodities: Tetraniliprole *plus* tetraniliprole-N-methylquinazolinone, expressed as tetraniliprole.

Definition of the residue for dietary risk assessment for animal commodities: Tetraniliprole + tetraniliprole-N-methylquinazolinone *plus* tetraniliprole-benzylalcohol, expressed as tetraniliprole.

The residue is not fat-soluble.