



منظمة الأغذية
والزراعة
للأمم المتحدة

联合国
粮食及
农业组织

Food
and
Agriculture
Organization
of
the
United
Nations

Organisation
des
Nations
Unies
pour
l'alimentation
et
l'agriculture

Organización
de las
Naciones
Unidas
para la
Agricultura
y la
Alimentación

FAO METHODOLOGY FOR THE MEASUREMENT OF FOOD DEPRIVATION

Updating the minimum dietary energy requirements

FAO Statistics Division

Rome, October 2008

TABLE OF CONTENTS

1. INTRODUCTION
2. METHODOLOGICAL FRAMEWORK
3. ESTIMATION PROCEDURES
4. MEANING, SIGNIFICANCE AND ADVANTAGES OF THE RESULTING ESTIMATES OF THE PREVALENCE OF UNDERNOURISHMENT

BIBLIOGRAPHY

1. INTRODUCTION

The FAO measure of food deprivation, referred as the prevalence of undernourishment, is based on a comparison of usual food consumption expressed in terms of dietary energy (kcal) with minimum energy requirement norms. The part of the population with food consumption below the minimum energy requirement is considered underfed.

FAO has been traditionally preparing estimates referring to the prevalence of food deprivation in connection with its World Food Survey reports, the last being The Sixth World Food Survey (FAO, 1996). The principal aim of the estimates in this context has been to provide information on the broad dimension of the hunger problem in the developing world. In fact, although the estimates have been worked out on a country-by-country basis, only the global and regional aggregates have been published. Furthermore, the focus has been on the long-term trends as the World Food Surveys were issued between periods of roughly ten years. However, monitoring needs have changed following recent major international Summits. The World Food Summit in 1996 fixed a hunger reduction target to be reached by 2015. The Millennium Declaration in 2000 integrated hunger and poverty reduction in one single goal, specifically as the first of the Millennium Development Goals.

For the purpose of monitoring progress towards the target of halving the number of undernourished, the need had arisen to regularly up-date such estimates at the global as well as country level. FAO has been undertaking this task in its annual report on “The State of Food Insecurity in the World” (SOFI), which was first issued in 1999. The estimates cover 106 countries published in SOFI.

The prevalence of undernourishment published in SOFI 2008 has updated the parameter on minimum dietary energy requirement used in the FAO methodology based on the report of a Joint FAO/ WHO/ UNU Expert Consultation on human energy requirements (<http://www.fao.org/docrep/007/y5686e/y5686e00.htm>) released in 2004 and WHO updated Body Mass Index (BMI¹) reference tables released in April 2006 (http://www.who.int/childgrowth/standards/bmi_for_age/en/index.html) for children aged less than five years and BMI reference tables released in 2007 (http://www.who.int/growthref/who2007_bmi_for_age/en/index.html) for children and adolescents from 5 to 19 years old.

In the following sections the basic methodological framework, the data sources and the procedures used by FAO for deriving the country estimates are described. The meaning and significance of the resulting estimates of prevalence of undernourishment are discussed. Then, the estimations at sub-national levels are also presented.

2. METHODOLOGICAL FRAMEWORK

The estimate of the proportion of the population below minimum level of dietary energy consumption has been defined within a probability distribution framework:

¹ The BMI refers to weight (kg) divided by height² (m).

$$P(U) = P(x < r_L) = \int_{x < r_L} f(x) dx = F_x(r_L)$$

where :

P(U) is the proportion of undernourished in total population

(x) refers to the dietary energy consumption


r_L is a cut-off point reflecting the minimum energy requirement

f(x) is the density function of dietary energy consumption

F_x is the cumulative distribution function

The next paragraphs discuss the estimation of **f(x)** and **r_L** and the graph below illustrates the methodological framework for the estimation procedures of the proportion of population undernourished, i.e. prevalence of undernourishment.

Theoretical distribution of dietary energy consumption



In the graph the curve **f(x)** depicts the proportion of the population corresponding to different per caput dietary energy consumption levels (**x**) represented by the horizontal line. The area

under the curve up to the minimum energy requirement which is a cut-off point in the curve, r_L , represents the proportion of the population undernourished, i.e. prevalence of undernourishment.

3. ESTIMATION PROCEDURES

The density function, $f(\mathbf{x})$, is assumed to be lognormal so that the parameters μ_x and σ_x^2 can be estimated on the basis of the mean, \bar{x} , and the coefficient of variation, $CV(\mathbf{x})$. A summarised description of the procedure for calculating the prevalence of undernourishment on the basis of \bar{x} , $CV(\mathbf{x})$ and r_L applied to an hypothetical country is given below.

3.1 Estimation of the mean and coefficient of variation of the density function $f(\mathbf{x})$

3.1.1 Estimation of the mean, \bar{x}

There are two options for estimating the mean: using Food Balance Sheet (FBS) data or Household Budget Survey (HBS) data. The first can be used to prepare annual estimates for monitoring progress in food security for the country as a whole. The second one allows the derivation of sub-national estimates. The latter estimates can not be prepared on a yearly basis, as they depend on the survey frequency, in general ranging from 5 to 10 years. The illustrative results are presented for both options, FBS and HBS.

a) Dietary energy consumption from the Food Balance Sheet

The mean is represented by the Dietary Energy Supply per person (**DES**) which refers to the food available for human consumption during the course of the reference period, expressed in terms of energy (kcal/person/day). The estimate is derived from the Food Balance Sheets compiled on the basis of data on the production (**PROD**) and trade (**IMP**orts and **EXP**orts) of food commodities. Using these data and the available information on stock changes (**STCH**), losses between the levels at which production is recorded and the household (**WASTE**) and types of utilization (**SEED**, **FEED**, **FOOD**, inputs for **PRO**Cessing derived products and **OTHER** uses) a supply/utilization account is prepared for each commodity in weight terms. The food component, which is usually derived as a balancing item, refers to the total amount of the commodity available for human consumption during the year. The DES is obtained by aggregating the food component of all commodities after conversion into energy values. The table below presents the standard Food Balance Sheet for the hypothetical country in 1999-2001.

The DES per person per day of 2414 (kcal) shown in the first row and last column of the table 1 is the figure used as the estimate of the mean \bar{x} for the hypothetical country, i.e.:

$$\bar{x} = 2414$$

b) Dietary energy consumption from Household Budget Survey

This option requires the conversion of quantities of the different food items consumed by the household into energy values. These data are usually collected through budget surveys using large scale samples which may allow mean estimates not only at the national level but also at sub-national levels such as geographic areas and socio-economic population groups.

Table 1. The Standard FBS for hypothetical country, 1999-2001

	PROD	+ IMP	+ STCH	- EXP	- FEED	- SEED	- PROC	WASTE	- OTHER	= FOOD	DES CALORIES PER PERSON / DAY (*) 2414
.....1000 MT / YEAR.....											
Grand Total											
Cereals (excl. Beer)	19973.7	1116.5	-355.7	6673.9	5211.8	434.7	407.5	969.4	9.7	7027.8	1114.2
Starchy Roots	16956.2	133.8	-1053.9	13525.9	0.4	0.9	143.7	1350.1	3.7	1011.4	45.2
Sugar crops	53406.6		-1333.3	0.3			43698.3	2753.7		5621.0	73.0
Sugar & Sweeteners	5267.7	11.3	-136.6	3360.6					13.0	1776.6	283.2
Pulses	269.5	5.7		37.9		21.9		8.2		207.3	31.5
Treenuts	54.0	2.2		15.8						40.5	6.5
Oil crops	2337.2	873.5	-198.7	38.5	1.0	14.3	1735.2	135.9		1087.5	100.1
Vegetable Oils	819.9	66.3	-149.9	116.5					272.8	348.7	137.9
Vegetables	2753.0	25.3		372.1			0.0	245.7		2163.9	26.8
Fruits (excl. Wine)	7270.5	55.9	0.2	1173.2			14.5	566.7		5574.7	114.4
Stimulants	78.1	21.4	-6.7	64.6						28.5	0.9
Spices	67.1	7.2		20.9				1.9		51.6	6.9
Alcoholic Beverages	2114.9	28.4		78.9					24.0	2040.4	163.4
Honey	3.0	0.2		1.6						1.6	0.2
Meat	1902.5	3.3		271.6				20.7		1614.8	150.9
Offals	75.7	2.8		0.4						77.7	3.8
Animal Fats	31.8	19.1		0.7					5.6	44.6	15.6
Milk (excl. Butter)	409.4	1095.4		81.1				12.3	12.0	1400.1	32.1
Eggs	812.0	1.3		6.8		137.4		40.6		628.4	42.6
Fish, Seafood	3458.0	532.1	1.7	809.3	1185.9					1996.5	62.2
Aquatic Products	30.1	0.5		14.5						16.1	0.2
Miscellaneous											2.1

(*) Food quantities converted into energy values and divided by total population and by 365 days.

3.1.2 Estimation of the coefficient of variation, CV(x)

The CV of the household daily per person dietary energy consumption is formulated as follows:

$$CV(x) = \sqrt{CV^2(x|v) + CV^2(x|r)}$$

Where CV(x) is the total CV of the household daily per person dietary energy consumption, CV(x/v) is the component due to household per person daily income (v) and CV(x/r) is the component due to energy requirement (r). CV(x/r) is considered to be a fixed component and

is estimated to correspond to about 0.20. $\mathbf{CV}(\mathbf{x}|\mathbf{v})$ is however estimated on the basis of household survey data.

For the purpose of estimation, $\mathbf{CV}(\mathbf{x}|\mathbf{v})$ is formulated as follows:

$$\mathbf{CV}(\mathbf{x}|\mathbf{v}) = \sigma(\mathbf{x}|\mathbf{v}) / \mu(\mathbf{x}).$$

The numerator of the ratio is derived as follows:

$$\sigma(x | v) = \sqrt{\left[\sum_{j=1}^k f_j (x | v)_j^2 - \left(\sum_{j=1}^k f_j (x | v)_j \right)^2 / n \right] / (n - 1)}$$

and the denominator, which is the overall average household daily per person dietary energy consumption, is derived as follows:

$$\mu(x) = \sum (x) / n$$

Where k is the number of income classes and f_j is the number of sampled households and $(x|v)_j$ is the average household daily per person dietary energy consumption of the j th income or expenditure class.

Thus, the data required for estimating $\mathbf{CV}(\mathbf{x}|\mathbf{v})$ are the averages of household daily per person dietary energy consumption by household income or expenditure classes from n households and the number of households in each class. The table below presents the average daily per person energy consumption by deciles of household per person total expenditure from a recent National Household Budget Survey conducted in the hypothetical country (sample of 2370 households).

Table 2. Average daily dietary energy consumption per person by household expenditure deciles

<u>Decile of household</u> <u>daily per person</u> <u>expenditure</u>	<u>Average dietary energy consumption</u> <u>(kcal per person/day)</u>
1	1554
2	1874
3	2066
4	2263
5	2413
6	2461
7	2530
8	2474
9	3093
10	3373

Using the data from the above table, $\mathbf{CV}(\mathbf{x}|\mathbf{v})$ is estimated as follows :

$$CV(x|v) = \sigma(x|v) / \mu(x) = 508 / 2410 = 0.211$$

Hence, given that $CV(x/r)$ corresponds to 0.20, we obtain

$$CV(x) = \sqrt{0.211^2 + 0.20^2} = 0.29$$

According to the sample size and design, the CV estimates can be broken-down by geographic areas and socio-economic groups.

3.2 Estimation of the minimum energy requirement (cut-off point), r_L

The procedure for estimating the minimum energy requirement by sex and age group begins with the specification of the reference body weight. After specifying the reference body weight the procedure for arriving at the corresponding energy requirement differs among population groups of different sex and age. Therefore the procedure for deriving the reference body weight for attained-height is handled first, followed by two separate subsections dealing with the derivation of minimum energy requirements for the various sex and age population groups and lastly a fourth subsection dealing with the derivation of the overall minimum daily per person energy requirement. Minimum energy requirements can be estimated for geographic and socioeconomic groups, using survey data on attained-heights and demographic structure.

3.2.1 Reference body- weight

The reference body weights by sex and age groups are based on the available weight-for-height in the Body Mass Index (BMI) reference tables published by the World Health Organization (WHO). Thus given an estimate of the actual attained-height the acceptable weight corresponding to this height is derived from these tables.

For **children below age 10** the reference body weight is fixed at the **median** of the range of weight-for-height given by the BMI reference tables (WHO, 2006 and 2007).

For **adults and children** of age 10 and above, the reference body weight is estimated on the basis of the **fifth** percentile of the distribution of the BMI (WHO, 1995 and 2007).

The actual attained-heights by sex and age used are those estimated by national anthropometric studies. The attained-height figures for the hypothetical country are given below.

3.2.2 Minimum energy requirements for children and adolescents below 18

The minimum energy requirement per person for children and adolescents is obtained by multiplying the reference body weight for attained-heights, as defined in section 3.2.1, by the recommended energy requirement per kilogram of body weight for each sex and age population group, using the total energy expenditure equations. The energy requirements per kilogram of body weight are based on the recommendations of the report of a Joint FAO/WHO/UNU Expert Consultation on human energy requirements (FAO/WHO/UNU, 2004).

Table 3: Average heights by age and sex

Actual height in cm					
Age (years)	Male	Female	Age (years)	Male	Female
0	66.5	61.5	10	128.8	129.0
1	73.5	71.5	11	133.5	135.7
2	81.5	79.5	12	138.6	142.4
3	87.5	85.5	13	144.3	146.8
4	96.5	95.5	14	150.9	149.1
5	102.5	100.5	15	157.6	151.0
6	108.5	108.5	16	162.8	152.9
7	113.5	113.5	17	165.0	153.8
8	118.5	117.5	18 +	166.0	154.4
9	122.5	122.5			

3.2.3 Minimum energy requirements for adults

The minimum energy requirements per person for adults are derived by first estimating the Basic Metabolic Rate (**BMR**) on the basis of the reference body weight, as defined in section 3.2.1, and using the sex and age specific regression parameters of the Schofield equations (James and Schofield, 1990). The Schofield equations were endorsed in the 2004 report on human energy requirements.

Then, the minimum energy requirements are derived by multiplying BMR by sex specific Physical Activity Level factors.

3.2.4 The overall minimum daily per person energy requirement

The overall minimum daily per person dietary energy requirement, which is used as the cut-off point, r_L , for estimating the prevalence of undernourishment, is derived by aggregating the sex-age requirements weighted by the proportion of each sex and age group in the total population.

Finally a pregnancy allowance (**PA**) in per person terms for the whole population is added to the overall requirement. The PA is estimated by multiplying the birth rate by 210 kilo-calories, assuming an estimated daily requirement of 280 kilo-calories during pregnancy over 75 per cent of the year.

The estimated country birth rate for the hypothetical country in 1999-2001 is 26 per thousand. Thus, the overall minimum daily per person energy requirement is derived as follows:

$$r_L = \sum_{ij} (\text{MER}_{ij} * P_{ij}) + \text{PA} = 1673 + 7 = 1680 \text{ kcal/day}$$

Where :

MER = minimum daily energy requirement per person

P_{ij} = proportion of each sex and age group in the total population

PA = pregnancy allowance

i = age group

j = sex

Details for computing this value are given in the annex.

3.3. Estimation of the proportion and number of undernourished

The density function of dietary energy consumption, $f(\mathbf{x})$, as indicated previously, is assumed to be lognormal with parameters μ_x and σ_x^2 . These parameters are estimated on the basis of the mean \bar{x} and coefficient of variation $CV(x)$ as follows:

$$\sigma_x = [\log_e (CV^2(x) + 1)]^{0.5} = [\log_e (0.29^2 + 1)]^{0.5} = 0.2842$$

and

$$\mu_x = \log_e \bar{x} - \sigma^2 / 2 = \log_e 2414 - 0.2842^2 / 2 = 7.7487.$$

The proportion of population below r_L is then evaluated as follows:

$$\Phi [(\log_e r_L - \mu) / \sigma] = \Phi [(\log_e 1680 - 7.7487) / 0.2842] = \Phi [-1.1335] = 0.1285$$

Where :

Φ = standard normal cumulative distribution.

Thus,

the percentage of the population undernourished = 13.

As the total population of the hypothetical country is 11 million, the number of undernourished is estimated as follows:

$$\text{Number of undernourished} = 11 * 0.1285 = 1.4 \text{ million.}$$

4. MEANING, SIGNIFICANCE AND ADVANTAGES OF THE ESTIMATES OF THE PREVALENCE OF UNDERNOURISHMENT

The data and approximations used to estimate the parameters of the density function of dietary energy consumption and the cut-off point have implications on the precise meaning and significance of the resulting estimate of the prevalence of food deprivation. These are discussed below.

4.1 *Concept of food consumption*

The daily per person DES refers to food acquired by (or available to) the households rather than the actual food intake of the individual household members.

4.2 *Time reference*

When Food Balance Sheet data are used, the daily per person DES taken as the mean of $f(\mathbf{x})$ corresponds to a three-year rather than annual average in order to even out the effect of errors in the annual food stocks data used in preparing the food balance sheets. On the other hand,

the daily per person food consumption from household budget surveys refer to a one year average period. Furthermore, for the purpose of deriving the $CV(x|v)$, only household survey data grouped according to income or total expenditure classes are used thus removing the effect of seasonal and other short-term variation that the household level data are subject to. As a consequence of these, the estimate refers to the average condition during the given one or three-year period and the effects of seasonal and other short-term variations in food availability are not considered.

4.3 *Use of concept of minimum energy requirement as cut-off point*

The cut-off point is derived by aggregating the sex-age specific minimum energy requirements using the proportion of the population in the different sex-age groups as weights. The specific minimum energy requirements for sex and age population groups are based on the total energy expenditure corresponding to the minimum acceptable limit of the range of body-weight for attained-height and the light physical activity norm. This approach of arriving at the cut-off point might give the impression that food deprivation is operationally defined as the state of having a food consumption level that is below that needed by an average individual for maintaining minimum acceptable body-weight and performing light physical activity. This is however, strictly speaking, not so. The minimal approach in establishing the cut-off point is a consequence of the consideration that, due to the effect of correlation between energy intake and requirement, the individuals with consumption falling within the range of variation of requirement are likely to be close to, if not exactly, matching their requirements. In other words their risk of food shortfall or excess is negligible if not exactly zero.

4.4 *Advantages of the use of food consumption estimates from Food Balance Sheets*

The procedure of using the daily per person DES derived from the food balance has some advantages as indicated below.

- The FAO daily per person DES database, which covers practically all countries of the world, is regularly revised and up-dated in connection with FAO's continuous work programme on supply and utilization accounts and food balance sheets. As a result the database represents a readily available source of information for the assessment and monitoring of the prevalence of food deprivation at the global, regional and country levels.
- The linkage of the daily per person DES with a measure of inequality within a probability framework provides a mechanism for assessing the effect of short-term changes in aggregate food availability as well as its components (production, import, etc.) on the distribution of dietary energy consumption and hence the prevalence of food deprivation. In addition, the use of a probability model – such as the log-normal function – facilitates the assessment of expected changes in the prevalence of food deprivation as a result of the combined effect of food supply increase and inequality reduction, as illustrated in the table below.

Mean Food Consumption (kcal/caput/day)	Food deprivation (%) at MDER of 1680 kilo-calories different levels of food consumption and inequality			
	<i>(CV = coefficient of variation)</i>			
	0.20	0.24	0.29	0.35
1700	52	53	54	55
2040	19	24	29	34
2450	4	7	12	17
2940	0	1	3	7

4.5. *Disaggregating estimates at sub-national levels*

There is of course an interest in obtaining information on the differences that may exist in the prevalence of undernourishment among individuals living in different areas within a country or belonging to different socioeconomic groups.

For global assessment purposes FAO has been estimating food deprivation for each country as a whole only as described in the previous section. However, for assisting countries, the FAO methodology has been applied to derive sub-national estimates of the prevalence of undernourishment, given that the mean and CV of the distribution of dietary energy consumption as well as the minimum energy requirement are estimated at representative sub-national areas from available household survey data of specific countries.

The data from Household Income and Expenditure Survey (HIES) allow to measure the urban and rural dietary energy consumption, minimum dietary energy requirements and CV of dietary energy consumption due to income, $CV(x|v)$. These estimations are illustrated below through a hypothetical example.

<i>Consumption statistics:</i>	<i>Urban</i>	<i>Rural</i>
Dietary energy consumption (kcal/person/day)	2380	2440
Minimum energy requirement (kcal/person/day)	1690	1650
Coefficient of variation of dietary energy consumption due to income (%)	25	20
Coefficient of variation of dietary energy consumption (%)	32.0	28.3
Food deprivation (%): Percentage of population with dietary energy consumption below the minimum energy requirement (indicator 1.9, target 1.C and goal 1 of the Millennium Development Goals)	17.4	10.2

BIBLIOGRAPHY

FAO, 1977. *The Fourth World Food Survey*, Rome

FAO, 1987. *The Fifth World Food Survey*, Rome

FAO, 1996. *The Sixth World Food Survey*, Rome

FAO, 2008. *The State of Food Insecurity in the World*, Rome

FAO/WHO/UNU, 1985. *Energy and Protein Requirements*. Report of a joint FAO/WHO/UNU ad hoc Expert Consultation. Geneva, WHO, WHO Tech. Rpt. Ser. 724.

FAO/WHO/UNU, 2004. *Human Energy Requirements*. Report of a Joint FAO/WHO/UNU Expert Consultation. Rome, FAO, FAO Food and Nutrition Tech. Rpt. Ser. 1.

James, W.P.T. & Schofield, E.C., 1990. Human energy requirements. Oxford, Oxford University Press.

Naiken, L., 1998. *On Certain Statistical Issues Arising from the Use of Energy Requirements in Estimating the Prevalence of Energy Inadequacy (Undernutrition)*. Journal of the Indian Society of Agricultural Statistics, Vol. L1, N. 2.3, pp. 113-128.

Naiken, L., 2002. *FAO methodology for estimating the prevalence of undernourishment* Key note paper presented at the International Scientific Symposium on Measurement of Food Deprivation and Undernutrition (26-28 June, 2002).

United Nations, 2001. Road Map towards the Implementation of the United Nations Millennium Declaration: Report of the Secretary - General. Document A/56/326.

United Nations, 2007. World Population Prospects: The 2006 Revision. United Nations, New York. Vol. I. Document ST/ESA/SER.A/261.

WHO, 1995. *Physical Status: the use and interpretation of anthropometry*. WHO Technical Report Series 854. Geneva.

WHO, 2006. *WHO Child Growth Standards: BMI for age tables*. WHO. Geneva. Available at http://www.who.int/childgrowth/standards/bmi_for_age/en/index.html

WHO, 2007. *WHO Growth reference 5-19 years: BMI for age tables*. WHO. Geneva. Available at http://www.who.int/growthref/who2007_bmi_for_age/en/index.html

Annex :

Computation of Minimum Dietary Energy Requirements

A Hypothetical Country

The Minimum Energy Requirement can be derived applying the algorithm described below using data on country attained-height and population structure for different age groups by sex. The algorithm is implemented using exogenous information on weight for attained-height based on the Body Mass Index (BMI) in the reference population for children less than five years of age (WHO 2006), for children and adolescents from five to less than 18 years of age (WHO 2007) and for adults (WHO 1995) as well as dietary energy requirements for weight (for attained-height) and for weight-gain by age groups (FAO/WHO/UNU 2004). Additional dietary energy allowance for pregnancy is estimated using country birth rate.

Algorithm for estimating the Minimum Dietary Energy Requirement (MDER)				
Age group	Weight for attained height (KG)	Lower Limit of Energy Requirement (LLER)		Total requirement per age group (TER)
		Male	Female	
		(kcal/person/day)		
less than 1		$(-99.4+88.6*KG50)+2*WG*Er_KG50$	$(-99.4+88.6*KG50)+2*WG*Er_KG50$	LLER*P _{ij}
1 to 1.9	KG50	$0.93*(310.2+63.3*KG50-0.263*KG502)+2*WG*Er_KG50$	$0.93*(263.4+65.3*KG50-0.454*KG502)+2*WG*Er_KG50$	
2 to 9.9		$(310.2+63.3*KG50-0.263*KG502)+WG*Er_KG50$	$(263.4+65.3*KG50-0.454*KG502)+WG*Er_KG50$	
10 to 17.9		$0.85*(310.2+63.3*KG5-0.263*KG52)+WG*Er_KG5$	$0.85*(263.4+65.3*KG5-0.454*KG52)+WG*Er_KG5$	
18 to 29.9		$1.55*(692.2+15.057*KG5)$	$1.55*(486.6+8.126*KG5)$	
30 to 59.9	KG5	$1.55*(873.1+11.472*KG5)$	$1.55*(845.6+8.118*KG5)$	
60 to 70 +		$1.55*(587.7+11.711*KG5)$	$1.55*(658.5+9.082*KG5)$	
		sum of		
		$KG50 = (BMI_{50th}*((H/100)^2))$	$MDER =$	TER age group
		$KG5 = (BMI_{5th}*((H/100)^2))$		

The reference weight for country attained-height correspond to the 50th percentile (median) for age-groups of individuals less than 10 years while the 5th percentile for age-groups of individuals of 10 years and above. The MDER is a weighted average of MDERs for the various age-groups and sex using the sex and age population structure provided by the UN Population Division (UN 2007).

The inputs for implementing the algorithm are given for the hypothetical country in the table below. The sex and age group population structure (Pop. ratio) adds up to unit for weighting the population MDER estimate. The attained-heights are estimated using anthropometric data collected in nutritional surveys.

Input data from reference tables and country statistics										
Age group	Pop. ratio		Attained HEIGHT (H) (cm)		BMI		Weight gain for age (WG) (kg)		Energy per KG of weight gain (Er_kg) (kcal)	
	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female
less than 1	0.0174	0.0168	66.5	61.5	17.30	16.90	16.16	15.07	4.1	4.4
1 to 1.9	0.0172	0.0166	73.5	71.5	16.10	15.70	6.58	6.58	2	2
2 to 2.9	0.0169	0.0164	81.5	79.5	15.80	15.50	6.30	6.30	2	2
3 to 3.9	0.0166	0.0161	87.5	85.5	15.40	15.30	5.75	5.21	2	2
4 to 4.9	0.0162	0.0158	96.5	95.5	15.30	15.30	5.48	4.66	2	2
5 to 5.9	0.0158	0.0154	102.5	100.5	15.26	15.25	5.48	4.93	2	2
6 to 6.9	0.0154	0.0150	108.5	108.5	15.38	15.32	6.03	6.30	2	2
7 to 7.9	0.0150	0.0146	113.5	113.5	15.60	15.52	6.58	8.22	2	2
8 to 8.9	0.0145	0.0142	118.5	117.5	15.89	15.87	7.67	10.14	2	2
9 to 9.9	0.0140	0.0138	122.5	122.5	16.23	16.34	9.04	10.96	2	2
10 to 10.9	0.0136	0.0134	128.8	129	14.29	14.13	6.30	7.95	2	2
11 to 11.9	0.0131	0.0129	133.5	135.7	14.67	14.62	8.22	8.77	2	2
12 to 12.9	0.0127	0.0125	138.6	142.4	15.14	15.19	10.41	9.86	2	2
13 to 13.9	0.0123	0.0123	144.3	146.8	15.69	15.77	13.15	9.86	2	2
14 to 14.9	0.0121	0.0121	150.9	149.1	16.27	16.28	14.25	8.77	2	2
15 to 15.9	0.0119	0.0118	157.6	151	16.82	16.67	13.42	6.58	2	2
16 to 16.9	0.0116	0.0116	162.8	152.9	17.32	16.94	10.68	3.84	2	2
17 to 17.9	0.0113	0.0113	165	153.8	17.75	17.10	6.58	1.64	2	2
18 to 18.9	0.0108	0.0110	166	154.4	18.10	17.19				
19 to 19.9	0.0103	0.0106	166	154.4	17.80	16.87				
20 to 24.9	0.0437	0.0471	166	154.4	18.66	17.38				
25 to 29.9	0.0335	0.0380	166	154.4	18.66	17.38				
30 to 34.9	0.0260	0.0303	166	154.4	18.66	17.38				
35 to 39.9	0.0213	0.0245	166	154.4	18.66	17.38				
40 to 44.9	0.0180	0.0201	166	154.4	18.66	17.38				
45 to 49.9	0.0165	0.0177	166	154.4	18.66	17.38				
50 to 54.9	0.0142	0.0151	166	154.4	18.66	17.38				
55 to 59.9	0.0110	0.0113	166	154.4	18.66	17.38				
60 to 64.9	0.0093	0.0095	166	154.4	18.66	17.38				
65 to 69.9	0.0077	0.0082	166	154.4	18.66	17.38				
70+	0.0115	0.0129	166	154.4	18.66	17.38				

The BMI values, 5th and 50th percentiles, are from WHO references (1995, 2006 and 2007). The weight gains and energy requirements for weight gains are from the FAO/WHO/UNU Joint Report of the Expert Consultation on Human Energy Requirements (2004).

The computation of the MDER for a hypothetical country is illustrated below. The first table refers to males contributing 892 kilo-calories and the second table refers to females contributing 781 kilo-calories; the pregnancy allowance is of 7 kilocalories. The MDER for the population of the hypothetical country is 1680 kilo-calories per person per day.

Estimating the Minimum Dietary Energy Requirement (MDER) in a hypothetical country: MALES

Age group	Attained HEIGHT (H) (cm)	BMI Weight (KG) for attained height $BMI*(height/100)^2$ (kg)	Weight gain (WG) for age (Ref. Tb) (kg)	Energy per KG of weight gain (kcal)	PAL (Physical Activity Level)	Lower Limit of Energy Requirement (LLER) (kcal/person/day)	Pop. ratio (p _{ij})	Total requirement per age group 2000 (kcal/person/day)	
less than 1	66.5	17.30	7.7	16.16	4.1	711	$(-99.4+88.6*KG)+2*WG*Er_kg$	0.017	12.39
1 to 1.9	73.5	16.10	8.7	6.58	2	808	$0.93*(310.2+63.3*KG-0.263*KG^2)+2*WG*Er_kg$	0.017	13.90
2 to 2.9	81.5	15.80	10.5	6.30	2	958	$(310.2+63.3*KG-0.263*KG^2)+WG*Er_kg$	0.017	16.21
3 to 3.9	87.5	15.40	11.8	5.75	2	1031		0.017	17.11
4 to 4.9	96.5	15.30	14.2	5.48	2	1170		0.016	18.97
5 to 5.9	102.5	15.26	16.0	5.48	2	1269		0.016	20.07
6 to 6.9	108.5	15.38	18.1	6.03	2	1382		0.015	21.28
7 to 7.9	113.5	15.60	20.1	6.58	2	1489		0.015	22.27
8 to 8.9	118.5	15.89	22.3	7.67	2	1607		0.015	23.30
9 to 9.9	122.5	16.23	24.4	9.04	2	1714		0.014	24.08
10 to 10.9	128.8	14.29	23.7	6.30	2	1426		$0.85*(310.2+63.3*KG-0.263*KG^2)+WG*Er_kg$	0.014
11 to 11.9	133.5	14.67	26.1	8.22	2	1534	0.013		20.06
12 to 12.9	138.6	15.14	29.1	10.41	2	1660	0.013		21.01
13 to 13.9	144.3	15.69	32.7	13.15	2	1809	0.012		22.33
14 to 14.9	150.9	16.27	37.0	14.25	2	1978	0.012		23.95
15 to 15.9	157.6	16.82	41.8	13.42	2	2148	0.012		25.46
16 to 16.9	162.8	17.32	45.9	10.68	2	2284	0.012		26.52
17 to 17.9	165	17.75	48.3	6.58	2	2355	0.011	26.57	
18 to 18.9	166	18.10	49.9		1.55	2237	$1.55*(692.2+15.057*KG)$	0.011	24.20
19 to 19.9	166	17.80	49.0		1.55	2218		0.010	22.77
20 to 24.9	166	18.66	51.4		1.55	2273		0.044	99.39
25 to 29.9	166	18.66	51.4		1.55	2273		0.033	76.11
30 to 34.9	166	18.66	51.4		1.55	2268	$1.55*(873.1+11.472*KG)$	0.026	58.97
35 to 39.9	166	18.66	51.4		1.55	2268		0.021	48.27
40 to 44.9	166	18.66	51.4		1.55	2268		0.018	40.79
45 to 49.9	166	18.66	51.4		1.55	2268		0.016	37.36
50 to 54.9	166	18.66	51.4		1.55	2268		0.014	32.11
55 to 59.9	166	18.66	51.4		1.55	2268		0.011	24.84
60 to 64.9	166	18.66	51.4		1.55	1844	$1.55*(587.7+11.711*KG)$	0.009	17.08
65 to 69.9	166	18.66	51.4		1.55	1844		0.008	14.13
70+	166	18.66	51.4		1.55	1844		0.011	21.19
Total									892.05

LLER*P_{ij}

Estimating the Minimum Dietary Energy Requirement (MDER) in a hypothetical country: FEMALES

Age group	Attained HEIGHT (H) (cm)	BMI for attained height BMI*(height/100) ² (kg)	Weight (KG) for attained height Weight (KG) (kg)	Weight gain (WG) for age (Ref. Tb) (kg)	Energy per KG of weight gain (kcal)	PAL (Physical Activity Level)	Lower Limit of Energy Requirement (LLER) (kcal/person/day)	Pop. ratio (P _{ij})	Total requirement per age group 2000 (kcal/person/day)
Female									
less than 1	61.5	16.90	6.4	15.07	4.4	600	$(-99.4+88.6*KG)+2*WG*Er_kg$	0.017	10.07
1 to 1.9	71.5	15.70	8.0	6.58	2	732	$0.93*(263.4+65.3*KG-0.454*KG^2)+2*WG*Er_kg$	0.017	12.15
2 to 2.9	79.5	15.50	9.8	6.30	2	872	$(263.4+65.3*KG-0.454*KG^2)+WG*Er_kg$	0.016	14.28
3 to 3.9	85.5	15.30	11.2	5.21	2	947		0.016	15.24
4 to 4.9	95.5	15.30	14.0	4.66	2	1096		0.016	17.27
5 to 5.9	100.5	15.25	15.4	4.93	2	1171		0.015	18.04
6 to 6.9	108.5	15.32	18.0	6.30	2	1306		0.015	19.62
7 to 7.9	113.5	15.52	20.0	8.22	2	1404		0.015	20.54
8 to 8.9	117.5	15.87	21.9	10.14	2	1497		0.014	21.28
9 to 9.9	122.5	16.34	24.5	10.96	2	1614		0.014	22.27
10 to 10.9	129	14.13	23.5	7.95	2	1331		$0.85*(263.4+65.3*KG-0.454*KG^2)+WG*Er_kg$	0.013
11 to 11.9	135.7	14.62	26.9	8.77	2	1456	0.013		18.83
12 to 12.9	142.4	15.19	30.8	9.86	2	1587	0.013		19.92
13 to 13.9	146.8	15.77	34.0	9.86	2	1684	0.012		20.66
14 to 14.9	149.1	16.28	36.2	8.77	2	1745	0.012		21.03
15 to 15.9	151	16.67	38.0	6.58	2	1789	0.012		21.16
16 to 16.9	152.9	16.94	39.6	3.84	2	1825	0.012		21.17
17 to 17.9	153.8	17.10	40.4	1.64	2	1841	0.011		20.86
18 to 18.9	154.4	17.19	41.0			1.55	$1.55*(486.6+8.126*KG)$	0.011	18.62
19 to 19.9	154.4	16.87	40.2			1.55		0.011	17.74
20 to 24.9	154.4	17.38	41.4			1.55		0.047	80.27
25 to 29.9	154.4	17.38	41.4			1.55		0.038	64.76
30 to 34.9	154.4	17.38	41.4			1.55	$1.55*(845.6+8.118*KG)$	0.030	55.52
35 to 39.9	154.4	17.38	41.4			1.55		0.025	44.91
40 to 44.9	154.4	17.38	41.4			1.55		0.020	36.74
45 to 49.9	154.4	17.38	41.4			1.55		0.018	32.50
50 to 54.9	154.4	17.38	41.4			1.55		0.015	27.60
55 to 59.9	154.4	17.38	41.4			1.55		0.011	20.74
60 to 64.9	154.4	17.38	41.4			1.55		$1.55*(658.5+9.082*KG)$	0.010
65 to 69.9	154.4	17.38	41.4			1.55	0.008		13.15
70+	154.4	17.38	41.4			1.55	0.013		20.73
Total									780.79

LLER*P_{ij}