

**Table 8.1 Feeding chart for common carp (*Cyprinus carpio*) cultured in ponds during summer (water temperature over 20 °C)<sup>1,2</sup>**

Individual weight of fish (g)	Density of fish per hectare													
	2 000–4 000		4 000–6 000		6 000–8 000		8 000–12 000		12 000–15 000		15 000–20 000		20 000–50 000	
	g	Protein content of feed (%)	g	Protein content of feed (%)	g	Protein content of feed (%)	g	Protein content of feed (%)	g	Protein content of feed (%)	g	Protein content of feed (%)	g	Protein content of feed (%)
20–50	1	12	2	12	2	12	3	12	3	12	3	18	3	18
50–100	2	12	3	12	4	12	4	18	4	18	4	18	4	18
100–200	6	12	6	18	9	18	9	18	9	18	9	25	9	25
200–300	10	12	11	18	11	18	12	25	12	25	12	25	12	25
300–400	11	18	13	25	14	25	15	25	15	25	15	25	15	25
400–500	14	18	15	25	16	25	17	25	17	30	17	30	17	30
500–600	15	25	17	25	18	25	19	30	19	30	19	30	19	30
600–700	15	25	18	25	19	30	19	30	19	30	19	30	19	30
700–800	16	25	18	25	20	30	19	30	19	30	19	30	19	30
800–900	17	25	18	25	18	30	20	30	20	30	20	30	20	30
900–1 000	17	25	19	25	20	30	21	30	21	30	21	30	21	30
1 000–1 100	18	25	20	30	21	30	22	30	22	30	22	30	22	30
1 100–1 200	18	25	20	30	21	30	22	30	22	30	22	30	22	30

<sup>1</sup>Daily amounts are given as grams per stocked fish.

<sup>2</sup>Source: Hephher (1988).