

Nutrition status	Health status	Food consumption (Individual)	Food access (Household)	Food availability (Household)	Care practices	Gender-related issues	Healthy Environment	
							WASH (not exhausted)	Access to health Services
Prevalence of wasting among pregnant and lactating women	No. Of consultation per person per year (proxy)	Individual/Women Dietary Diversity Score (IDDS/WDDS)	Household Food consumption score (FCS)	Level of production	Nutrition Knowledge	Childcare practices	Access to drinking water	Coverage of health services
Prevalence of wasting among children under 5 (and under 2)	HIV/AIDS	No. Of meals per day (Adults/children)	Household dietary diversity score (HDDS)	Availability of micronutrient-rich food in the market.	Immediate breastfeeding	Mothers time/workload	Distance to safe water source	No. of Health workers per person person/people
Prevalence of stunting among children under 5	Measles	Consumption of iron- rich or iron-fortified food	% of household consuming specific food groups	Availability of different food groups in the market (Seasonality)	Exclusive breastfeeding until 6 months	No. of men and women participating on project decision making and implementation		Distance to the health facilities
Prevalence of micronutrient deficiencies	Diarrhoea	Minimum acceptable diet (for children 6- 23 yrs)	Coping strategy Index (CSI) and Reduced copying strategy index and (rCSI)	No. of days covered by the current food stock	Minimum acceptable diet-dietary diversity+ meal frequency	Division of labour at household level		No. of health facilities
Body Mass Index (for adults)	Malaria	Minimum meal frequency (for children 6-23 yrs)	Household hunger scale (HHS)		Early childhood stimulation	Women control of household cash		No. of skilled health personnel
Low Birth Weight	Morbidity rate	Minimum dietary diversity (for children 6-23 yrs)	Household food insecurity access scale (HFIAS)		Preventive health seeking behaviour	No. of labour saving devices introduced (poshomills.)		% of budget allocated to health
	Mortality rate (maternal, under 5, infant)	Early initiation of breastfeeding	Household income		Hand washing	.		% of people accessing health
		Exclusive breastfeeding under 6 months	% Expenditure on food					% of cases referred
		Continued beastfeeding at 2 yrs	Price of staples in the market (e.g. compared to 10 yrs average)					
			Terms of trade for livestock					
			Price of micronutrient-rich food in market					