

DICLAZURIL

**First draft prepared by
Dr. D. Arnold
Federal Institute for Health Protection of
Consumer and Veterinary Medicine
Berlin, Germany**

**ADDENDUM
to the Diclazuril monograph prepared by
the forty-fifth meeting of the Committee and published in
FAO Food and Nutrition Paper 41/8, Rome 1996**

At its forty-fifth meeting the Committee had recommended temporary MRLs for residues of diclazuril in certain tissues of food animals expressed as parent compound. The MRL's were temporary because the ADI was temporary.

No additional residue data were provided to the Committee for consideration at its current meeting. Since the final ADI established by the Committee at its current meeting was not lower than the temporary ADI allocated at the 45th meeting and since the temporary MRLs already reflected good practice in the use of this veterinary drug the Committee decided to delete the temporary qualification and recommend MRLs at the same level for the same species/tissue combinations as had been recommended at its forty-fifth session.

The recommended MRLs expressed as parent compound, are:

| Species | Animal Tissue | MRL (µg/kg) |
|----------------|----------------------|--------------------|
| Sheep, rabbits | Liver | 3000 |
| | Kidney | 2000 |
| | Fat | 1000 |
| | Muscle | 500 |
| Poultry | Liver | 3000 |
| | Kidney | 2000 |
| | Skin/Fat | 1000 |
| | Muscle | 500 |